Published in final edited form as:

Chem Commun (Camb). 2011 October 14; 47(38): 10614–10616. doi:10.1039/c1cc14165a.

Mesoionic Thiazol-5-ylidenes as Ligands for Transition Metal Complexes

Daniel Mendoza-Espinosa, Gaël Ung, Bruno Donnadieu, and Guy Bertrand

Guy Bertrand: guy.bertrand@ucr.edu

^aUCR-CNRS Joint Research Chemistry Laboratory (UMI 2957), Department of Chemistry, University of California, Riverside CA 92521-0403 (USA). Fax (+1) 202 354 5267

Abstract

The first examples of thiazol-5-ylidene complexes featuring group 9, 10 and 11 metal centers, have been prepared by deprotonation of a series of 2,3,4-triaryl-substituted thiazolium salts in the presence of the corresponding transition metal precursor.

In 2001, Crabtree *et al.* reported the preparation of complex **B**,¹ featuring an imidazol-5-ylidene **C** as ligand. It was quickly demonstrated that this type of ligands² are even stronger electron-donors than their imidazol-2-ylidene isomers, the so-called NHCs **A**.^{3,4} (Figure 1). Recently, it has been shown that derivatives of type **C**,⁵ as well as pyrazolin-4-ylidenes **D**⁶ and 1,2,3-triazol-5-ylidenes **E**,⁷ can be isolated. Since no reasonable canonical resonance forms showing a carbene can be drawn without additional charges, and since they are mesoionic compounds, they can be named mesoionic carbenes (MICs).⁸ Importantly, in contrast to regular carbenes, no obvious dimerization pathway can be foreseen, and therefore it was reasonable to believe that unhindered MICs with various heteroatom-containing skeletons could be stable. To test this hypothesis, we attempted the preparation of 1,3-dithiol-5-ylidenes **F**, but observed a spontaneous ring opening yielding the corresponding ethynylcarbamodithioate **G**.⁹ The formation of **G** is reminiscent of the ring-opening reaction observed in the deprotonation of isothiazolium salts,¹⁰ and therefore is probably due to the weakness of the sulfur-carbon bond. These results prompted us to investigate the preparation of thiazol-5-ylidenes **H**, since a similar ring-opening process would imply the cleavage of a carbon-nitrogen bond. Moreover, although 1,3-dithiol-2-ylidenes **I** have never been observed, it is noteworthy that 1,3-thiazol-2-ylidenes **J** have been used for decades as organocatalysts,¹¹ beginning with the seminal work by Breslow,¹² before their isolation by Arduengo.¹³ Here we report the synthesis of a series of 2,3,4-aryl-substituted thiazolium salts, and their use for the preparation of thiazol-5-ylidene transition metal complexes.

2,3,4-Triaryl-substituted thiazolium salts **3a–c** are conveniently prepared in two steps from thioamides **1a–c** (Scheme 1). Addition of triethylamine in the presence of bromoacetophenone affords ketones **2a–c** in 55 to 76% yield. Then, cyclization under acidic

Correspondence to: Guy Bertrand, guy.bertrand@ucr.edu.

[†]Electronic Supplementary Information (ESI) available: [Preparation and spectroscopic data for all compounds, selected X-ray data for compounds **3b** and **5b**. CCDC 833533 and 833534. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b000000x/]

[‡]Crystal data: **3b**: C₂₇H₃₀BrNO₂S, *M* = 512.49, *monoclinic*, *a* = 9.4960(14), *b* = 28.054(4), *c* = 9.6399(14), α = 90, β = 104.693(2), γ = 90, *V* = 2484.1(6) Å³, *T* = 100(2) K, *space group* *P*2₁/c, *Z* = 4, 20859 reflections measured, 6277 unique (*R*_{int} = 0.0276), which were used in all calculations. The final *R* [*I* > 2σ(*I*)] was 0.0319. **5b**: C₂₄H₂₁AuClNS, *M* = 587.89, *monoclinic*, *a* = 18.170(4), *b* = 9.7695(15), *c* = 24.734(4), α = 90, β = 99.134(3), γ = 90, *V* = 4334.9(12) Å³, *T* = 200(2) K, *space group* *C*2/c, *Z* = 8, 11994 reflections measured, 4607 unique (*R*_{int} = 0.0501), which were used in all calculations. The final *R* [*I* > 2σ(*I*)] was 0.0648.

conditions produces the desired thiazolium salts **3a–c** in 42 to 60% yields. In order to increase their solubility, a salt metathesis reaction with silver trifluoromethanesulfonate was carried out, producing **4a–c** in excellent yields. The formation of the thiazolium salts was confirmed by the characteristic downfield signal between 8.64 and 9.20 ppm (thiazolium C–H) in the ^1H NMR spectra and, in the case of salt **3b**, by a single crystal X-ray diffraction study (Figure 2, left).

All attempts to deprotonate salts **3** and **4** with a variety of bases [MHMDS (M = Li, Na, K), $^n\text{BuLi}$, KO^tBu and LDA] resulted in complex mixtures. Even monitoring the deprotonation reaction by NMR spectroscopy, at $-50\text{ }^\circ\text{C}$, did not allow for the observation of the desired thiazol-5-ylidenes of type **H**. However, when salts **4a–c** were treated at $-78\text{ }^\circ\text{C}$ with an equimolar amount of LiHMDS, in the presence of (THT)AuCl or $[\text{Pd}(\text{Allyl})\text{Cl}]_2$, the corresponding thiazol-5-ylidene transition metal complexes were obtained and isolated in moderate to good yields (51 to 90%) (Scheme 2). The ^{13}C NMR chemical shifts of the carbene carbon of Au^{I} (149.5 to 151.4 ppm) and Pd^{II} (156.0 to 159.5 ppm) complexes are in the range observed for previously reported metal complexes featuring MICs **C–F**,^{5,9,14} and as expected, at much higher field than those of classical NHCs.¹⁵

To unambiguously confirm the existence of the new thiazol-5-ylidene ligands, single crystals of complex **5b**, obtained by slow diffusion of hexane into a concentrated chloroform solution, were subjected to an X-ray diffraction analysis (Figure 2, right). Complex **5b** crystallizes in the $C2/c$ space group displaying a monomeric unit with a Au(1)–C(1) bond distance of 2.005(14) Å, which is slightly longer than those observed for the (1,3-dithiol-5-ylidene)AuCl [1.978(4) Å]⁹ and (imidazol-5-ylidene)AuCl [1.98 Å].⁵ The carbene bond angle in **5b** [110.0(10)°] is slightly more acute than the corresponding angle of thiazolium salt **3b** [112.0(4)°]. This feature is consistent with the increased s-character of the σ lone-pair orbital on the carbene atom in **5b** as compared with the C–H⁺ bond orbital in **3b**. Finally, as expected, the Au(I) center is in a linear environment with an C(1)–Au(1)–Cl(1) angle of 179.2(4)°.

To evaluate the donor properties of thiazol-5-ylidene ligands, the corresponding rhodium(I) dicarbonyl chloride complexes **7a–c** were prepared by the *in situ* deprotonation of salts **4a–c**, at $-78\text{ }^\circ\text{C}$, with LiHMDS, in the presence of half an equivalent of $[\text{Rh}(\text{COD})\text{Cl}]_2$, followed by treatment with an excess of carbon monoxide (Scheme 3).

The average CO vibration frequency for **7a–c** ($\nu_{\text{av}} = 2034\text{ cm}^{-1}$) indicates that the electron-donor capabilities of thiazol-5-ylidenes are slightly superior to those of conventional NHCs ($\nu_{\text{av}} = 2039\text{--}2041\text{ cm}^{-1}$),¹⁶ similar to those of 1,3-dithiol-5-ylidene **F** ($\nu_{\text{av}} = 2030\text{ cm}^{-1}$)⁹ and cyclic (alkyl)(amino)carbenes ($\nu_{\text{av}} = 2036\text{ cm}^{-1}$),¹⁷ but inferior to those of MICs **C–E** ($\nu_{\text{av}} = 2016\text{--}2025\text{ cm}^{-1}$).^{2d}

Although thiazol-5-ylidenes cannot be observed, the readily available 2,3,4-triaryl-substituted thiazolium salts are modular precursors for the synthesis of a variety of thiazol-5-ylidenes transition metal complexes. This novel type of MIC ligand features stronger donor properties than classical NHCs, and gives rise to complexes, which are thermally- and air-stable. Research in our laboratory is currently underway to extend the scope of mesoionic carbenes, and explore the catalytic activity of the corresponding complexes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH (R01 GM 68825) and DOE (DE-FG02-09ER16069).

Notes and references

1. Gründemann S, Kovacevic A, Albrecht M, Faller JW, Crabtree RH. *Chem Commun.* 2001;2274.
2. For reviews on imidazol-5-ylidenes and related species, see: Schuster O, Yang L, Raubenheimer HG, Albrecht M. *Chem Rev.* 2009; 109:3445. [PubMed: 19331408] Albrecht M. *Chem Commun.* 2008; 31:3601. Arnold PL, Pearson S. *Coord Chem Rev.* 2007; 251:596. Albrecht M. *Chimia.* 2009; 63:105. Melaimi M, Soleilhavoup M, Bertrand G. *Angew Chem Int Ed.* 2010; 49:8810.
3. For recent reviews on NHCs, see for examples: Dröge T, Glorius F. *Angew Chem Int Ed.* 2010; 49:6940. Vougioukalakis GC, Grubbs RH. *Chem Rev.* 2010; 110:1746. [PubMed: 20000700] Lin JCY, Huang RTW, Lee CS, Bhattacharyya A, Hwang WS, Lin IJB. *Chem Rev.* 2009; 109:3561. [PubMed: 19361198] Arnold PL, Casely IJ. *Chem Rev.* 2009; 109:3599. [PubMed: 19358527] Díez-González S, Marion N, Nolan SP. *Chem Rev.* 2009; 109:3612. [PubMed: 19588961] Poyatos M, Mata JA, Peris E. *Chem Rev.* 2009; 109:3677. [PubMed: 19236010] Samojłowicz C, Bieniek M, Grela K. *Chem Rev.* 2009; 109:3708. [PubMed: 19534492] van Otterlo WAL, de Koning CB. *Chem Rev.* 2009; 109:3743. [PubMed: 19618929] Monfette S, Fogg DE. *Chem Rev.* 2009; 109:3783. [PubMed: 19537778] Alcaide B, Almendros P, Luna A. *Chem Rev.* 2009; 109:3817. [PubMed: 19572654] Hahn FE, Jahnke MC. *Angew Chem Int Ed.* 2008; 47:3122. Bourissou D, Guerret O, Gabbai FP, Bertrand G. *Chem Rev.* 2000; 100:39. [PubMed: 11749234]
4. For recent books on NHCs, see: Díez-González S. *N-Heterocyclic Carbenes, From laboratory Curiosities to Efficient Synthetic Tools.* Royal Society of Chemistry Publishing 2011. Glorius FA. *N-Heterocyclic Carbenes in Transition Metal Catalysis.* Springer-Verlag 2007. Nolan SP. *N-Heterocyclic Carbenes in Synthesis.* Wiley-VCH 2006
5. a) Aldeco-Perez E, Rosenthal AJ, Donnadiou B, Parameswaran P, Frenking G, Bertrand G. *Science.* 2009; 326:556. [PubMed: 19900893] b) Ung G, Bertrand G. *Chem Eur J.* 2011; 17:8269. [PubMed: 21656597] c) Wang Y, Xie Y, Abraham MY, Wei P, Schaefer HF III, Schleyer PvR, Robinson GH. *J Am Chem Soc.* 2010; 132:14370. [PubMed: 20863065]
6. a) Lavallo V, Dyker CA, Donnadiou B, Bertrand G. *Angew Chem Int Ed.* 2008; 47:5411. b) Fernández I, Dyker CA, DeHope A, Donnadiou B, Frenking G, Bertrand G. *J Am Chem Soc.* 2009; 131:11875. [PubMed: 19645451]
7. a) Guisado-Barrios G, Bouffard J, Donnadiou B, Bertrand G. *Angew Chem Int Ed.* 2010; 49:4759. b) Bouffard J, Keitz BK, Tonner R, Guisado-Barrios G, Frenking G, Grubbs RH, Bertrand G. *Organometallics.* 2011; 30:2617. [PubMed: 21572542] c) Keitz BK, Bouffard J, Bertrand G, Grubbs RH. *J Am Chem Soc.* 2011; 133:8498. [PubMed: 21574621]
8. For the origin of the acronym MIC, see: Araki S, Wanibe Y, Uno F, Morikawa A, Yamamoto K, Chiba K, Butsugan Y. *Chem Ber.* 1993; 12:1149.
9. a) Ung G, Mendoza-Espinosa D, Bouffard J, Bertrand G. *Angew Chem Int Ed.* 2011; 50:4215. b) Ung G, Frey GD, Schoeller WW, Bertrand G. accepted for publication.
10. Dehope A, Lavallo V, Donnadiou B, Schoeller WW, Bertrand G. *Angew Chem Int Ed.* 2007; 46:6922.
11. See for recent examples: Hirano K, Biju AT, Piel I, Glorius F. *J Am Chem Soc.* 2009; 131:14190. [PubMed: 19807174] Lebeuf R, Hirano K, Glorius F. *Org Lett.* 2008; 10:4243. [PubMed: 18763794] Biju AT, Glorius F. *Angew Chem Int Ed.* 2010; 49:9761. Piel I, Steinmetz M, Hirano K, Frohlich R, Grimme S, Glorius F. *Angew Chem Int Ed.* 2011; 50:4983.
12. Breslow R. *J Am Chem Soc.* 1958; 80:3719.
13. Arduengo AJ III, Goerlich JR, Marshal WJ. *Liebigs, Ann Recl.* 1997; 365
14. a) Mathew P, Neels A, Albrecht M. *J Am Chem Soc.* 2008; 130:13534. [PubMed: 18785741] b) John A, Shaik MM, Ghosh P. *Dalton Trans.* 2009:10581. [PubMed: 20023883] c) Xu X, Xu B, Li Y, Hong SH. *Organometallics.* 2010; 29:6343.
15. Tapu D, Dixon DA, Roe C. *Chem Rev.* 2009; 109:3385. [PubMed: 19281270]

16. Fürstner A, Alcarazo M, Krause H, Lehmann CW. J Am Chem Soc. 2007; 129:12676. [PubMed: 17910460]
17. Lavallo V, Canac Y, Dehope A, Donnadiou B, Bertrand G. Angew Chem Int Ed. 2005; 44:7236.

Figure 1.

NHCs (**A**), Crabtree's imidazol-5-ylidene complex (**B**), types of MIC that have previously been isolated (**C–E**), transient 1,3-dithiol-5-ylidene (**F**) and its ring opened isomer (**G**), the targeted thiazol-5-ylidenes (**H**), the never-observed 1,3-dithiol-2-ylidenes **I**, and the isolated 1,3-thiazol-2-ylidenes **J**

Figure 2.

Crystal structures of thiazolium salt **3b** (left) and gold complex **5b** (right) with ellipsoids shown at 50% probability. Hydrogen atoms, except for the CH of **3b**, are omitted for clarity. Selected bond distances [Å] and angles [°]: **3b**: C(1)-S(1) = 1.716(6), C(1)-C(3) = 1.363(6), C(3)-N(1) = 1.397(3), N(1)-C(2) = 1.361(3), C(2)-S(1) = 1.684(2), C(3)-C(1)-S(1) = 112.0(4); **5b**: C(1)-S(1) = 1.697(14), C(1)-C(3) = 1.364(16), C(3)-N(1) = 1.414(15), N(1)-C(2) = 1.351(15), C(2)-S(1) = 1.695(14), Au(1)-C(1) = 2.005(14), Au(1)-Cl(1) = 2.293(4), C(3)-C(1)-S(1) = 110.0(10).

Scheme 1.

Synthesis of thiazolium salts **3a-c** and **4a-c** [Ar = Phenyl (**a**); 2,4,6-trimethylphenyl (**b**); 2,6-diisopropylphenyl (**c**)]

Scheme 2.

Synthesis of metal complexes **5a-c** and **6a-c** [Ar = Phenyl (**a**); 2,4,6-trimethylphenyl (**b**); 2,6-diisopropylphenyl (**c**)]

Scheme 3.

Synthesis of the rhodium(I) complexes **7a-c** [Ar = Phenyl (**a**); 2,4,6-trimethylphenyl (**b**); 2,6-diisopropylphenyl = Dipp (**c**)]