Published in final edited form as:

Curr Opin Cell Biol. 2012 June ; 24(3): 333–340. doi:10.1016/j.ceb.2012.03.006.

Small Non-coding RNAs Mount a Silent Revolution in Gene Expression

Antti P. Aalto^{1,#} and Amy E. Pasquinelli^{2,*}

¹Institute of Biotechnology and Department of Biosciences, University of Helsinki, Helsinki, Finland ²Division of Biological Sciences, University of California, San Diego, La Jolla, California, USA

Abstract

During the last decade, it has become evident that small non-coding RNAs (ncRNAs) participate in widespread and essential regulatory mechanisms in most eukaryotic cells. Novel classes of small RNAs, their biogenesis pathways and cellular effects are continuously being described, and new properties of already established ncRNAs are still being discovered. As the list of small RNA molecules and their roles becomes more and more extensive, one can get lost in the midst of new information. In this review, we attempt to bring order to the small ncRNA transcriptome by covering some of the major milestones of recent years. We go through many of the new properties that have been attributed to already familiar RNA molecules, and introduce some of the more recent novel classes of tiny ncRNAs.

Introduction

The discovery of microRNAs (miRNAs) and small interfering RNAs (siRNAs) established the field of small RNA (sRNA) research in animals at the turn of the 21st century. The first described small ncRNAs, miRNAs, siRNAs and Piwi-interacting RNAs (piRNAs), are classified by distinct biogenesis pathways and regulatory mechanisms, but they have in common the use of sequence complementarity to recognize target genes and silence their expression. Recent technical developments have enabled scientists to scan the small RNA transcriptome more thoroughly, which has revealed a multitude of novel types of RNA molecules that do not fit into the well-established classes. Here we review some of the new results that have emerged from large-scale sequencing studies as well as from more mechanistic approaches, and strive to categorize these recent advances in a meaningful context. The classes of sRNAs discussed in this review are listed in Table 1. As this review primarily focuses on animal small ncRNAs, the reader is directed to [1] for a discussion of plant ncRNAs.

© 2012 Elsevier Ltd. All rights reserved.

*Corresponding author: apasquinelli@ucsd.edu, 9500 Gilman Drive MC 0349, UCSD, Biology, La Jolla, CA 92093-0349, 858-822-3006 (phone), 858-822-3021 (FAX).

#Current address: Department of Biological Sciences, University of California, San Diego, La Jolla, California, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

miRNAs: Diverse biogenesis and regulatory pathways

Since their discovery in *C. elegans* over a decade ago, miRNAs have emerged as one of the most abundant and important classes of small ncRNAs. Mis-regulation of specific miRNAs has been linked to cancer, heart ailments, diabetes and neurological defects in humans [2]. The growing number of miRNAs is curated at the miRBase database (<http://www.mirbase.org/>) [3]. The latest release (18) has 18,226 entries that give rise to 21,643 mature miRNAs from 168 species including animals, plants, unicellular algae and even viruses. Some miRNAs, such as let-7, are perfectly conserved across diverse species. Others share identical 5' end sequences (typically nucleotides 2–7 called the seed region) and comprise families of miRNAs. While some individual miRNAs have proven to be essential, the redundancy of miRNAs within a family can often compensate for the loss of single members. However, it is unclear if the functional overlap stems from related miRNAs recognizing identical target sites or regulating genes in a common pathway.

The details of miRNA biogenesis have been extensively reviewed and the general pathway is summarized in Figure 1a [4]. Recent studies have revealed unexpected alternatives to the canonical pathway. The generation of mature miRNAs has been regarded to depend on the catalytic activity of Dicer with Ago proteins directing the repression of mRNAs recognized by the miRNAs. However, this view was challenged by several reports examining the biogenesis of vertebrate miR-451 [5–7]. The precursor of miR-451 bypasses Dicer processing and, instead, undergoes final maturation by Ago2 the only mammalian Argonaute that displays catalytic activity. miR-451 is co-transcribed with another miRNA, miR-144. Although both miRNAs share a common primary transcript, because of its structure, only the miR-451 precursor is subject to Ago2-dependent maturation. Reasons for this alternative miRNA processing pathway are currently unknown, but one of its consequences may be the preservation of a catalytically competent Argonaute during mammalian evolution [5–7].

Other variations in miRNA processing pathways add to the complexity of miRNA biogenesis (Figure 1a). Some miRNAs bypass Drosha-mediated cleavage altogether, but undergo normal digestion by Dicer in the cytoplasm. The initial processing of this class of miRNAs is spliceosomal, and the RNAs have been assigned as “mirtrons” for their miRNA- and intron-like features [8–10]. In addition, some miRNAs are Dicer-dependent but Microprocessor- and splicing-independent. These miRNAs originate from RNAs that have the capacity to form extensive stem-loop structures and are known as short hairpin-derived miRNAs [10]. Moreover, many miRNAs display variation in their annotated 5' or 3' termini. These isomers (isomiRs) may arise from products of alternative 5' processing or from the addition of non-templated nucleotides to the 3' ends of miRNAs [11,12]. In addition, the 5' and 3' arms of some *Ciona intestinalis* and human pre-miRNAs give rise to miRNA-offset RNAs (moRNAs) that apparently are products of Drosha processing [13,14]. Thus, miRNAs share the feature of deriving from intramolecular stem-loop structures capable of giving rise to ~22 nt RNA products but are not classified by a uniform processing pathway.

Mature miRNAs guide the RNA induced silencing complex (miRISC), which consists of Argonaute and other effector proteins, to target mRNAs through antisense base-pairing. In animals, most miRNAs only partially base-pair with their targets. The most common motif from computational predictions is perfect pairing of the miRNA 5' end, called the seed region, usually to mRNA 3'UTR sequences [15]. Validating predicted target sites *in vivo*, however, has until recently only been feasible through reporter gene analyses. An advance in identifying endogenous miRNA target sites has been made possible through CLIP-seq (cross-linking immunoprecipitation coupled with high-throughput sequencing, also known

as HITS-CLIP). This method has been used to isolate and map the sequences bound by Ago *in vivo* [16–19]. Results from CLIP-seq studies revealed that, in addition to 3' UTR sequences, coding exon sites make up a large fraction of Ago bound regions. Furthermore, seed pairing capacity was enriched in many but not all of the sequences associated with Ago, suggesting that other types of complementarity guide a considerable portion of miRISC interactions. While CLIP-seq narrows the miRNA target site to less than 100 nt and in some cases reveals the bases in contact with the Ago protein, the actual sequences paired to the miRNA cannot yet be resolved. Nonetheless, the extensive datasets of endogenous RISC binding activity provide a much needed experimental basis for further improvement of miRNA target prediction methods.

Originally, partial pairing of a miRNA with its target was thought to repress gene expression by inhibiting translation without inducing mRNA decay. It has been proposed that the miRNA complex can inhibit translation initiation or ribosome subunit joining, induce premature degradation of the nascent polypeptide chain and increase ribosome drop off (Figure 2a) [20]. Other studies have demonstrated that regulation by miRNAs can induce target mRNA deadenylation and destabilization (Figure 2a,b) [20]. A recent investigation suggests that miRNA-dependent mRNA destabilization may be a much more widely employed regulatory mechanism than previously appreciated [21]. Here, the authors used ribosome profiling to determine the exact positions of ribosomes on mRNAs, and observed that at least in HeLa cells miRNA-mediated target repression is primarily due to decreased mRNA levels and translational repression alone is a rare event. However, others argue that the mRNA decay may be merely an outcome of translational inhibition [22]. At this point, it remains to be shown which of the several models of miRNA-mediated target regulation most accurately reflects *in vivo* conditions. It is likely that the regulatory outcome can differ depending on the cell type, developmental stage, cellular condition and context of the miRNA target site.

piRNAs: Silencers of the transposable genome

Piwi-interacting RNAs (piRNAs) are characterized by their specific association with Argonaute proteins of the Piwi-clade, the absence of dsRNA precursors and independence of Dicer [23]. piRNAs have been shown to be crucial for early development, epigenetic regulation, gametogenesis and silencing of transposable elements (TEs) as well as some protein coding genes. Although the specifics of the pathways differ to some extent depending on the species, piRNAs have been mainly studied in *Drosophila*, where they were originally called repeat-associated siRNAs (rasiRNAs), *C. elegans*, where they are also called 21U-RNAs, and mouse. piRNAs were long considered to be confined in animal germline cells, but recently piRNAs have been detected in somatic tissues [24,25]. For a more thorough review of piRNA pathways, we recommend [23].

The piRNA pathway can be divided into initiation, effector and adaptation phases [26]. During the initiation phase, primary piRNAs that are mostly antisense to TEs are produced from piRNA clusters (Figure 1b). These RNAs are processed into 24–30 nt piRNAs that subsequently associate with specific Piwi proteins. In the piRNA pathway effector phase, the piRNA-loaded piRISCs base-pair to sense transposon transcripts and cleave them endonucleolytically. In *Drosophila* germline cells there is additionally an adaptation phase (a.k.a. the ping-pong amplification cycle) that depends partly on Ago3 and replenishes the piRNA pool while consuming the TE transcripts [23,24]. An intact piRNA pathway is continuously required for keeping expression of TEs at bay [26].

In addition to cleaving RNA transcripts directly, piRNA pathways induce TE silencing by establishing DNA methylation patterns within the genome (Figure 2b,c) [27]. piRNAs have

also been shown to contribute to deadenylation and decay of maternal mRNAs in the *Drosophila* embryo [28]. Subcellular localization seems to be important for these processes, and components of the piRNA pathway are known to reside both in the nucleus and specific cytoplasmic foci. The mechanistic details of how piRNA complexes sort to particular subcellular locations and direct transcriptional and post-transcriptional silencing of TEs and other types of transcripts are yet to be deciphered.

Endogenous siRNAs: A natural RNAi pathway

Animal endogenous siRNAs (endo-siRNAs) are reminiscent of piRNAs in the sense that they often target TEs or repetitive elements. Indeed, mammalian and fly endo-siRNAs are derived from dsRNA intermediates that result from “natural” sources such as bidirectional transcription, inverted repeats or pseudogenes hybridized to mRNAs (Figure 1c) [29–33]. The dsRNA is cleaved by Dicer and the siRNAs associate mostly with the catalytically competent Ago2. Presumably, the siRNA loaded Ago2 complex can direct cleavage of targets that pair perfectly with the endo-siRNA, although direct evidence for this mode of action in animal cells is limited (Figure 2b). Endo-siRNAs are also abundant in *C. elegans*, but their mechanism of biosynthesis is unique. Most nematode endo-siRNAs are secondary and result from unprimed RNA-dependent RNA polymerase (RdRP) activity, and Dicer-dependent primary endo-siRNAs are scarce (Figure 1c) [34,35]. In *C. elegans* and *Drosophila*, endo-siRNAs also have a role in chromatin regulation and thus can act as co-transcriptional silencers of gene expression (Figure 2c) [36,37].

New tricks for tRNAs: tRFs and tsRNAs

High-throughput sequencing efforts of small RNAs always yield hits to abundant RNA species, such as rRNAs, tRNAs and snoRNAs, that are usually regarded as random degradation products and omitted from the final analyses. However, recent studies have shown that these housekeeping RNAs may give rise to biologically relevant small RNAs. Sequencing the small RNAs of 17 to 26 nt from two prostate cancer cell lines revealed reads that matched mature tRNAs or pre-tRNAs [38]. These tRNA-derived RNA fragments (tRFs) were the second-most copious class of small RNAs in these cell types, and they displayed distinct expression patterns. Interestingly, blocking the expression of an abundantly expressed tRF (tRF-1001) by siRNAs reduced cell viability and proliferation. These effects were rescued by synthetic tRF-1001, which suggests that tRFs may be essential for certain biological processes [38]. Another study identified two types of tRNA-derived small RNAs (tsRNAs): type I tsRNAs are Dicer-dependent, whereas type II tsRNAs are Dicer-independent but dependent on RNaseZ and RNA polymerase III transcription termination (Figure 1d) [39]. The two types of tsRNAs seemed to associate with separate sets of Ago proteins, which affected their ability to induce silencing of a reporter gene (Figure 2b). The physiological functions of these tRFs/tsRNAs remain to be determined, but the authors suggest the possibilities of Dicer-mediated cleavage of tRNAs as a regulatory or quality control mechanism, canonical RNA silencing, or regulation of cellular homeostasis by fine-tuning the small RNA pool and Dicer/Ago occupancy [38–40].

Transcription meets silencing: PASRs, TASRs, TSSa-RNAs, tiRNAs and spliRNAs

Recent studies have revealed new types of small RNA species that originate from different regions of annotated protein-coding genes. Promoter-associated sRNAs (PASRs) and termini-associated sRNAs (TASRs) are 22 to 200 nt in length, and are apparently derived from longer precursor RNAs [41]. Transcription start site-associated RNAs (TSSa-RNAs) have the mean length of 20 nt and are of both sense and antisense orientation [42], while

transcription initiation RNAs (tiRNAs) are about 18 nt long and originate largely from the same genomic strand as the TSS [43]. TSSa-RNAs are thought to be products of RNA polymerase II (RNAPII) divergent transcription and pausing. Productive RNAPII transcription elongation occurs unidirectionally, and divergent transcription and TSSa-RNAs might be a transcription regulatory mechanism [42]. Another group of RNAs that may be related to RNAPII backtracking is splice-site RNAs (spliRNAs) [13]. SpliRNAs are small RNAs whose 3' termini map precisely to the splice donor site, are enriched in the nucleus and are present at internal exons. The authors suggest that the accumulation of both tiRNAs and spliRNAs might reflect the relation between RNAPII activity and nucleosome positioning [13]. A recent study suggests that TSSa-RNAs/tiRNAs are unlikely to be derived from RNAPII backtracking, but instead emerge as a result of RNAPII protecting nascent RNAs from degradation [44]. Similarly, in this study spliRNAs are considered as remnants of splicing intermediates [44].

Other newly established small RNA classes

In addition to the small RNA species presented above, several distinct classes of sRNAs have been described lately. Small nucleolar RNAs (snoRNAs) are normally needed for alternative splicing and RNA modifications, but they also give rise to snoRNA-derived RNAs (sdRNAs) [45,46]. The snoRNA ACA45 is a precursor for a Dicer-dependent sdRNA that acts as a miRNA in a reporter assay, indicating that sdRNAs may function in gene expression regulation (Figure 2a) [45]. In the filamentous fungus *Neurospora crassa*, DNA damage induces the synthesis of a class of RNAs known as QDE-2-interacting small RNAs (qiRNAs) [47]. qiRNAs match almost exclusively to rDNA loci, and are probably essential components of the DNA damage response pathway of the fungus. Small RNA sequencing in *Neurospora* has also recently detected fungal miRNA-like small RNAs (milRNAs) [48]. milRNAs are synthesized by at least four distinct pathways and are able to regulate the levels of specific target genes (Figure 2a). Vault particles are conserved organelles found in most animals that are composed of several proteins and three vault RNAs (vRNAs). Small RNAs matching to vRNAs (svRNAs) were discovered in human breast cancer samples [49]. svRNAs are synthesized by a Dicer-dependent pathway, and they are capable of guiding sequence-specific target RNA cleavage *in vitro*. These various examples indicate that high-throughput sequencing projects continuously add complexity to the already extensive list of small RNAs.

Perspectives

During the recent years, next-generation sequencing efforts have resulted in a plethora of small RNA reads. The analysis of these data has shown that regulation of gene expression is an exceedingly complex phenomenon and we are still in the midst of discovering the executors of this process. Therefore, caution should be exercised in categorizing these sequence reads. It may well be that many reads of abundant housekeeping RNAs are not unspecific degradation products, but represent *bona fide*, biologically relevant RNA species. On the other hand, as the libraries originate from various organisms, cell types and developmental or physiological stages, there may be functional or mechanistic overlap that is not immediately evident at the sequence level. It is possible that seemingly novel classes of small RNAs might actually represent previously described molecules. Thus, scientists should be conservative in assigning novel names for newly described small RNAs, as this will eventually lead to an overload of terminology.

Many of the small noncoding RNAs presented above show similar characteristics. As an example, miRNAs, tRFs, sdRNAs, svRNAs, qiRNAs and some endo-siRNAs seem to be Dicer-dependent, but their Microprocessor-dependence, subcellular localization and affinity

to Argonautes may differ. Moreover, miRNA-like characteristics are continuously described in small RNAs that are not canonical miRNAs (e.g. mirtrons, sdRNAs, svRNAs, milRNAs). It is conceivable that the abundance of small RNAs is a way to control the silencing machinery by regulating Dicer/Drosha activity and Ago occupancy. In *S. pombe*, there is a surveillance mechanism that prevents unwanted small RNA molecules from entering the silencing pathway [50]. Other organisms may accomplish the same goal by “titration” of the silencing components.

Remarkably, the predominant function of sRNAs is to induce silencing of gene expression. While there are some examples to the contrary, base-pairing of a sRNA to a target sequence typically results in repressed gene expression through a variety of mechanisms (Figure 2) [20]. It remains to be discovered if in the context of specific proteins or cellular conditions sRNAs might also have widespread roles in gene activation.

All in all, regulation of gene expression in a eukaryotic cell appears to be largely dictated by the composition of its small RNA repertoire. As the roster of small RNAs keeps on growing, there is a dire need for a set of guidelines for their systematic classification. We are only beginning to appreciate the amazing complexity of these tiny molecules and the discernment of common features will surely accelerate our understanding of the biological functions of specific sRNAs.

Acknowledgments

This work was supported by the US National Institutes of Health (GM071654), Keck, and Peter Gruber Foundations (AEP). APA was supported by the Academy of Finland Centre of Excellence Program in Virus Research and the Sigrid Jusélius Foundation. The contents of this review are solely the responsibility of the authors and do not necessarily represent the official views of the funding sources.

References

1. Vazquez F, Legrand S, Windels D. The biosynthetic pathways and biological scopes of plant small RNAs. *Trends Plant Sci.* 2010; 15:337–345. [PubMed: 20427224]
2. Sayed D, Abdellatif M. MicroRNAs in development and disease. *Physiol Rev.* 2011; 91:827–887. [PubMed: 21742789]
3. Kozomara A, Griffiths-Jones S. miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res.* 2011; 39:D152–157. [PubMed: 21037258]
4. Krol J, Loedige I, Filipowicz W. The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet.* 2010; 11:597–610. [PubMed: 20661255]
- 5. Cifuentes D, Xue H, Taylor DW, Patnode H, Mishima Y, Cheloufi S, Ma E, Mane S, Hannon GJ, Lawson ND, et al. A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science.* 2010; 328:1694–1698. These three papers reveal a novel miRNA processing pathway by showing that the maturation of pre-miR-451 does not require Dicer processing but is dependent on the catalytic activity of Ago2. [PubMed: 20448148]
- 6. Cheloufi S, Dos Santos CO, Chong MM, Hannon GJ. A dicer-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature.* 2010; 465:584–589. These three papers reveal a novel miRNA processing pathway by showing that the maturation of pre-miR-451 does not require Dicer processing but is dependent on the catalytic activity of Ago2. [PubMed: 20424607]
- 7. Yang JS, Maurin T, Robine N, Rasmussen KD, Jeffrey KL, Chandwani R, Papapetrou EP, Sadelain M, O'Carroll D, Lai EC. Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc Natl Acad Sci U S A.* 2010; 107:15163–15168. These three papers reveal a novel miRNA processing pathway by showing that the maturation of pre-miR-451 does not require Dicer processing but is dependent on the catalytic activity of Ago2. [PubMed: 20699384]
- 8. Okamura K, Hagen JW, Duan H, Tyler DM, Lai EC. The mirtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell.* 2007; 130:89–100. Epub 2007 Jun 28. In these two

studies, mirtrons were described as miRNAs that do not follow the canonical processing pathway but originate from intron splicing and bypass Drosha cleavage. [PubMed: 17599402]

- 9. Ruby JG, Jan CH, Bartel DP. Intronic microRNA precursors that bypass Drosha processing. *Nature*. 2007; 448:83–86. Epub 2007 Jun 24. In these two studies, mirtrons were described as miRNAs that do not follow the canonical processing pathway but originate from intron splicing and bypass Drosha cleavage. [PubMed: 17589500]
10. Babiarz JE, Ruby JG, Wang Y, Bartel DP, Blelloch R. Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes Dev*. 2008; 22:2773–2785. [PubMed: 18923076]
11. Berezikov E, Robine N, Samsonova A, Westholm JO, Naqvi A, Hung JH, Okamura K, Dai Q, Bortolamiol-Becet D, Martin R, et al. Deep annotation of *Drosophila melanogaster* microRNAs yields insights into their processing, modification, and emergence. *Genome Res*. 2011; 21:203–215. [PubMed: 21177969]
12. Wyman SK, Knouf EC, Parkin RK, Fritz BR, Lin DW, Dennis LM, Krouse MA, Webster PJ, Tewari M. Post-transcriptional generation of miRNA variants by multiple nucleotidyl transferases contributes to miRNA transcriptome complexity. *Genome Res*. 2011
13. Taft RJ, Simons C, Nahkuri S, Oey H, Korbie DJ, Mercer TR, Holst J, Ritchie W, Wong JJ, Rasko JE, et al. Nuclear-localized tiny RNAs are associated with transcription initiation and splice sites in metazoans. *Nat Struct Mol Biol*. 2010; 17:1030–1034. [PubMed: 20622877]
14. Shi W, Hendrix D, Levine M, Haley B. A distinct class of small RNAs arises from pre-miRNA-proximal regions in a simple chordate. *Nat Struct Mol Biol*. 2009; 16:183–189. [PubMed: 19151725]
15. Bartel DP. MicroRNAs. target recognition and regulatory functions. *Cell*. 2009; 136:215–233. [PubMed: 19167326]
- 16. Chi SW, Zang JB, Mele A, Darnell RB. Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature*. 2009; 460:479–486. These papers introduce HITS-CLIP, CLIP-seq and PAR-CLIP as complementary, high-throughput methods to analyze Argonaute-bound sequences and miRNA-mRNA interactions. The results from these studies pave the way towards understanding miRNA targeting *in vivo*. [PubMed: 19536157]
- 17. Zisoulis DG, Lovci MT, Wilbert ML, Hutt KR, Liang TY, Pasquinelli AE, Yeo GW. Comprehensive discovery of endogenous Argonaute binding sites in *Caenorhabditis elegans*. *Nat Struct Mol Biol*. 2010; 17:173–179. These papers introduce HITS-CLIP, CLIP-seq and PAR-CLIP as complementary, high-throughput methods to analyze Argonaute-bound sequences and miRNA-mRNA interactions. The results from these studies pave the way towards understanding miRNA targeting *in vivo*. [PubMed: 20062054]
- 18. Leung AK, Young AG, Bhutkar A, Zheng GX, Bosson AD, Nielsen CB, Sharp PA. Genome-wide identification of Ago2 binding sites from mouse embryonic stem cells with and without mature microRNAs. *Nat Struct Mol Biol*. 2011; 18:237–244. These papers introduce HITS-CLIP, CLIP-seq and PAR-CLIP as complementary, high-throughput methods to analyze Argonaute-bound sequences and miRNA-mRNA interactions. The results from these studies pave the way towards understanding miRNA targeting *in vivo*. [PubMed: 21258322]
- 19. Hafner M, Landthaler M, Burger L, Khorshid M, Hausser J, Berninger P, Rothballer A, Ascano M Jr, Jungkamp AC, Munschauer M, et al. Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell*. 2010; 141:129–141. These papers introduce HITS-CLIP, CLIP-seq and PAR-CLIP as complementary, high-throughput methods to analyze Argonaute-bound sequences and miRNA-mRNA interactions. The results from these studies pave the way towards understanding miRNA targeting *in vivo*. [PubMed: 20371350]
20. Huntzinger E, Izaurralde E. Gene silencing by microRNAs. contributions of translational repression and mRNA decay. *Nat Rev Genet*. 2011; 12:99–110. [PubMed: 21245828]
- 21. Guo H, Ingolia NT, Weissman JS, Bartel DP. Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature*. 2010; 466:835–840. The authors employ ribosome profiling and proteomics analyses to elegantly show that the majority of miRNA-mediated downregulation in HeLa cells occurs through mRNA destabilization. [PubMed: 20703300]
22. Djuranovic S, Nahvi A, Green R. A parsimonious model for gene regulation by miRNAs. *Science*. 2011; 331:550–553. [PubMed: 21292970]

23. Siomi MC, Sato K, Pezic D, Aravin AA. PIWI-interacting small RNAs: the vanguard of genome defence. *Nat Rev Mol Cell Biol.* 2011; 12:246–258. [PubMed: 21427766]
24. Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R, Hannon GJ. Specialized piRNA pathways act in germline and somatic tissues of the *Drosophila* ovary. *Cell.* 2009; 137:522–535. Epub 2009 Apr 2023. [PubMed: 19395010]
25. Li C, Vagin VV, Lee S, Xu J, Ma S, Xi H, Seitz H, Horwich MD, Syrzycka M, Honda BM, et al. Collapse of germline piRNAs in the absence of Argonaute3 reveals somatic piRNAs in flies. *Cell.* 2009; 137:509–521. Epub 2009 Apr 2023. [PubMed: 19395009]
26. Haase AD, Fenoglio S, Muerdter F, Guzzardo PM, Czech B, Pappin DJ, Chen C, Gordon A, Hannon GJ. Probing the initiation and effector phases of the somatic piRNA pathway in *Drosophila*. *Genes Dev.* 2010; 24:2499–2504. [PubMed: 20966049]
27. Aravin AA, Sachidanandam R, Bourc'his D, Schaefer C, Pezic D, Toth KF, Bestor T, Hannon GJ. A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol Cell.* 2008; 31:785–799. [PubMed: 18922463]
28. Rouget C, Papin C, Boureux A, Meunier AC, Franco B, Robine N, Lai EC, Pelisson A, Simonelig M. Maternal mRNA deadenylation and decay by the piRNA pathway in the early *Drosophila* embryo. *Nature.* 2010; 467:1128–1132. [PubMed: 20953170]
29. Ghildiyal M, Seitz H, Horwich MD, Li C, Du T, Lee S, Xu J, Kittler EL, Zapp ML, Weng Z, et al. Endogenous siRNAs derived from transposons and mRNAs in *Drosophila* somatic cells. *Science.* 2008; 320:1077–1081. [PubMed: 18403677]
30. Tam OH, Aravin AA, Stein P, Girard A, Murchison EP, Cheloufi S, Hodges E, Anger M, Sachidanandam R, Schultz RM, et al. Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature.* 2008; 453:534–538. Epub 2008 Apr 2010. [PubMed: 18404147]
31. Watanabe T, Totoki Y, Toyoda A, Kaneda M, Kuramochi-Miyagawa S, Obata Y, Chiba H, Kohara Y, Kono T, Nakano T, et al. Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature.* 2008; 453:539–543. Epub 2008 Apr 2010. [PubMed: 18404146]
32. Okamura K, Chung WJ, Ruby JG, Guo H, Bartel DP, Lai EC. The *Drosophila* hairpin RNA pathway generates endogenous short interfering RNAs. *Nature.* 2008; 453:803–806. [PubMed: 18463630]
33. Kawamura Y, Saito K, Kin T, Ono Y, Asai K, Sunohara T, Okada TN, Siomi MC, Siomi H. *Drosophila* endogenous small RNAs bind to Argonaute 2 in somatic cells. *Nature.* 2008; 453:793–797. [PubMed: 18463636]
- 34. Pak J, Fire A. Distinct populations of primary and secondary effectors during RNAi in *C. elegans*. *Science.* 2007; 315:241–244. Epub 2006 Nov 2023. These studies show that most *C. elegans* siRNAs are secondary to Dicer-cleaved primary siRNAs and products of RNA-dependent RNA polymerases. [PubMed: 17124291]
- 35. Sijen T, Steiner FA, Thijssen KL, Plasterk RH. Secondary siRNAs result from unprimed RNA synthesis and form a distinct class. *Science.* 2007; 315:244–247. These studies show that most *C. elegans* siRNAs are secondary to Dicer-cleaved primary siRNAs and products of RNA-dependent RNA polymerases. [PubMed: 17158288]
36. Burkhart KB, Guang S, Buckley BA, Wong L, Bochner AF, Kennedy S. A pre-mRNA-associating factor links endogenous siRNAs to chromatin regulation. *PLoS Genet.* 2011; 7:e1002249. [PubMed: 21901112]
37. Fagegaltier D, Bouge AL, Berry B, Poisot E, Sismeiro O, Coppee JY, Theodore L, Voinnet O, Antoniewski C. The endogenous siRNA pathway is involved in heterochromatin formation in *Drosophila*. *Proc Natl Acad Sci U S A.* 2009; 106:21258–21263. [PubMed: 19948966]
38. Lee YS, Shibata Y, Malhotra A, Dutta A. A novel class of small RNAs. tRNA-derived RNA fragments (tRFs). *Genes Dev.* 2009; 23:2639–2649. [PubMed: 19933153]
- 39. Haussecker D, Huang Y, Lau A, Parameswaran P, Fire AZ, Kay MA. Human tRNA-derived small RNAs in the global regulation of RNA silencing. *RNA.* 2010; 16:673–695. By combining deep sequencing data and biochemical assays, these three papers show that biologically functional sRNAs are derived from tRNAs. [PubMed: 20181738]

- 40. Cole C, Sobala A, Lu C, Thatcher SR, Bowman A, Brown JW, Green PJ, Barton GJ, Hutvagner G. Filtering of deep sequencing data reveals the existence of abundant Dicer-dependent small RNAs derived from tRNAs. *RNA*. 2009; 15:2147–2160. By combining deep sequencing data and biochemical assays, these three papers show that biologically functional sRNAs are derived from tRNAs. [PubMed: 19850906]
- 41. Kapranov P, Cheng J, Dike S, Nix DA, Dutttagupta R, Willingham AT, Stadler PF, Hertel J, Hackermuller J, Hofacker IL, et al. RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science*. 2007; 316:1484–1488. [PubMed: 17510325]
- 42. Seila AC, Calabrese JM, Levine SS, Yeo GW, Rahl PB, Flynn RA, Young RA, Sharp PA. Divergent transcription from active promoters. *Science*. 2008; 322:1849–1851. [PubMed: 19056940]
- 43. Taft RJ, Glazov EA, Cloonan N, Simons C, Stephen S, Faulkner GJ, Lassmann T, Forrest AR, Grimmond SM, Schroder K, et al. Tiny RNAs associated with transcription start sites in animals. *Nat Genet*. 2009; 41:572–578. [PubMed: 19377478]
- 44. Valen E, Preker R, Andersen PR, Zhao X, Chen Y, Ender C, Dueck A, Meister G, Sandelin A, Jensen TH. Biogenic mechanisms and utilization of small RNAs derived from human protein-coding genes. *Nat Struct Mol Biol*. 2011
- 45. Ender C, Krek A, Friedlander MR, Beitzinger M, Weinmann L, Chen W, Pfeffer S, Rajewsky N, Meister G. A human snoRNA with microRNA-like functions. *Mol Cell*. 2008; 32:519–528. [PubMed: 19026782]
- 46. Taft RJ, Glazov EA, Lassmann T, Hayashizaki Y, Carninci P, Mattick JS. Small RNAs derived from snoRNAs. *RNA*. 2009; 15:1233–1240. [PubMed: 19474147]
- 47. Lee HC, Chang SS, Choudhary S, Aalto AP, Maiti M, Bamford DH, Liu Y. qiRNA is a new type of small interfering RNA induced by DNA damage. *Nature*. 2009; 459:274–277. [PubMed: 19444217]
- 48. Lee HC, Li L, Gu W, Xue Z, Crosthwaite SK, Pertsemlidis A, Lewis ZA, Freitag M, Selker EU, Mello CC, et al. Diverse pathways generate microRNA-like RNAs and dicer-independent small interfering RNAs in fungi. *Mol Cell*. 2010; 38:803–814. The authors show that in contrast to previous analyses, miRNAs are found also in the fungal kingdom. The miRNAs of *Neurospora crassa* are synthesized by at least four distinct pathways and function as miRNAs in a reporter assay. [PubMed: 20417140]
- 49. Persson H, Kvist A, Vallon-Christersson J, Medstrand P, Borg A, Rovira C. The non-coding RNA of the multidrug resistance-linked vault particle encodes multiple regulatory small RNAs. *Nat Cell Biol*. 2009; 11:1268–1271. [PubMed: 19749744]
- 50. Buhler M, Spies N, Bartel DP, Moazed D. TRAMP-mediated RNA surveillance prevents spurious entry of RNAs into the *Schizosaccharomyces pombe* siRNA pathway. *Nat Struct Mol Biol*. 2008; 15:1015–1023. [PubMed: 18776903]
- 51. Batista PJ, Ruby JG, Claycomb JM, Chiang R, Fahlgren N, Kasschau KD, Chaves DA, Gu W, Vasale JJ, Duan S, et al. PRG-1 and 21U-RNAs interact to form the piRNA complex required for fertility in *C. elegans*. *Mol Cell*. 2008; 31:67–78. Epub 2008 Jun 20. [PubMed: 18571452]

Figure 1. Many ways to make a silencer

Biogenesis pathways of small RNAs (sRNAs). **(a)** Following transcription of a primary miRNA (pri-miRNA) transcript, the Drosha complex or spliceosome releases the hairpin precursor miRNA (pre-miRNA) from typical or mirtron type miRNAs, respectively. The pre-miRNA or transcribed short hairpin (shRNA) miRNA transcripts undergo final processing by Dicer before incorporation into Argonaute complexes. The pre-miR-451 is an exception that utilizes Argonaute instead of Dicer in the final cleavage step. **(b)** Primary piRNAs from genomic clusters are loaded onto Piwi-related proteins and target homologous sequences for cleavage and generation of secondary piRNAs. These piRNAs are bound by distinct Piwi family proteins, which catalyze the generation of additional piRNAs through the cleavage and processing of RNAs with antisense sequences. **(c)** Endo-siRNAs derive from intramolecular hairpins or sense-antisense transcripts that base-pair and are cleaved by Dicer. The ~21 nt endo-siRNAs are bound by Argonaute proteins and in some organisms initiate the production of secondary endo-siRNAs by RNA-dependent RNA polymerases (RdRPs) that use cleaved target RNAs as templates. **(d)** Precursor tRNAs can be cleaved by RNaseZ to produce Type II tsRNAs from the 3' termini. Following 3'-end CCA-addition, Dicer processes some tRNAs, potentially those that are misfolded, to produce Type I tsRNAs.

Figure 2. Many roads to silencing

Mechanisms used by sRNAs to silence gene expression. **(a)** Typically, miRNAs, and possibly miRNAs and sdRNAs, partially pair to mRNA target sequences and promote mRNA degradation through the recruitment of deadenylases by GW182 proteins bound to Argonaute. Alternatively, miRISC inhibits translation initiation, stalls translation elongation or stimulates proteolysis of nascent peptides encoded by the target mRNA. **(b)** Perfect pairing of many types of sRNAs results in cleavage of the target mRNA by certain Argonaute proteins. **(c)** Endo-siRNAs, piRNAs and possibly many of the sRNAs derived from protein-coding regions of the genome direct DNA and histone modifications (red stars) that regulate transcriptional activity.

Table 1

Classes of small non-coding RNAs.

ncRNA	Abbreviation	Model organism(s)	References
microRNA	miRNA	Most eukaryotic organisms except fungi; viruses	[4,20]
miRNA-offset RNA	moRNA	<i>Ciona intestinalis</i> , mammals	[13,14]
short hairpin-derived miRNA	shRNA-derived miRNA	<i>Mus musculus</i>	[10]
miRNA-like small RNA	miRNA	<i>Neurospora crassa</i>	[48]
mirtron	-	<i>D. melanogaster</i> , <i>C. elegans</i> , mammals	[8–10]
Piwi-interacting RNA	piRNA	<i>D. melanogaster</i> , mammals, <i>Danio rerio</i> , <i>Xenopus laevis</i> , <i>Bombyx mori</i>	[23]
21U-RNA	-	<i>C. elegans</i>	[51]
endogenous small interfering RNA	endo-siRNA	<i>D. melanogaster</i> , <i>C. elegans</i> , mammals	[29–35]
tRNA-derived RNA fragment, tRNA-derived small RNA	tRF, tsRNA	Mammals	[38–40]
promoter-associated small RNA, terminus-associated small RNA	PASR, TASR	Mammals	[41]
transcription start site-associated RNA, transcription initiation RNA	TSSa-RNA, tiRNA	Mammals, <i>Gallus gallus</i> , <i>D. melanogaster</i>	[42–44]
splice-site RNA	spliRNA	Metazoans	[13,44]
snoRNA-derived RNA	sdRNA	Metazoans, <i>Arabidopsis thaliana</i> , <i>Schizosaccharomyces pombe</i>	[13,45,46]
QDE-2-interacting small RNA	qiRNA	<i>N. crassa</i>	[47]
small vault RNA	svRNA	<i>Homo sapiens</i>	[49]