Published in final edited form as:

J Exp Child Psychol. 2012 August ; 112(4): 361–377. doi:10.1016/j.jecp.2012.03.007.

Seven-month-old infants chunk items in memory

Mariko Moher^{a,1}, Arin S. Tuerk^{a,1}, and Lisa Feigenson^a

^aDepartment of Psychological & Brain Sciences, Johns Hopkins University 3400 N. Charles St., Baltimore, MD 21218 U.S.A.

Abstract

Although working memory has a highly constrained capacity limit of 3 to 4 items, both adults and toddlers can increase the total amount of stored information by chunking object representations in memory. To examine the developmental origins of chunking, we used a violation-of-expectation procedure to ask whether 7-month-old infants, whose working memory capacity is still maturing, also can chunk items in memory. In Experiment 1 we found that in the absence of chunking cues, infants failed to remember 3 identical hidden objects. In Experiments 2 and 3 we found that infants successfully remembered 3 hidden objects when provided with overlapping spatial and featural chunking cues. In Experiment 4 we found that infants did not chunk when provided with either spatial or featural chunking cues alone. Finally, in Experiment 5 we found that infants also failed to chunk when spatial and featural cues specified different chunks (i.e., were pitted against each other). Taken together, these results suggest that chunking is available before working memory capacity has matured, but still may undergo important development over the first year of life.

Keywords

working memory; cognitive development; chunking; infants

Introduction

Thinking about things that are not directly perceptually accessible requires memory. Although both infants (Bauer, 2007; Oakes & Bauer, 2007; Rovee-Collier, 1999) and adults (Squire, 2009; Tulving, 2002) can store durable memory representations over long periods of time, they also must have access to a form of memory that can create and manipulate representations rapidly and on the fly. Working memory, which allows for the temporary storage of information, has been shown to serve this role across the lifespan (Cowan, 1997; Oakes & Bauer, 2007). A hallmark of working memory is its highly limited capacity, whereby only a few items can be represented at any given time. However, adults, children, and even 14-month-old toddlers have been shown to mentally reorganize or *chunk* information in working memory, thereby increasing the total amount of remembered

© 2012 Elsevier Inc. All rights reserved.

Corresponding author: Mariko Moher, mmoher@wjh.harvard.edu Phone: 1(617)496-2695, Fax: 1(617)495-3728.
atuerk@wjh.harvard.edu, feigenson@jhu.edu.

¹Present address: Department of Psychology, Harvard University William James Hall, 33 Kirkland St., Cambridge, MA 02138 U.S.A.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

material. Here we asked whether this type of chunking of items is also available to younger, pre-verbal infants whose working memory has not yet matured to adult-like capacity.

Working memory capacity limits in infancy

From early in life infants represent hidden objects in memory, an ability critical to learning about the world. Consider the everyday challenges infants face, like watching a favorite toy become covered by a blanket or seeing one's mother disappear into another room. In order to lift the blanket and retrieve the hidden object, or in order to crawl after a parent, infants must rely on representations stored in memory. Laboratory tasks have demonstrated that infants can represent such absent or occluded objects. For example, 5-month-old infants who saw a doll hidden behind a screen, then saw another doll added behind the same screen, looked longer when the screen was lifted to reveal unexpected outcomes of one or three dolls than at the expected outcome of two dolls (Wynn, 1992). Because the two dolls were never shown simultaneously during the hiding event, this looking preference suggests that infants maintained a representation of the first object in memory, then mentally updated it to reflect the addition of the second object (see also Feigenson, Carey, & Spelke, 2002; Koechlin, Dehaene & Mehler, 1997; Simon, Hespos, & Rochat, 1995; Uller, Carey, Huntley-Fenner, & Klatt, 1999).

The representations that allow infants to remember a toy hidden under a blanket, or to represent the outcome when two dolls are serially hidden, need not reside in working memory. For example, these could be long-term representations with greater longevity and robustness. However, one piece of evidence suggesting that infants may rely on working memory in tasks like Wynn's comes from the striking cases in which infants fail to remember. Across a variety of paradigms, adults have been shown to concurrently represent a maximum of only three or four visual items over brief durations (e.g., Alvarez & Cavanagh, 2004; Broadbent, 1975; Cowan, 2001; Luck & Vogel, 1997; Song & Jiang, 2006; Y. Xu, 2002; Y. Xu & Chun, 2006). This limit on the number of remembered items has been taken as a signature of visual short-term memory or working memory representations. A similar limit has emerged from tasks measuring the number of hidden objects infants can remember over durations ranging from less than a second to several seconds. In one series of studies, the memory capacity of 12- to 20-month-old infants was measured by hiding varying numbers of objects in a box, allowing infants to retrieve either all or just a subset of the objects, and then asking whether infants continued searching the box for any remaining objects. When three or fewer objects were hidden and they had retrieved only a subset, infants successfully continued searching for the missing objects. However, when more than three objects were hidden, infants failed to continue searching (Barner, Thalwitz, Wood, & Carey, 2007; Feigenson & Carey, 2003; 2005; Feigenson & Halberda, 2004, 2008). These results suggest that infants can successfully represent one, two or three hidden objects, but fail to remember four or more.

A similar capacity limit has been revealed when measuring infants' visual short-term memory (VSTM) for very briefly presented items, using methods more similar to those used with adults (e.g., Luck & Vogel, 1997). In that experiment, 10- and 13-month-old infants saw two flickering streams of colored squares. In the Changing Stream, one of the squares changed its color between each 500 ms flicker while the other squares maintained their colors. In the Non-Changing Stream, all of the squares' colors stayed constant across flickers. Infants looked longer at the Changing than the Non-Changing Stream when the array contained 1, 2, 3, or 4 squares, suggesting that they had maintained representations of the squares, compared these in memory, and noticed the color change. However, they failed to look longer at the Changing Stream when the array contained six squares (Ross-Sheehy, Oakes, & Luck, 2003), suggesting that infants could not remember the features of this many items. This similarity in the capacity limits observed with 10- to 20-month-old children and

with adults, spanning a range of methods, suggests that working memory capacity may be in some ways consistent across development, from durations ranging from 500 ms to the longer 4- to 8-second durations that have been demonstrated in adult working memory (Noles, Scholl, & Mitroff, 2008; Zhang & Luck, 2009).

Chunking items in memory

Although working memory is limited in the number of items it can maintain, it also appears flexible in terms of what can count as an item. In particular, adults have been shown to store representations not just of individual items, but also of groups – or chunks – of items. Such chunking appears to expand the total amount of information that can be remembered. For example, adults might have difficulty remembering the 9-letter sequence LAXJFKSFO, yet easily remember the 3-acronym sequence of airport codes: LAX (Los Angeles), JFK (John F. Kennedy), and SFO (San Francisco). Adults have been shown to chunk using knowledge about race times (Chase & Ericsson, 1981; Ericsson, Chase & Faloon, 1980), the mathematical constant π (Ericsson, Delaney, Weaver & Mahadevan, 2004), chess configurations (Chase & Simon, 1973), word sequences (Simon, 1974), and ordinal relations (Mathy & Feldman, 2012), all of which rely on information stored in long-term memory. Adults also can chunk using visual statistics learned during a short laboratory session. Observers showed a memory advantage for items from arrays in which particular items tended to co-occur, compared to arrays in which no such regularities were present (Brady, Konkle & Alvarez, 2009; see also Fiser & Aslin, 2005), and were better at recalling lists containing recently learned word-pairings than at recalling individually learned words (Cowan, Chen & Rouder, 2004). In all of these cases, strong associations between individual items allowed the items to be grouped in memory, thereby creating a more efficiently stored representation.

What are the developmental origins of chunking? One possibility is that chunking is an acquired strategy learned through formal instruction. Alternatively, chunking may be a foundational memory process that emerges early, without explicit training. One way to test these possibilities is to examine young children who are unlikely to have explicit control over memory strategies or to have received instruction in such strategies. This approach has shown that, like adults, 14-month-old toddlers can increase memory via chunking. Toddlers who saw an array of four identical, equally spaced objects and then watched the experimenter hide all four objects inside a box failed just a few seconds later to remember how many objects had been hidden. That is, toddlers did not continue searching for the missing objects when four objects had been hidden and any subset of these was retrieved (Feigenson & Carey, 2003, 2005). However when the four identical objects initially were presented in two spatially separated groups of two (e.g., two objects on the left side of a hiding location, and two objects on the right), toddlers now searched successfully (Feigenson & Halberda, 2004; Rosenberg & Feigenson, submitted). This simple manipulation of spatial separation apparently allowed toddlers to remember two groups, each containing two objects, in the face of their failure to remember a single group of four. Like adults, toddlers can use several types of information as a basis for chunking. In addition to the spatiotemporal chunking cues just described, 14-month-olds can use conceptual knowledge of familiar object kinds (e.g., two cats and two cars), perceptual similarity between objects, and shared verbal labels to chunk objects in memory (Feigenson & Halberda, 2008).

Although the above studies show that young children chunk without instruction, several questions remain regarding the origins of this ability. First, it is unclear whether children require linguistic competence in order to chunk (non-human animals have been shown to chunk, but only after many hundreds of training trials – see Terrace, 1987; 1991; Terrace & Chen, 1991). Although variable in their linguistic abilities, by the time they are 14 months

old children typically comprehend and produce tens and even hundreds of words (Fenson, Dale, Reznick, Bates, Thal & Pethick, 1994). The finding that 14-month-old toddlers can use shared novel labels to chunk objects together (Feigenson & Halberda, 2008) demonstrates that linguistic information can drive chunking. The question remains whether linguistic coding is necessary for chunking, at least in humans.

Second, it is unknown whether children younger than 14 months have access to chunking. Mounting evidence suggests that working memory capacity undergoes significant development during the first year of life. By 10 to 14 months infants can remember both the existence and at least some of the features of up to three objects across a range of tasks (Feigenson & Carey, 2003; 2005; Feigenson, Carey & Hauser, 2002; Rose, Feldman, & Jankowski, 2001; Ross-Sheehy et al., 2003). However, younger infants appear to have a smaller working memory capacity, at least for object features (Káldy & Leslie, 2005; Kibbe & Leslie, 2011; Ross-Sheehy et al., 2003). These studies suggest that infants' capacity for remembering the presence of hidden objects may not asymptote until about 10 months of age, and that infants' memory capacity for the features of hidden objects may lag even further.

The finding that working memory capacity undergoes important development over the first year of life raises the question of whether chunking is available during this time of change. Chunking may not be performed until working memory has reached more adult-like capacity, at approximately 10 months. Alternatively, chunking might be available throughout the lifespan, even while working memory capacity is still increasing. Testing for the deployment of chunking in infants younger than those previously studied can help address both this question and the question of whether chunking requires linguistic abilities.

Finally, evidence for untrained chunking in very young children is currently limited to a single experimental paradigm: manual search (Feigenson & Halberda, 2004, 2008; Rosenberg & Feigenson, submitted). The claim that chunking is available very early in life would be bolstered by evidence using different methods and measuring a different behavior.

The current experiments

Here we asked whether children younger than 10 months, whose working memory capacity has not yet matured, can use chunking to expand the amount of remembered information. Estimates of working memory capacity at any age will depend on the methods of the task and on the evidence taken to indicate successful storage of an item. For example, studies that test for infants' memory for an object's presence (e.g., Wynn, 1992) yield evidence that infants remember more items than do studies that test for infants' memory of particular object features (e.g., Kibbe & Leslie, 2011; Ross-Sheehy et al., 2003). In the present investigation our goal was not to make claims about the absolute number of items 7-month-old infants can remember, but rather to test the influence of chunking cues within the context of a single task. That is, we asked whether infants provided with chunking cues remember more objects than infants not provided with such cues.

We tested 7-month-olds because current evidence suggests that for infants of this age, working memory capacity has not yet reached the mature three to four item limit (Káldy & Leslie, 2005; Rose et al., 2001; Ross-Sheehy et al., 2003), and because 7-month-olds are considerably younger than the 14-month-old toddlers with whom chunking has already been demonstrated (Feigenson & Halberda, 2004, 2008). Our first step was to ask whether 7-month-old infants can remember the presence of three hidden objects in the absence of chunking cues—a task which we predicted 7-month-old infants would fail, based on the findings reviewed earlier. We showed infants three objects sequentially hidden behind a

screen, then lifted the screen to reveal either the expected outcome of three objects or the unexpected outcome of two objects.

After establishing that 7-month-old infants fail to remember three hidden objects when no chunking cues were provided (Experiment 1), we asked whether providing different types of chunking cues would allow infants to succeed. In Experiments 2 and 3 we asked whether 7-month-old infants would remember three hidden objects when given both spatial and featural cues to group objects into chunks. In Experiment 4 we asked whether infants would remember three hidden objects when given only spatial *or* featural cues, in isolation. Finally, in Experiment 5 we asked whether infants would remember three hidden objects when spatial and featural chunking cues suggested opposing parses of the array (i.e., when spatial and featural chunking cues were pitted against each other).

Experiment 1

In Experiment 1 we first sought to establish the limits of 7-month-old infants' memory capacity within the context of our task. Although previous research has found that infants of roughly this age have difficulty maintaining the features of more than one item in working memory (Káldy & Leslie, 2005; Ross-Sheehy et al., 2003), the limits of their ability to represent the *existence* of three objects (regardless of whether they remember the objects' features) remains unknown. Because by 5 months of age infants can remember the existence of at least two hidden objects (Kibbe & Leslie, 2011; Koechlin et al., 1995; Wynn, 1992), we began by asking whether 7-month-old infants also can remember three hidden objects.

Method

Participants—Twenty healthy, full-term infants from the Baltimore, MD area participated (10 boys; 5 Black or African American, 13 Caucasian, and 2 Other Race). Their mean age was 7 months, 2 days (range: 6 months, 18 days to 7 months, 13 days). Eight additional infants were tested but excluded from analysis (5 for fussiness, 1 for experimental error, 2 for parent or sibling interference). Infants were recruited by telephone and by mail, and received a small gift (shirt, book, or stuffed toy) to thank them for their participation.

Apparatus and Stimuli—Infants sat in a highchair approximately 95 cm from a black wooden puppet stage (40 cm high, 129 cm wide, and 50 cm deep). A black curtain could be lowered to cover the entire stage.

A video camera mounted within the stage provided a direct view of infants' gaze, and a wall-mounted camera behind infants captured a view of the stage. Both images were recorded digitally and viewed live in an adjacent room by a trained observer who did not know the experimental condition in which infants were being tested (the portion of the video screen showing the stage was covered during coding). An experimenter hidden behind the stage inserted objects onto and removed objects from the stage, and could observe whether infants were watching these events via a hidden television monitor.

A black foam core screen (16 cm high by 80 cm wide) with a hidden ledge (10 cm deep) was used to hide the stimulus objects. The objects were three identical green and white balls (9 cm diameter) decorated with colorful dots. Squeezing the balls produced a squeaking sound that was used to draw infants' attention to each object presentation.

Procedure—The basic procedure was the same across all experiments. Parents sat out of view behind infants and were instructed to refrain from interacting with infants. Classical music was played at a low volume throughout the experiment.

Baseline trials: To measure infants' baseline preference to look at two versus three objects, the experiment began with one 2-Object baseline trial and one 3-Object baseline trial. The same stimulus objects were used throughout the baseline and test trials. For each baseline trial the experimenter said, "Up goes the curtain," and raised the curtain to reveal the black screen already on the stage. The experimenter made sure that infants were looking (tapping the screen if they were not), then lifted the screen to reveal either two or three objects. Infants' looking was measured from the moment the screen was completely out of view. The objects remained in place until either a) infants looked away for at least 2 continuous seconds, or b) infants looked for a total of 120 seconds, as indicated by the observer in an adjacent room who pressed a button to signal the end of each trial. When each trial ended, the experimenter said, "Down goes the curtain," and lowered the curtain. For this and all subsequent experiments, whether infants saw the 2-Object or the 3-Object baseline trial first was counterbalanced across infants.

Test trials: On each of the eight test trials the experimenter said, "Up goes the curtain," while raising the curtain to reveal an empty stage. The experimenter then placed the occluding screen onto the middle of the stage. Next, she lowered a ball directly above the center of the screen, squeaked it once, then hid it behind the screen. Throughout this object presentation the experimenter's arm remained visible above the screen so that infants could infer the final hiding location of the ball. This procedure was repeated until three balls were hidden, at which point the experimenter lifted the screen and the measurement period began. On half of the test trials the screen lifted to reveal the expected outcome (three balls), and on the other half the screen lifted to reveal the unexpected outcome (two balls, with the third ball surreptitiously removed from the stage on the concealed ledge of the screen) (Figure 1). On unexpected outcome trials the remaining balls were centered behind the screen. For this and all subsequent experiments, expected and unexpected outcomes alternated across the eight test trials, with test trial order (expected outcome shown first or second) counterbalanced across infants.

All testing sessions were later re-coded by two trained observers who were unaware of the experimental condition and order of trial types. Their inter-rater agreement averaged 95.1% across all participants. Looking times greater than 2.5 standard deviations from the group mean for that trial were replaced by the same-trial average looking times of participants in the same condition. This resulted in the replacement of 3.8% of all trials.

Results

Baseline Trials—To assess infants' baseline preference to look at two versus three objects, we first analyzed looking times from the two baseline trials in a 2 (Outcome: 2 Objects vs. 3 Objects) $\times$ 2 (Trial Order: 2-Object vs. 3-Object array shown first) ANOVA. We observed no effect of Outcome; infants exhibited no significant baseline preference for either the 2-Object (9.79 seconds) or 3-Object array (7.65 seconds) ($F(1, 16) = 1.873, p = .188$) (Figure 2A).

Test Trials—We next analyzed infants' looking times from the test trials in a 2 (Outcome: 2 Objects vs. 3 Objects) $\times$ 4 (Test Pair: 1st, 2nd, 3rd, or 4th test pair) $\times$ 2 (Trial Order: 2-Object vs. 3-Object outcome shown first) ANOVA¹. This revealed a significant main effect of Test Pair ($F(3, 54) = 7.010, p < .001, \eta_p^2 = .280$); infants' looking times were longer on the first three pairs (8.34 seconds, 9.05 seconds, and 7.71 seconds, respectively) than on the last pair (5.76 seconds). There was also a significant Outcome $\times$ Test Pair interaction ($F(3,$

¹In this and all other experiments we first analyzed the data including Gender as a factor. No significant main effects or interactions involving Gender emerged in Experiment 1 or in any of the other experiments.

54) = 3.288, $p < .001$, $\eta_p^2 = .154$), driven by infants looking longer at the unexpected outcome on the second test pair, but looking equally to both outcomes on all other pairs. No other main effects or interactions were observed. Critically, there was no significant main effect of Outcome ($F(1, 18) = 0.359$, $p = .557$). Averaged across the four test pairs, infants did not look longer at unexpected 2-Object outcomes (7.55 seconds) than at expected 3-Object outcomes (7.87 seconds) (Figure 2A).

Finally, we asked whether infants' preference to look at two versus three objects in the baseline trials differed from their preference during the test trials. A 2 (Trial Type: Baseline vs. Test Trials) $\times$ 2 (Outcome) ANOVA revealed no significant main effects or interactions.

Discussion

In Experiment 1 we found that 7-month-old infants failed to remember three identical objects when they were hidden sequentially behind a single screen. This suggests that, in addition to their limited representation of specific object features (Káldy & Leslie, 2005; Ross-Sheehy et al., 2003; Kibbe & Leslie, 2011; Simon et al., 1995), infants of this age may also be limited in the number of object representations they can maintain in working memory, at least in the context of this task. It is an open question whether infants of this age might succeed at representing three hidden objects if the objects were hidden simultaneously rather than sequentially (for evidence that this type of "object first" presentation can improve infants' performance see Uller et al., 1999), or if the retention interval were shortened by presenting the objects more quickly (Diamond, 1990). However, our primary interest was whether, given a constant sequence of events, providing chunking cues would improve infants' performance.

Experiment 2

By 14 months of age toddlers can use spatial cues to chunk identical objects (Feigenson & Halberda, 2004) and can also use featural cues to chunk equally spaced objects (Feigenson & Halberda, 2008). Here, in order to maximize the chance that these much younger, 7-month-old infants would also show successful chunking, we began by presenting infants with arrays containing both spatial and featural chunking cues. First, we spatially separated the objects into two groups by hiding them behind two different screens rather than the single screen used in Experiment 1. One object was hidden behind one screen, and two objects were hidden behind the other. In the Contrasting Screens condition, we further emphasized the distinctiveness of the screens by outlining the border of one screen in pink and the other in blue. In the Identical Screens condition the two screens were identical but clearly spatially separated. Second, in both the Contrasting and Identical Screens conditions we presented objects with contrasting features. One object was a leopard-print block with a painted face, and the other two objects were the green and white balls used in Experiment 1. The number of objects hidden and the timing and duration of the hiding events were identical to those in Experiment 1.

Method

Participants—Forty healthy full-term infants from the Baltimore, MD area participated, with 20 infants in the Contrasting Screens condition and 20 in the Identical Screens condition (19 boys; 1 Asian, 4 Black or African American, 30 Caucasian, and 5 Other Race). Their mean age was 6 months, 29 days (range: 6 months, 14 days to 7 months, 15 days). Ten additional infants were tested but excluded from the final analysis (3 for fussiness, 5 for experimenter error, and 2 for parental interference).

Apparatus and Stimuli—The apparatus and stimuli were identical to those used in Experiment 1, except for two changes. First, two smaller black foam core screens (16 cm high $\times$ 25 cm wide) replaced the single screen used in Experiment 1. In the Contrasting Screens condition, one screen had a pink border and the other had a blue border (1.6 cm wide) around all four sides. In the Identical Screens condition, both screens were black. Second, one of the green and white balls used in Experiment 1 was replaced with a leopard-print block (12.5 cm high $\times$ 6.5 cm wide $\times$ 6.5 cm deep) with a schematic smiley face.

Procedure—The procedure was identical in both conditions. As in Experiment 1, infants first saw two baseline trials consisting of one 2-Object array and one 3-Object array, with order counterbalanced across infants. The curtain first was raised to reveal the two hiding screens already on place on the stage, approximately 30 cm apart (so that their endpoints were separated by the same distance as the endpoints of the single screen used in Experiment 1). On the 2-Object trial, the two screens were lifted simultaneously to reveal one ball behind one screen and one block behind the other screen. On the 3-Object trial, the screens were simultaneously lifted to reveal two balls behind one screen and one block behind the other. The side on which the ball(s) or the block appeared was counterbalanced across infants. The 3-Object array always contained two balls and one block (never two blocks and one ball).

Infants then saw eight test trials, as in Experiment 1. On each test trial the curtain first was raised to reveal an empty stage. The experimenter simultaneously lowered the two hiding screens to their resting locations 30 cm apart in the center of the stage. Infants then saw two balls sequentially hidden behind one screen and one block hidden behind the other (with side and presentation order counterbalanced across trials). Because the block did not emit a squeak, the experimenter instead squeaked an extra ball (concealed behind the stage) each time an object was presented (whether it was a ball or a block) in order to attract infants' attention. As in Experiment 1, the test trials alternately revealed the unexpected 2-object outcome and the expected 3-object outcome. The unexpected outcome always contained just one ball behind one screen and one block behind the other, and the expected outcome always contained two balls behind one screen and one block behind the other (Figure 1).

As in Experiment 1, looking times greater than 2.5 standard deviations from the group mean for that trial were replaced by the same-trial average looking times of participants in the same condition. This resulted in the replacement of 5.0% of all trials.

Results

Baseline Trials—A 2 (Outcome) $\times$ 2 (Trial Order) $\times$ 2 (Chunking Cue: Contrasting vs. Identical Screens) ANOVA revealed a marginally significant main effect of Chunking Cue ($F(1, 36) = 3.573, p = .067, \eta_p^2 = .090$). Infants in the Identical Screens condition looked longer overall (14.66 seconds) than infants in the Contrasting screens condition (9.00 seconds). However, there was no main effect of Outcome ($F(1, 36) = 1.137, p = .293$); infants looked equally at the 2-Object (11.03 seconds) and 3-Object (12.64 seconds) arrays. There were no other significant main effects or interactions.

Test Trials—We next analyzed looking times from the test trials in a 2 (Outcome) $\times$ 4 (Test Pair) $\times$ 2 (Trial Order) $\times$ 2 (Chunking Cue) ANOVA. As predicted, this revealed a significant main effect of Outcome ($F(1, 36) = 4.460, p = .042, \eta_p^2 = .110$). Averaged across all test trials, infants looked significantly longer at unexpected 2-Object outcomes (10.46 seconds) than at expected 3-Object outcomes (9.38 seconds) (Figure 2B). This preference for the 2-Object outcome was observed both for infants in the Contrasting Screens condition (9.56 seconds vs. 8.14 seconds; $t(19) = 2.329, p = .031, d = 0.591$) and infants in the

Identical Screens condition (11.69 seconds vs. 10.94 seconds), though in the Identical Screens condition the preference was not statistically significant; $t(19) = 0.938$, $p = .360$. There was also a significant main effect of Test Pair ($F(3, 108) = 3.488$, $p = .018$, $\eta_p^2 = .088$), driven by infants looking longer at the first (11.49 seconds) and second (10.50 seconds) test pairs than the third (8.73 seconds) and fourth test pairs (8.96 seconds), regardless of outcome. Finally, there was a main effect of Chunking Cue ($F(1, 36) = 4.359$, $p = .044$, $\eta_p^2 = .108$), driven by infants in the Identical Screens condition looking longer across all trials than infants in the Contrasting Screens condition, regardless of outcome. There were no other main effects or interactions. The lack of an interaction between Outcome and Chunking Cue ($F(1, 36) = 0.055$, $p = .817$) suggests that infants looked longer at the unexpected outcome whether or not the hiding screens were marked by contrasting borders.

As in Experiment 1, we asked whether infants' preference to look at two versus three objects differed between the baseline and test trials. A 2 (Trial Type) $\times$ 2 (Outcome) $\times$ 2 (Chunking Cue) ANOVA revealed a significant main effect of Chunking Cue ($F(1, 38) = 4.766$, $p = .035$, $\eta_p^2 = .111$), with infants in the Identical Screens condition looking longer across all trials (12.79 seconds) than infants in the Contrasting Screens condition (8.97 seconds). In addition, there was a marginally significant Trial Type $\times$ Outcome interaction ($F(1, 38) = 3.184$, $p = .082$, $\eta_p^2 = .077$), with infants looking longer at the 2-Object outcome only during the test trials, when it was the unexpected result of the hiding sequence (Figure 2B). There was no Trial Type $\times$ Outcome $\times$ Chunking Cue interaction ($F(1, 38) = 1.599$, $p = .214$), indicating that this marginal preference for the 2-Object outcome only during Test Trials was observed for infants in both Chunking Cue conditions.

Discussion

The results of Experiment 2 suggest that 7-month-old infants successfully represented three objects in memory. When provided with multiple redundant chunking cues including distinctive hiding locations and contrasting object features, 7-month-old infants who saw three objects sequentially hidden looked longer when two objects were revealed than when 3 objects were revealed, in the face of their failure to do so when these chunking cues were absent (Experiment 1). To our knowledge, this is the first evidence that infants younger than 14 months old may be able to expand the number of items held in memory through chunking.

Because little previous work has explored infants' ability to chunk objects, and because infants' success in the Identical Screens condition of Experiment 2 did not reach statistical significance, we wished to confirm infants' chunking abilities before exploring the particular cues underlying performance. In Experiment 3 we aimed to replicate the finding of successful chunking in a separate group of infants, using the Identical Screens condition from Experiment 2 (which offered fewer chunking cues and therefore provided a more stringent test of replication).

Experiment 3

Method

Participants—Twenty healthy full-term infants from the Baltimore, MD area participated (7 boys; 1 Asian, 4 Black or African American, 12 Caucasian, and 3 Other Race). Their mean age was 6 months, 26 days (range: 6 months, 14 days to 7 months, 14 days). Seven additional infants tested but excluded from the final analysis (1 for fussiness, 4 for experimenter error, 2 for parental interference).

Apparatus, Stimuli, and Procedure—The apparatus, stimuli, and procedure were identical to those from the Identical Screens condition of Experiment 2. As in Experiments 1 and 2, looking times greater than 2.5 standard deviations from the group mean for that trial were replaced by the same-trial average looking times of participants in the same condition. This resulted in the replacement of 6% of all trials.

Results

Baseline Trials—A 2 (Outcome) $\times$ 2 (Trial Order) ANOVA revealed no significant main effects or interactions, including a main effect of Outcome ($F(1, 16) = 1.738, p = .206$). Infants did not look significantly longer at either the 2-Object (7.35 seconds) or the 3-Object outcome (9.61 seconds).

Test Trials—We next analyzed looking times from the test trials in a 2 (Outcome) $\times$ 4 (Test Pair) $\times$ 2 (Trial Order) ANOVA. This yielded only a significant main effect of Outcome ($F(1, 18) = 5.671, p = .028, \eta_p^2 = .240$). Infants looked longer at unexpected 2-Object outcomes (9.38 seconds) than at expected 3-Object outcomes (7.66 seconds) (Figure 2C).

We also asked whether infants' preference to look at two versus three objects differed between the baseline and test trials. A 2 (Trial Type) $\times$ 2 (Outcome) ANOVA revealed a marginally significant interaction ($F(1, 19) = 3.906, p = .063, \eta_p^2 = .171$). Infants preferred to look longer at the 2-Object than the 3-Object outcome, but only during the test trials when it was the unexpected result of a hiding sequence.

Discussion

Experiment 3 replicated our previous finding that 7-month-old infants can use spatial and feature chunking cues to increase the amount of remembered information. As in Experiment 2, we found that infants who saw three objects hidden sequentially behind two screens looked longer at an unexpected outcome of two objects than an expected outcome of three objects. Given that this pattern of success was not observed in Experiment 1, in which no chunking cues were provided, these results suggest that infants in Experiment 3 were able to chunk representations of the hidden objects in memory.

Infants' success in Experiments 2 and 3 raises the question of which cues drove infants' chunking. Fourteen-month-old toddlers have been shown to chunk on the basis of spatiotemporal cues alone (Feigenson & Halberda, 2004, 2008) and featural cues alone (Feigenson & Halberda, 2008). Therefore in our next experiment we asked whether spatial separation between hiding locations or contrasting object features would independently support 7-month-old infants' chunking, as they appear to do for older children (Feigenson & Halberda, 2004, 2008).

Experiment 4

In Experiment 4 we examined 7-month-old infants' ability to use spatial or featural cues alone as a basis for chunking. In the Spatial Cues condition infants saw three identical objects hidden behind two identical spatially separated screens. In the Feature Cues condition infants saw three objects, two of which had identical features, hidden behind a single screen.

Method

Participants—Forty healthy, full-term infants from the Baltimore, MD area participated (17 boys; 2 Black or African American, 33 Caucasian, and 5 Other Race). Their mean age

was 6 months, 29 days (range: 6 months, 16 days to 7 months, 13 days). Eleven additional infants were tested but excluded from the final analysis (3 for fussiness, 4 for experimenter error, and 4 for parental interference).

Apparatus and Stimuli—Infants in the Spatial Cues condition saw three identical green and white balls (the same objects used in Experiment 1) hidden behind two identical black screens (the same screens used in the Identical Screens condition of Experiment 2, and in Experiment 3). Infants in the Feature Cues condition saw one leopard-print block and two green and white balls (the same objects used in Experiments 2 and 3) hidden behind a single large screen (the same screen used in Experiment 1).

Procedure—Infants saw two baseline trials and eight test trials that were similar to those in Experiments 1-3. In the baseline trials of the Spatial Cues condition, the two screens were lifted simultaneously to reveal one ball behind each screen (2-Object trial), or two balls behind one screen and one ball behind the other (3-Object trial). In the baseline trials of the Feature Cues condition, the screen was lifted to reveal one ball and one leopard-print block (2-Object trial) or two balls and one block (3-Object trial).

During the test trials, infants always saw three objects hidden sequentially behind the screen(s) (Figure 1). In the Spatial Cues condition, infants saw three identical balls hidden behind two screens. The side on which two objects were hidden, and whether one or two balls hidden were behind the first screen were counterbalanced across trials for each infant. In the Feature Cues condition, infants saw two balls and one block hidden behind a single screen, with the two balls always hidden in succession (ball-ball-block or block-ball-ball). Whether infants first saw the balls or the block hidden was counterbalanced across trials for each infant.

As in the previous experiments, looking times greater than 2.5 standard deviations from the group mean for that trial were replaced by the same-trial average looking times of participants in the same condition. This resulted in the replacement of 5.0% of all trials.

Results

Baseline Trials—A 2 (Outcome) $\times$ 2 (Trial Order) $\times$ 2 (Chunking Cue: Spatial vs. Feature) ANOVA revealed a significant Outcome $\times$ Trial Order interaction ($F(1, 36) = 4.133$, $p = .049$, $\eta_p^2 = .103$) driven by infants looking longer at the first trial they saw, regardless of whether it contained two or three objects (Trial 1: 8.86 seconds, Trial 2: 5.84 seconds). There were no other main effects or interactions. Infants did not look longer at the 2-Object (6.81 seconds) or the 3-Object outcome (7.97 seconds), $F(1, 36) = 0.624$, $p = .435$.

Test Trials—We next analyzed looking times from the test trials in a 2 (Outcome) $\times$ 4 (Test Pair) $\times$ 2 (Trial Order) $\times$ 2 (Chunking Cue) ANOVA. This yielded a significant main effect of Pair ($F(3, 108) = 3.886$, $p = .011$, $\eta_p^2 = .093$), driven by infants looking longer at the first three test pairs (8.08 seconds, 8.43 seconds, and 8.83 seconds, respectively) than the last pair (6.29 seconds). There was also a main effect of Chunking Cue ($F(1, 36) = 8.713$, $p = .005$, $\eta_p^2 = .187$), driven by infants in the Spatial Cue condition looking longer across all trials (9.35 seconds) (Figure 2D) than infants in the Feature Cue condition (6.47 seconds) (Figure 2E). However, there was no effect of Outcome ($F(1, 36) = 0.250$, $p = .620$). Averaged across all test pairs, infants did not look longer at unexpected 2-Object outcomes (8.02 seconds) than at expected 3-Object outcomes (7.79 seconds). This lack of a significant preference for either outcome was observed both for infants in the Spatial condition (6.69 seconds vs. 6.25 seconds) and Feature condition (9.36 seconds vs. 9.33 seconds) (Figures 2 D and E).

We also asked whether infants' preference to look at two versus three objects differed between the baseline and test trials. A 2 (Trial Type) $\times$ 2 (Outcome) $\times$ 2 (Chunking Cue) ANOVA revealed no significant main effects or interactions. Critically, there was no Trial Type $\times$ Outcome interaction ($F(1,38) = 0.703, p = .407$), suggesting that infants' preference for two versus three objects did not change when these were the unexpected and expected results of an object hiding sequence.

Discussion

Fourteen-month-old toddlers chunk arrays of identical objects when provided with spatial grouping cues (Feigenson & Halberda, 2004), and chunk equi-spaced arrays of objects when provided with featural or conceptual grouping cues (Feigenson & Halberda, 2008). The results of Experiment 4 suggest that for younger, 7-month-old infants, neither spatial nor featural cues are independently sufficient to support object chunking, at least in the context of the present task. Although our finding does not preclude the possibility that a single cue other than those we tested here could suffice to support chunking at this age, it does raise the possibility that these younger infants may require multiple cues in order to chunk successfully, and that chunking may undergo developmental change between the ages of 7 and 14 months (see General Discussion).

However, an alternative to the claim that 7-month-old infants can chunk if provided with multiple cues is that infants succeeded in Experiments 2 and 3 simply because the stimulus arrays in those experiments were more perceptually interesting, thereby attracting infants' attention more strongly, and making it more likely that the objects would be encoded and remembered. The stimulus arrays in Experiments 2 and 3 contained two distinctive hiding locations (demarcated by two screens), and also contained objects with high featural contrast (a smiling, leopard-print block versus spotted balls). In comparison, the stimulus arrays in Experiments 1 and 4 either contained just a single screen/hiding location (Experiment 1 and the Featural Cues condition of Experiment 4) or contained all identical objects (Experiment 1 and the Spatial Cues condition of Experiment 4). One might therefore worry that infants paid more attention to the arguably more complex arrays in Experiments 2 and 3—the experiments in which they succeeded—than to the simpler arrays in Experiments 1 and 4—the experiments in which they failed.

To test this possibility, in Experiment 5 we presented infants with arrays that contained exactly the same amount of spatial and featural information as in Experiments 2 and 3. Infants saw objects hidden behind two spatially separated screens, and saw the same featurally contrasting objects as in Experiments 2 and 3. However, in Experiment 5 we reorganized the arrays such that spatial and featural cues no longer specified the same chunks—we hid a ball and a block behind one screen, and another ball behind the other screen. If infants require multiple redundant cues to support chunking—that is, if spatial and featural cues must specify the same chunks—then infants in this experiment should fail to remember the three hidden objects.

Experiment 5

Method

Participants—Twenty healthy, full-term infants from the Baltimore, MD area participated (9 boys; 1 Asian, 2 Black or African American, 13 Caucasian, and 4 Other Race). Their mean age was 6 months, 29 days (range: 6 months, 15 days to 7 months, 13 days). Eleven additional infants were tested but excluded from the final analysis (7 for fussiness, 3 for experimenter error, and 1 for parental interference).

Apparatus and Stimuli—The apparatus and stimuli were the same as those in the Identical Screens condition of Experiment 2 and Experiment 3 (two identical black hiding screens and two green and white balls plus one leopard-print block).

Procedure—In the two baseline trials infants saw a 2-Object array in which one ball was revealed from behind one screen and one block was revealed from behind the other, and a 3-Object array in which one ball was revealed from behind one screen and one ball and one block were revealed from behind the other (Figure 1).

In the eight test trials infants always saw a sequential presentation in which one ball was hidden behind one screen and one ball and one block were hidden behind the other. Infants always saw the block as either the first or last object hidden so that the two balls were always presented in temporal succession. Expected outcomes always contained two balls and one block, and unexpected outcomes always contained just one ball and one block (Figure 1).

As in the previous experiments, looking times greater than 2.5 standard deviations from the group mean for that trial were replaced by the same-trial average looking times of participants in the same condition. This resulted in the replacement of 5.0% of all trials.

Results

Baseline Trials—A 2 (Outcome) $\times$ 2 (Trial Order) ANOVA yielded no significant main effects or interactions. Infants did not look longer at either the 2-Object (9.74 seconds) or the 3-Object outcome (8.59 seconds) ($F(1, 16) = 0.732, p = .403$).

Test Trials—We next analyzed looking times from the test trials in a 2 (Outcome) $\times$ 4 (Test Pair) $\times$ 2 (Trial Order) ANOVA. This revealed only a significant main effect of Test Pair ($F(3, 54) = 3.340, p = .026, \eta_p^2 = .157$). Infants looked longer on the second (9.97 seconds) and third (9.96 seconds) test pairs than on the first (7.52 seconds) and fourth (6.90 seconds) test pairs. Critically, there was no main effect of Outcome ($F(1, 18) = 0.227, p = .640$). Averaged across all test pairs, infants did not look significantly longer at unexpected 2-Object outcomes (9.74 seconds) than at expected 3-Object outcomes (8.59 seconds) (Figure 2F).

We also asked whether infants' preference to look at two versus three objects differed between the baseline and test trials. A 2 (Trial Type) $\times$ 2 (Outcome) ANOVA revealed no significant main effects or interactions. Importantly, there was no Trial Type $\times$ Outcome interaction ($F(1, 19) = 0.243, p = .628$), indicating that infants' preference for two versus three objects did not change when these were the unexpected and expected results of an object hiding sequence.

Discussion

The results from Experiment 5 strengthen our claim that infants' success at remembering three objects in Experiments 2 and 3 reflects chunking. Simply presenting infants with multiple hiding locations and contrasting object features did not cause infants to remember the hidden objects. Instead, it appears that the spatial and featural cues had to redundantly specify the same chunks in order for infants to benefit.

General Discussion

Previous studies demonstrated that 10- to 20-month-old infants can only store three to four items concurrently in working memory (Barner et al., 2007; Feigenson & Carey, 2003;

2005; Feigenson, Carey & Hauser, 2002; Rose et al., 2001; Ross-Sheehy et al., 2003), paralleling similar limits in adults (Alvarez & Cavanagh, 2004; Awh et al., 2007; Cowan, 2001; Feigenson, 2008; Halberda et al., 2006; Luck & Vogel, 1997; Scolari, Vogel & Awh, 2008; Song & Jiang, 2006; Todd & Marois, 2004; Vogel & Machizawa, Y. Xu & Chun, 2006; Zhang & Luck, 2008). Previous studies also have shown that toddlers, older children, and adults can increase the total amount of stored information by binding representations of individual items into chunks (Bower, 1972; Chase & Ericsson, 1981; Chase & Simon, 1973; Cowan et al., 2004; Ericsson et al., 1980; Feigenson & Halberda, 2004, 2008; Mathy & Feldman, 2012; Rosner, 1971; Simon, 1974). In the present series of experiments we explored the developmental origins of this ability. We found that younger, 7-month-old infants, who are pre-linguistic and whose working memory has not yet reached mature capacity, also can remember more items by chunking. This suggests that chunking may be a foundational computation—one used from infancy onward.

To recap our findings, Experiment 1 found that 7-month-old infants did not remember three identical objects sequentially hidden in a single location, consistent with previous suggestions that working memory capacity is not yet mature at this age (Káldy & Leslie, 2005; Rose et al., 2001; Ross-Sheehy et al., 2003). Experiments 2 and 3 demonstrated that when provided with multiple redundant chunking cues, including spatially separated hiding locations and featurally contrasting objects, infants successfully looked longer when three objects were hidden and just two were revealed. However, Experiment 4 revealed that neither spatial separation of the hiding locations nor contrasting object features alone were sufficient chunking cues. Infants failed to remember all three objects when given either spatial or featural chunking cues individually. Finally, Experiment 5 showed that infants' performance in Experiments 2 and 3 could not be attributed to the fact that the stimuli were more perceptually complex, and that spatial and featural cues had to specify consistent chunks in order for 7-month-olds to benefit.

Many factors influence memory, making it more or less likely that observers will remember the items in a scene. For example, previous research suggests that infants are more successful at remembering objects over shorter durations (Diamond, 1990), when given sustained or repeated presentations that allow for the potential recruitment of long-term memory resources (Canfield & Smith, 1996; Sharon & Wynn, 1998), when the number of required memory updates decreased (Uller et al., 1999), when items are presented in direct succession (Feigenson & Yamaguchi, 2009), when objects are presented later rather than earlier in a hiding sequence (Káldy & Leslie, 2005), and when given with the opportunity to compare multiple objects during encoding (Oakes, Kovack-Lesh, & Horst, 2009). Here we show that in addition to the above factors, the extent to which an object array supports chunking also affects infants' memory, with other aspects of the presentation held constant.

Our finding that 7-month-old infants only chunked when given redundant cues contrasts with previous findings that 14-month-old toddlers needed just a single cue in order to chunk. Fourteen-month-olds demonstrated successful chunking when shown spatially interleaved arrays of four familiar objects, such as cat, car, cat, car (Feigenson & Halberda, 2008). Because the cats and cars were not spatially grouped (i.e., identical objects were non-adjacent), spatial adjacency and featural information specified conflicting parses of the array. The finding that 14-month-olds successfully chunked when spatial and featural cues did not overlap (Feigenson & Halberda, 2008), but that 7-month-olds did not (Experiment 5 in the present series) suggests that chunking may undergo developmental change between 7 and 14 months of age, with infants becoming increasingly adept at using more subtle cues to support chunking. However, we note that the methods used to test chunking differed across the two age groups (manual search for 14-month-olds versus violation-of-expectation

looking time for 7-month-olds). Therefore, future investigations of developmental change in chunking abilities should test infants of different ages using a single paradigm.

An additional question for future work concerns the specificity of infants' object representations. Although our findings show that infants used spatial and featural information in conjunction to remember hidden objects, it remains unknown whether infants actually remembered both types of information. Indeed, recent work suggests that infants who remember the existence of an object may remember little about the object's features (Kibbe & Leslie, 2011). This raises the possibility that infants in our experiments may have used spatial and featural cues to encode or maintain object representations, but then failed to retain this featural information in memory or to use it at retrieval (see Wang & Baillargeon, 2006; Wang & Mitroff, 2009). A related question is whether, even if infants do remember spatial and featural information, they correctly bind this information in memory. For example, infants might correctly remember that there were two objects behind the left screen and one behind the right, but misremember which objects were where (Káldy & Leslie, 2003; Oakes, Ross-Sheehy, & Luck, 2006). More clearly characterizing the contents of infants' representations of object chunks remains a prospect for future investigation.

Finally, the role of recent experience in guiding infants' chunking remains to be explored. In particular, it is an open question whether infants' visual exposure to the baseline arrays in Experiments 2 and 3 played a role in their later chunking of the test arrays. It is possible that seeing baseline arrays, with objects grouped by spatial location and by feature, helped infants later use these cues to chunk. The performance of infants given only test trials would bear on this question.

To summarize, our results highlight the possibility of both developmental continuity and change in early memory. Here we found evidence that 7-month-old infants, whose working memory capacity is still developing, can chunk items together to increase total information storage. However, this ability appears to be somewhat fragile in these younger infants; 7-month-olds required multiple redundant cues in order to chunk, whereas older toddlers were able to use spatial and featural cues independently. Future work should continue to characterize both these continuous and discontinuous properties of a fundamental aspect of memory processing.

Acknowledgments

This work was supported by a National Science Foundation Graduate Student Fellowship to M.M. (as M. Yamaguchi), and by a McDonnell Scholar Award and a National Institute of Child Health and Human Development Grant HD054416 to L.F. We thank Rachel Austin, Rebecca Etkin, Catherine Juravich Roe, Nick Minar, Lenae Stansky, Andrea Stevenson and Allison Wessel for help with data collection, and the families in the greater Baltimore area for their participation.

References

- Alvarez GA, Cavanagh P. The capacity of visual short-term memory is set both by visual information load and by number of objects. *Psychological Science*. 2004; 15(2):106–111. [PubMed: 14738517]
- Awh E, Barton B, Vogel EK. Visual working memory represents a fixed number of items regardless of complexity. *Psychological Science*. 2007; 18(7):622–628. [PubMed: 17614871]
- Barner D, Thalwitz D, Wood J, Yang SJ, Carey S. On the relation between the acquisition of singular-plural morpho-syntax and the conceptual distinction between one and more than one. *Developmental Science*. 2007; 10(3):365–373. [PubMed: 17444976]
- Bauer, PJ. Remembering the times of our lives: Memory in infancy and beyond. Laurence Erlbaum Associates; Mahwah, NJ: 2007.

- Bower GH. Perceptual groups as coding units in immediate memory. *Psychonomic Science*. 1972; 27(4):217–219.
- Brady TF, Konkle T, Alvarez GA. Compression in visual working memory: Using statistical regularities to form more efficient memory representations. *Journal of Experimental Psychology: General*. 2009; 138(4):487–502. [PubMed: 19883132]
- Broadbent, DE. The magic number seven after fifteen years.. In: Kennedy, A.; Wilkes, A., editors. *Studies in Long-Term Memory*. Wiley; New York: 1975. p. 3-18.
- Canfield RL, Smith EG. Number-based expectations and sequential enumeration by 5-month-old infants. *Developmental Psychology*. 1996; 32:269–279.
- Chase WG, Ericsson KA. Bower GH. Skill and working memory. *The psychology of learning and motivation*. 1981; 16:1–58.
- Chase WG, Simon HA. Perception in chess. *Cognitive Psychology*. 1973; 4:55–81.
- Cowan, N. The development of memory in infancy and childhood. Psychology Press; Hove, U.K.: 1997.
- Cowan N. The magical number 4 in short-term memory: A reconsideration of mental storage capacity. *Behavioral and Brain Sciences*. 2001; 24(1):87–114. [PubMed: 11515286]
- Cowan N, Chen Z, Rouders JN. Constant capacity in an immediate serial-recall task. *Psychological Science*. 2004; 15:634–640. [PubMed: 15327636]
- Diamond A. The development and neural bases of memory functions as indexed by the AB and delayed response tasks in human infants and infant monkeys. *Annals of the New York Academy of Sciences*. 1990; 608:267–317. [PubMed: 2127511]
- Ericsson KA, Chase WG, Faloon S. Acquisition of a memory skill. *Science*. 1980; 208:1181–1182. [PubMed: 7375930]
- Ericsson KA, Delaney PF, Weaver G, Mahadevan R. Uncovering the structure of a memorist's superior "basic" memory capacity. *Cognitive Psychology*. 2004; 49:181–237.
- Feigenson L. Parallel non-verbal enumeration is constrained by a set-based limit. *Cognition*. 2008; 107:1–18. [PubMed: 17761158]
- Feigenson L, Carey S. Tracking individuals via object-files: Evidence from infants' manual search. *Developmental Science*. 2003; 6:568–584.
- Feigenson L, Carey S. On the limits of infants' quantification of small object arrays. *Cognition*. 2005; 97:295–313. [PubMed: 16260263]
- Feigenson L, Carey S, Hauser M. The representations underlying infants' choice of more: Object-files versus analog magnitudes. *Psychological Science*. 2002a; 13:150–156. [PubMed: 11933999]
- Feigenson L, Carey S, Spelke ES. Infants' discrimination of number versus continuous extent. *Cognitive Psychology*. 2002b; 44:33–66. [PubMed: 11814309]
- Feigenson L, Halberda J. Infants chunk object arrays into sets of individuals. *Cognition*. 2004; 91:173–190. [PubMed: 14738772]
- Feigenson L, Halberda J. Conceptual knowledge increases infants' memory capacity. *Proceedings of the National Academy of Sciences*. 2008; 105(29):9926–9930.
- Feigenson L, Yamaguchi M. Limits on infants' ability to dynamically update object representations. *Infancy*. 2009; 14(2):244–262.
- Fenson L, Dale PS, Reznick JS, Bates E, Thal DJ, Pethick SJ. Variability in early communicative development. *Monographs of the Society for Research in Child Development*. 1994; 59(5) Serial No. 242.
- Fiser J, Aslin RN. Encoding multielement scenes: Statistical learning of visual feature hierarchies. *Journal of Experimental Psychology: General*. 2005; 134(4):521–537. [PubMed: 16316289]
- Halberda J, Sires SF, Feigenson L. Multiple spatially overlapping sets can be enumerated in parallel. *Psychological Science*. 2006; 17(7):572–576. [PubMed: 16866741]
- Káldy Z, Leslie AM. Identification of objects in 9-month-old infants: Integrating 'what' and 'where' information. *Developmental Science*. 2003; 6:360–373.
- Káldy Z, Leslie AM. A memory span of one? Object identification in 6.5-month-old infants. *Cognition*. 2005; 97:153–177. [PubMed: 16226561]

- Kibbe M, Leslie AM. What do infants remember when they forget? Location and identity in 6-month old infants' memory for objects. *Psychological Science*. 2011; 22(11)
- Koechlin E, Dehaene S, Mehler J. Numerical transformations in five-month-old human infants. *Mathematical Cognition*. 1997; 3:89–104.
- Luck SJ, Vogel EK. The capacity of visual working memory for features and conjunctions. *Nature*. 1997; 390:279–281. [PubMed: 9384378]
- Mathy F, Feldman J. What's magic about magic numbers? Chunking and data compression in short-term memory. *Cognition*. 2012; 122:346–362. [PubMed: 22176752]
- Noles NS, Scholl BJ, Mitroff SR. The persistence of object file representations. *Perception and Psychophysics*. 2008; 67(2):324–334. [PubMed: 15973783]
- Oakes, LM.; Bauer, PJ. Short- and long-term memory in infancy and early childhood: Taking the first steps toward remembering. Oxford University Press; New York: 2007.
- Oakes LM, Kovack-Lesh KA, Horst JS. Two are better than one: Comparison influences infants' visual recognition memory. *Journal of Experimental Child Psychology*. 2009; 104:124–131. [PubMed: 18951555]
- Oakes LM, Ross-Sheehy S, Luck SJ. Rapid development of feature binding in visual short-term memory. *Psychological Science*. 2006; 17(9):781–787. [PubMed: 16984295]
- Rose SA, Feldman JF, Jankowski JJ. Visual short-term memory in the first year of life: Capacity and recency effects. *Developmental Psychology*. 2001; 4(37):539–549. [PubMed: 11444489]
- Rosner SR. The effects of rehearsal and chunking instructions on children's multitrial free recall. *Journal of Experimental Child Psychology*. 1971; 11(1):93–105.
- Ross-Sheehy S, Oakes LM, Luck SJ. The development of visual short-term memory capacity in infants. *Child Development*. 2003; 74(6):1807–1922. [PubMed: 14669897]
- Rovee-Collier CK. The development of infant memory. *Current Directions in Psychological Science*. 1999; 8:80–85.
- Scolari M, Vogel EK, Awh E. Perceptual expertise enhances the resolution but not the number of representations in working memory. *Psychonomic Bulletin and Review*. 2008; 15(1):215–222. [PubMed: 18605506]
- Sharon T, Wynn K. Individuation of actions from continuous motion. *Psychological Science*. 1998; 9:357–362.
- Simon HA. How big is a chunk? *Science*. 1974; 183:482–488. [PubMed: 17773029]
- Simon T, Hespos SJ, Rochat P. Do infants understand simple arithmetic? A replication of Wynn (1992). *Cognitive Development*. 1995; 10:253–269.
- Song J-H, Jiang Y. Visual working memory for simple and complex features: an fMRI study. *Neuroimage*. 2006; 30:963–972. [PubMed: 16300970]
- Squire LR. Memory and brain systems: 1969–2009. *The Journal of Neuroscience*. 2009; 29(41):12711–12716. [PubMed: 19828780]
- Terrace HS. Chunking by a pigeon in a serial learning task. *Nature*. 1987; 325:149–151. [PubMed: 3808071]
- Terrace HS. Chunking during serial learning by a pigeon: I. Basic evidence. *Journal of Experimental Psychology: Animal Behavior Processes*. 1991; 17(1):81–93. [PubMed: 2002309]
- Terrace HS, Chen S. Chunking during serial learning by a pigeon: III. What are the necessary conditions for establishing a chunk? *Journal of Experimental Psychology: Animal Behavior Processes*. 1991; 17(1):107–118. [PubMed: 2002304]
- Todd JJ, Marois R. Capacity limit of visual short-term memory in human posterior parietal cortex. *Nature*. 2004; 428(6984):751–754. [PubMed: 15085133]
- Tulving E. Episodic memory: From mind to brain. *Annual Review of Psychology*. 2002; 53:1–25.
- Uller C, Carey S, Huntley-Fenner G, Klatt L. What representations might underlie infant numerical knowledge. *Cognitive Development*. 1999; 14:1–36.
- Wang S, Baillargeon R. Infants' physical knowledge affects their change detection. *Developmental Science*. 2006; 9:173–181. [PubMed: 16472318]
- Wang S, Mitroff SR. Preserved visual representations despite change blindness in 11-month-old infants. *Developmental Science*. 2009; 12:681–687. [PubMed: 19702760]

- Wynn K. Addition and subtraction by human infants. *Nature*. 1992; 358:749–750. [PubMed: 1508269]
- Xu Y. Limitations of object-based feature encoding in visual short-term memory. *Journal of Experimental Psychology: Human Perception and Performance*. 2002; 28(2):458–468. [PubMed: 11999866]
- Xu Y, Chun MM. Dissociable neural mechanisms supporting visual short-term memory for objects. *Nature*. 2006; 440:91–95. [PubMed: 16382240]
- Zhang W, Luck SJ. Discrete fixed-resolution representations in visual working memory. *Nature*. 2008; 453:233–235. [PubMed: 18385672]
- Zhang W, Luck SJ. Sudden death and gradual decay in visual working memory. *Psychological Science*. 2009; 20(4):423–428. [PubMed: 19320861]

Highlights

1. Adults and toddlers can chunk to increase the total amount of remembered information
2. We asked whether pre-verbal 7-month old infants also can chunk
3. With no chunking cues, 7-month old infants failed to remember 3 hidden objects
4. With redundant spatial and featural cues, infants succeeded
5. Chunking appears to be a foundational aspect of memory throughout the lifespan

	Unexpected outcome	Expected outcome
Experiment 1
Experiment 2 Contrasting screens
Experiment 2 Identical screens
Experiment 3 Identical screens
Experiment 4 Spatial cues only
Experiment 4 Feature cues only
Experiment 6 Conflicting cues

Figure 1.

Depiction of the Unexpected and Expected test outcomes for each experiment. The black borders around the objects depict the screens, which occluded the objects as they were being sequentially hidden, and then were lifted to reveal the objects. These same arrays served as the baseline arrays within each experiment.

Figure 2.

Infants' average looking times to two versus three objects in the Baseline and Test arrays. Bars depict standard error. (* $p < .05$, + $p < .01$)