Published in final edited form as:

J Psychoactive Drugs. 2011 ; 43(1): 20–26. doi:10.1080/02791072.2011.566493.

Predictors of Membership in Alcoholics Anonymous in a Sample of Successfully Remitted Alcoholics

Amy R. Krentzman, Ph.D.^{*}, Elizabeth A. R. Robinson, Ph.D.^{*}, Brian E. Perron, Ph.D.^{**}, and James A. Cranford, Ph.D.^{*}

^{*}University of Michigan Addiction Research Center, 4250 Plymouth Road, Ann Arbor, MI 48109-5740, U.S.A.

^{**}University of Michigan School of Social Work, 1080 S. University Avenue, Ann Arbor, MI 48109, U.S.A.

Abstract

This study identifies factors associated with Alcoholics Anonymous (AA) membership in a sample of 81 persons who have achieved at least one year of total abstinence from drugs and alcohol. Forty-four were AA members, 37 were not. Logistic regression was used to test the cross-sectional associations of baseline demographic, substance-related, spiritual and religious, and personality variables with AA membership. Significant variables from the bivariate analyses were included in a multivariate model controlling for previous AA involvement. Having more positive views of God and more negative consequences of drinking were significantly associated with AA membership. This information can be used by clinicians to identify clients for whom AA might be a good fit, and can help others overcome obstacles to AA or explore alternative forms of abstinence support.

Keywords

Alcoholism; Alcoholics Anonymous; predictors of Alcoholics Anonymous membership

Introduction

A number of studies have shown membership in Alcoholics Anonymous (AA) to be among the most important predictors of positive outcomes and sustained recovery from alcoholism (Bond, Kaskutas, & Weisner, 2003; Morgenstern, Labouvie, McCrady, Kahler, & Frey, 1997; Vaillant, 2003). Thus, identifying characteristics of AA participants can inform the development of effective interventions. To our knowledge, to date, no previous study has examined predictors of membership in AA among remitted alcoholics. The current study

Corresponding author: Amy R. Krentzman, Ph.D., University of Michigan Addiction Research Center, 4250 Plymouth Rd., Ann Arbor, MI 48109-5740, U.S.A., Telephone: 718-570-2706, Fax: 734.998.7992, amykrent@med.umich.edu.

Contributors

Amy R. Krentzman designed the current study, conducted the literature review and statistical analyses, and had primary responsibility for writing the first drafts of the manuscript. Author Elizabeth A.R. Robinson designed the parent study, wrote the methods section, and provided editorial leadership. Author Brian E. Perron shaped the direction of the paper and provided editorial leadership. Author James Cranford provided statistical direction and consultation and made editorial contributions. All authors contributed to and have approved the final manuscript.

Conflict of Interest

All authors declare that they have no conflicts of interest.

was designed to address this gap in our knowledge by identifying factors associated with AA membership in a sample of remitted alcoholics.

Previous research has followed the 12-step behavior of substance abusers following treatment. Differences in demographics, substance use severity, spiritual and religious activity, and social involvement have been observed among those who do and do not participate in AA. One study reported no demographic differences between those who attend AA and those who do not (Humphreys, Mavis, & Stofflemayr, 1994). Other studies found that certain demographic groups, such as women (Humphreys, Mavis, and Stofflemayr, 1991), those with higher levels of education (Terra et al., 2007), and African Americans (Humphreys et al., 1991; Kelly & Moos, 2003) are more likely to attend or less likely to drop out of AA. Higher substance use severity and more family, social and psychological problems have been shown to be associated with later AA participation (Emrick, Tonigan, Motgomery, & Little, 1993; Humphreys et al., 1991). Studies have also found that individuals who participate in more spiritual and religious activity are more likely to later affiliate with AA (Emrick et al., 1993; Kelly & Moos, 2003). Certain personality characteristics of those who affiliate with AA have been identified. Janowsky, Boone, Morter, and Howe (1999) found those who had attended meetings had higher levels of extraversion and lower levels of “shyness with strangers.” Kelly and Moos (2003) similarly found that individuals who were less likely to dropout of AA reported more interpersonal social involvement with other people in their daily lives.

This study examines predictors of AA membership in a longitudinal sample of alcoholics who 3 years after intake were abstinent for at least one year and either identified as members of AA or did not. This study has two aims. The first is to identify correlates of AA participation among successful remitters. The specific variables we investigate are those found to be predictive of AA participation in previous literature, including education, gender, race, spirituality, religiousness, extroversion, and addiction severity. The second aim is to examine the relative strength of the correlates within a multivariate framework.

Method

Data from this study are derived from the Life Transition Study (LTS), a longitudinal survey that followed 364 alcohol-dependent individuals for 3 years (Robinson, Krentzman, Pierce, Webb, & Brower, 2010). The LTS was designed to explore spiritual and religious change and its relationship to drinking outcomes and recovery efforts in a diverse sample of alcohol-dependent individuals. Diagnoses of alcohol dependence were based on the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID; First et al., 1997). To ensure demographic, socioeconomic, and substance-use diversity, study respondents were recruited from four sources in a Midwestern city: a university outpatient treatment program (n=157), a VA outpatient treatment program (n=80), an outpatient moderation program designed to help individuals reduce drinking (n=34), and untreated individuals recruited from the local community (n=93). Individuals were interviewed in person every 6 months and by telephone during the intervening 3 month periods.

The current study focuses on 81 of the 87 original respondents who had achieved a year of abstinence at their final interview and were not involved in other forms of mutual-help groups beyond AA. Six individuals were abstinent and attending other self-help organizations such as Women in Recovery and S.M.A.R.T. Recovery. Abstinence was based on drinking and drug use data assessed by the TimeLine FollowBack Interview (TLFB; Sobell & Sobell, 1992, 1996). The sample was then divided into AA membership status using the following question from the Alcoholics Anonymous Involvement scale (AAI; Tonigan, Connors, & Miller, 1996): “Do you consider yourself to be a member of

Alcoholics Anonymous?” At the final interview, 44 (54.3%) answered “yes.” This group formed the AA group. The remaining 37 (45.7%) individuals were neither members of AA nor involved in any other mutual-help abstinence-support groups in the 90 days prior to the 3-year follow-up. This sample of 81 came from all four of the original recruitment sites: the university outpatient treatment program (n=47, 58.0%), the VA outpatient treatment program (n=22, 27.2%), the moderation program (n=3, 3.7%), and the community sample (n=9, 11.1%). All of the individuals recruited from the community received professional treatment either before or during the study.

Measures

Demographic Variables—Demographic variables included education (in years), race (white or other), and gender (male or female).

Spirituality and Religiousness Variables—Measures of spirituality and religiousness were chosen based on their hypothesized ability to measure openness to the spiritual aspects of the AA program and included measures of belief in God, perception of God as loving, and the experience of being raised in a religious tradition. *Belief in God* was measured with the first item of the Religious Background and Behaviors scale (RBB; Connors et al., 1996). This single item asks, “Which of the following best describes you at the present time?” The 5-point response options were 1 = I do not believe in God (atheist); 2 = I believe we really can’t know about God (agnostic); 3 = I don’t know what to believe about God (unsure); 4 = I believe in God, but I’m not religious (spiritual); and 5 = I believe in God and practice religion (religious). In the current study, belief in God was defined as endorsement of items 4 or 5, non-belief by endorsing items 1–3.

Perceptions of God were assessed with the Loving God scale (Benson & Spilka, 1973). This 5-item semantic differential measure uses opposite words as anchors at either end of a seven-point scale. Examples of item pairs are *rejecting-accepting*, *unforgiving-forgiving*, and *approving-disapproving*. The Loving God subscale measures the degree the respondent finds God to be loving (or accepting, saving, forgiving, approving) ($\alpha = .79$).

Religious tradition was measured using the following single item; “Would you say you were brought up in a religious tradition?” (yes / no).

2.1.3 Personality Variable: Extraversion—The NEO Five Factor Inventory (Costa & McCrae, 1985, 1992) measures the five major domains of personality: neuroticism, extraversion, openness, agreeableness, and conscientiousness. This study used 12 items from the NEO that measures extraversion. These items used a 5-point Likert-type response format (1 = *strongly disagree* to 5 = *strongly agree*; $\alpha = .82$).

Substance-Use Variables—*Drinking severity* was measured by the *Short Index of Problems* scale (SIP; Miller et al., 1995). The SIP is a 15-item measure which asks about several negative consequences of drinking. Sample items are “I have felt guilty or ashamed because of my drinking” and “My family has been hurt by my drinking.” The scale uses a 4-point Likert-type response format (0 = *never*, 3 = *daily or almost daily*; $\alpha = .91$).

Alcoholics Anonymous Variables—*Alcoholics Anonymous involvement* refers to involvement in AA beyond meeting attendance, including celebrating an AA sobriety birthday, having a sponsor, and being a sponsor. The construct was measured using a modified version of the Alcoholics Anonymous Involvement index (AAI; Tonigan et al., 1996). Baseline values of this variable were included to control for previous AA involvement. Scores were rendered by summing the positive responses of seven of the

scale's items which used a dichotomous response set, resulting in a range of possible scores from 0 to 7 ($\alpha = .83$).

Alcoholics Anonymous attendance was measured by asking participants to calculate the total number of AA meetings they had attended throughout their lives.

2.2 Analysis Plan

Associations between study variables and AA attendance were compared using unadjusted logistic regression analysis. Associations that were significant at the bivariate level were included in a multivariate logistic regression model. Odds ratios (OR) were computed for all unadjusted and adjusted models and were considered statistically significant if the 95% confidence interval (CI) did not bound the value 1.0.

Results

Summary of Demographic and Clinical Variables

Table 1 summarizes the demographic and clinical variables of respondents who had been abstinent for at least one year at their final interview ($N = 81$). The majority of respondents were either married/living with a significant other (38.3%) or separated, widowed, or divorced (40.7%), White (85.2%), and male (69.1%). The mean age was 47.2 ($SD = 11.7$), and the years of education was 14.4 ($SD = 2.16$). Almost half the sample earned less than \$30,000 per year (46.9%) and were employed (49.4%).

AA Attendance and Involvement Prior to Baseline

Table 1 indicates a high level of previous AA attendance in the sample. This is seen in the average number of total lifetime meetings attended ($M=233$), although considerable heterogeneity was observed based on the large standard deviation ($SD=500.7$). The mean score on the AA involvement index was 2.21 ($SD = 2.29$).

Bivariate Analyses

Table 2 summarizes bivariate analyses between AA membership and other study variables. Women with one year of abstinence were three times more likely to be in the AA group than men with one year of abstinence ($OR = 3.0$, 95% $CI = 1.07-8.22$). While believing in God was not predictive of AA membership in this sample, believing in a loving God was ($OR = 1.1$, 95% $CI = 1.03-1.28$). Individuals raised in a religious tradition were almost three times more likely to become AA members than those who were not ($OR=2.8$).

Multivariate Analyses

Table 3 displays significant correlates of AA membership using multivariate logistic regression. All significant variables from bivariate correlates were entered into the model simultaneously. The overall model exhibited a good fit with the data ($\chi^2=24.01$, $p < .001$, pseudo R -square=.36). Three variables were statistically significant predictors of AA membership at the final interview: The Loving God subscale ($OR = 1.2$, 95% $CI = 1.01-1.32$), SIP measure of drinking consequences ($OR = 1.1$, 95% $CI = 1.00-1.12$) and previous AA involvement ($OR = 1.3$, 95% $CI = 1.00-1.801$). Gender and being raised in a religious tradition were not significantly associated with AA membership at the final interview after controlling for other potentially confounding variables.

Discussion

To our knowledge, this is the first study that explored correlates of AA membership in a sample of AA and non-AA abstinent alcoholics. We found that a number of factors found in previous literature to predict AA participation in samples of alcoholics whose drinking was allowed to vary were also found in this study to predict AA membership in a sample of abstinent alcoholics. In this study, higher drinking consequences and higher scores on the Loving God subscale were associated with subsequent AA membership. These findings are consistent with prior research which analyzed samples of alcoholics whose drinking was allowed to vary. Specifically, our study confirmed that even in a sample of successfully remitted alcoholics drinking severity and positive disposition toward religion and spirituality are independently predictive of AA membership. These significant findings make a contribution to the literature on the characteristics predictive of AA membership among individuals successful at achieving abstinence. Following the work of Avants, Beitel, and Margolin (2005), included throughout the following discussion are the words of those respondents included in this analysis as recorded during the baseline interview. These quotes were chosen to verbalize examples of baseline constructs associated with later AA membership.

While female gender was a predictor of AA membership in the bivariate analysis, it was not significant in the multivariate model. Humphreys et al. (1991) reported that women would be more likely to affiliate with AA than men despite a recent Alcoholics Anonymous membership survey that found that 67% of AA members are male (Alcoholics Anonymous, 2008). Further research using larger samples would aid in clarifying the role of gender for AA membership.

Findings in the literature related to extraversion suggest that people-oriented, social individuals might be more comfortable at AA meetings and in AA culture. Speaking in front of others, making new friends, calling others on the phone, doing service, going out for coffee, and helping others are all highly interpersonal staples in the AA culture. While these activities are voluntary, the social nature of AA may be more desirable for some people than others. In our study, extraversion was not significantly associated with AA membership. This is contrary to research by Kelly and Moos (2003) and Janowsky et al. (1999). One possibility for this negative finding may have been the circumscribed sample (i.e., all abstinent or within-group heterogeneity). While extraversion was not significantly associated with AA membership, supplemental data from qualitative interviews suggest that it remains an important area of future study. For example, a 41-year-old white woman who had previously participated in AA was asked the following question, “What do you think has been most helpful in dealing with some of the problems of alcohol?” She said, “Going to AA and getting involved with people because I’m a very social person so I do like people around me and I like to be active and do stuff.” Her response indicates that she responded positively to AA in part because she is social and outgoing.

Negative consequences of drinking were predictive of AA membership among abstinent alcoholics in both the bivariate and multivariate analyses. Similar findings have been reported by Emrick et al. (1993) and by Tonigan, Bogenschutz, and Miller (2006), who found that Type B (more severe) alcoholics were more likely to have sustained attendance in AA than less severe alcoholics. More severe drinking consequences may be predictive of later AA membership as the AA program encourages an admission of defeat and surrender in the battle with alcohol. Step 1 of the AA program involves admitting that one has become “powerless over alcohol” and that life has become “unmanageable” (AA, 1986, p. 5). AA members routinely self-identify as alcoholics. It might take a greater number of negative

consequences to bring an individual to come to accept one's powerlessness over alcohol and to embrace the stigma of the alcoholic label.

Interestingly in the bivariate model, belief in God was not predictive of AA membership in this sample but belief in a loving God was. Further, having been raised in a religious tradition was predictive of AA membership. The Loving God subscale remained significant even when controlling for other variables in the multivariate model. These findings add to evidence found in previous studies that spiritual and religious constructs are predictive of AA participation or retention (Emrick et al., 1993; Kelly & Moos, 2003). The current study furthers the understanding of this dimension by controlling for previous AA involvement, ruling out that the findings are byproducts of previous AA involvement. This suggests that a generally favorable view of God and previous experience with religiosity in childhood may be characteristic of individuals who later affiliate with AA. Those who relate to ideas of religion and spirituality in a positive way might be more open to trying and joining a spiritually-based recovery program such as AA.

A 45-year-old Black male respondent with high scores on measures of spirituality and religiousness found the spirituality of AA to be highly resonant with his attitude and world view:

[The AA founders] got a knowledge [*sic*] of human nature. You know, and it's spiritual. I like to read spiritual a lot, and AA is actually like a next Bible for me... You know, they talk about it being a way of living, not just a program... It's a way to get along with people, and treat people right. You know, it always tells you: you have to look at yourself, which is a spiritual principle. Turn it, turn it, have an inventory. The inventory is you, not somebody else's, you turn it around, and that's what caught me with AA.

>

This participant has a positive disposition toward spirituality and religiousness. He finds self examination, a way of being with others, and human nature itself to be implicitly spiritual. In these ways the spirituality of the AA program seems to be a good match with his pre-existing view of spirituality in human life. He is sober and identifies as an AA member at the 3-year follow-up.

Other respondents were more ambivalent. A 32-year-old white female participant, who at baseline had not yet attended her first AA meeting but who at 3-year follow-up was an AA member, conveyed that while she was critical of religion, there were aspects of a spiritual community that she yearned for. She stated:

Religion--it's bad for me and I think it's bad for a lot of people but I think some people really need the structure of it and really need the faith to get through horrible crises. In a way I am jealous of that. I wish I had that because I think it's really helpful, when people die, you have all that support from church but I'd sort of like to be like that but also still be intellectual and smart and know about everything else that's out there.

While she feels religion is bad for her, her desire for structure and the social support of a faith community may portend her later affiliation with Alcoholics Anonymous. Clinicians can look for this orientation in clients and incorporate it when making referrals to AA.

Taken together, our findings indicate that having a more favorable view of God and more negative consequences of drinking may indicate a propensity to be amenable to AA membership, even in a sample of abstinent alcoholics controlling for previous AA

involvement. These predictors suggest that certain individuals may find AA a more comfortable and synchronistic experience. This has important clinical implications. Counselors can explore these concepts with clients and can encourage those with these characteristics to give AA a try as such clients may find AA to be a good fit. Practitioners may also work with individuals who do not have the characteristics associated with AA membership to help them explore whether these obstacles might be overcome or find other sources of support for sobriety with which they would feel more comfortable. The spiritual component of the AA program is a barrier to some. But some clients with negative views of spirituality or religion may in fact be ambivalent. This ambivalence could be resolved using clinical interventions such as motivational interviewing in order to reduce obstacles to trying AA.

Future Research and Limitations

This study is comprised of a sample of individuals who achieved stable long-term abstinence, it controlled for previous AA involvement, and it used a high standard for classifying AA membership. However, several limitations are important to note. First, the sample size is relatively small. This was a consequence of choosing a sample who had strictly achieved one year of total abstinence by the 3-year follow-up period. While this makes a contribution by controlling for favorable drinking outcomes, it is a limitation in terms of sample size. Therefore, results of this study should be interpreted with caution and replicated in larger samples. In addition, there are limitations related to generalizability because participants were recruited in one small Midwestern city.

While there is a literature on untreated alcoholics who experience natural recovery (Sobell, Ellingstad, & Sobel, 2000), little is known about treated alcoholics who achieve total abstinence without AA participation. Future research on this group can explore the resources used by Non-AA members to support their abstinence. This work could help to expand options for sobriety for individuals who are not comfortable in AA meetings.

Conclusion

Prior studies have examined predictors of AA involvement or dropout, but to our knowledge this is the first that focuses on correlates of AA membership among a sample of successfully remitted alcohol-dependent individuals. Some findings of the current study replicate those from previous research and further showed that variables reflecting spirituality and severity of alcoholism are uniquely associated with AA membership. Results enhance understanding of the factors associated with AA participation and suggest that greater attention to spirituality-related constructs might improve the translational impact of research on AA.

Acknowledgments

The first author thanks Jaclyn Christine Bradley and Arielle Sherman for their assistance in documenting the selected quotations and Kirk Brower for his helpful review of the manuscript.

Role of Funding Sources

Funding for this study was provided by NIAAA Grant R01 AA014442 and T32 AA007477-21. NIAAA had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

References

- Alcoholics Anonymous. Alcoholics Anonymous 2007 Membership Survey. New York: 2008. Author. Retrieved February 27, 2010 from http://www.aa.org/pdf/products/p-48_07survey.pdf

- Alcoholics Anonymous. Twelve Steps and Twelve Traditions. New York: Author; 1986.
- Avants SK, Beitel M, Margolin A. Making the shift from 'addict self' to 'spiritual self': Results from a Stage I study of Spiritual Self-Schema (3-S) therapy for the treatment of addiction and HIV risk behavior. *Mental Health, Religion & Culture*. 2005; 8(3):167–177.
- Benson P, Spilka B. God image as a function of self-esteem and locus of control. *Journal for the Scientific Study of Religion*. 1973; 12:297–310.
- Bond J, Kaskutas LA, Weisner C. The persistent influence of social networks and Alcoholics Anonymous on abstinence. *Journal of Studies on Alcohol*. 2003; 64(4):579–588. [PubMed: 12921201]
- Connors GJ, Tonigan JS, Miller WR. A measure of religious background and behavior for use in behavior change research. *Psychology of Addictive Behaviors*. 1996; 10(2):90–96.
- Costa, PT., Jr; McCrae, RR. The NEO Personality Inventory manual. Odessa, FL: Psychological Assessment Resources; 1985.
- Costa, PT., Jr; McCrae, RR. Revised NEO Personality Inventory (NEO-PI-R) and NEO Five-Factor Inventory (NEO-FFI) manual. Odessa, FL: Psychological Assessment Resources; 1992.
- Crumbaugh JC, Maholick LT. An experimental study in existentialism: The psychometric approach to Frankl's concept of "noogenic" neurosis. *Journal of Clinical Psychology*. 1964; 20:200–207. [PubMed: 14138376]
- Emrick, CD.; Tonigan, JS.; Montgomery, HA.; Little, L. Alcoholics Anonymous: What is currently known?. In: McCrady, BS.; Miller, WR., editors. *Research on Alcoholics Anonymous: Opportunities and alternatives*. New Brunswick, NJ: Rutgers Center on Alcohol Studies; 1993. p. 41-76.
- Fetzer Monograph: Multidimensional Measurement of Religiousness/Spirituality for Use in Health Research: A Report of the Fetzer Institute/National Institute on Aging Workshop Group. Kalamazoo, MI: John E. Fetzer Institute; 1999.
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I), Clinician Version: User's Guide. Washington, DC: American Psychiatric Press; 1997.
- Humphreys K, Mavis B, Stoffelmayr B. Factors predicting attendance at self-help groups after substance abuse treatment: preliminary findings. *Journal of Consulting & Clinical Psychology*. 1991; 59(4):591–593. [PubMed: 1918563]
- Humphreys K, Mavis B, Stoffelmayr B. Are twelve step programs appropriate for disenfranchised groups? Evidence from a study of posttreatment mutual help involvement. *Prevention in Human Services*. 1994; 11:65–179.
- Janowsky DS, Boone A, Morter S, Howe L. Personality and alcohol/substance-use disorder patient relapse and attendance at self-help group meetings. *Alcohol & Alcoholism*. 1999; 34(3):359–369. [PubMed: 10414612]
- Kelly JF, Moos R. Dropout from 12-step self-help groups: Prevalence, predictors, and counteracting treatment influences. *Journal of Substance Abuse Treatment*. 2003; 24:241–250. [PubMed: 12810145]
- Miller, WR.; Tonigan, JS.; Longabaugh, R. The Drinker Inventory of Consequences (DrInC): An Instrument for Assessing Adverse Consequences of Alcohol Abuse: Test manual. Project MATCH Monograph Series. Vol. Volume 4. Rockville, MD: National Institute on Alcohol Abuse & Alcoholism; 1995.
- Morgenstern J, Labouvie E, McCrady B, Kahler CW, Frey RM. Affiliation with Alcoholics Anonymous after treatment: A study of its therapeutic effects and mechanisms of action. *Journal of Consulting and Clinical Psychology*. 1997; 65(5):768–777. [PubMed: 9337496]
- Robinson EAR, Krentzman AR, Pierce LS, Webb J, Brower KJ. Six-month changes in spirituality and religiousness in alcoholics predict drinking outcomes at nine months. 2010 Manuscript submitted for publication.
- Sobell LC, Ellingstad TP, Sobell MB. Natural recovery from alcohol and drug problems: Methodological review of the research with suggestions for future directions. *Addiction*. 2000; 95:749–764. [PubMed: 10885050]

- Sobell, LC.; Sobell, MD. Timeline Follow-back: A technique for assessing self-reported alcohol consumption. In: Litten, R.; Allen, J., editors. *Measuring Alcohol Consumption: Psychosocial and Biochemical Methods*. Totowa, NJ: Humana Press; 1992.
- Sobell LC, Brown J, Leo GI, Sobell MB. The reliability of the Alcohol Timeline Follow-back when administered by telephone and by computer. *Drug and Alcohol Dependence*. 1996; 42(1):49–54. [PubMed: 8889403]
- Terra MB, Barros HMT, Stein AT, Figueira I, Athayde LD, Palermo LH, Tergolina LP, Rovani JS. Predictors of engagement in the Alcoholics Anonymous group or to psychotherapy among Brazilian alcoholics: A six-month follow-up study. *European Archives of Psychiatry and Clinical Neuroscience*. 2007; 257(4):237–244. [PubMed: 17401731]
- Tonigan JS, Bogenschutz MP, Miller WR. Is alcoholism typology a predictor of both Alcoholics Anonymous affiliation and disaffiliation after treatment? *Journal of Substance Abuse Treatment*. 2006; 30(4):323–330. [PubMed: 16716847]
- Tonigan JS, Connors GJ, Miller WR. The Alcoholics Anonymous Involvement (AAI) Scale: Reliability and norms. *Psychology of Addictive Behaviors*. 1996; 10(2):75–80.
- Underwood LG, Teresi JA. The daily spiritual experience scale: Development, theoretical description, reliability, exploratory factor analysis, and preliminary construct validity using health-related data. *Annals of Behavioral Medicine*. 2002; 24(1):22–33. [PubMed: 12008791]
- Vaillant G. A 60-year follow-up of alcoholic men. *Addiction*. 2003; 98(8):1043–1051. [PubMed: 12873238]

Table 1

Demographic and AA-Related Variables at Baseline for Individuals Who Achieved at least One Year of Abstinence with and without Alcoholics Anonymous in a Subset of the Life Transition Study (N = 81) Observations, Percentages or Means, Standard Deviations

	AA Member* (n=44)	Not an AA Member* (n=37)	Total Sample (n=81)
Demographics			
Marital Status			
Never married	9 (20.5%)	8 (21.6%)	17 (21.0%)
Married/living with sign. Other	19 (43.2%)	12 (32.4%)	31 (38.3%)
Separated, divorced, widowed	16 (36.4%)	17 (45.9%)	33 (40.7%)
Race			
White	36 (81.8%)	33 (89.2%)	69 (85.2%)
Other**	8 (10.8%)	4 (18.2%)	12 (14.8%)
Household Income			
< \$30,000	19 (43.2%)	19 (51.4%)	38 (46.9%)
\$30,001-\$60,000	11 (25.0%)	8 (21.6%)	19 (23.4%)
\$60,000 +	14 (31.8%)	10 (27.0%)	24 (29.6%)
Employment			
Employed	24 (54.5%)	16 (43.2%)	40 (49.4%)
Not employed	20 (45.5%)	21 (56.8%)	41 (50.6%)
Gender			
Male	26 (59.1%)	30 (81.1%)	56 (69.1%)
Female	18 (40.9%)	7 (18.9%)	25 (30.9%)
Age	47.95 (11.44)	46.32 (12.10)	47.21 (11.70)
Education	14.52 (2.06)	14.30 (2.28)	14.42 (2.16)
Spiritual and Religious			
Raised in a religious tradition	32 (72.7%)	18 (48.6%)	50 (61.7%)
Belief in God	37 (84.1%)	25 (67.6%)	62 (76.5%)
Loving God	25.95 (3.80)	23.06 (5.47)	24.66 (4.81)
Substance-Related			
Drinking Consequences (SIP)	25.64 (9.47)	20.57 (11.62)	23.32 (10.74)
Total Lifetime # of AA Meetings Attended	331.77 (590.52)	114.70 (337.99)	232.62 (500.68)
Alcoholics Anonymous Involvement	2.80 (2.31)	1.51 (1.56)	2.21 (2.09)
Personality			
Extraversion	38.66 (7.33)	35.27 (7.99)	37.11 (7.78)

* "In AA" refers to individuals who, at 3-year follow up, had achieved at least 1 year of abstinence and considered themselves members of Alcoholics Anonymous. "Not in AA" refers to individuals who, at 3-year follow-up, had achieved at least 1 year of abstinence and did not participate in Alcoholics Anonymous or any other abstinence support group.

** Other includes African Americans (n=9), Native Americans (n=2), and multiracial individuals (n=2).

Table 2

Results from bivariate logistic regression analyses for individual baseline predictors of AA membership (N = 81)

		95% CI	
Individual predictors	OR	Lower	Upper
Demographics			
Education ₁₀₋₁₉ (in years)	1.050	.856	1.289
Gender (female = 1, male = 0)	2.967	1.071	8.218
Race (white = 1, other* = 0)	.545	.150	1.981
Spiritual and Religious			
Loving God ₁₁₋₃₀	1.147	1.029	1.278
I believe in God (yes = 1, no = 0)	2.537	.878	7.333
Raised in a Religious Tradition (yes = 1, no = 0)	2.815	1.116	7.099
Personality			
Extraversion ₁₋₄₅	1.062	.998	1.129
Substance-Related			
Drinking Consequences (SIP) ₁₋₄₄	1.047	1.003	1.094
Alcoholics Anonymous Involvement ₀₋₇	1.394	1.090	1.782

Note: Each line represents a separate bivariate logistic regression model. OR = Odds ratio. CI = Confidence interval. Values in bold are statistically significant at an alpha of .05. Figures in subscript are the range of values for continuous variables.

* Other includes African Americans (n=9), Native Americans (n=1), and multiracial individuals (n=2).

Table 3

Multiple Logistic Regression Analysis of Longitudinal Predictors of AA Membership

Predictors	OR	95% CI	
		lower	upper
AA Involvement	1.345	1.002	1.805
Gender (female = 1, male = 0)	2.795	.795	9.827
Loving God	1.155	1.009	1.320
Raised in a religious tradition	2.619	.842	8.143
Drinking Consequences (SIP)	1.060	1.003	1.119

Note: OR = Odds ratio. CI = Confidence interval. Values in bold are statistically significant based on a 95% CI that does not bound 1.0.