Published in final edited form as:

Psychiatry Res. 2012 July 30; 198(2): 212–215. doi:10.1016/j.psychres.2011.08.014.

Inflammatory Markers in Antipsychotic-naïve Patients with Nonaffective Psychosis and Deficit vs. Nondeficit Features

Clemente Garcia-Rizo^{a,*}, Emilio Fernandez-Egea^{b,c}, Cristina Oliveira^a, Azucena Justicia^b, Miguel Bernardo^{a,d,e}, and Brian Kirkpatrick^f

^aSchizophrenia Program, Department of Psychiatry, Neuroscience Institute, Hospital Clinic, University of Barcelona, Barcelona, Spain

^bDepartment of Psychiatry, University of Cambridge, Addenbrookes Hospital, CB2 0QQ Cambridge, UK

^cCambridgeshire and Peterborough NHS Foundation Trust, Huntingdon PE29 3RJ, UK

^dInstitute of Biomedical Research Agusti Pi i Sunyer (IDIBAPS), Barcelona, Spain

^eCIBERSAM, Madrid, Spain

^fDepartment of Psychiatry, Texas A&M University College of Medicine and Scott & White Healthcare, Temple, TX, USA

Abstract

Newly diagnosed, antipsychotic-naïve patients with nonaffective psychosis appear to have increases in pro-inflammatory cytokines. Patients characterized by primary, enduring negative symptoms (deficit symptoms) differ from patients without such features with regard to course of illness, treatment response, risk factors and metabolic disturbances. We hypothesized that they would also differ on the inflammatory markers, interleukin-6 (IL6) and C-reactive protein (CRP) concentrations. Newly diagnosed, antipsychotic-naïve patients with nonaffective psychosis were categorized into deficit ($N=20$) and nondeficit ($N=42$) groups, and were matched on age, gender, body mass index, smoking, cortisol level, socioeconomic status, and the severity of psychotic symptoms. Fasting concentrations of IL6 levels were significantly higher in deficit (mean [SD]) (8.0 pg/ml [12.7]) than nondeficit patients (0.3 pg/ml [1.3]) CRP levels were also significantly higher in the deficit patients (0.3 mg/dl [0.4]) vs. (0.2 mg/dl [0.4]), respectively. In contrast, 2-h glucose concentrations (2HG) in a glucose tolerance test were lower in the deficit than the nondeficit group. Our results show a double dissociation with regard to glucose intolerance and inflammation: the deficit group has greater inflammation, but less severe glucose intolerance. These results provide further evidence for the validity of the deficit/nondeficit categorization.

Keywords

Deficit schizophrenia; interleukin 6; C-reactive protein; glucose tolerance; negative symptoms; first episode psychosis

© 2011 Elsevier Ireland Ltd. All rights reserved.

*Corresponding Author: Clemente Garcia-Rizo, Hospital Clinic Schizophrenia Program, Department of Psychiatry, Villarroel 170, 08036 Barcelona, Spain, +34932275547 (phone), +34932275548 (fax), cgarcia3@clinic.ub.es.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Schizophrenia subtypes have long been the topic of research. One proposed subtype is deficit schizophrenia, which is characterized by primary (or idiopathic), enduring negative symptoms. Deficit/nondeficit differences in course of illness, signs and symptoms other than negative symptoms, risk factors, biological correlates, and treatment response have all been found (Kirkpatrick et al., 2001). In some studies, despite the presence of primary negative symptoms, the deficit group did not have a more severe form of the same abnormality found in nondeficit schizophrenia. Instead, with regard to some variables, the deficit group was either less severely affected than the nondeficit group, or neither group was more severely affected, but both differed from control subjects (e.g., excess winter birth in nondeficit schizophrenia vs. excess summer births in deficit schizophrenia) (Messias et al., 2004).

A number of studies have found increased pro-inflammatory markers in patients with schizophrenia (Miller et al., 2011). Inflammation is governed by the interaction of multiple specific mediators including cytokines, prostaglandins, and chemokines. One of these mediators, c-reactive protein (CRP), an acute-phase protein, is a general marker of inflammation. Increased levels of CRP have been associated with increased risk of inflammatory cardiovascular disorders (Lowe, 2005). In schizophrenia patients, CRP has been associated with cognitive impairment (Dickerson et al., 2007), more severe psychopathology (Fan et al., 2007), and certain clinical phenotypes (Akanji et al., 2009). Although CRP values were found to be increased in treated schizophrenia patients (Carrizo et al., 2008), the difference was not replicated in a drug-naïve sample compared with matched control subjects (Fernandez-Egea et al., 2009).

Interleukin-6 (IL6) is a multifunctional cytokine with biological activities that include regulation of immune response, inflammation, and haematopoiesis. IL6 induces T-cell proliferation, systemic inflammatory symptoms such as fever, generalized fatigue, and anorexia and angiogenesis by augmenting the production of vascular endothelial growth factor. It also has both neurotrophic and neurotoxic effects in different neuronal types and at different developmental stages (Deverman and Patterson, 2009). One of these effects is stimulation of CRP production (Papanicolaou and Vgontzas, 2000). We had previously shown that glucose tolerance differs between deficit and nondeficit schizophrenia patients (Kirkpatrick et al., 2009). In the current study, we tested the hypothesis that patients with psychosis and deficit features would also differ from patients without these features (nondeficit patients) with regard to IL6 and CRP concentrations.

2. Materials and methods

2.1. Subjects

Newly diagnosed, antipsychotic-naïve patients with nonaffective psychosis were assessed at the time of their first contact with psychiatric services. In addition to patients with schizophrenia, patients with schizophreniform disorder, delusional disorder, and psychosis not otherwise specified were included because these disorders share clinical and genetic factors with schizophrenia, and most newly diagnosed individuals with these disorders receive a diagnosis of schizophrenia within the first year after first clinical contact (Addington et al., 2005). Subjects with schizoaffective disorder were not included. The patients had a maximum cumulative lifetime exposure to antipsychotics of 1 week, and no antipsychotic use in the 30 days before study entry.

The setting was an academic medical center in Barcelona that provides health services for residents of the surrounding catchment area as part of the Spanish national health care system. The catchment area is a relatively homogeneous, middle class/upper middle class

neighbourhood in the center of Barcelona. Further details on this study have been presented (Fernandez-Egea et al., 2009).

Additional inclusion and exclusion criteria for all subjects were as follows: 1) age from 18 to 64 years, 2) no history of diabetes or other serious medical or neurological condition associated with glucose intolerance or insulin resistance (e.g. Cushing's disease), 3) not taking a medication associated with insulin resistance (hydrochlorothiazide, furosemide, ethacrynic acid, metolazone, chlorthalidone, beta blockers, glucocorticoids, phenytoin, nicotinic acid, cyclosporine, pentamidine, or narcotics), and 4) no history of cocaine use in the previous 30 days.

2.2. Assessments

All subjects were interviewed using the Spanish translation of the Structured Clinical Interview for DSM-IV Axis I Disorders, clinician version (SCID-I). They were also administered the Dartmouth Assessment of Lifestyle Inventory to quantify substance abuse. Socioeconomic status (SES) was assessed using the Hollingshead–Redlich scale (Hollingshead and Redlich, 1958)

2.3. Procedure

All participants were given a 2-h, 75 g oral glucose tolerance test, which began between 08.00 and 09.00h after an overnight fast. Samples for IL6, CRP, and cortisol blood concentrations were also taken before the patients' ingested glucose. They were allowed to receive anti-anxiety medication (lorazepam) the night before blood was drawn to a maximum of 3 mg, but not on the day of assessment. No correlation was found between lorazepam administration and the variables studied. Height, weight, and waist and hip circumference, when wearing underwear and without shoes, were recorded between the baseline and two blood samples. The determination of IL6 was performed with a commercial enzyme immunoassay technique (Biosource, Nivelles, Belgium). Cortisol was measured using a radioimmunoassay (ImmuChem, Ivoz-Ramet, Belgium).

2.4 Deficit/nondeficit categorization

Patients were categorized into groups with and without deficit features using the Proxy for the Deficit Syndrome, or PDS (Arango et al., 2011; Chemerinski et al., 2006; Dickerson et al., 2006; Kirkpatrick et al., 1996a; Kirkpatrick et al., 1996b; Kirkpatrick et al., 1993; Kirkpatrick et al., 2009; Kirkpatrick and Galderisi, 2008; Kirkpatrick et al., 2002a; Kirkpatrick et al., 2003; Kirkpatrick et al., 1998; Kirkpatrick et al., 1996c; Kirkpatrick et al., 2000; Kirkpatrick et al., 2002b; Messias and Bienvenu, 2003; Messias and Kirkpatrick, 2001; Messias et al., 2004; Tek et al., 2001; Wang et al., 2008). Patients' ($N=95$) deficit/nondeficit symptomatology was assessed with the Positive and Negative Syndrome Scale (PANSS) item scores; the PANSS rating was completed on the day of the glucose tolerance test, before any of the blood measures were available to the investigators. Each patient was assigned a PDS score as defined by PANSS item scores as follows: PDS = blunted affect + lack of spontaneity and flow of conversation – (hostility + guilt + anxiety + depressive mood). This score quantified the combination of prominent negative symptoms and a lack of distress that is characteristic of deficit compared with nondeficit patients (Kirkpatrick et al., 1993, 1994), with high scores reflecting deficit-like features (high negative symptom scores and an absence of dysphoria) and low scores reflecting their absence. These scores ranged from –11 to 5. Patients with high PDS scores (below –1) were assigned to the putative deficit group ($N=24$), while the other patients (PDS scores over 0) were designated nondeficit ($N=71$).

The deficit and nondeficit groups were matched, without knowledge of outcome variables, on age, gender, smoking, body mass index (BMI), and socioeconomic status (SES) of the family of origin by omitting nondeficit patients until these variables were very similar for the two groups. Due to lack of matched data on several patients, our total number decreased from 95 to 62. The validity of these two matched groups, deficit ($N=20$) and nondeficit ($N=42$), was tested by comparing the features of the two groups with those found in deficit and nondeficit groups in other studies (Kirkpatrick et al., 2001). In addition to high negative symptoms and an absence of dysphoria (present in that group by definition, because of the use of the PDS), a valid deficit group, compared with nondeficit subjects, would not have a greater length of illness or more severe psychotic symptoms. In our examination of psychotic symptoms, the sum of the delusions and hallucinatory behaviour items on the PANSS was analyzed separately from the conceptual disorganization item. Physical examination and other laboratory tests did not reveal the presence of infection in these patients, and in this relatively young group, no one met criteria for the metabolic syndrome.

2.5. Statistical analysis

We conducted two sets of analyses. In the first set of analyses, deficit and nondeficit groups were matched for age, gender, body mass index, and smoking (the last variable was highly correlated with all drug abuse). This matching made it necessary to omit some nondeficit patients from the first set of analyses. Only those subjects with data on all of the relevant variables were included, yielding matched groups of 20 deficit and 42 nondeficit subjects. Matching was done without knowledge of IL6 and CRP concentrations. Because the IL6 and CRP values were not normally distributed, these variables were log-transformed for all analyses. In order to assess the robustness of our results with the matched sample, we also conducted logistic regression analysis among patients who were not matched. In bivariate correlations, age had a correlation < 0.20 with both IL6 and CRP (both log-transformed), whereas gender, smoking, BMI, and SES all had a correlation > 0.40 . Given these results, we conducted two analyses; in each, deficit vs. nondeficit categorization was the dependent variable. In one of these analyses, log-transformed IL6 and age were the independent variables, while in the other analysis, log-transformed CRP and age were the independent variables. Because of some missing data, there were 21 deficit and 67 nondeficit subjects in the IL6 analysis versus 23 deficit and 64 nondeficit subjects in the CRP analysis.

Statistical tests were performed using version 17.0 for Windows of SPSS (Statistical Package for the Social Sciences).

3. Results

The validity of the categorization was good, as demonstrated by the clinical and demographic features of the matched deficit and nondeficit groups (see Table 1). Both groups were overwhelmingly composed of Spanish Caucasians. By definition, the deficit group had more severe negative symptoms but less dysphoria than the nondeficit group, but there was no significant difference in demographic characteristics or positive psychotic symptoms.

Both IL6 and CRP concentrations were significantly higher in the deficit than the nondeficit group (see Table 1). In contrast, 2-h glucose concentrations in the glucose tolerance test were significantly higher in the nondeficit than the deficit group. Cortisol values were similar in the two groups. The effect size for log-transformed IL6 was 1.1, much greater than the normal definition of a large effect size; the effect size for the log-transformed CRP values was 0.86, which also represents a large effect size.

In the logistic regression analyses, consistent with the results in the smaller, matched sample, the deficit group had significantly higher IL6 and CRP values (respective p values <0.001 and <0.03 ; data not shown). There was no relationship between the inflammatory markers and duration of untreated psychosis (data not shown).

4. Discussion

In this study, newly diagnosed, antipsychotic-naïve patients with nonaffective psychosis and features of deficit schizophrenia were found to have higher IL6 and CRP concentrations than did matched patients without deficit features. These findings could not be attributed to confounding by age, gender, SES of the family of origin, BMI, smoking status, cortisol concentrations, positive psychotic symptoms, antipsychotic use (as the subjects were antipsychotic-naïve), or antidepressant or mood-stabilizing drugs (no subject had previous use of an antidepressant or a mood-stabilizer). In contrast, the group with deficit features had lower glucose concentrations than did those without these characteristics, as we had previously shown in a sample that overlapped substantially with the current subjects (Kirkpatrick et al., 2009).

4.1. Study limitations

A limitation of our study is that the number of subjects with deficit features was limited. However, the limited sample size should have biased toward a negative result because of poor statistical power. Another limitation was that our putative deficit/nondeficit categorization was made using the PDS, rather than the Schedule for the Deficit Syndrome, the standard instrument used to distinguish patients with and without primary negative symptoms (Kirkpatrick et al., 1989). The categorizations in this study were validated by comparison to the clinical features of deficit and nondeficit groups diagnosed in the literature; the groups in this study were very similar in their characteristics to those diagnosed by the Schedule for the Deficit Syndrome. Moreover, the Proxy for the Deficit Syndrome has been validated in a number of previous publication (Arango et al., 2011; Chemerinski et al., 2006; Dickerson et al., 2006; Kirkpatrick et al., 1996a; Kirkpatrick et al., 1996b; Kirkpatrick et al., 1993; Kirkpatrick et al., 2009; Kirkpatrick and Galderisi, 2008; Kirkpatrick et al., 2002a; Kirkpatrick et al., 2008; Kirkpatrick et al., 2003; Kirkpatrick et al., 1998; Kirkpatrick et al., 1996c; Kirkpatrick et al., 2000; Kirkpatrick et al., 2002b; Messias and Bienvenu, 2003; Messias and Kirkpatrick, 2001; Messias et al., 2004; Tek et al., 2001; Wang et al., 2008). The use of the PDS also should have biased toward a negative study because of miscategorizations compared to use of the SDS. The large effect sizes we found suggest these difference are robust, but replication is needed. We only examined IL6 and CRP as inflammatory markers; examination of others, such as tumor necrosis factor-alpha would be desirable.

4.2. Conclusions

Our results represent a double dissociation in metabolic measures, with greater inflammation but less severe glucose intolerance in the deficit group. (We have previously shown that both patient groups differ from matched control subjects.) Higher CRP concentrations in the deficit group are consistent with a previous study in which treated patients with high CRP values had higher scores on the negative symptom subscale of the PANSS (Fan et al., 2007). Our results are also consistent with other studies in which elevated concentrations of IL6 were found in patients with predominantly negative symptoms and/or poor therapy outcome (Schwarz et al., 2001), longer duration and an unfavourable course (Müller et al., 2000) or treatment resistance (Lin et al., 1998), as the characteristics of deficit patients are similar to these correlates. A marginally statistically association between the severity of negative symptoms and a polymorphism of the IL6 gene (IL6-1 rs1800797) has also been described

in schizophrenia (Liu et al., 2010). Some (Akiyama, 1999; Fernandez-Egea et al., 2009; Ganguli et al., 1994) but not all studies (Singh et al., 2009) have found an increased IL6 in drug-naïve patients. The described abnormality in each of these markers provides a degree of validation of the abnormality in the other.

Whether these abnormalities in cytokines are part of the pathophysiology or even etiology of schizophrenia, or a result of the stress associated with the disorder, is not known (Corcoran et al., 2003). Another complicating factor for many studies (although not the present study) is that antipsychotic treatment may also modulate the production of cytokines (Szuster-Ciesielska et al., 2004). Inflammation may prove to be an important target for the development of therapeutics, as some studies suggest they may increase the speed of recovery from a relapse (Akhondzadeh et al., 2007; Bresee et al., 2006; Laan et al., 2007; Müller et al., 2004, 2005, 2010).

We previously hypothesized that deficit schizophrenia is a separate disease within the syndrome of schizophrenia, based on studies that had shown that deficit and nondeficit schizophrenia differed with regard to signs and symptoms, risk factors, course of illness, biological correlates, and treatment response (Kirkpatrick et al., 2001). The current double dissociation in glucose tolerance and inflammation supports this hypothesis. Two other double dissociations have also been found. One is in event-related potentials: N1 has been found to be abnormal in deficit but not nondeficit patients, whereas P3 was affected only in nondeficit patients (Mucci et al., 2007). Another double dissociation is in season of birth: deficit schizophrenia is associated with summer birth (Messias et al., 2004) whereas schizophrenia as a whole (and therefore nondeficit schizophrenia) is associated with winter birth (Torrey et al., 1997).

Deficit and nondeficit patients have been found to have different risk factors. Differences that suggest these two groups have differences in etiopathophysiology. The metabolic differences we report here may therefore reflect differences in prenatal metabolic programming. Although replication would be needed, our present results support the hypothesis of a separate disease.

Acknowledgments

This work was supported by grant RO1 DK069265 from the National Institute of Diabetes and Digestive and Kidney Diseases (Dr. Kirkpatrick), NARSAD (Dr. Fernandez-Egea) and by the Government of Catalonia, Comissionat per Universitats i Recerca del Departament d'Innovació, Universitats i Empresa (DIUE) 2009SGR1295 (Dr. Bernardo). The views stated in this article represent those of the authors and are not official statements of the National Institute of Diabetes and Digestive and Kidney Diseases or the National Institutes of Health. The authors thank Eduard Parellada and Linh Nguyen for their assistance.

References

- Addington J, Saeedi H, Addington D. The course of cognitive functioning in first episode psychosis: changes over time and impact on outcome. *Schizophr Research*. 2005; 78:35–43.
- Akanji AO, Ohaeri JU, Al-Shammri S, Fatania HR. Association of blood levels of C-reactive protein with clinical phenotypes in Arab schizophrenic patients. *Psychiatry Research*. 2009; 169:56–61. [PubMed: 19619902]
- Akiyama K. Serum levels of soluble IL-2 receptor alpha, IL-6 and IL-1 receptor antagonist in schizophrenia before and during neuroleptic administration. *Schizophrenia Research*. 1999; 37:97–106. [PubMed: 10227112]
- Akhondzadeh S, Tabatabaee M, Amini H, Ahmadi Abhari SA, Abbasi SH, Behnam B. Celecoxib as adjunctive therapy in schizophrenia: a double-blind, randomized and placebo-controlled trial. *Schizophrenia Research*. 2007; 90:179–185. [PubMed: 17208413]

- Arango C, Bobes J, Kirkpatrick B, Garcia-Garcia M, Rejas J. Psychopathology, coronary heart disease and metabolic syndrome in schizophrenia spectrum patients with deficit versus non-deficit schizophrenia: findings from the CLAMORS study. *European Neuropsychopharmacology*. 2011
- Bresee CJ, Delrahim K, Maddux RE, Dolnak D, Ahmadpour O, Rapaport MH. The effects of celecoxib augmentation on cytokine levels in schizophrenia. *International Journal of Neuropsychopharmacology*. 2006; 9:343–348. [PubMed: 16035956]
- Carrizo E, Fernandez V, Quintero J, Connell L, Rodriguez Z, Mosquera M, Acosta A, Baptista T. Coagulation and inflammation markers during atypical or typical antipsychotic treatment in schizophrenia patients and drug-free first-degree relatives. *Schizophr Research*. 2008; 103:83–93.
- Chemerinski E, Reichenberg A, Kirkpatrick B, Bowie CR, Harvey PD. Three dimensions of clinical symptoms in elderly patients with schizophrenia: prediction of six-year cognitive and functional status. *Schizophrenia Research*. 2006; 85:12–19. [PubMed: 16624531]
- Corcoran C, Walker E, Huot R, Mittal V, Tessner K, Kestler L, Malaspina D. The stress cascade and schizophrenia: etiology and onset. *Schizophrenia Bulletin*. 2003; 29:671–692. [PubMed: 14989406]
- Deverman BE, Patterson PH. Cytokines and CNS development. *Neuron*. 2009; 64:61–78. [PubMed: 19840550]
- Dickerson F, Kirkpatrick B, Boronow J, Stallings C, Origoni A, Yolken R. Deficit schizophrenia: association with serum antibodies to cytomegalovirus. *Schizophrenia Bulletin*. 2006; 32:396–400. [PubMed: 16166610]
- Dickerson F, Stallings C, Origoni A, Boronow J, Yolken R. C-reactive protein is associated with the severity of cognitive impairment but not of psychiatric symptoms in individuals with schizophrenia. *Schizophrenia Research*. 2007; 93:261–265. [PubMed: 17490859]
- Fan X, Pristach C, Liu EY, Freudenreich O, Henderson DC, Goff DC. Elevated serum levels of C-reactive protein are associated with more severe psychopathology in a subgroup of patients with schizophrenia. *Psychiatry Research*. 2007; 149:267–271. [PubMed: 17112596]
- Fernandez-Egea E, Bernardo M, Donner T, Conget I, Parellada E, Justicia A, Esmatjes E, Garcia-Rizo C, Kirkpatrick B. Metabolic profile of antipsychotic-naïve individuals with non-affective psychosis. *British Journal of Psychiatry*. 2009; 194:434–438. [PubMed: 19407273]
- Ganguli R, Yang Z, Shurin G, Chengappa KN, Brar JS, Gubbi AV, Rabin BS. Serum interleukin-6 concentration in schizophrenia: elevation associated with duration of illness. *Psychiatry Research*. 1994; 51:1–10. [PubMed: 7910974]
- Kirkpatrick B, Buchanan RW, Ross DE, Carpenter WT Jr. A separate disease within the syndrome of schizophrenia. *Archives of General Psychiatry*. 2001; 58:165–171. [PubMed: 11177118]
- Kirkpatrick B, Fernandez-Egea E, Garcia-Rizo C, Bernardo M. Differences in glucose tolerance between deficit and nondeficit schizophrenia. *Schizophr Research*. 2009; 107:122–127.
- Kirkpatrick B, Amador XF, Flaum M, Yale SA, Gorman JM, Carpenter WT Jr, Tohen M, McGlashan T. The deficit syndrome in the DSM-IV Field Trial: I. Alcohol and other drug abuse. *Schizophrenia Research*. 1996a; 20:69–77. [PubMed: 8794495]
- Kirkpatrick B, Amador XF, Yale SA, Bustillo JR, Buchanan RW, Tohen M. The deficit syndrome in the DSM-IV Field Trial. Part II. Depressive episodes and persecutory beliefs. *Schizophrenia Research*. 1996b; 20:79–90. [PubMed: 8794496]
- Kirkpatrick B, Buchanan RW, Breier A, Carpenter WT Jr. Case identification and stability of the deficit syndrome of schizophrenia. *Psychiatry Research*. 1993; 47:47–56. [PubMed: 8516416]
- Kirkpatrick B, Buchanan RW, Breier A, Carpenter WT Jr. Depressive symptoms and the deficit syndrome of schizophrenia. *Journal of Nervous and Mental Disease*. 1994; 182:452–455. [PubMed: 8040655]
- Kirkpatrick B, Buchanan RW, McKenney PD, Alphs LD, Carpenter WT Jr. The Schedule for the Deficit syndrome: an instrument for research in schizophrenia. *Psychiatry Research*. 1989; 30:119–123. [PubMed: 2616682]
- Kirkpatrick B, Galderisi S. Deficit schizophrenia: an update. *World Psychiatry*. 2008; 7:143–147. [PubMed: 18836581]
- Kirkpatrick B, Herrera Castanedo S, Vazquez-Barquero JL. Summer birth and deficit schizophrenia: Cantabria, Spain. *Journal of Nervous and Mental Disease*. 2002a; 190:526–532. [PubMed: 12193837]

- Kirkpatrick B, Messias E, LaPorte D. Schizoid-like features and season of birth in a nonpatient sample. *Schizophrenia Research*. 2008; 103:151–155. [PubMed: 18261884]
- Kirkpatrick B, Messias EM, Tek C. Substance abuse and the heterogeneity of schizophrenia a population-based study. *Schizophrenia Research*. 2003; 62:293–294. [PubMed: 12837529]
- Kirkpatrick B, Ram R, Amador XF, Buchanan RW, McGlashan T, Tohen M, Bromet E. Summer birth and the deficit syndrome of schizophrenia. *American Journal of Psychiatry*. 1998; 155:1221–1226. [PubMed: 9734546]
- Kirkpatrick B, Ram R, Bromet E. The deficit syndrome in the Suffolk County Mental Health Project. *Schizophrenia Research*. 1996c; 22:119–126. [PubMed: 8958595]
- Kirkpatrick B, Ross DE, Walsh D, Karkowski L, Kendler KS. Family characteristics of deficit and nondeficit schizophrenia in the Roscommon Family Study. *Schizophrenia Research*. 2000; 45:57–64. [PubMed: 10978873]
- Kirkpatrick B, Tek C, Allardyce J, Morrison G, McCreadie RG. Summer birth and deficit schizophrenia in Dumfries and Galloway, southwestern Scotland. *American Journal of Psychiatry*. 2002b; 159:1382–1387. [PubMed: 12153832]
- Laan W, Seltén JP, Grobbee DE, Smeets H, Kahn RS, Burger H. Non-steroidal anti-inflammatory drugs and the risk of psychosis. *European Neuropsychopharmacology*. 2007; 17:309–311. [PubMed: 17097858]
- Lin A, Kenis G, Bignotti S, Tura GJ, De Jong R, Bosmans E, Pioli R, Altamura C, Scharpe S, Maes M. The inflammatory response system in treatment-resistant schizophrenia: increased serum interleukin-6. *Schizophrenia Research*. 1998; 32:9–15. [PubMed: 9690329]
- Liu YL, Liu CM, Fann CS, Yang WC, Chen YH, Tseng LJ, Liu SK, Hsieh MH, Hwang TJ, Chan HY, Chen JJ, Chen WJ, Hwu HG. Genetic variants of IL-6 and its receptor are not associated with schizophrenia in Taiwan. *Neuroscience Letters*. 2010; 468(3):330–333. [PubMed: 19914334]
- Lowe GD. Circulating inflammatory markers and risks of cardiovascular and non-cardiovascular disease. *Journal of Thrombosis and Haemostasis*. 2005; 3:1618–1627.
- Messias E, Bienvenu OJ. Suspiciousness and alcohol use disorders in schizophrenia. *Journal of Nervous and Mental Disease*. 2003; 191:387–390. [PubMed: 12826920]
- Messias E, Kirkpatrick B. Summer birth and deficit schizophrenia in the epidemiological catchment area study. *Journal of Nervous and Mental Disease*. 2001; 189:608–612. [PubMed: 11580004]
- Messias E, Kirkpatrick B, Bromet E, Ross D, Buchanan RW, Carpenter WT Jr, Tek C, Kendler KS, Walsh D, Dollfus S. Summer birth and deficit schizophrenia: a pooled analysis from 6 countries. *Archives of General Psychiatry*. 2004; 61:985–989. [PubMed: 15466671]
- Miller BJ, Buckley P, Seabolt W, Mellor A, Kirkpatrick B. Meta-analysis of cytokine alterations in schizophrenia: clinical status and antipsychotic affects. *Biological Psychiatry*. 2011 Electronic publication 11 June 2011. 10.1016/j.biopsych.2011.04.013
- Mucci A, Galderisi S, Kirkpatrick B, Bucci P, Volpe U, Merlotti E, Centanaro F, Catapano F, Maj M. Double dissociation of N1 and P3 abnormalities in deficit and nondeficit schizophrenia. *Schizophrenia Research*. 2007; 92:252–261. [PubMed: 17363220]
- Müller N, Krause D, Dehning S, Musil R, Schennach-Wolff R, Obermeier M, Möller HJ, Klauss V, Schwarz MJ, Riedel M. Celecoxib treatment in an early stage of schizophrenia: results of a randomized, double-blind, placebo-controlled trial of celecoxib augmentation of amisulpride treatment. *Schizophrenia Research*. 2010; 121:118–124. [PubMed: 20570110]
- Müller N, Riedel M, Ackenheil M, Schwarz MJ. Cellular and humoral immune system in schizophrenia: a conceptual re-evaluation. *World Journal of Biological Psychiatry*. 2000; 1:173–179. [PubMed: 12607212]
- Müller N, Riedel M, Schwarz MJ, Engel RR. Clinical effects of COX-2 inhibitors on cognition in schizophrenia. *European Archives Psychiatry Clinical Neuroscience*. 2005; 255:149–151.
- Müller N, Strassnig M, Schwarz MJ, Ulmschneider M, Riedel M. COX-2 inhibitors as adjunctive therapy in schizophrenia. *Expert Opinion Investigational Drugs*. 2004; 13:1033–1044.
- Papanicolaou DA, Vgontzas AN. Interleukin-6: the endocrine cytokine. *Journal of Clinical Endocrinology Metabolism*. 2000; 85:1331–1333. [PubMed: 10720086]

- Schwarz MJ, Müller N, Riedel M, Ackenheil M. The Th2-hypothesis of schizophrenia: a strategy to identify a subgroup of schizophrenia caused by immune mechanisms. *Medical Hypotheses*. 2001; 56:483–486. [PubMed: 11339852]
- Singh B, Bera NK, Nayak CR, Chaudhuri TK. Decreased serum levels of interleukin-2 and interleukin-6 in Indian Bengalee schizophrenic patients. *Cytokine*. 2009; 47:1–5. [PubMed: 19502081]
- Szuster-Ciesielska A, Slotwinska M, Stachura A, Marmurowska-Michalowska H, Kandefer-Szerszen M. Neuroleptics modulate cytokine and reactive oxygen species production in blood leukocytes of healthy volunteers. *Archives of Immunology and Experimental Therapy (Warsz)*. 2004; 52:59–67.
- Tek C, Kirkpatrick B, Buchanan RW. A five-year followup study of deficit and nondeficit schizophrenia. *Schizophrenia Research*. 2001; 49:253–260. [PubMed: 11356586]
- Torrey EF, Miller J, Rawlings R, Yolken RH. Seasonality of births in schizophrenia and bipolar disorder: a review of the literature. *Schizophrenia Research*. 1997; 28:1–38. [PubMed: 9428062]
- Wang X, Yao S, Kirkpatrick B, Shi C, Yi J. Psychopathology and neuropsychological impairments in deficit and nondeficit schizophrenia of Chinese origin. *Psychiatry Research*. 2008; 158:195–205. [PubMed: 18243336]

Table

Demographic, Clinical, and Metabolic Characteristics of Patients With and Without Deficit Features

	Deficit (N=20)	Non Deficit (N=42)	p value
Age	29.8 (9.8)	27.7 (6.5)	.40
% Gender (Male)	67	61	.58
Cigarettes/day	4.7 (6.9)	6.8 (7.9)	.32
Body Mass Index ¹	23.4 (2.5)	22.5 (4.3)	.41
SES ²	5.0 (2.2)	5.3 (2.3)	.68
PANSS Items Scores:			
Disorganization	3.9 (1.8)	3.5 (1.4)	.34
Reality distortion ³	8.1 (3.3)	8.2 (2.2)	.82
IL-6 (pg/mL) ⁴	8.0 (12.7)	0.3 (1.3)	<.001
CRP (mg/dL) ⁴	0.3 (0.4)	0.2 (0.4)	.01
2HG (mg/dL)	99.1 (25.1)	119.2 (36.9)	.03
Cortisol	18.7 (5.9)	17.4 (5.9)	.43

Values are mean (SD), except for % male

P values are from T-test, except for % male (chi-square) and cigarettes (Mann-Whitney U).

¹N =19 for the deficit group

²For deficit N=14 and nondeficit N=37

³Reality distortion scores = scores for hallucinations + delusions

⁴L-6 and CRP values were analyzed using log-transformed data