Published in final edited form as:

Cell Tissue Res. 2012 July ; 349(1): 27–37. doi:10.1007/s00441-012-1388-6.

NEUROTROPHIC FACTORS IN COMBINATORIAL APPROACHES FOR SPINAL CORD REGENERATION

Julianne McCall¹, Norbert Weidner¹, and Armin Blesch^{1,2,*}

¹Spinal Cord Injury Center, Heidelberg University Hospital, 69118 Heidelberg, Germany

²Department of Neurosciences, University of California, San Diego, La Jolla, CA 92093, USA

Abstract

Axonal regeneration is inhibited by a plethora of different mechanisms in the adult central nervous system (CNS). While neurotrophic factors have been shown to stimulate axonal growth in numerous animal models of nervous system injury, a lack of suitable growth substrates, an insufficient activation of neuron-intrinsic regenerative programs and extracellular inhibitors of regeneration limit the efficacy of neurotrophic factor delivery for anatomical and functional recovery after spinal cord injury. Thus, growth-stimulating factors will likely have to be combined with other treatment approaches to tap into the full potential of growth factor therapy for axonal regeneration. In addition, the temporal and spatial distribution of growth factors have to be tightly controlled to achieve biologically active concentrations, to allow for the chemotropic guidance of axons and to prevent adverse effects related to the widespread distribution of neurotrophic factors. Here, we will review the rationale for combinatorial treatments in axonal regeneration and summarize some recent progress in promoting axonal regeneration in the injured CNS using such approaches.

Keywords

neurotrophin; gene therapy; axonal regeneration; spinal cord injury; cell transplantation

INTRODUCTION

In vitro and in vivo studies conducted over the last 30 years have identified several key issues that need to be considered to improve axonal regeneration in the adult mammalian central nervous system (CNS). There is ample evidence that axon growth-inhibitory molecules present in the adult injured CNS contribute to an unfavorable environment for axonal regeneration (Silver and Miller, 2004, Xie and Zheng, 2008). Glial and inflammatory reactions after injury further add to the adverse milieu of the injured spinal cord and diminish the prospects for axonal regeneration (Bethea and Dietrich, 2002, Fitch and Silver, 2008, Popovich and McTigue, 2009). Molecules and guidance cues present in the developing nervous system are not preserved throughout adulthood or have different functions, resulting in an insufficient stimulation of axonal growth and limited means to direct axons towards appropriate targets (Giger, et al., 2010). Neuronal and glial cell death at a site of injury cannot be easily replenished from the intrinsic neural stem cell pool, and cystic degeneration resulting in the formation of fluid-filled cavities at sites of spinal cord injury (SCI) require the provision of a substrate for axons to bridge across a lesion site.

*To whom correspondence should be addressed: Armin Blesch, Ph.D., Spinal Cord Injury Center, Heidelberg University Hospital, Schlierbacher Landstrasse 200 a, 69118 Heidelberg, Germany, Phone: +49-6221-969200, Fax: +49-6221-969308, armin.blesch@med.uni-heidelberg.de.

Finally, differentiated neurons in the adult CNS do not activate transcriptional programs that are adequate and sufficiently sustained to allow for regeneration to occur.

No single experimental treatment can tackle all of these factors and will therefore be limited in its efficacy to augment axonal regeneration. During development and axon regeneration in the injured peripheral nervous system (PNS), multiple mechanisms interact to contribute to efficient and controlled axonal growth. Physical guidance by appropriate cellular orientation and provision of a growth-permissive extracellular matrix clearly contribute to directed axonal extension. In both circumstances, neurons have an intrinsically high capacity for axonal growth either based on their developmental status (Blackmore, et al., 2010, Moore, et al., 2009) or the initiation of appropriate changes in gene expression (Michailevski, et al., 2010). At the axon tip, growth is also stimulated by levels of growth factors higher than those present in the adult CNS, and expression patterns of growth factors are tightly regulated in a spatial and temporal manner. Thus, in both situations, multiple mechanisms act synergistically to result in significant axonal growth. The above-mentioned examples from development and PNS injury and the empirical evidence about limited effects of single treatments on axonal regeneration in the injured spinal cord corroborate a need to address more than one factor in axon regeneration failure. In this review, we aim to summarize recent data on regeneration, sprouting and plasticity of injured axons following SCI. We will focus on studies that have combined means to stimulate axonal growth, including neurotrophic factor delivery, cellular grafts, neutralization or digestion of growth-inhibitory molecules and activation of the intrinsic growth capacity of injured neurons.

Bridging the lesion site

The significance of the environment for adult CNS regeneration became for the first time clearly evident in a landmark study by Aguayo and colleagues, showing that brainstem and spinal cord neurons can extend axons into an implanted peripheral nerve graft (David and Aguayo, 1981, Richardson, et al., 1980). At the same time, these studies also demonstrated that adult neurons in the CNS retain some capacity for regeneration if they are provided with a favorable milieu, challenging earlier beliefs that these neurons completely lack the ability to activate any intrinsic regenerative program.

These studies stimulated the search for a cellular or acellular matrix that could mimic some aspects of a peripheral nerve graft or provide an even superior environment. Cellular grafts comprised of fibroblasts, bone marrow stromal cells (Ankeny, et al., 2004, Hofstetter, et al., 2002), Schwann cells (Li and Raisman, 1994, Tuszynski, et al., 1998, Weidner, et al., 1999, Xu, et al., 1997, Xu, et al., 1995, Xu, et al., 1994), olfactory ensheathing cells (Li, et al., 1997, Lu, et al., 2006, Ramon-Cueto, et al., 2000, Ramon-Cueto and Nieto-Sampedro, 1994, Richter and Roskams, 2008), stem cells (Lepore and Fischer, 2005, Lu, et al., 2003, Pfeifer, et al., 2004) and embryonic tissue (Bregman, et al., 1993, Mori, et al., 1997, Stokes and Reier, 1992, Tessler, 1991, Tessler, et al., 1997) are some of the cellular substrates that have been examined in this context (Tetzlaff, et al., 2011). However, a uniform cellular graft lacks important aspects found in the regenerating PNS. Upon injury in the PNS, profound changes in the expression patterns of neurotrophic factors can be observed, changes which have proven critical to successful peripheral nerve regeneration by studies using knockout animals and neutralization of growth factors (English, et al., 2005, Geremia, et al., 2010, Zhang, et al., 2000, Zhong, et al., 1999). Schwann cells upregulate expression of nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), glial cell line-derived neurotrophic factor (GDNF), ciliary neurotrophic factor (CNTF) and other growth factors in particular in the distal nerve stump, resulting in chemotropic growth factor gradients. Upon reinnervation of the distal nerve, expression of growth factors declines (Funakoshi, et al., 1993, Heumann, et al., 1987, Naveilhan, et al., 1997, Sendtner, et al., 1992, Trupp, et al., 1997). When grafted into the injured spinal cord, available data support that a peripheral

nerve may retain patterns of neurotrophic factor upregulation observed after PNS injury, whereby NGF levels exhibit two peaks, immediately at the time of injury and some days following the injury, while BDNF levels gradually increase and peak at later time points (Kuo, et al., 2011, Meyer, et al., 1992). Thus, a regulated spatial and temporal expression pattern of growth factors contributes to effective and rapid regeneration in the PNS.

Studies combining cell transplantation with neurotrophic factors using genetically modified cells have aimed to mimic some components of the growth-promoting environment of the PNS. Indeed, robust axonal growth into cellular transplants can be induced by neurotrophic factor production from grafts of genetically engineered cells. In summary, NGF and GDNF can induce growth of peptidergic and non-peptidergic nociceptive axons, respectively (Blesch and Tuszynski, 2003, Tuszynski, et al., 1996), growth of reticulospinal, raphespinal, rubrospinal and propriospinal axons is augmented by BDNF and NT-4/5 (Blesch, et al., 2004, Liu, et al., 1999, Lu, et al., 2005, Menei, et al., 1998) and NT-3 enhances growth of ascending sensory axons (Bradbury, et al., 1999, Taylor, et al., 2006). NT-3 delivery in the injured spinal cord also induces sprouting of corticospinal axons (Blits, et al., 2000, Fortun, et al., 2009, Grill, et al., 1997, Schnell, et al., 1994, Zhou, et al., 2003). While studies on cellular growth factor delivery indicated substantial growth responses for many axonal populations into the lesion site, corticospinal axons failed to penetrate most cellular grafts examined to date. Instead, corticospinal axon sprouting was mainly observed in remaining gray matter in partial lesion models. Given the importance of corticospinal projections in fine motor control in humans and primates, the quest for an appropriate substrate for the corticospinal tract therefore remains a high priority.

Equally important, axons responding to cellular delivery of neurotrophic factors rarely exit cellular grafts to regenerate across the lesion site, limiting the usefulness of this approach. Only when growth factors are delivered beyond the lesion site in combination with cell or tissue transplantation have regenerating axons been shown to bridge across a site of SCI (see below).

Guidance and directional growth of regenerating axons

After injury in the PNS, a regulated spatial and temporal expression pattern of neurotrophic factors contributes to successful regeneration through context-dependent survival, growth and guidance signaling. Numerous studies have described the neurotrophin family, consisting of NGF, BDNF, NT-3 and NT-4/5, and their tyrosine kinase receptors TrkA, -B, and -C and the low-affinity p75 neurotrophin receptor, which are integral to regenerative processes (Boyd and Gordon, 2003, Cui, 2006). Upon neurotrophin binding to their appropriate Trk receptor along the axon, the activated complex is endocytosed and retrogradely transported via microtubule/dynein machinery to the cell soma to propagate signaling cascades that underlie neuronal survival and growth (Neet and Campenot, 2001). In addition to promoting neuronal survival, neurotrophins are also strong stimulators of axonal growth, specifically when receptors in the axon tip are activated (Campenot, 1982, Kimpinski, et al., 1997, Zhou and Snider, 2006). Neurotrophin stimulation at the growth cone can also induce chemotropic guidance, *in vitro* (Gundersen and Barrett, 1979) and *in vivo* in the PNS (Hu, et al., 2010) and CNS (Alto, et al., 2009, Taylor, et al., 2006).

Assuming that injured axons in the CNS are as responsive to growth factor gradients as they are during PNS regeneration, delivery of growth factors only into the SCI lesion site may actually prevent axons from exiting the lesion site and reentering the distal host spinal cord. Indeed, studies examining growth factor-expressing cellular grafts have rarely reported axon growth across the lesion site. Only if neurotrophin delivery is modified such that a growth factor gradient is established with highest concentrations beyond the lesion site, axons can bridge a site of SCI and reenter the host spinal cord. This was demonstrated using a

combination of bone marrow stromal cell grafts at the lesion site and lentiviral NT-3 gene transfer distal to a spinal cord lesion site, allowing for ascending dorsal column sensory axons to extend across a lesion site (Taylor, et al., 2006). In contrast, delivery of NT-3 only within the lesion leads to growth into but not beyond a lesion in the injured spinal cord (Lu, et al., 2003). Similarly, transplanted dorsal root ganglion (DRG) neurons can cross a lesion in the corpus callosum towards a viral NGF source (Jin, et al., 2008). Regeneration and guidance of sensory axons across the dorsal root entry zone has also been demonstrated towards areas of highest neurotrophin expression (Romero, et al., 2000). These responses can be further refined by combining chemoattractive and chemorepulsive guidance cues, thereby restricting the reinnervation pattern of sensory axons in the spinal cord or directing the growth of transplanted DRG neurons (Tang, et al., 2004, Ziemba, et al., 2008). Taken together, these studies support the hypothesis that injured adult neurons can respond to growth-promoting and growth-inhibiting guidance factors during axonal regeneration. By combining appropriate substrates with gradients of attractive and repulsive cues, more directed axonal growth can be achieved.

Directional growth of axons has also been demonstrated for grafts of neural precursors. Upon grafting a mixture of embryonic neural and glial restricted precursors into a spinal cord lesion site, axons originating from grafted neurons extend preferentially towards a source of BDNF established by lentiviral gene transfer (Bonner, et al., 2010). If progenitor grafts are close to the dorsal column nuclei, they can form synapses and serve as a relay for injured dorsal column sensory axons (Bonner, et al., 2011).

In addition to an appropriate spatial distribution of growth factors, the temporal availability of axon growth-promoting stimuli might be equally important. Cellular growth factor delivery in a lesion site has the potential to increase the pool of axons available to extend from a lesion into the distal spinal cord, but continuous expression in a lesion site certainly does not promote axonal growth into the inhibitory environment present beyond the lesion site. On the contrary, long-term overexpression of neurotrophic factors by genetically modified cells can lead to a gradual increase in graft size at least partially due to the invasion and proliferation of Schwann cells (Blesch and Tuszynski, 2003, Blesch, et al., 2004). Experiments with regulated neurotrophic factor expression further indicate that high levels of trophic factors are only necessary to induce growth into a lesion, whereas low levels are sufficient to sustain axons that have extended into a cellular graft (Blesch and Tuszynski, 2007). Clearly, there is an urgent need for efficient and safe means to regulate neurotrophic factor expression for axonal regeneration in vivo or, alternatively, vectors that automatically turn off within a defined time period due to epigenetic silencing.

Approaches to reproduce the physical guidance properties found in the PNS have also been described in recent years using different forms of biomaterials. Cellular grafts lacking orientation and alignment of grafted cells along the longitudinal axis of the spinal cord result in randomly directed axonal growth, thereby effectively limiting the size of a lesion that axons can successfully bridge. This is of particular importance when translating findings to the injured human spinal cord. The larger extent of lesions in the injured human spinal cord will require axons to bridge over significantly longer distances than any rodent studies have shown to date. Increasing the distance and rostro-caudal orientation of axonal growth therefore remains an important objective for translating axonal regeneration in the foreseeable future. Using linear scaffolds with defined pore sizes that can be filled with a cellular substrate, a more directional, linear axonal growth pattern can be achieved (Prang, et al., 2006, Stokols, et al., 2006, Straley, et al., 2010). When combined with neurotrophin gene transfer, the vast majority of axons entering a scaffold can extend throughout this bridge and are found to exit distally (Gros, et al., 2010). The latter study also activated a regenerative program in injured neurons, as discussed below.

Activating cell body responses to injury

Upon injury in a distal axon segment, a retrograde signal travels for extended distances (possibly more than one meter), manifesting in responses in injured neuronal cell bodies (Hanz, et al., 2003). This response has marked influence on the success or failure of axon regeneration. It can involve substantial, long-term changes in the expression of thousands of genes until target innervation is achieved (Costigan, et al., 2002, Goldberg, et al., 2002, Michaelievski, et al., 2010, Stam, et al., 2007, Xiao, et al., 2002) or it can lead to cellular atrophy (Kwon, et al., 2002) or, in the worst case, to neuronal cell death (Giehl and Tetzlaff, 1996). The latter was initially thought to be prominent in corticospinal motor neurons after SCI. However, more recent studies have indicated that corticospinal neurons do not die but become atrophic (Brock, et al., 2010, Nielson, et al., 2011), a reaction that has also been observed for neurons in the red nucleus following spinal cord lesions (Kobayashi, et al., 1997).

Preventing neuronal atrophy and activating regenerative programs in injured neurons might be a prerequisite for long-distance axonal regeneration. One of the most characterized systems for the activation of regenerative programs in injured neurons is the sensory system and so-called conditioning lesions (Richardson and Issa, 1984). If the central branch of dorsal column sensory axons is injured in the spinal cord, regeneration and sprouting does not occur. However, if a lesion of the peripheral branch precedes lesions of the central branch, neurons are pre-conditioned, and axons activate a regenerative program. This results in sprouting into or around a spinal cord lesion site or into a peripheral nerve graft (Neumann and Woolf, 1999, Richardson, et al., 1984). The signaling cascades leading to the activation of this regenerative program remain incompletely understood, but increases in intracellular second messengers, modulation of cytoskeletal signaling and upregulation of cytokines and growth factors have been implicated in this response. In particular, elevation of cAMP (Cai, et al., 2001, Gao, et al., 2004, Hannila and Filbin, 2008, Lu, et al., 2004, Neumann, et al., 2002, Qiu, et al., 2002), suppression of Rho signaling or constitutively activated growth cone components (Dergham, et al., 2002, Jain, et al., 2011) and activation of stat3 via neurotrophic cytokines (IL-6/LIF/CNTF) (Cafferty, et al., 2004, Cao, et al., 2006, Qiu, et al., 2005, Wu, et al., 2006) seem to play major roles in enhancing growth in an inhibitory CNS environment. Whereas a conditioning lesion preceding a central lesion has been clearly shown to enhance localized sprouting at a lesion site, a more significant effect is apparent when combined with neurotrophic factors to stimulate axonal growth. Provision of bone marrow stromal cells in a dorsal column lesion site combined with lentiviral NT-3 gene transfer rostral to the lesion site and a conditioning lesion increases the number and distance of sensory axons bridging across the lesion site (Alto, et al., 2009). Enhanced growth of sensory axons in such a combination is also observed if pre-conditioning lesions are replaced by elevations of cAMP (Lu, et al., 2004), although it is likely that cAMP is only partially responsible for effects observed after conditioning lesions. Importantly, even if peripheral neurons are conditioned after the central lesion in a chronic model of SCI, sensory axons extend across the grafted cellular bridge into the distal host spinal cord (Blesch, et al., 2012, Kadoya, et al., 2009). Although the *in vivo* regenerative response is diminished compared to an acute lesion (Blesch, et al., 2012), *in vitro* neurite growth assays and changes in DRG gene expression suggest that the intrinsic growth program is equally activated.

Much less is known about the activation of regenerative programs in CNS neurons. One of the best-studied CNS systems in regeneration is that of retinal ganglion neurons after optic nerve injury. Lens injury activates a signaling cascade that enhances optic nerve regeneration, and the mechanisms underlying this cascade have been partially identified. Activation of stat3 via CNTF/LIF appears to be one major factor (Muller, et al., 2009). In addition, PTEN knockout and siRNA studies in mice suggest that the activation of PI3-

kinase and the mTOR pathway can substantially enhance optic nerve regeneration (Park, et al., 2008). If activation of PI3-kinase via PTEN knockout is combined with activation of stat3 via SOCS3 knockout, the number and distance of regenerating axons is even further enhanced (Sun, et al., 2011). Activation of the mTOR pathway can also enhance corticospinal sprouting in mice, at least if it is activated prior to SCI (Liu, et al., 2010). As both pathways have been shown to contribute to the conditioning lesion effect in sensory axon regeneration, it appears that general principles in the activation of intrinsic regenerative programs exist.

Practical means to activate growth programs in injured CNS neurons with projections to the spinal cord are limited, and only few have been identified to date. Infusions and viral gene delivery of neurotrophins have been examined for several neuronal populations. In the red nucleus, provision of BDNF at the cell soma has been shown to ameliorate neuronal atrophy even at chronic time points up to 1 year post-lesion (Kwon, et al., 2007, Kwon, et al., 2002, Ruitenberg, et al., 2004). In addition, BDNF also increases expression of regeneration-associated genes, like alpha-tubulin and GAP43 (Kobayashi, et al., 1997), promoting growth of rubrospinal axons into peripheral nerve grafts placed in the lesioned spinal cord (Kobayashi, et al., 1997, Kwon, et al., 2002).

Infusion of BDNF into the motor cortex can apparently also activate regenerative responses, evident in enhanced corticospinal collateral sprouting in the injured spinal cord (Vavrek, et al., 2006). Although BDNF can reverse corticospinal neuronal atrophy in rodents and primates even when provided in a spinal cord lesion site (Brock, et al., 2010), BDNF or NT-4/5 delivered by genetically modified fibroblasts in the spinal cord fail to increase corticospinal sprouting after a midthoracic injury (Blesch, et al., 2004, Lu, et al., 2001). Indeed, BDNF delivery to corticospinal axons appears to only enhance regeneration into a transplant when the cognate receptor, trkB, is overexpressed by corticospinal neurons and BDNF-expressing grafts are placed in a subcortical lesion site (Hollis, et al., 2009).

Other studies have focused on genes important in CNS development to enhance the innate regenerative capacity (Blackmore, et al., 2010, Moore, et al., 2009, Wong, et al., 2006, Yip, et al., 2006, Yip, et al., 2010) or cell adhesion molecules (Andrews, et al., 2009), but these treatments have not been combined with substrates for axonal growth or other means to stimulate axon regeneration.

Target innervation and functional integration

If axons regenerate across a lesion in the injured spinal cord, finding appropriate targets and establishing functional synapses will be a prerequisite for behavioral recovery. From hundreds of neurons present in the vicinity of a regenerated axon, proper connections need to be formed to have functional implications. Spontaneous recovery of function after SCI can at least partially be attributed to localized axon sprouting and the formation of new connections. For example, corticospinal and reticulospinal axons have been shown to sprout towards propriospinal neurons and motor neurons after SCI (Ballermann and Fouad, 2006, Bareyre, et al., 2004, Fouad, et al., 2001, Girgis, et al., 2007, Vavrek, et al., 2006, Weidner, et al., 2001, Z'Graggen, et al., 2000). Extensive collateral sprouting that is accompanied by behavioral recovery is also observed in the primate spinal cord (Rosenzweig, et al., 2010). Propriospinal neurons can serve as relays to allow for supraspinal control of motor function without any direct connections beyond a lesion in the spinal cord (Courtine, et al., 2008).

Taken together, these studies indicate that some natural mechanisms exist that support the formation of appropriate new connections after SCI. However, rather little is known about the molecular basis and the guiding principles of these changes. Use- and disuse-related Hebbian mechanisms might be at least partially involved in these processes. Recent data

suggest that inherent mechanisms of sprouting and synapse formation might be strengthened by rehabilitative training. Rehabilitation and training, possibly via upregulation of neurotrophic factors including BDNF and insulin-like growth factor 1 by specific neuronal populations, could provide one means to attract or stabilize synaptic connections (Gomez-Pinilla, et al., 2002, Hutchinson, et al., 2004, Vaynman and Gomez-Pinilla, 2005, Ying, et al., 2005).

Combining neurotrophins with degradation of inhibitory molecules

Inhibitory extracellular matrix including chondroitinsulfate proteoglycans (CSPGs) is one key factor limiting axonal regeneration in the injured mammalian CNS. Within 24 hours post-injury, reactive astrocytes begin to deposit high amounts of CSPGs at the lesion site, establishing a long-lasting inhibitory boundary that has been comprehensively shown in vitro and in vivo to hinder regenerative growth of adult CNS axons and to inhibit synaptogenesis (Jones, et al., 2003, Silver and Miller, 2004). Degradation of the glycosaminoglycan (GAC) side chains of CSPG molecules by local administration of the enzyme chondroitinase ABC (ChABC) can enhance local sprouting and facilitate axonal regeneration, in part by reducing the inhibitory components of the glial scar and disrupting the CSPG perineuronal nets which envelope neuronal somata (Bartus, et al., 2011, Galtrey, et al., 2008). Without a cellular graft or trophic support, ChABC treatment upon acute or subchronic SCI induces moderate plasticity of surrounding tissue and some regeneration of select axonal tracts (Alilain, et al., 2011, Bradbury, et al., 2002), even when coupled with acute or delayed rehabilitative training (Garcia-Alias, et al., 2009, Wang, et al., 2011) or a systemic pharmacological neuroprotective agent (Yick, et al., 2004). When administered in parallel with a graft at the lesion site, ChABC has been shown to increase the number of axons regenerating across the lesion site into the distal host spinal cord (Chau, et al., 2004, Fouad, et al., 2005, Houle, et al., 2006, Tom and Houle, 2008). In combination with GDNF delivered directly to and around the lesion site to activate regenerative responses in chronically injured neurons, ChABC and a peripheral nerve graft applied four weeks post-injury stimulates descending axons to regenerate into and through a PNS graft and exit to form new connections (Tom, et al., 2009). The fast rate of inactivation of ChABC in vivo at physiological temperatures might be one limiting factor in ChABC-mediated effects on morphological and functional recovery. A recently described thermostabilized form of ChABC (ChABC/trehalose) results in suppressed CS-CAG levels for 6 weeks after SCI, three-fold longer than previously shown (Lee, et al., 2010). When ChABC/trehalose was employed in parallel with NT-3 in an agarose gel scaffold in the lesion site, a significant increase of serotonergic axon sprouting was observed adjacent to the lesion compared to control groups. Using a similar combination of ChABC (microinjection) and NT-3 (lentiviral delivery), the growth of transplanted adult DRG neurons into the CSPG-rich dorsal column nuclei can be increased 10-fold compared to ChABC or NT-3 treatment alone, strongly suggesting a cooperative interaction between mechanisms of disinhibition and a boost in growth-promoting signal (Massey, et al., 2008). Of clinical relevance, several studies consistently demonstrate that pathways facilitating pain perception appear not to be influenced by ChABC treatment (Barritt, et al., 2006, Karimi-Abdolrezaee, et al., 2010). Taken together, removal of inhibitory extracellular matrix components in conjunction with neurotrophin delivery appears to synergistically enhance axonal regeneration.

Conclusions

Over the last two decades, it has become increasingly clear that the lack of axonal regeneration in the mammalian CNS is not due to a single inhibitory mechanism or entirely to the reticence of CNS axons to regenerate. The complexity of SCI and the numerous factors influencing regeneration provide a clear rationale for combinatorial treatments, and several studies have now successfully demonstrated that effects of neurotrophic factors can

be further increased if growth factors are combined with other treatments, and vice versa. However, in vivo experiments examining multiple treatment parameters are laborious and demanding, requiring a large number of controls to test each single treatment and possible combination. A regressive analysis with control groups that receive systematically fewer treatments might be one means to collect evidence for improvements without initial testing of all necessary controls. In addition, interactions between treatments could not only change the dose response profile of each therapy but lead to negative consequences and functional deterioration. A better understanding of the mechanisms of each single treatment and more localized targeting of therapeutic approaches might be helpful to avoid adverse outcomes. Numerous challenges remain, but empirical evidence from several experimental treatments suggests that improved outcomes are possible with combinatorial strategies, justifying the further pursuit of multi-mechanistic approaches.

Acknowledgments

Supported by grants from NIH/NINDS (NS054883), Wings for Life, International Foundation for Research in Paraplegia (IFP/IRP), the EU (IRG268282) and a fellowship to J.M. (Landesgraduiertenförderung).

References

- Alilain WJ, Horn KP, Hu H, Dick TE, Silver J. Functional regeneration of respiratory pathways after spinal cord injury. *Nature*. 2011; 475:196–200. [PubMed: 21753849]
- Alto LT, Havton LA, Conner JM, Hollis ER II, Blesch A, Tuszynski MH. Chemotropic guidance facilitates axonal regeneration and synapse formation after spinal cord injury. *Nature neuroscience*. 2009; 12:1106–1113.
- Andrews MR, Czitkovich S, Dasse E, Vogelaar CF, Faissner A, Blits B, Gage FH, French-Constant C, Fawcett JW. Alpha9 integrin promotes neurite outgrowth on tenascin-C and enhances sensory axon regeneration. *J Neurosci*. 2009; 29:5546–5557. [PubMed: 19403822]
- Ankeny DP, McTigue DM, Jakeman LB. Bone marrow transplants provide tissue protection and directional guidance for axons after contusive spinal cord injury in rats. *Exp Neurol*. 2004; 190:17–31. [PubMed: 15473977]
- Ballermann M, Fouad K. Spontaneous locomotor recovery in spinal cord injured rats is accompanied by anatomical plasticity of reticulospinal fibers. *Eur J Neurosci*. 2006; 23:1988–1996. [PubMed: 16630047]
- Bareyre FM, Kerschensteiner M, Raineteau O, Mettenleiter TC, Weinmann O, Schwab ME. The injured spinal cord spontaneously forms a new intraspinal circuit in adult rats. *Nature neuroscience*. 2004; 7:269–277.
- Barritt AW, Davies M, Marchand F, Hartley R, Grist J, Yip P, McMahon SB, Bradbury EJ. Chondroitinase ABC promotes sprouting of intact and injured spinal systems after spinal cord injury. *J Neurosci*. 2006; 26:10856–10867. [PubMed: 17050723]
- Bartus K, James ND, Bosch KD, Bradbury EJ. Chondroitin sulphate proteoglycans: Key modulators of spinal cord and brain plasticity. *Exp Neurol*. 2011 in press.
- Bethea JR, Dietrich WD. Targeting the host inflammatory response in traumatic spinal cord injury. *Curr Opin Neurol*. 2002; 15:355–360. [PubMed: 12045737]
- Blackmore MG, Moore DL, Smith RP, Goldberg JL, Bixby JL, Lemmon VP. High content screening of cortical neurons identifies novel regulators of axon growth. *Mol Cell Neurosci*. 2010; 44:43–54. [PubMed: 20159039]
- Blesch A, Lu P, Tsukada S, Taylor L, Roet K, Coppola G, Geschwind D, Tuszynski MH. Conditioning lesions before or after spinal cord injury recruit broad genetic mechanisms that sustain axonal regeneration: superiority to cAMP-mediated effects. *Exp Neurol*. 2012 in press.
- Blesch A, Tuszynski MH. Cellular GDNF delivery promotes growth of motor and dorsal column sensory axons after partial and complete spinal cord transection and induces remyelination. *J Comp Neurol*. 2003; 467:403–417. [PubMed: 14608602]

- Blesch A, Tuszynski MH. Transient growth factor delivery sustains regenerated axons after spinal cord injury. *J Neurosci*. 2007; 27:10535–10545. [PubMed: 17898225]
- Blesch A, Yang H, Weidner N, Hoang A, Otero D. Axonal responses to cellularly delivered NT-4/5 after spinal cord injury. *Mol Cell Neurosci*. 2004; 27:190–201. [PubMed: 15485774]
- Blits B, Dijkhuizen PA, Boer GJ, Verhaagen J. Intercostal nerve implants transduced with an adenoviral vector encoding neurotrophin-3 promote regrowth of injured rat corticospinal tract fibers and improve hindlimb function. *Exp Neurol*. 2000; 164:25–37. [PubMed: 10877912]
- Bonner JF, Blesch A, Neuhuber B, Fischer I. Promoting directional axon growth from neural progenitors grafted into the injured spinal cord. *J Neurosci Res*. 2010; 88:1182–1192. [PubMed: 19908250]
- Bonner JF, Connors TM, Silverman WF, Kowalski DP, Lemay MA, Fischer I. Grafted neural progenitors integrate and restore synaptic connectivity across the injured spinal cord. *J Neurosci*. 2011; 31:4675–4686. [PubMed: 21430166]
- Boyd JG, Gordon T. Neurotrophic factors and their receptors in axonal regeneration and functional recovery after peripheral nerve injury. *Molecular neurobiology*. 2003; 27:277–324. [PubMed: 12845152]
- Bradbury EJ, Khemani S, Von R, King, Priestley JV, McMahon SB. NT-3 promotes growth of lesioned adult rat sensory axons ascending in the dorsal columns of the spinal cord. *Eur J Neurosci*. 1999; 11:3873–3883. [PubMed: 10583476]
- Bradbury EJ, Moon LD, Popat RJ, King VR, Bennett GS, Patel PN, Fawcett JW, McMahon SB. Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature*. 2002; 416:636–640. [PubMed: 11948352]
- Bregman BS, Kunkel-Bagden E, Reier PJ, Dai HN, McAtee M, Gao D. Recovery of function after spinal cord injury: mechanisms underlying transplant-mediated recovery of function differ after spinal cord injury in newborn and adult rats. *Exp Neurol*. 1993; 123:3–16. [PubMed: 8405277]
- Brock JH, Rosenzweig ES, Blesch A, Moseanko R, Havton LA, Edgerton VR, Tuszynski MH. Local and remote growth factor effects after primate spinal cord injury. *J Neurosci*. 2010; 30:9728–9737. [PubMed: 20660255]
- Cafferty WB, Gardiner NJ, Das P, Qiu J, McMahon SB, Thompson SW. Conditioning injury-induced spinal axon regeneration fails in interleukin-6 knock-out mice. *J Neurosci*. 2004; 24:4432–4443. [PubMed: 15128857]
- Cai D, Qiu J, Cao Z, McAtee M, Bregman BS, Filbin MT. Neuronal cyclic AMP controls the developmental loss in ability of axons to regenerate. *J Neurosci*. 2001; 21:4731–4739. [PubMed: 11425900]
- Campanot RB. Development of sympathetic neurons in compartmentalized cultures. II Local control of neurite growth by nerve growth factor. *Developmental biology*. 1982; 93:1–12. [PubMed: 7128926]
- Cao Z, Gao Y, Bryson JB, Hou J, Chaudhry N, Siddiq M, Martinez J, Spencer T, Carmel J, Hart RB, Filbin MT. The cytokine interleukin-6 is sufficient but not necessary to mimic the peripheral conditioning lesion effect on axonal growth. *J Neurosci*. 2006; 26:5565–5573. [PubMed: 16707807]
- Chau CH, Shum DK, Li H, Pei J, Lui YY, Wirthlin L, Chan YS, Xu XM. Chondroitinase ABC enhances axonal regrowth through Schwann cell-seeded guidance channels after spinal cord injury. *Faseb J*. 2004; 18:194–196. [PubMed: 14630702]
- Costigan M, Belfort K, Karchewski L, Griffin RS, D'Urso D, Allchorne A, Sitariski J, Mannion JW, Pratt RE, Woolf CJ. Replicate high-density rat genome oligonucleotide microarrays reveal hundreds of regulated genes in the dorsal root ganglion after peripheral nerve injury. *BMC Neurosci*. 2002; 3:16. [PubMed: 12401135]
- Courtine G, Song B, Roy RR, Zhong H, Herrmann JE, Ao Y, Qi J, Edgerton VR, Sofroniew MV. Recovery of supraspinal control of stepping via indirect propriospinal relay connections after spinal cord injury. *Nat Med*. 2008; 14:69–74. [PubMed: 18157143]
- Cui Q. Actions of neurotrophic factors and their signaling pathways in neuronal survival and axonal regeneration. *Molecular neurobiology*. 2006; 33:155–179. [PubMed: 16603794]

- David S, Aguayo AJ. Axonal elongation into peripheral nervous system “bridges” after central nervous system injury in adult rats. *Science*. 1981; 214:931–933. [PubMed: 6171034]
- Dergham P, Ellezam B, Essagian C, Avedissian H, Lubell WD, McKerracher L. Rho signaling pathway targeted to promote spinal cord repair. *J Neurosci*. 2002; 22:6570–6577. [PubMed: 12151536]
- English AW, Meador W, Carrasco DI. Neurotrophin-4/5 is required for the early growth of regenerating axons in peripheral nerves. *Eur J Neurosci*. 2005; 21:2624–2634. [PubMed: 15926911]
- Fitch MT, Silver J. CNS injury, glial scars, and inflammation: Inhibitory extracellular matrices and regeneration failure. *Exp Neurol*. 2008; 209:294–301. [PubMed: 17617407]
- Fortun J, Puzis R, Pearse DD, Gage FH, Bunge MB. Muscle injection of AAV-NT3 promotes anatomical reorganization of CST axons and improves behavioral outcome following SCI. *Journal of neurotrauma*. 2009; 26:941–953. [PubMed: 19275471]
- Fouad K, Pedersen V, Schwab ME, Brosamle C. Cervical sprouting of corticospinal fibers after thoracic spinal cord injury accompanies shifts in evoked motor responses. *Curr Biol*. 2001; 11:1766–1770. [PubMed: 11719218]
- Fouad K, Schnell L, Bunge MB, Schwab ME, Liebscher T, Pearse DD. Combining Schwann cell bridges and olfactory-ensheathing glia grafts with chondroitinase promotes locomotor recovery after complete transection of the spinal cord. *J Neurosci*. 2005; 25:1169–1178. [PubMed: 15689553]
- Funakoshi H, Frisen J, Barbany G, Timmusk T, Zachrisson O, Verge VM, Persson H. Differential expression of mRNAs for neurotrophins and their receptors after axotomy of the sciatic nerve. *J Cell Biol*. 1993; 123:455–465. [PubMed: 8408225]
- Galtrey CM, Kwok JC, Carulli D, Rhodes KE, Fawcett JW. Distribution and synthesis of extracellular matrix proteoglycans, hyaluronan, link proteins and tenascin-R in the rat spinal cord. *Eur J Neurosci*. 2008; 27:1373–1390. [PubMed: 18364019]
- Gao Y, Deng K, Hou J, Bryson JB, Barco A, Nikulina E, Spencer T, Mellado W, Kandel ER, Filbin MT. Activated CREB is sufficient to overcome inhibitors in myelin and promote spinal axon regeneration in vivo. *Neuron*. 2004; 44:609–621. [PubMed: 15541310]
- Garcia-Alias G, Barkhuysen S, Buckle M, Fawcett JW. Chondroitinase ABC treatment opens a window of opportunity for task-specific rehabilitation. *Nature neuroscience*. 2009; 12:1145–1151.
- Geremia NM, Pettersson LM, Hasmatali JC, Hryciw T, Danielsen N, Schreyer DJ, Verge VM. Endogenous BDNF regulates induction of intrinsic neuronal growth programs in injured sensory neurons. *Exp Neurol*. 2010; 223:128–142. [PubMed: 19646438]
- Giehl KM, Tetzlaff W. BDNF and NT-3, but not NGF, prevent axotomy-induced death of rat corticospinal neurons in vivo. *Eur J Neurosci*. 1996; 8:1167–1175. [PubMed: 8752586]
- Giger RJ, Hollis ER 2nd, Tuszynski MH. Guidance molecules in axon regeneration. *Cold Spring Harb Perspect Biol*. 2010; 2:a001867. [PubMed: 20519341]
- Girgis J, Merrett D, Kirkland S, Metz GA, Verge V, Fouad K. Reaching training in rats with spinal cord injury promotes plasticity and task specific recovery. *Brain*. 2007; 130:2993–3003. [PubMed: 17928316]
- Goldberg JL, Klassen MP, Hua Y, Barres BA. Amacrine-signaled loss of intrinsic axon growth ability by retinal ganglion cells. *Science*. 2002; 296:1860–1864. [PubMed: 12052959]
- Gomez-Pinilla F, Ying Z, Roy RR, Molteni R, Edgerton VR. Voluntary exercise induces a BDNF-mediated mechanism that promotes neuroplasticity. *J Neurophysiol*. 2002; 88:2187–2195. [PubMed: 12424260]
- Grill R, Murai K, Blesch A, Gage FH, Tuszynski MH. Cellular delivery of neurotrophin-3 promotes corticospinal axonal growth and partial functional recovery after spinal cord injury. *J Neurosci*. 1997; 17:5560–5572. [PubMed: 9204937]
- Gros T, Sakamoto JS, Blesch A, Havton LA, Tuszynski MH. Regeneration of long-tract axons through sites of spinal cord injury using templated agarose scaffolds. *Biomaterials*. 2010; 31:6719–6729. [PubMed: 20619785]
- Gundersen RW, Barrett JN. Neuronal chemotaxis: chick dorsal-root axons turn toward high concentrations of nerve growth factor. *Science*. 1979; 206:1079–1080. [PubMed: 493992]

- Hannila SS, Filbin MT. The role of cyclic AMP signaling in promoting axonal regeneration after spinal cord injury. *Exp Neurol*. 2008; 209:321–332. [PubMed: 17720160]
- Hanz S, Perlson E, Willis D, Zheng JQ, Massarwa R, Huerta JJ, Koltzenburg M, Kohler M, van-Minnen J, Twiss JL, Fainzilber M. Axoplasmic importins enable retrograde injury signaling in lesioned nerve. *Neuron*. 2003; 40:1095–1104. [PubMed: 14687545]
- Heumann R, Lindholm D, Bandtlow C, Meyer M, Radeke MJ, Misko TP, Shooter E, Thoenen H. Differential regulation of mRNA encoding nerve growth factor and its receptor in rat sciatic nerve during development, degeneration, and regeneration: role of macrophages. *Proc Natl Acad Sci U S A*. 1987; 84:8735–8739. [PubMed: 2825206]
- Hofstetter CP, Schwarz EJ, Hess D, Widenfalk J, El Manira A, Prockop DJ, Olson L. Marrow stromal cells form guiding strands in the injured spinal cord and promote recovery. *Proc Natl Acad Sci U S A*. 2002; 99:2199–2204. [PubMed: 11854516]
- Hollis ER 2nd, Jamshidi P, Low K, Blesch A, Tuszynski MH. Induction of corticospinal regeneration by lentiviral trkB-induced Erk activation. *Proc Natl Acad Sci U S A*. 2009; 106:7215–7220. [PubMed: 19359495]
- Houle JD, Tom VJ, Mayes D, Wagoner G, Phillips N, Silver J. Combining an autologous peripheral nervous system “bridge” and matrix modification by chondroitinase allows robust, functional regeneration beyond a hemisection lesion of the adult rat spinal cord. *J Neurosci*. 2006; 26:7405–7415. [PubMed: 16837588]
- Hu X, Cai J, Yang J, Smith GM. Sensory axon targeting is increased by NGF gene therapy within the lesioned adult femoral nerve. *Exp Neurol*. 2010; 223:153–165. [PubMed: 19733564]
- Hutchinson KJ, Gomez-Pinilla F, Crowe MJ, Ying Z, Basso DM. Three exercise paradigms differentially improve sensory recovery after spinal cord contusion in rats. *Brain*. 2004; 127:1403–1414. [PubMed: 15069022]
- Jain A, McKeon RJ, Brady-Kalnay SM, Bellamkonda RV. Sustained delivery of activated Rho GTPases and BDNF promotes axon growth in CSPG-rich regions following spinal cord injury. *PloS one*. 2011; 6:e16135. [PubMed: 21283639]
- Jin Y, Ziemba KS, Smith GM. Axon growth across a lesion site along a preformed guidance pathway in the brain. *Exp Neurol*. 2008; 210:521–530. [PubMed: 18261727]
- Jones LL, Margolis RU, Tuszynski MH. The chondroitin sulfate proteoglycans neurocan, brevican, phosphacan, and versican are differentially regulated following spinal cord injury. *Exp Neurol*. 2003; 182:399–411. [PubMed: 12895450]
- Kadota K, Tsukada S, Lu P, Coppola G, Geschwind D, Filbin M, Blesch A, Tuszynski MH. Combined intrinsic and extrinsic neuronal mechanisms facilitate bridging axonal regeneration one year after spinal cord injury. *Neuron*. 2009; 64:165–172. [PubMed: 19874785]
- Karimi-Abdolrezaee S, Eftekharpour E, Wang J, Schut D, Fehlings MG. Synergistic effects of transplanted adult neural stem/progenitor cells, chondroitinase, and growth factors promote functional repair and plasticity of the chronically injured spinal cord. *J Neurosci*. 2010; 30:1657–1676. [PubMed: 20130176]
- Kimpinski K, Campenot RB, Mearow K. Effects of the neurotrophins nerve growth factor, neurotrophin-3, and brain-derived neurotrophic factor (BDNF) on neurite growth from adult sensory neurons in compartmented cultures. *Journal of neurobiology*. 1997; 33:395–410. [PubMed: 9322157]
- Kobayashi NR, Fan DP, Giehl KM, Bedard AM, Wiegand SJ, Tetzlaff W. BDNF and NT-4/5 prevent atrophy of rat rubrospinal neurons after cervical axotomy, stimulate GAP-43 and α -tubulin mRNA expression, and promote axonal regeneration. *J Neurosci*. 1997; 17:9583–9595. [PubMed: 9391013]
- Kuo HS, Tsai MJ, Huang MC, Chiu CW, Tsai CY, Lee MJ, Huang WC, Lin YL, Kuo WC, Cheng H. Acid fibroblast growth factor and peripheral nerve grafts regulate Th2 cytokine expression, macrophage activation, polyamine synthesis, and neurotrophin expression in transected rat spinal cords. *J Neurosci*. 2011; 31:4137–4147. [PubMed: 21411654]
- Kwon BK, Liu J, Lam C, Plunet W, Oschipok LW, Hauswirth W, Di Polo A, Blesch A, Tetzlaff W. Brain-derived neurotrophic factor gene transfer with adeno-associated viral and lentiviral vectors

- prevents rubrospinal neuronal atrophy and stimulates regeneration-associated gene expression after acute cervical spinal cord injury. *Spine*. 2007; 32:1164–1173. [PubMed: 17495772]
- Kwon BK, Liu J, Messerer C, Kobayashi NR, McGraw J, Oschipok L, Tetzlaff W. Survival and regeneration of rubrospinal neurons 1 year after spinal cord injury. *Proc Natl Acad Sci U S A*. 2002; 99:3246–3251. [PubMed: 11867727]
- Lee H, McKeon RJ, Bellamkonda RV. Sustained delivery of thermostabilized chABC enhances axonal sprouting and functional recovery after spinal cord injury. *Proc Natl Acad Sci U S A*. 2010; 107:3340–3345. [PubMed: 19884507]
- Lepore AC, Fischer I. Lineage-restricted neural precursors survive, migrate, and differentiate following transplantation into the injured adult spinal cord. *Exp Neurol*. 2005; 194:230–242. [PubMed: 15899260]
- Li Y, Field M, Raisman G. Repair of adult rat corticospinal tract by transplants of olfactory ensheathing cells. *Science*. 1997; 277:2000–2002. [PubMed: 9302296]
- Li Y, Raisman G. Schwann cells induce sprouting in motor and sensory axons in the adult rat spinal cord. *J Neurosci*. 1994; 14:4050–4063. [PubMed: 8027762]
- Liu K, Lu Y, Lee JK, Samara R, Willenberg R, Sears-Kraxberger I, Tedeschi A, Park KK, Jin D, Cai B, Xu B, Connolly L, Steward O, Zheng B, He Z. PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nature neuroscience*. 2010; 13:1075–1081.
- Liu Y, Kim D, Himes BT, Chow SY, Schallert T, Murray M, Tessler A, Fischer I. Transplants of fibroblasts genetically modified to express brain-derived neurotrophic factor promote regeneration of adult rat rubrospinal axons and recovery of forelimb function. *J Neurosci*. 1999; 19:4370–4387. [PubMed: 10341240]
- Lu P, Blesch A, Tuszynski MH. Neurotrophism without neurotropism: BDNF promotes survival but not growth of lesioned corticospinal neurons. *J Comp Neurol*. 2001; 436:456–470. [PubMed: 11447589]
- Lu P, Jones LL, Snyder EY, Tuszynski MH. Neural stem cells constitutively secrete neurotrophic factors and promote extensive host axonal growth after spinal cord injury. *Exp Neurol*. 2003; 181:115–129. [PubMed: 12781986]
- Lu P, Jones LL, Tuszynski MH. BDNF-expressing marrow stromal cells support extensive axonal growth at sites of spinal cord injury. *Exp Neurol*. 2005; 191:344–360. [PubMed: 15649491]
- Lu P, Yang H, Culbertson M, Graham L, Roskams AJ, Tuszynski MH. Olfactory ensheathing cells do not exhibit unique migratory or axonal growth-promoting properties after spinal cord injury. *J Neurosci*. 2006; 26:11120–11130. [PubMed: 17065452]
- Lu P, Yang H, Jones LL, Filbin MT, Tuszynski MH. Combinatorial therapy with neurotrophins and cAMP promotes axonal regeneration beyond sites of spinal cord injury. *J Neurosci*. 2004; 24:6402–6409. [PubMed: 15254096]
- Massey JM, Amps J, Viapiano MS, Matthews RT, Wagoner MR, Whitaker CM, Alilain W, Yonkof AL, Khalyfa A, Cooper NG, Silver J, Onifer SM. Increased chondroitin sulfate proteoglycan expression in denervated brainstem targets following spinal cord injury creates a barrier to axonal regeneration overcome by chondroitinase ABC and neurotrophin-3. *Exp Neurol*. 2008; 209:426–445. [PubMed: 17540369]
- Menei P, Montero-Menei C, Whittemore SR, Bunge RP, Bunge MB. Schwann cells genetically modified to secrete human BDNF promote enhanced axonal regrowth across transected adult rat spinal cord. *Eur J Neurosci*. 1998; 10:607–621. [PubMed: 9749723]
- Meyer M, Matsuoka I, Wetmore C, Olson L, Thoenen H. Enhanced synthesis of brain-derived neurotrophic factor in the lesioned peripheral nerve: different mechanisms are responsible for the regulation of BDNF and NGF mRNA. *J Cell Biol*. 1992; 119:45–54. [PubMed: 1527172]
- Michailevski I, Segal-Ruder Y, Rozenbaum M, Medzihradsky KF, Shalem O, Coppola G, Horn-Saban S, Ben-Yaakov K, Dagan SY, Rishal I, Geschwind DH, Pilpel Y, Burlingame AL, Fainzilber M. Signaling to transcription networks in the neuronal retrograde injury response. *Sci Signal*. 2010; 3:ra53. [PubMed: 20628157]
- Moore DL, Blackmore MG, Hu Y, Kaestner KH, Bixby JL, Lemmon VP, Goldberg JL. KLF family members regulate intrinsic axon regeneration ability. *Science*. 2009; 326:298–301. [PubMed: 19815778]

- Mori F, Himes BT, Kowada M, Murray M, Tessler A. Fetal spinal cord transplants rescue some axotomized rubrospinal neurons from retrograde cell death in adult rats. *Exp Neurol*. 1997; 143:45–60. [PubMed: 9000445]
- Muller A, Hauk TG, Leibinger M, Marienfeld R, Fischer D. Exogenous CNTF stimulates axon regeneration of retinal ganglion cells partially via endogenous CNTF. *Mol Cell Neurosci*. 2009; 41:233–246. [PubMed: 19332123]
- Naveilhan P, ElShamy WM, Ernfors P. Differential regulation of mRNAs for GDNF and its receptors Ret and GDNFR alpha after sciatic nerve lesion in the mouse. *Eur J Neurosci*. 1997; 9:1450–1460. [PubMed: 9240402]
- Neet KE, Campenot RB. Receptor binding, internalization, and retrograde transport of neurotrophic factors. *Cellular and molecular life sciences : CMLS*. 2001; 58:1021–1035. [PubMed: 11529495]
- Neumann S, Bradke F, Tessier-Lavigne M, Basbaum AI. Regeneration of sensory axons within the injured spinal cord induced by intraganglionic cAMP elevation. *Neuron*. 2002; 34:885–893. [PubMed: 12086637]
- Neumann S, Woolf CJ. Regeneration of dorsal column fibers into and beyond the lesion site following adult spinal cord injury. *Neuron*. 1999; 23:83–91. [PubMed: 10402195]
- Nielson JL, Strong MK, Steward O. A reassessment of whether cortical motor neurons die following spinal cord injury. *J Comp Neurol*. 2011; 519:2852–2869. [PubMed: 21618218]
- Park KK, Liu K, Hu Y, Smith PD, Wang C, Cai B, Xu B, Connolly L, Kramvis I, Sahin M, He Z. Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway. *Science*. 2008; 322:963–966. [PubMed: 18988856]
- Pfeifer K, Vroemen M, Blesch A, Weidner N. Adult neural progenitor cells provide a permissive guiding substrate for corticospinal axon growth following spinal cord injury. *Eur J Neurosci*. 2004; 20:1695–1704. [PubMed: 15379990]
- Popovich P, McTigue D. Damage control in the nervous system: beware the immune system in spinal cord injury. *Nat Med*. 2009; 15:736–737. [PubMed: 19584863]
- Prang P, Muller R, Eljaouhari A, Heckmann K, Kunz W, Weber T, Faber C, Vroemen M, Bogdahn U, Weidner N. The promotion of oriented axonal regrowth in the injured spinal cord by alginate-based anisotropic capillary hydrogels. *Biomaterials*. 2006; 27:3560–3569. [PubMed: 16500703]
- Qiu J, Cafferty WB, McMahon SB, Thompson SW. Conditioning injury-induced spinal axon regeneration requires signal transducer and activator of transcription 3 activation. *J Neurosci*. 2005; 25:1645–1653. [PubMed: 15716400]
- Qiu J, Cai D, Dai H, McAtee M, Hoffman PN, Bregman BS, Filbin MT. Spinal axon regeneration induced by elevation of cyclic AMP. *Neuron*. 2002; 34:895–903. [PubMed: 12086638]
- Ramon-Cueto A, Cordero MI, Santos-Benito FF, Avila J. Functional recovery of paraplegic rats and motor axon regeneration in their spinal cords by olfactory ensheathing glia. *Neuron*. 2000; 25:425–435. [PubMed: 10719896]
- Ramon-Cueto A, Nieto-Sampedro M. Regeneration into the spinal cord of transected dorsal root axons is promoted by ensheathing glia transplants. *Exp Neurol*. 1994; 127:232–244. [PubMed: 8033963]
- Richardson PM, Issa VM. Peripheral injury enhances central regeneration of primary sensory neurones. *Nature*. 1984; 309:791–793. [PubMed: 6204205]
- Richardson PM, Issa VM, Aguayo AJ. Regeneration of long spinal axons in the rat. *Journal of Neurocytology*. 1984; 13:165–182. [PubMed: 6707710]
- Richardson PM, McGuinness UM, Aguayo AJ. Axons from CNS neurons regenerate into PNS grafts. *Nature*. 1980; 284:264–265. [PubMed: 7360259]
- Richter MW, Roskams AJ. Olfactory ensheathing cell transplantation following spinal cord injury: hype or hope? *Exp Neurol*. 2008; 209:353–367. [PubMed: 17643431]
- Romero MI, Rangappa N, Li L, Lightfoot E, Garry MG, Smith GM. Extensive sprouting of sensory afferents and hyperalgesia induced by conditional expression of nerve growth factor in the adult spinal cord. *J Neurosci*. 2000; 20:4435–4445. [PubMed: 10844012]
- Rosenzweig ES, Courtine G, Jindrich DL, Brock JH, Ferguson AR, Strand SC, Nout YS, Roy RR, Miller DM, Beattie MS, Havton LA, Bresnahan JC, Edgerton VR, Tuszynski MH. Extensive spontaneous plasticity of corticospinal projections after primate spinal cord injury. *Nature neuroscience*. 2010; 13:1505–1510.

- Ruitenbergh MJ, Blits B, Dijkhuizen PA, te Beek ET, Bakker A, van Heerikhuizen JJ, Pool CW, Hermens WT, Boer GJ, Verhaagen J. Adeno-associated viral vector-mediated gene transfer of brain-derived neurotrophic factor reverses atrophy of rubrospinal neurons following both acute and chronic spinal cord injury. *Neurobiol Dis.* 2004; 15:394–406. [PubMed: 15006710]
- Schnell L, Schneider R, Kolbeck R, Barde YA, Schwab ME. Neurotrophin-3 enhances sprouting of corticospinal tract during development and after adult spinal cord lesion. *Nature.* 1994; 367:170–173. [PubMed: 8114912]
- Sendtner M, Stockli KA, Thoenen H. Synthesis and localization of ciliary neurotrophic factor in the sciatic nerve of the adult rat after lesion and during regeneration. *J Cell Biol.* 1992; 118:139–148. [PubMed: 1618901]
- Silver J, Miller JH. Regeneration beyond the glial scar. *Nat Rev Neurosci.* 2004; 5:146–156. [PubMed: 14735117]
- Stam FJ, MacGillavry HD, Armstrong NJ, de Gunst MC, Zhang Y, van Kesteren RE, Smit AB, Verhaagen J. Identification of candidate transcriptional modulators involved in successful regeneration after nerve injury. *Eur J Neurosci.* 2007; 25:3629–3637. [PubMed: 17610582]
- Stokes BT, Reier PJ. Fetal grafts alter chronic behavioral outcome after contusion damage to the adult rat spinal cord. *Exp Neurol.* 1992; 116:1–12. [PubMed: 1559561]
- Stokols S, Sakamoto J, Breckon C, Holt T, Weiss J, Tuszynski MH. Templated agarose scaffolds support linear axonal regeneration. *Tissue Eng.* 2006; 12:2777–2787. [PubMed: 17518647]
- Straley KS, Foo CW, Heilshorn SC. Biomaterial design strategies for the treatment of spinal cord injuries. *Journal of neurotrauma.* 2010; 27:1–19. [PubMed: 19698073]
- Sun F, Park KK, Belin S, Wang D, Lu T, Chen G, Zhang K, Yeung C, Feng G, Yankner BA, He Z. Sustained axon regeneration induced by co-deletion of PTEN and SOCS3. *Nature.* 2011 in press.
- Tang XQ, Tanelian DL, Smith GM. Semaphorin3A inhibits nerve growth factor-induced sprouting of nociceptive afferents in adult rat spinal cord. *J Neurosci.* 2004; 24:819–827. [PubMed: 14749426]
- Taylor L, Jones L, Tuszynski MH, Blesch A. Neurotrophin-3 gradients established by lentiviral gene delivery promote short-distance axonal bridging beyond cellular grafts in the injured spinal cord. *J Neurosci.* 2006; 26:9713–9721. [PubMed: 16988042]
- Tessler A. Intraspinal transplants. *Ann Neurol.* 1991; 29:115–123. [PubMed: 2012382]
- Tessler A, Fischer I, Giszter S, Himes BT, Miya D, Mori F, Murray M. Embryonic spinal cord transplants enhance locomotor performance in spinalized newborn rats. *Adv Neurol.* 1997; 72:291–303. [PubMed: 8993706]
- Tetzlaff W, Okon EB, Karimi-Abdolrezaee S, Hill CE, Sparling JS, Plemel JR, Plunet WT, Tsai EC, Baptiste D, Smithson LJ, Kawaja MD, Fehlings MG, Kwon BK. A systematic review of cellular transplantation therapies for spinal cord injury. *Journal of neurotrauma.* 2011; 28:1611–1682. [PubMed: 20146557]
- Tom VJ, Houle JD. Intraspinal microinjection of chondroitinase ABC following injury promotes axonal regeneration out of a peripheral nerve graft bridge. *Exp Neurol.* 2008; 211:315–319. [PubMed: 18353313]
- Tom VJ, Sandrow-Feinberg HR, Miller K, Santi L, Connors T, Lemay MA, Houle JD. Combining peripheral nerve grafts and chondroitinase promotes functional axonal regeneration in the chronically injured spinal cord. *J Neurosci.* 2009; 29:14881–14890. [PubMed: 19940184]
- Trupp M, Belluardo N, Funakoshi H, Ibanez CF. Complementary and overlapping expression of glial cell line-derived neurotrophic factor (GDNF), c-ret proto-oncogene, and GDNF receptor-alpha indicates multiple mechanisms of trophic actions in the adult rat CNS. *J Neurosci.* 1997; 17:3554–3567. [PubMed: 9133379]
- Tuszynski MH, Gabriel K, Gage FH, Suhr S, Meyer S, Rosetti A. Nerve growth factor delivery by gene transfer induces differential outgrowth of sensory, motor, and noradrenergic neurites after adult spinal cord injury. *Exp Neurol.* 1996; 137:157–173. [PubMed: 8566208]
- Tuszynski MH, Weidner N, McCormack M, Miller I, Powell H, Conner J. Grafts of genetically modified Schwann cells to the spinal cord: survival, axon growth, and myelination. *Cell Transplant.* 1998; 7:187–196. [PubMed: 9588600]

- Vavrek R, Girgis J, Tetzlaff W, Hiebert GW, Fouad K. BDNF promotes connections of corticospinal neurons onto spared descending interneurons in spinal cord injured rats. *Brain*. 2006; 129:1534–1545. [PubMed: 16632552]
- Vaynman S, Gomez-Pinilla F. License to run: exercise impacts functional plasticity in the intact and injured central nervous system by using neurotrophins. *Neurorehabil Neural Repair*. 2005; 19:283–295. [PubMed: 16263961]
- Wang D, Ichiyama RM, Zhao R, Andrews MR, Fawcett JW. Chondroitinase combined with rehabilitation promotes recovery of forelimb function in rats with chronic spinal cord injury. *J Neurosci*. 2011; 31:9332–9344. [PubMed: 21697383]
- Weidner N, Blesch A, Grill RJ, Tuszynski MH. Nerve growth factor-hypersecreting Schwann cell grafts augment and guide spinal cord axonal growth and remyelinate central nervous system axons in a phenotypically appropriate manner that correlates with expression of L1. *J Comp Neurol*. 1999; 413:495–506. [PubMed: 10495438]
- Weidner N, Ner A, Salimi N, Tuszynski MH. Spontaneous corticospinal axonal plasticity and functional recovery after adult central nervous system injury. *Proc Natl Acad Sci U S A*. 2001; 98:3513–3518. [PubMed: 11248109]
- Wong LF, Yip PK, Battaglia A, Grist J, Corcoran J, Maden M, Azzouz M, Kingsman SM, Kingsman AJ, Mazarakis ND, McMahon SB. Retinoic acid receptor beta2 promotes functional regeneration of sensory axons in the spinal cord. *Nature neuroscience*. 2006; 9:243–250.
- Wu D, Zhang Y, Bo X, Huang W, Xiao F, Zhang X, Miao T, Magoulas C, Subang MC, Richardson PM. Actions of neurotrophic cytokines and cyclic AMP in regenerative conditioning of rat primary sensory neurons. *Exp Neurol*. 2006; 204:66–76. [PubMed: 17112514]
- Xiao HS, Huang QH, Zhang FX, Bao L, Lu YJ, Guo C, Yang L, Huang WJ, Fu G, Xu SH, Cheng XP, Yan Q, Zhu ZD, Zhang X, Chen Z, Han ZG, Zhang X. Identification of gene expression profile of dorsal root ganglion in the rat peripheral axotomy model of neuropathic pain. *Proc Natl Acad Sci U S A*. 2002; 99:8360–8365. [PubMed: 12060780]
- Xie F, Zheng B. White matter inhibitors in CNS axon regeneration failure. *Exp Neurol*. 2008; 209:302–312. [PubMed: 17706966]
- Xu XM, Chen A, Guenard V, Kleitman N, Bunge MB. Bridging Schwann cell transplants promote axonal regeneration from both the rostral and caudal stumps of transected adult rat spinal cord. *J Neurocytol*. 1997; 26:1–16. [PubMed: 9154524]
- Xu XM, Guenard V, Kleitman N, Aebischer P, Bunge MB. A combination of BDNF and NT-3 promotes supraspinal axonal regeneration into Schwann cell grafts in adult rat thoracic spinal cord. *Exp Neurol*. 1995; 134:261–272. [PubMed: 7556546]
- Xu XM, Guenard V, Kleitman N, Bunge MB. Axonal regeneration into Schwann cell-seeded guidance channels grafted into transected adult rat spinal cord. *J Comp Neurol*. 1994; 351:145–160. [PubMed: 7896937]
- Yick LW, So KF, Cheung PT, Wu WT. Lithium chloride reinforces the regeneration-promoting effect of chondroitinase ABC on rubrospinal neurons after spinal cord injury. *Journal of neurotrauma*. 2004; 21:932–943. [PubMed: 15307905]
- Ying Z, Roy RR, Edgerton VR, Gomez-Pinilla F. Exercise restores levels of neurotrophins and synaptic plasticity following spinal cord injury. *Exp Neurol*. 2005; 193:411–419. [PubMed: 15869943]
- Yip PK, Wong LF, Pattinson D, Battaglia A, Grist J, Bradbury EJ, Maden M, McMahon SB, Mazarakis ND. Lentiviral vector expressing retinoic acid receptor beta2 promotes recovery of function after corticospinal tract injury in the adult rat spinal cord. *Hum Mol Genet*. 2006; 15:3107–3118. [PubMed: 16984961]
- Yip PK, Wong LF, Sears TA, Yanez-Munoz RJ, McMahon SB. Cortical overexpression of neuronal calcium sensor-1 induces functional plasticity in spinal cord following unilateral pyramidal tract injury in rat. *PLoS Biol*. 2010; 8:e1000399. [PubMed: 20585375]
- Z'Graggen WJ, Fouad K, Raineteau O, Metz GA, Schwab ME, Kartje GL. Compensatory sprouting and impulse rerouting after unilateral pyramidal tract lesion in neonatal rats. *J Neurosci*. 2000; 20:6561–6569. [PubMed: 10964961]

- Zhang JY, Luo XG, Xian CJ, Liu ZH, Zhou XF. Endogenous BDNF is required for myelination and regeneration of injured sciatic nerve in rodents. *Eur J Neurosci*. 2000; 12:4171–4180. [PubMed: 11122329]
- Zhong J, Dietzel ID, Wahle P, Kopf M, Heumann R. Sensory impairments and delayed regeneration of sensory axons in interleukin-6-deficient mice. *J Neurosci*. 1999; 19:4305–4313. [PubMed: 10341234]
- Zhou FQ, Snider WD. Intracellular control of developmental and regenerative axon growth. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2006; 361:1575–1592.
- Zhou L, Baumgartner BJ, Hill-Felberg SJ, McGowen LR, Shine HD. Neurotrophin-3 expressed in situ induces axonal plasticity in the adult injured spinal cord. *J Neurosci*. 2003; 23:1424–1431. [PubMed: 12598631]
- Ziembra KS, Chaudhry N, Rabchevsky AG, Jin Y, Smith GM. Targeting axon growth from neuronal transplants along preformed guidance pathways in the adult CNS. *J Neurosci*. 2008; 28:340–348. [PubMed: 18184776]

Table 1
Selected combinatorial therapies for axonal regeneration after experimental spinal cord injury.

Axonal population	Cell soma stimulation	Substrate at the lesion site	Neutralization of inhibition	Growth factor	References
Serotonergic, noradrenergic, corticospinal axons	-	E14 Fetal graft	-	BDNF +/- NT-3, NT-4 protein NT-3/BDNF infusion	(Bregman et al. 1997) (Cummings et al. 2001)
Serotonergic propriospinal, vestibulospinal, reticulospinal, raphespinal, and other bulbospinal axons	-	Schwann cells + olfactory ensheathing cells	ChABC	-	(Fouad et al. 2005; Vavrek et al. 2007)
Serotonergic axons	cAMP + rolipram	Schwann cells	cAMP + Rolipram	-	(Pearse et al. 2004) but (Sharp et al. 2010)
Dorsal column sensory axons	cAMP	Bone marrow stromal cells	-	NT-3 protein	(Lu et al. 2004)
Dorsal column sensory axons	Conditioning lesion	Bone marrow stromal cells	-	NT-3 lentivirus	(Alto et al. 2009; Blesch et al. 2012; Kadoya et al. 2009)
Reticulospinal, raphespinal, rubrospinal, propriospinal coeruleospinal, axons	-	Peripheral nerve graft	ChABC	-	(Houle et al. 2006)
Descending axons	-	Peripheral nerve graft	ChABC	GDNF infusion	(Tom et al. 2009)
Corticospinal, serotonergic axons	-	Neural stem cells	ChABC	EGF+ bFGF+ PDGF-AA infusion	(Karimi-Abdolrezaee et al. 2010)
Serotonergic axons	Rolipram (cAMP)	Fetal graft	Rolipram	-	(Nikulina et al. 2004)
Transplanted DRG neurons	-	-	ChABC	NT-3 lentivirus	(Massey et al. 2008)
Serotonergic axons	-	-	Thermo- stabilized ChABC/ trehalose	NT-3 protein	(Lee et al. 2010)
Rubrospinal axons	LiCl-induced suppression of GSK-3 β /upregulation of Bcl-2	-	ChABC	-	(Yick et al. 2004)
Serotonergic and descending bulbospinal axons	-	Peripheral nerve graft	ChABC	-	(Allain et al. 2011)