Published in final edited form as:

Vaccine. 2010 December 10; 29(1): 51–57. doi:10.1016/j.vaccine.2010.10.022.

Higher human CD4 T cell response to novel *Mycobacterium tuberculosis* latency associated antigens Rv2660 and Rv2659 in latent infection compared with tuberculosis disease

Lerisa Govender¹, Brian Abel¹, E. Jane Hughes¹, Thomas J. Scriba¹, Benjamin M. N. Kagina¹, Marwou de Kock¹, Gerhard Walzl², Gillian Black², Ida Rosenkrands³, Gregory D Hussey¹, Hassan Mahomed¹, Peter Andersen³, and Willem A. Hanekom^{1,*}

¹ South African Tuberculosis Vaccine Initiative, Institute of Infectious Diseases and Molecular Medicine and School of Child and Adolescent Health, University of Cape Town, South Africa

² Department of Biochemistry, University of Stellenbosch, Tygerberg, South Africa

³ Statens Serum Institut, Copenhagen, Denmark

Abstract

One third of the world's population is infected with *Mycobacterium tuberculosis* (*M.tb*). A vaccine that would prevent progression to TB disease will have a dramatic impact on the global TB burden. We propose that antigens of *M.tb* that are preferentially expressed during latent infection will be excellent candidates for post-exposure vaccination. We therefore assessed human T cell recognition of two such antigens, Rv2660 and Rv2659. Expression of these was shown to be associated with non-replicating persistence *in vitro*. After six days incubation of PBMC from persons with latent tuberculosis infection (LTBI) and tuberculosis (TB) disease, Rv2660 and Rv2659 induced IFN- γ production in a greater proportion of persons with LTBI, compared with TB diseased patients. Persons with LTBI also had increased numbers of viable T cells, and greater specific CD4⁺ T cell proliferation and cytokine expression capacity. Persons with LTBI preferentially recognize Rv2659 and Rv2660, compared with patients with TB disease. These results suggest promise of these antigens for incorporation into post-exposure TB vaccines.

Keywords

Mycobacterium tuberculosis; LTBI; TB disease; latency antigens; post-infection vaccine

1. Introduction

The success of *Mycobacterium tuberculosis* (*M.tb*) as a human pathogen may be ascribed to its ability to persist for long periods in asymptomatic people, a state known as latency [1]. Healthy latently infected individuals have approximately 10% lifetime risk of developing

*Correspondence should be addressed to Willem A. Hanekom, University of Cape Town Health Sciences, Anzio Road, Observatory, 7925, South Africa, Tel. +27 21 406 6080, Fax. +27 21 406 6693, Willem.Hanekom@uct.ac.za.

Conflict of interest

P.A. is the inventor of TB vaccines that incorporates the RV2659/60 antigens. All rights have been assigned to the Statens Serum Institut. The rest of the authors declare no financial or commercial conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

active TB disease [2]. This risk increases to approximately 10% per year in HIV-infected persons not on antiretroviral therapy [3]. Approximately one third of the world's population is infected with *M.tb*, 9 million develop TB disease and 1.8 million die annually [4,5]. To control this pandemic, an effective vaccine is urgently needed; however, this requires extensive insight into the mechanisms underlying protective immunity against *M.tb*.

The low bacterial burden associated with a latent infection has proven to be a major obstacle in characterizing the mechanisms by which *M.tb* persists and reactivates in the host [2]. In latent granulomatous lesions, *M.tb* is successfully contained and has to adapt to a hypoxic and nutritionally compromised environment [6]. It has recently been hypothesized that this immune pressure drives *M.tb* into a different metabolic state, compared with actively replicating organisms in lesions that characterize early *M.tb* infection or active disease [1]. This is supported by findings from *in vitro* studies which mimic the granuloma environment, where *M.tb* was shown to upregulate sets of genes that are distinct from those upregulated in actively replicating organisms; e.g., in nutrient deficient medium, a particular set of genes that were termed the "starvation stimulon" were upregulated [7]. We focused on 2 proteins of the starvation stimulon, Rv2660 and Rv2659, and hypothesized that these antigens will be preferentially recognized by persons with latent *M.tb* infection (LTBI), compared with persons with TB disease. Rv2660 and Rv2659 belong to the RD11 region encoded in *M.tb* [8], however the function of these proteins has not been determined [7].

The rationale for these studies is that latency-associated antigens might in the future be incorporated into novel post-exposure TB vaccine candidates, which could protect infected individuals from developing TB disease [1].

2. Materials and Method

2.1 Study participants

Participants were recruited from the Western Cape region of South Africa. Persons with LTBI were included if they had a positive QuantiFERON®-TB Gold In-Tube (QFT™) test, or a positive ELISPOT test for ESAT-6/CFP-10 (>17 spots/10⁶ PBMC after background subtraction), or a positive Mantoux skin test (induration >10mm), for greater than 1 year at time of recruitment. Any clinical signs or symptoms suggestive of TB disease, a history of TB treatment, HIV infection, any other acute or chronic medical conditions, receipt of immunosuppressive medication, or a hemoglobin level <9g/dL resulted in exclusion.

Patients with TB disease were also included, if they had 2 positive sputum smears and/or a positive culture, with clinical features consistent with TB disease. Patients with extra-pulmonary TB disease were excluded, as were patients with exclusion criteria mentioned for LTBI above. Informed consent was obtained from all participants prior to study participation. The study was approved by the research ethics committees of the University of Cape Town and the University of Stellenbosch.

2.2 Whole blood incubation

Whole blood was collected in heparinised syringes. Blood was diluted in RPMI-1640 medium (Sigma) containing 1% L-glutamine (Sigma, final dilution 1:10), and incubated with an ESAT-6 peptide pool (17mers, overlapping by 10, final concentration 1 µg/mL/peptide), CFP-10 peptide pool (17mers, overlapping by 10, final concentration 1 µg/mL/peptide), TB10.4 protein (10 µg/mL), Rv2660 peptide pool (20mers, overlapping by 10, final concentration 1 µg/mL/peptide), or Rv2659 peptide pool (20mers, overlapping by 10, final concentration 1 µg/mL/peptide), in 200 µL volumes in 96-well U-bottom plates (Nunc, Cat No 163320), for 6 days at 37°C in 5% CO₂. All antigens were obtained from the Statens Serum Institut, Denmark. Blood incubated without antigen and blood incubated with

phytohaemagglutinin (PHA, 5µg/mL) were used as negative and positive controls, respectively. After incubation, 150µL culture supernatant was removed and stored at -80°C until further analysis.

2.3 PBMC incubation

PBMC were isolated by density gradient centrifugation and cryopreserved in 10% DMSO (Sigma-Aldrich) in heat-inactivated fetal calf serum (Adcock Ingram). PBMC were thawed in 12.5% AB serum/RPMI containing 2.5µg/mL DNase (Sigma-Aldrich) and stained with 10µg/mL of CellTrace Oregon Green 488 (Molecular Probes, Invitrogen) per 1×10^7 cells. The PBMC were resuspended in 12.5% AB serum in RPMI, and 200,000 cells in 200µL medium were rested overnight at 37°C, 5% CO₂ in a 96 well plate. Duplicate wells containing cells (i.e. 400,000 PBMC per condition) were stimulated with BCG (1331 Danish BCG) at an MOI of 0.01, PPD (0.5µg/mL), ESAT-6/CFP10 peptide pool (2µg/mL/peptide), TB10.4 peptide pool (15mers, overlapping by 10, final concentration 1µg/mL/peptide), Rv2660 peptide pool (20mers, overlapping by 10, final concentration 1µg/mL/peptide), Rv2659 peptide pool (20mers, overlapping by 10, final concentration 1µg/mL/peptide), SEB (0.05µg/mL, positive control, Sigma-Aldrich) or no antigen (negative control), for 6 days at 37°C with 5% CO₂. The optimum concentrations of Rv2660 and Rv2659 peptide pools were determined in preliminary studies to be 1µg/mL/peptide. All antigens were obtained from the Statens Serum Institut, Denmark. After incubation, plasma was collected and stored at -80°C until further use. PBMC were then incubated with phorbol 12-myristate 13-acetate (PMA, 50ng/mL) and ionomycin (250ng/mL), to induce cytokine production, and Brefeldin A (10µg/mL) was added while incubation continued for a further 5 hours. PBMC were harvested, stained for viability with LIVE/DEAD Fixable Violet Dead Cell Stain (ViViD (Invitrogen), fixed with FACS Lysing Solution (BD Biosciences), and cryopreserved until later analysis.

2.4 Flow cytometric analysis of processed PBMC

Fixed, cryopreserved PBMC were thawed and permeabilised with Perm/Wash Solution (BD Biosciences) for 10 minutes. PBMC were washed with PBS and stained with the following antibodies: anti-CD3 Qdot605 (clone UCHT1) anti-CD8 PerCPCy5.5 (SK-1), anti-IFN-γ AlexaFluor700 (B27), anti-IL2 APC (MQ1-17H12) and anti-TNF-α Cy7.PE (MAb11) (all from BD Biosciences) for 1 hour at 4°C. After washing, cells were acquired on a LSRII flow cytometer (BD Biosciences), configured with 3 lasers and 10 detectors, using the FACS Diva 6.1 software. Compensation settings were defined using anti-mouse kappa Comp Beads (BD Biosciences) stained with each fluorochrome-conjugated antibody. Because the cellular dyes Oregon Green and ViViD do not bind Comp Beads, compensation of these dyes was done using FITC and Pacific Blue conjugated antibody-bound Comp Beads, respectively. Data was analyzed using Flowjo 8.8.4 (Treestar). Frequencies of proliferating and cytokine-expressing CD4 or CD8 T cells were determined after exclusion of dead cells (ViViD^{high} events). Boolean gating was applied to generate combinations of cytokine expressing CD4⁺ and CD8⁺ T cell subsets.

2.5 IFN-γ measurement in supernatants

IFN-γ was quantified in culture supernatants by sandwich ELISA. The ELISA was performed as previously described [9], with the following modifications: OPD Fast (Sigma) was utilized as substrate, and plates were read at 405 nm on a Versamax ELISA plate reader using Softmax Pro software Version 4.7.1. The limit of detection of the ELISA was determined to be 15.6pg/mL. A 1:2 dilution of supernatants was used, which was found optimal in preliminary experiments.

2.6 Data analysis

For soluble IFN- γ analysis, antigen-specific cytokine levels were determined by subtracting levels from unstimulated conditions. A positive IFN- γ response was defined as 31.2pg/mL, after background subtraction.

For the flow cytometric assay, proliferation and cytokine expression levels from the unstimulated condition (restimulated with PMA and Ionomycin) were subtracted from levels obtained after antigen-specific stimulation. Individuals were excluded from analysis for samples with <100 viable CD3⁺ T cells, or if the positive control yielded a negative result. The Mann-Whitney U test was used to assess differences between persons in the two groups. The Fisher's exact test was used to assess differences between proportions of responders. P-values <0.05 were considered to be significant.

3. Results

3.1 Participant Characteristics

We examined immune recognition of Rv2660 and Rv2659, measured by IFN- γ expression in whole blood in a pilot study on 21 adults with LTBI. The median age was 32 years (interquartile range: 28–40); 90% were female. All participants had been vaccinated with BCG at birth, and all were from a mixed ethnic background. Following the pilot study, PBMC proliferation and cytokine production were studied in 25 additional persons with LTBI, as well as 25 patients with TB disease. The median ages of the LTBI and TB diseased groups were 19 years (18.5–31.5) and 31 years (23.5–42), respectively ($p=0.003$); 64% of individuals with LTBI were female, compared with 24% in the TB group ($p=0.044$). Due to the difference in age and gender we compared the outcomes with respect to these co-variables. Only polyfunctional CD4⁺ T cells, expressing IFN- γ , TNF- α and IL-2 were different in males and females. Participants excluded from analysis, with criteria for exclusion, are shown in Supplementary Figure 1. Twenty-eight % of TB patients were excluded due either culture contamination or <100 viable CD3⁺ T cells; 8% of LTBI participants were excluded due to a negative SEB result.

3.2 Rv2660 and Rv2659 are recognized by T cells of persons with established latency

To assess whether Rv2660 and Rv2659 are recognized during established human latent *M.tb* infection, diluted whole blood was incubated with the antigens for 6 days, and IFN- γ measured in the supernatant. Rv2660 and Rv2659 induced detectable IFN- γ production in a high proportion of persons with LTBI (Figure 1A). The immunodominant control antigen, TB10.4, present in both BCG and in *M.tb*, and the *M.tb*-specific antigens ESAT-6 and CFP-10, were recognized in substantially greater proportions of individuals (Figure 1A). The positive control, SEB induced a response in all individuals (data not shown).

To compare immune recognition of Rv2660 and Rv2659 during established latent infection and TB disease, we incubated PBMC with these antigens for 6 days, followed by measurement of IFN- γ in supernatants. A greater proportion of persons with LTBI recognized Rv2660 and Rv2659, compared with patients with TB disease (Figure 1B). A similar proportion responded to BCG; however, more persons with LTBI responded to ESAT-6/CFP-10, compared with patients with TB disease. Further, levels of IFN- γ in response to all antigens were higher in the LTBI group, compared with the TB diseased group (Figure 1C).

3.3 Decreased viability of CD3⁺ T cells during TB disease

We hypothesized that lower Rv2660 and Rv2659 recognition during TB disease is due to lesser capacity of peripheral blood T cells of TB patients to survive, compared with cells

from persons with LTBI. After 6 days of incubation of PBMC in conditions as aforementioned, cell viability was assessed with flow cytometry following staining with the Violet amine reactive dye, ViViD (Figure 2A). The LTBI group had significantly higher frequencies of viable CD3⁺ T cells, for all conditions tested (Figure 2B). The median number of viable CD3⁺ T cells was 15,771 (IQR, 4,912–99,654; Supplementary Figure 2).

3.4 Increased frequency of proliferating Rv2660- and Rv2659-specific CD4⁺ T cells during established latency

We further characterized viable CD4⁺ T cell responses to Rv2660 and Rv2659 during established latent infection and TB disease by assessing antigen-specific proliferation with a dye dilution assay, and the cytokine producing capacity of proliferating cells (Figure 3A and B, respectively). Frequencies of proliferating antigen-specific CD4⁺ T cells were higher in LTBI, compared with TB disease, for all antigens tested but the difference was clearly more pronounced for BCG and the ESAT6/CFP10 antigens (Figures 4C–F, left panels).

The cytokine expression profile of proliferating CD4⁺ T cells was assessed by adding a polyclonal stimulant for the last 5 hours of incubation. Upon stimulation with Rv2660 and Rv2659 antigens, frequencies of proliferating CD4⁺ T cells were higher in the LTBI group for all cytokine producing subsets, except for the total IL-2⁺ population (Figures 4C–D, right panels, respectively). In addition, Rv2660 and Rv2659 induced more polyfunctional (co-expression of IFN- γ ⁺, TNF- α ⁺ and IL-2⁺ in the same cell) cytokine expression in persons with LTBI, compared with TB patients. In response to BCG- and to ESAT-6/CFP-10, specific (proliferating) CD4⁺ T cells were also more likely to be polyfunctional in persons with LTBI, compared with patients with TB disease (Figure 4E–F); total IFN- γ ⁺, total TNF- α ⁺ and total IL-2 were also higher in persons with latent infection (Figures 4E–F, right panels).

CD8⁺ T cell proliferation and cytokine expression in response to Rv2660 and Rv2659 was very low, and no difference was demonstrated between the 2 groups (Figures 4A–B). However, persons with LTBI had greater frequencies of CD8⁺ T cells that proliferated in response to ESAT-6/CFP10 and BCG, compared with frequencies in patients with TB disease. Proliferating cells were also more likely to be polyfunctional in persons with LTBI (Figures 4C–D). The positive control, SEB induced a response in all individuals (data not shown).

4. Discussion

Mathematical modeling has indicated that an effective post-infection vaccine that targets latently infected individuals, would have a dramatic impact on the TB epidemic [10]. Our study suggests that Rv2660 and Rv2659 are good antigenic candidates for inclusion in such a vaccine; we show that T cells of persons with LTBI preferentially recognize these two latency-associated antigens, compared with T cells from patients with TB disease. This is the first clinical report of human immune recognition of the starvation stimulon gene products, Rv2660 and Rv2659.

Interestingly, the same pattern of preferential recognition in LTBI, compared with TB disease, was observed for the crude mycobacterial antigens BCG, PPD, and the immunodominant/early antigens ESAT-6/CFP-10 and TB10.4 (data for PPD and TB10.4 not shown). In agreement with our findings, Schuck *et al.* recently reported higher T cell responses upon stimulation of PBMC with PPD and ESAT-6/CFP10, and with other latency antigens, such as Rv1733 and Rv0140, in individuals with LTBI, compared with TB patients [11]. This decreased T cell response in peripheral blood may result from migration of specific T cells to the lung during active disease [[12]], or, with respect to Rv2659 and

Rv2660, may relate to a preferential expression of these two antigens during latency. Alternatively, aberrant immune regulation during disease, mediated by regulatory T cells [11], anti-inflammatory cytokines such as IL-10 and TGF- β [13,14], or Th2 cytokines like IL-4 and IL-13 may be responsible for the differences noted [15]. Our observation that T cells from patients with TB disease were strikingly less likely to survive in a 6-day culture, compared with T cells from persons with LTBI, suggest that T cell exhaustion may also be responsible for lower responses in the diseased group. In chronic infections such as HIV and CMV, a persistently high antigenic load drives specific T cell exhaustion and dysfunction [16,17]. These cells upregulate markers like PD-1, and are more prone to apoptosis [18]. We postulate that this also occurs during active TB disease. We have recently shown that PD-1 expression is increased on *M.tb*-specific CD4⁺ T cells in TB diseased patients, compared to persons with LTBI (C. Day, unpublished observations).

Recent advances in imaging have led to the observation that granulomatous lesions in persons with LTBI reflect a spectrum of differential ability to support or suppress the persistence of viable bacteria [19]. This spectrum is supported by our data of similar preferential recognition of latency-associated and immunodominant antigens in persons with LTBI, compared with active TB. A post-exposure vaccine would thus ideally contain both immunodominant and latency-associated antigens, respectively, to target replicating or persisting bacteria in progressing lesions and dormant bacteria in successful lesions.

LTBI individuals had elevated frequencies of Th1 cytokine expression among proliferating CD4⁺ T cells, compared with those in TB patients. The pattern of cytokine production also differed: specific T cells from persons with LTBI were more likely to be “polyfunctional”, i.e., able to co-express IFN- γ , TNF- α and IL-2. This characteristic has been proposed to be associated with long-term memory and efficient protection against intracellular infections like *M.tb* [20–23]. We therefore propose that robust polyfunctional responses to latency and immunodominant antigens may prevent LTBI individuals from progressing to TB disease. The lesser cytokine production in TB disease may also be due to T cell exhaustion, leading to a loss of function in particular for antigens like ESAT6/CFP10 that predominate during the early stages of infection, and where continued exposure may lead to exhaustion [16]. Therefore, because latency associated antigens would not be expressed in early stages of infection, T cells directed to these antigens may not be subjected to regulation by the established effector and/or regulatory T cells primed in the earlier stages of infection.

CD8⁺ T cell responses to immunodominant antigens could readily be detected, and were higher in LTBI compared with TB disease. However, CD8⁺ T cell responses to Rv2659 and Rv2660 were very low, which may be the reason why no difference could be demonstrated between the participant groups. We cannot exclude that differences in antigen presentation between CD4⁺ and CD8⁺ T cells could have affected these results.

A limitation of our study was the different age and gender ratios in the LTBI and TB groups. Ideally, participants in such groups should be matched for these co-variates. However, given the fact that only polyfunctional CD4⁺ T cells and no other outcomes were different in males and females, the unequal gender ratio was unlikely to have markedly confounded our results.

The prophylactic TB subunit vaccines currently under development are almost exclusively based on early or immunodominant antigens secreted by replicating *M.tb*, which are recognized in the first stage of infection, and are designed as BCG boosters to prevent primary TB disease [24]. Addition of latency associated antigens to well-established prophylactic vaccines (i.e., the vaccine candidates under clinical testing today) has been suggested as the basis for a future generation of multi-stage TB vaccines with activity against all stages of infection [1]. Our finding of immune recognition of these novel antigens

in persons with LTBI has contributed to the development of a new subunit vaccine, H56, by the Statens Serum Institut. This vaccine is based upon the H1 vaccine currently in clinical trials, which contains Ag85B and ESAT-6, and the Rv2660 antigen, and will be the first to enter clinical trials as a post-infection vaccine candidate [25]. We hypothesize that inducing robust T cell responses to latency and immunodominant antigens may enhance immunological control of latent TB, and prevent reactivation. Much more research is needed to determine whether candidate latency-, starvation-, or resuscitation-associated antigens other than Rv2659 or Rv2660 would be needed for inclusion in an optimal post-infection vaccine.

Ultimately, a combined strategy that includes primary vaccination with a live attenuated vaccine followed by a multistage subunit boost vaccine; may have maximum impact on the TB epidemic.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Acknowledgements/Funding

The work described in this report was supported by the Bill & Melinda Gates Foundation through a Grand Challenge grant: Preclinical and clinical evaluation of a post-exposure TB vaccine.

W. A. H. is supported by the NIH (RO1-AI065653 and NO1-AI70022).

L.G. is a recipient of The National Research (NRF) Prestigious and Equity Scholarship (SFH2007081400194) and the South African Tuberculosis and AIDS Training Award (SATBAT) (1U2RTW007370).

Abbreviations

<i>M.tb</i>	<i>Mycobacterium tuberculosis</i>
LTBI	latent tuberculosis infection
OG	Oregon green

5. References

1. Andersen P. Vaccine strategies against latent tuberculosis infection. Trends Microbiol. 2007 Jan; 15(1):7–13. [PubMed: 17141504]
2. Flynn JL, Chan J. Immunology of tuberculosis. Annu Rev Immunol. 2001; 19:93–129. [PubMed: 11244032]
3. Corbett EL, Watt CJ, Walker N, Maher D, Williams BG, Raviglione MC, et al. The growing burden of tuberculosis: global trends and interactions with the HIV epidemic. Arch Intern Med. 2003 May 12; 163(9):1009–21. [PubMed: 12742798]
4. Organisation WH. Global Tuberculosis Control-surveillance, planning, financing. World Health Organisation; Geneva, Switzerland: 2008.
5. Baumann S, Nasser Eddine A, Kaufmann SH. Progress in tuberculosis vaccine development. Curr Opin Immunol. 2006 Aug; 18(4):438–48. [PubMed: 16777396]
6. Sherman DR, Voskuil M, Schnappinger D, Liao R, Harrell MI, Schoolnik GK. Regulation of the Mycobacterium tuberculosis hypoxic response gene encoding alpha -crystallin. Proc Natl Acad Sci U S A. 2001 Jun 19; 98(13):7534–9. [PubMed: 11416222]

7. Betts JC, Lukey PT, Robb LC, McAdam RA, Duncan K. Evaluation of a nutrient starvation model of *Mycobacterium tuberculosis* persistence by gene and protein expression profiling. *Mol Microbiol*. 2002 Feb; 43(3):717–31. [PubMed: 11929527]
8. Gordon SV, Eiglmeier K, Garnier T, Brosch R, Parkhill J, Barrell B, et al. Genomics of *Mycobacterium bovis*. *Tuberculosis (Edinb)*. 2001; 81(1–2):157–63. [PubMed: 11463237]
9. Black GF, Fine PEM, Warndorff DK, Floyd S, Weir RE, Blackwell JM, et al. Relationship between IFN-gamma and skin test responsiveness to *Mycobacterium tuberculosis* PPD in healthy, non-BCG-vaccinated young adults in Northern Malawi. *Int J Tuberc Lung Dis*. 2001 Jul; 5(7):664–72. [PubMed: 11467373]
10. Abu-Raddad LJ, Sabatelli L, Achterberg JT, Sugimoto JD, Longini IM Jr, Dye C, et al. Epidemiological benefits of more-effective tuberculosis vaccines, drugs, and diagnostics. *Proc Natl Acad Sci U S A*. 2009 Aug 18; 106(33):13980–5. [PubMed: 19666590]
11. Schuck SD, Mueller H, Kunitz F, Neher A, Hoffmann H, Franken KL, et al. Identification of T-cell antigens specific for latent mycobacterium tuberculosis infection. *PLoS One*. 2009; 4(5):e5590. [PubMed: 19440342]
12. Tully G, Kortsik C, Hohn H, Zehbe I, Hitzler WE, Neukirch C, et al. Highly focused T cell responses in latent human pulmonary *Mycobacterium tuberculosis* infection. *J Immunol*. 2005 Feb 15; 174(4):2174–84. [PubMed: 15699149]
13. Fiorenza G, Rateni L, Farroni MA, Bogue C, Dlugovitzky DG. TNF-alpha, TGF-beta and NO relationship in sera from tuberculosis (TB) patients of different severity. *Immunol Lett*. 2005 Apr 15; 98(1):45–8. [PubMed: 15790507]
14. Guyot-Revol V, Innes JA, Hackforth S, Hinks T, Lalvani A. Regulatory T cells are expanded in blood and disease sites in patients with tuberculosis. *Am J Respir Crit Care Med*. 2006 Apr 1; 173(7):803–10. [PubMed: 16339919]
15. Ordway DJ, Martins MS, Costa LM, Freire MS, Arroz MJ, Dockrell HM, et al. [Increased IL-4 production in response to virulent *Mycobacterium tuberculosis* in tuberculosis patients with advanced disease]. *Acta Med Port*. 2005 Jan–Feb; 18(1):27–36. [PubMed: 16202332]
16. Barber DL, Wherry EJ, Masopust D, Zhu B, Allison JP, Sharpe AH, et al. Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature*. 2006 Feb 9; 439(7077):682–7. [PubMed: 16382236]
17. Day CL, Kaufmann DE, Kiepiela P, Brown JA, Moodley ES, Reddy S, et al. PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature*. 2006 Sep 21; 443(7109):350–4. [PubMed: 16921384]
18. van Grevenynghe J, Procopio FA, He Z, Chomont N, Riou C, Zhang Y, et al. Transcription factor FOXO3a controls the persistence of memory CD4(+) T cells during HIV infection. *Nat Med*. 2008 Mar; 14(3):266–74. [PubMed: 18311149]
19. Barry CE 3rd, Boshoff HI, Dartois V, Dick T, Ehrh S, Flynn J, et al. The spectrum of latent tuberculosis: rethinking the biology and intervention strategies. *Nat Rev Microbiol*. 2009 Dec; 7(12):845–55. [PubMed: 19855401]
20. Forbes EK, Sander C, Ronan EO, McShane H, Hill AV, Beverley PC, et al. Multifunctional, high-level cytokine-producing Th1 cells in the lung, but not spleen, correlate with protection against *Mycobacterium tuberculosis* aerosol challenge in mice. *J Immunol*. 2008 Oct 1; 181(7):4955–64. [PubMed: 18802099]
21. Lindstrom T, Agger EM, Korsholm KS, Darrah PA, Aagaard C, Seder RA, et al. Tuberculosis subunit vaccination provides long-term protective immunity characterized by multifunctional CD4 memory T cells. *J Immunol*. 2009 Jun 15; 182(12):8047–55. [PubMed: 19494330]
22. Makedonas G, Betts MR. Polyfunctional analysis of human t cell responses: importance in vaccine immunogenicity and natural infection. *Springer Semin Immunopathol*. 2006 Nov; 28(3):209–19. [PubMed: 16932955]
23. Seder RA, Darrah PA, Roederer M. T-cell quality in memory and protection: implications for vaccine design. *Nat Rev Immunol*. 2008 Apr; 8(4):247–58. [PubMed: 18323851]
24. Andersen P. Tuberculosis vaccines - an update. *Nat Rev Microbiol*. 2007 Jul; 5(7):484–7. [PubMed: 17571458]

25. van Dissel JT, Arend SM, Prins C, Bang P, Tingskov PN, Lingnau K, et al. Ag85B-ESAT-6 adjuvanted with IC31((R)) promotes strong and long-lived Mycobacterium tuberculosis specific T cell responses in naive human volunteers. *Vaccine*. Mar 11.

Figure 1. Recognition of novel TB antigens in persons with LTBI and with TB disease

(A) Whole blood from persons with LTBI was incubated for 6 days with mycobacterial antigens and IFN- γ levels in supernatants measured by ELISA. The proportion of persons with a positive response, defined as an IFN- γ level above 31.2pg/mL, following subtraction of the IFN- γ level from the unstimulated control, is shown. (B) PBMC from a different cohort of persons with LTBI or TB disease were incubated with the antigens for 6 days with mycobacterial antigens and IFN- γ levels in supernatants measured by ELISA. Proportion of persons with a detectable response, defined as mentioned above, is shown. The Fisher's exact test was used to assess differences between proportions of responders. (C) IFN- γ response (pg/mL) to the antigens indicated. The medians are represented by the grey horizontal bars. A positive response was defined as an IFN- γ level above 31.2pg/mL after background subtraction. The Mann-Whitney U test was used to assess differences between the 2 groups.

Figure 2. CD3⁺ T cell Viability

Comparison of viable CD3⁺ T cells in PBMC of LTBI and TB individuals stimulated with BCG and *M.tb*-specific antigens for 6 days. (A) Viable T cells were selected by gating on CD3⁺ T cells against ViViD. (B) Frequency of viable CD3⁺ T cells. The medians are represented by the grey horizontal bar. The Mann-Whitney U test was used to assess differences between the 2 groups.

Figure 3. Gating strategy to analyze the cytokine expression profiles of proliferating CD4⁺ and CD8⁺ T cells

PBMC were stained with Oregon Green and stimulated with mycobacterial antigens, SEB, or were left unstimulated for 6 days, and analyzed by flow cytometry. (A) Doublet cells were excluded by gating on forward scatter-height (FSC-H) against forward scatter-area (FSC-A), and viable T cells were then selected by gating on CD3⁺ against ViViD. CD3⁺ T cells were further differentiated into proliferating (OG^{low}) CD8⁻ and CD8⁺ T cell subsets by staining for CD8 (CD3⁺CD8⁻ = CD4⁺ T cells, CD3⁺CD8⁺ = CD8⁺ T cells), and then cytokine expression patterns were analyzed in proliferating T cells. (B) Representative dot plots of cytokine co-expression patterns in unstimulated or antigen-specific proliferating CD4⁺ T cells. (C–F): Comparison of T cell proliferation and intracellular cytokine expression in individuals with LTBI or TB disease. CD4⁺ T cell antigen-specific

proliferation upon incubation with Rv2660 (C), Rv2659 (D), BCG (E), and ESAT-6/CFP-10 (F), detected by gating on Oregon Green^{low} CD4⁺ T cells. Frequencies of CD4⁺ T cell cytokine expression profiles; IFN- γ TNF- α ⁺IL2⁺, total IFN- γ , total TNF- α ⁺ and total IL-2⁺, of proliferating cells following incubation of PBMC with antigens (A–D, right panels), detected by intracellular cytokine staining. The medians are represented by the grey horizontal bar. The Mann-Whitney U test was used to assess differences between the 2 groups.

Figure 4. Rv2660- and Rv2659-specific CD8⁺ T Lymphocyte Proliferation and Cytokine expression profiles

Comparison of T cell proliferation and intracellular cytokine expression in individuals with LTBI or TB disease. CD8⁺ T cell antigen-specific proliferation upon incubation with Rv2660 (A), Rv2659 (B), BCG (C), and ESAT-6/CFP-10 (D), detected by gating on Oregon Green^{low} CD4⁺ T cells. Frequencies of CD8⁺ T cell cytokine expression profiles; IFN- γ TNF- α +IL-2⁺, Total IFN- γ , Total TNF- α + and Total IL-2⁺, of proliferating cells following incubation of PBMC with antigens (A–D, right panels), and detected by intracellular cytokine staining. The medians are represented by the grey horizontal bar. The Mann-Whitney U test was used to assess between the 2 groups.