police or social services may become the “lead” agency, but the paediatrician and primary healthcare team continue their involvement with the family.

Multiagency meeting to discuss the case

Finally, two to three months after the death a meeting is held to discuss the case, which includes all professionals who were involved with the family. This gives an opportunity to review the classification of the death, identify any contributory factors (fig 2), debrief all parties participating in the care of the family, and plan for continuing support of the family, including informing them of the assessment of the cause of the infant's death. The family is given a written explanation in plain English of what is understood of the cause of death and the results of the pathology investigations, and a meeting is arranged for the family with the paediatrician and general practitioner to answer their questions and identify continuing needs for support.

This joint approach ensures that all necessary information is collected sensitively and promptly, with a minimum of repetition. The broad experience of normal childcare practices in the community that is brought by the paediatrician and child protection team reduces the risk of attributing death or injury to normal variants in patterns of child care. The continuing participation of the paediatricians in research into current childcare practices in the community further helps to inform their interpretation of information obtained after infant deaths.

Summary

The sudden unexpected death of an infant or young child warrants careful, evidence based investigation involving child protection teams, healthcare staff, and social services, to include a careful review of the history, examination of the scene of the death, postmortem examination to an evidence based protocol, and a multiagency review meeting. The aim is to provide help and support for the bereaved family, to identify where possible the cause of the death, to help prevent further such deaths if possible, and to ensure that future children are protected from avoidable environmental hazards, medical conditions (for example, metabolic conditions), and abuse.

Contributors: PS, PJF, and PSB jointly conducted the literature review. PS, PJF, and TH jointly developed the multiagency protocol.

Competing interests: PJF has given expert evidence in both civil and criminal court proceedings on the subject of the investigation of unexpected deaths of infants.

- 1 Byard RW, Kohle SD. *Sudden death in infancy, childhood and adolescence*. Cambridge: Cambridge University Press, 1994.
- 2 Fleming PJ, Blair P, Bacon C, Berry PJ. *Sudden unexpected deaths in infancy. The CESDI SUDI studies 1993-1996*. London: Stationery Office, 2000.
- 3 Centers for Disease Control and Prevention. Guidelines for death scene investigation of sudden unexplained infant deaths: recommendations of the interagency panel on sudden infant death syndrome. *Morb Mortal Wkly Rep MMWR* 1996;45(RR-10):1-22.
- 4 Bass M, Kavath RE, Glass L. Death-scene investigation in sudden infant death. *N Engl J Med* 1986;315:100-5.
- 5 American Academy of Pediatrics. Committee on Child Abuse and Neglect. Distinguishing sudden infant death syndrome from child abuse fatalities. *Pediatrics* 2001;107:437-41.
- 6 Stanton J, Simpson A. Murder misdiagnosed as SIDS: a perpetrator's perspective. *Arch Dis Child* 2001;85:454-9.
- 7 Dyer C. Group to review babies' deaths. *BMJ* 2003;327:10.
- 8 Walkins SJ. Conviction by mathematical error? *BMJ* 2000;320:2-3.
- 9 Meadow R. A case of murder and the BMJ. *BMJ* 2002;324:41-3.
- 10 Dent A, Condon L, Blair P, Fleming PJ. A study of bereavement care after a sudden and unexpected death. *Arch Dis Child* 1996;74:522-6.
- 11 Morentin B, Aguilera B, Garamendi PM, Suarez-Mier MP. Sudden unexpected non-violent death between 1 & 19 years in north Spain. *Arch Dis Child* 2000;82:456-61.
- 12 Durfee M, Tilton-Durfee D. Multiagency child death review teams: experiences in the United States. *Child Abuse Rev* 1995;4:377-81.
- 13 Arnestad M, Vege A, Rognum TO. Evaluation of diagnostic tools applied in the examination of sudden unexpected deaths in infancy and early childhood. *Forensic Sci Int* 2002;125:262-8.
- 14 Rognum T, Lier L. Police investigation and SIDS: an improved system of co-operation between health personnel, forensic pathologists and the police. In: Rognum T, ed. *Sudden infant death syndrome: new trends in the nineties*. Oslo: Scandinavian University Press, 1995:289-92.
- 15 Gilbert RE, Rudd PT, Berry PJ, Fleming PJ, Hall E, White DG, et al. Combined effect of infection and heavy wrapping on the risk of sudden infant death. *Arch Dis Child* 1992;67:272-7.
- 16 Willinger M, James S, Catz C. Defining the SIDS: deliberations of an expert panel convened by the National Institute of Child Health and Human Development. *Pediatr Pathol* 1991;11:677-84.
- 17 Dyer C. Paediatricians did not have duty of care to patient's mother. *BMJ* 2002;325:1321.
- 18 Working Party of the Royal College of Paediatrics and Child Health. *Fabricated or induced illness by carers. Report of the working party of the Royal College of Paediatrics and Child Health*. London: RCPCH, 2002.
- 19 Department of Health. *Safeguarding children in whom illness is fabricated or induced*. London: DoH, 2002.
- 20 Mitchell E, Krous HF, Donald T, Byard RW. An analysis of the usefulness of specific stages in the pathologic investigation of sudden infant death. *Am J Forensic Med Pathol* 2000;21:395-400.
- 21 Krous HF. The international standardised autopsy protocol for sudden unexpected infant death. In: Rognum T, ed. *Sudden infant death syndrome: new trends in the nineties*. Oslo: Scandinavian University Press, 1995:81-98.
- 22 Wigglesworth JS, Keeling JW, Rushton DI, Berry PJ. Pathological investigations in cases of sudden infant death. *J Clin Pathol* 1987;40:1481-3.
- 23 Mitchell EA, Stewart AW, Scragg R, Ford RPK, Taylor BJ, Becroft DMO, et al. Ethnic differences in mortality from sudden infant death syndrome in New Zealand. *BMJ* 1993;306:13-6.
- 24 Creighton SJ. Fatal child abuse—how preventable is it? *Child Abuse Rev* 1995;4:318-28.

Pepys and blood transfusion?

On 21 November 1667, Samuel Pepys was in discussion with colleagues at the Tavern in St. Clements Churchyard and:

they discourse of a man that is a little frantic (that hath been a kind of minister, Dr. Wilkins saying that he hath read for him in his church) that is poor and a debauched man, that the College have hired for 20s to have some of the blood of a sheep let into his body; and it is to be done on Saturday next. They purpose to let in about twelve ounces, which they compute is what will be let in in a minutes time by a watch. They differ in the opinion they have of the effects of it; some think that it may have a good effect upon him as a frantic man, by cooling his blood; others, that it will not have any effect at all. But the man is a healthy man, and by this means will be able to give an account what alteration, if any, he doth find in himself, and so may be usefull.

On 30 November, while in “good discourse” Pepys writes:

But here above all, I was pleased to see the person who had his blood taken out. He speaks well, and did this day give the Society a relation thereof in Latin, saying that he finds himself much better since, and as a new man. But he is cracked a little in the head, though he speaks very reasonably and very well. He had but 20s for his suffering it, and is to have the same again tried upon him - the first sound man that ever had it tried on him in England, and but one that hear of in France, which was a porter hired by the virtuosi. Here all the afternoon till within night.

Pepys's report (*The Shorter Pepys*, by Robert Latham, Folio Society, 1985) suggests an even earlier interest in transfusion than that reported by Schmidt and Leacock (*BMJ* 2002;322:1485-7).

D F Scott *retired clinical neurophysiologist, London*