Long term mortality after starvation during the Leningrad siege

Crucial confounding factor was overlooked

EDITOR—Sparén et al retrospectively analysed outcomes, in terms of cardiovascular mortality, in siege survivors and have selected as controls men of similar age who did not experience the siege.¹ The data show that survivors have a higher risk of cardiovascular disease. The authors conclude that the process of starvation has adversely affected the physiology of the men who experienced food shortages at a critical stage of their development.

This conclusion could be drawn if the mortality of the starvation process itself could be shown to have been applicable to all physiological types equally. People with greater fat reserves seem likely to outlive people with less in periods of famine, and that ability to withstand protracted famine is probably related to an efficient metabolism, which is substantially genetically inherited. This cohort of siege survivors may owe its survival to the possibility that its members, if not overweight themselves at the start of the siege, nevertheless might have been the children of overweight parents.

These parents, with better fat reserves themselves, would have been in a better position to survive the siege than their thinner peers and hence in a better position to ensure the survival of their children. This would have been true whether or not these children had inherited their parents' metabolic type, but especially true if they had. The bread ration for a family would have been distributed in the household as the parents saw fit, and parents with greater fat reserves in the winter of 1941-2 would have been able to hand over a greater proportion of the bread ration to their children.

This might imply that siege survivors in Leningrad represent a selected survival group with the metabolic characteristic of obesity, favourable in a time of famine. It would therefore be surprising if this group, when followed up into old age, did not show the increased risk of cardiovascular disease that we all know they should have.

Michael D Croft *salaried general practitioner*
Gateways Surgery, Shenley, Hertfordshire
WD7 9LP
michael.croft@hertsmere-pct.nhs.uk

Competing interests: None declared.

1 Sparén P, Vägerö D, Shestov DB, Plavinskaja S, Parfenova N, Hoptiar V, et al. Long term mortality after severe starvation during the siege of Leningrad: prospective cohort study. *BMJ* 2004;328:11. (3 January.)

No evidence that starvation around puberty causes later cardiovascular disease

EDITOR—In their study of survivors of the Leningrad siege Sparén et al present no evidence for a causal link between starvation around the age of puberty and later cardiovascular disease.¹

They allocate subjects to age groups 8 years or younger, 9-15 years, and 16-26 years on the basis of the entire 29 month siege. However, Pavlov's data show that there was only a three month window (November 1941 to January 1942) during which energy and nutritional intakes were extremely low.²

In Russia in 1941 pubertal changes would have not been likely much before the age of 12. More importantly, from their 12th birthday

boys were classed as adult dependants rather than children and in consequence their rations were cut substantially over this period. Therefore, linking para-pubertal starvation and later morbidity would rely on identifying specific outcome differences between those aged 12-15 over the winter of 1941-2 and other age groups.

Despite throwaway allusions that stress might have contributed to cardiovascular changes, the authors effectively dismiss this since they state repeatedly that their analysis indicates an effect of starvation. But persistent stress is well known to raise sympathetic

tone and cortisol secretion, with long term increase in blood pressure; 2.5 years under siege bombardment definitely constitutes a stressful environment. In addition, Pavlov mentions that boys aged 12 or over were assigned to defence duties such as bomb disposal.

In the absence of supporting data, it seems redundant to suggest nutritional involvement in any increased cardiovascular morbidity.

C Bell *head, department of physiology*
Trinity College Dublin, Dublin 2, Republic of Ireland
cbell@tcd.ie

Competing interests: None declared.

1 Sparén P, Vägerö D, Shestov DB, Plavinskaja S, Parfenova N, Hoptiar V, et al. Long term mortality after severe starvation during the siege of Leningrad: prospective cohort study. *BMJ* 2004;328:11. (3 January.)

2 Pavlov DV. *Leningrad 1941: the blockade*. Chicago, IL: University of Chicago Press, 1965.

Authors' reply

EDITOR—Croft makes the point that children of fat parents would have had a better chance of surviving the Leningrad siege: obese parents would be more likely to survive and protect their children. This may not have been the case. One of us who survived the siege (DS) recalls a common impression that fat people were among the first to die. Observing Russian prisoners of war, Leyton found "a big well-built man standing up to shortage of food less well than his smaller brother."¹

Siege survivors were not fatter than other war survivors when both groups were examined in 1975-7 (table 1 in the long version of our paper on bmj.com), and adjustment for body mass index did not reduce our estimates of siege effects. Croft's hypothesis, moreover, would be unable to account for the interaction with age.

Bell suggests that the stress of war may have caused hypertension and circulatory mortality 30-60 years later. We agree that prolonged bombardment constitutes a considerable stressor, as do food shortage, starvation, and threat of starvation; most deaths were indeed due to starvation.² The endocrine profile of severe starvation in several aspects mimics that of reaction to stress. In the short version of our paper we said that "starvation is entangled with the trauma of the siege." Whatever the biological pathway, the fact remains that the effect on cardiovascular mortality of exposure to siege strongly interacted with (peri-pubertal) age and was partly mediated by hypertension.

bmj.com

Letters appearing here are an edited selection of rapid responses originally posted on bmj.com

We ask for all letters to the editor to be submitted as rapid responses via bmj.com

For advice see: bmj.com/rapidresponses

Nutritional intake was extremely low for a period even after January 1942, when bread rations went up to 250 g.² Six months of severe food shortage, followed by hard physical work during refeeding triggered the "Leningrad blockade hypertension epidemic."³ We suggest that this hypertension epidemic was particularly pronounced among peri-adolescent boys since energy restriction or deficiency of certain nutrients around puberty may be more likely to interfere with blood pressure regulation.⁴

Raised blood pressure was still visible when subjects were examined in 1975-7. Surely it is not "redundant" to suggest that the starvation episode also played a part in the increased mortality from circulatory disease some 20 years later.

Denny Vägerö *professor*
Centre for Health Equity Studies, CHESS,
Stockholm University/Karolinska Institute,
Stockholm, Sweden

Dimitri Shestov *professor*
Institute of Experimental Medicine, Russian
Academy of Medical Sciences, St Petersburg, Russia

Rosaria Galanti *senior researcher*
Unit of Clinical Epidemiology, Department of
Medicine, Karolinska Hospital, Stockholm

Pär Sparén *senior researcher*
Department of Medical Epidemiology and
Biostatistics, Karolinska Institute, Stockholm
Par.Sparen@meb.ki.se

Competing interests: None declared.

- 1 Leyton GB. Effect of slow starvation. *Lancet* 1946;248:i:73-9.
- 2 Pavlov DV. *Leningrad 1941: the blockade*. Chicago, IL: University of Chicago Press, 1965:125.
- 3 Keys A, Brozek J, Henschel A, Mickelsen O, Taylor HL. *The biology of human starvation*. Vol 1. Minneapolis, MN: University of Minnesota Press, 1950:621.
- 4 Falkner B, Sherif K, Michel S, Kushner H. Dietary nutrients and blood pressure in urban minority adolescents at risk for hypertension. *Arch Pediatr Adolesc Med* 2000;154: 918-22.

Critical care outreach team's effect on patient outcome

Other conclusions are possible

EDITOR—In their paper Ball et al show that the intervention of a critical care outreach team increased survival to discharge and rate of readmissions to critical care.¹ Despite the reduced probability of inhospital mortality in the intervention group of patients, the relative risk ratio was 1.00-1.18; it therefore includes 1 (unless a further figure greater than zero behind the two zeros was omitted) and does not seem to indicate significance (a P value was not published).

The reduction of readmissions by nine patients over 12 months is likely to be explained by a relative increase of survivors in critical care by 34% in the intervention period, with an increase of median length of stay of 0.9 days. Unless the number of critical care beds was increased in the intervention period or the non-survivors died at an earlier stage during their admission, these two figures make the availability of empty beds for readmissions less likely. The increased length of stay in intensive care is also likely to reduce the need for readmission. Multivariate analysis with the number

of empty beds per day as a covariable may be able to show this.

The five senior critical care nurses and the nurse consultant participated in many activities other than following up patients discharged from critical care. The real effects of the critical care outreach team may be seen if the quality of care received by patients not admitted to intensive care could be assessed better. However, these data are more difficult to collect and subject to multiple confounding factors.

Christian P Subbe *advanced trainee in intensive care medicine*
Intensive Therapy Unit, Morriston Hospital,
Swansea SA6 6NL
csubbe@hotmail.com

Competing interests: CPS received grants for evaluation of early warning scores from the North East Wales NHS Trust.

- 1 Ball C, Kirkby M, Williams S. Effect of the critical care outreach team on patient survival to discharge from hospital and readmission to critical care: non-randomised population based study. *BMJ* 2003;327:1014-6. (1 November.)

More information is needed

EDITOR—The average hospital mortality in patients admitted to intensive care units in England is 30%, of which around 8% occurs after discharge from intensive care.¹ In the study by Ball et al the retrospective control group had a median predicted hospital mortality of 20.4%, and yet 19% died after discharge from intensive care.² No details are given of mortality in the intensive care unit, and only the results in the subgroup who survived to discharge from intensive care were reported. Clinicians can therefore only question why mortality after intensive care seems so much higher in a population of patients with an unremarkable predicted mortality.

Outcome prediction using the acute physiology and chronic health evaluation (APACHE II) score predicts hospital mortality rather than mortality in intensive care³ because discharge criteria from intensive care may vary between units and even within units as a function of demand (therefore influencing mortality in intensive care). By the same argument variations in mortality after intensive care might have been influenced by variation in discharge practices rather than variation in the quality of ward care. According to Ball et al, the difference between the two groups of survivors of critical care on admission to intensive care did not reach significance, but they cannot conclude that the groups were equivalent on discharge from intensive care.

If the outreach team has had a genuine effect on mortality, a reduction in the standardised mortality ratio to hospital discharge for all patients admitted to intensive care may be expected. If, however, the

standardised mortality ratio has not changed, a reduction in ward mortality may simply reflect changes in discharge policy of the intensive care unit.

Michael R Parker *consultant in intensive care medicine*
Manchester Royal Infirmary, Manchester M13 9WL
michael.r.parker@man.ac.uk

Competing interests: None declared.

- 1 Audit Commission. *Critical to success: the place of efficient and effective critical care services within the acute hospital*. London: Audit Commission, 1999.
- 2 Ball C, Kirkby M, Williams S. Effect of the critical care outreach team on patient survival to discharge from hospital and readmission to critical care: non-randomised population based study. *BMJ* 2003;327:1014-6. (1 November.)
- 3 Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: A severity of disease classification system. *Crit Care Med* 1985;13:818-29.

Authors' reply

EDITOR—The significance for the relative risk ratio, for patients surviving to discharge, between the two groups was $P = 0.0452$.

In relation to our readmission data, the increase in survival and increase in median length of stay before and after the introduction of the outreach team might explain the published results. After discussion with consultant colleagues we think that patients could be discharged to the ward with more confidence with the introduction of the outreach team. Therefore, the increase in length of stay in the unit was associated with patients' need for level 3 care rather than a conscious decision to keep patients in the unit for a longer period to avoid possible readmission.¹

We cannot claim the subgroups were equivalent on discharge. For some time we have been considering the utility of collecting acute physiology and chronic health evaluation (APACHE II) data on discharge from intensive care as this would help us to examine severity of illness on discharge to the ward. The table shows hospital mortality for all patients admitted to the intensive care unit. Our standardised mortality ratio for all patients admitted to intensive care eligible for mortality standardisation by APACHE II was 1.81 (95% CI 1.65 to 1.96) before the introduction of the outreach team ($n = 315$) and 1.38 (1.23 to 1.52) afterwards ($n = 354$). The difference in the ratio between the two groups implies that the critical care outreach team has had a genuine effect in reducing hospital mortality.

Carol Ball *consultant nurse*
carol.ball@royalfree.nhs.uk

Margaret Kirkby *senior sister critical care outreach team*

Susan Williams *clinical audit manager*
Critical Care Unit, Royal Free Hampstead NHS Trust, London NW3 2QG

Competing interests: None declared.

- 1 Daly K, Beale R, Chang RWS. Reduction in mortality after inappropriate early discharge from critical care unit: logistic regression triage model. *BMJ* 2001;322:1-5.

Hospital mortality. Values are numbers (percentages)

Hospital outcome	Before critical outreach team	After critical outreach team	Total
Alive	168 (48)	235 (62)	403 (55)
Dead	181 (52)	145 (38)	326 (45)
Total	349	380	729

Treatment of unilateral visual impairment on preschool vision screening

Preschool vision screening should continue, perhaps earlier

EDITOR—The report by Clarke et al showed that mild visual loss in preschool children benefited little by treatment and that current management of this group ought to be reviewed.¹

Their conclusion advocating “the use of relatively later screening” seems incorrect. This study and others show the benefits of treatment for greater degrees of vision loss.² Conversely, Simons and Preslan show the consequences of failure of treatment.³ To dismiss the findings of the study of Simons and Preslan as merely an inevitability of being a non-compliant group seems harsh.

Preschool screening should continue for all, perhaps even at an earlier age.² The results of the current study, however, imply that later referral and eventual expectant management of mild visual loss may be appropriate. Certainly, newer technologies to improve screening accuracy would be helpful.

As an aside, the report does not indicate whether any underlying eye disease was fortuitously diagnosed by the ophthalmological referrals.

Gerard O'Brien *general practitioner*
38 Cecil Street, Limerick, Republic of Ireland
obrienrta@yahoo.com

Competing interests: None declared.

- 1 Clarke MP, Wright CM, Hrisos S, Anderson JD, Henderson J, Richardson SR. Randomised controlled trial of treatment of unilateral visual impairment detected at preschool vision screening. *BMJ* 2003;327:251-4. (29 November.)
- 2 Williams C, Northstone K, Harrad RA, Sparrow J, Harvey I. Amblyopia treatment outcomes after screening before or at 3 years: follow up from randomised trial. *BMJ* 2002;324:1549.
- 3 Simons K, Preslan M. Natural history of amblyopia untreated due to lack of compliance. *Br J Ophthalmol* 1999;83:582-7.

Study leaves questions unanswered

EDITOR—Clarke et al provide evidence that deferring treatment for amblyopia from age 4 to 5 is not harmful as visual acuity remains the same after treatment.¹ Deferring makes it easier to target the population, mobilise resources, and motivate children to wear glasses and patches. They also recommend that fewer children need patching. Hence, it is an important article that could change current practice.

The efficacy of patching, however, depends on several variables such as total duration of patching, severity of amblyopia, initial visual acuity, the cause of amblyopia, and compliance.^{2,3}

Although two thirds of the children for whom patching was advised wore patches for

a mean of 20 weeks, the paper is not clear about the number of hours advised daily and the compliance. Cleary showed that maximal improvement occurs in response to 400 hours of occlusion wear or less, and to full time occlusion.³ She defines full time occlusion as wearing more than 8 hours a day (8-12 hours prescribed) and part time wear as less than 7 hours (2-6 hours prescribed). Improvement in acuity with six hours of prescribed daily patching is similar to that with prescribed fulltime patching in severe amblyopia.⁴ Shorter times of occlusion (such as 1 hour daily) fail to produce a clinically or statistically significant improvement in vision.⁵

I wonder whether the patching regimen and outcome in children with microtropia was similar to that in those without a squint.

Hirekatur V Srinivas *ophthalmologist*
Aditya Eye Clinic, Bangalore
560 050, India
srinivashv@excite.com

Competing interests: None declared.

- 1 Clarke MP, Wright CM, Hrisos S, Anderson JD, Henderson J, Richardson SR. Randomised controlled trial of treatment of unilateral visual impairment detected at preschool vision screening. *BMJ* 2003;327:251-4. (29 November.)
- 2 Woodruff G, Hiscox F, Thompson JR, Smith LK. Factors affecting the outcome of children treated for amblyopia. *Eye* 1994;8:627-31.
- 3 Cleary M. Efficacy of occlusion for strabismic amblyopia: can an optimal duration be identified? *Br J Ophthalmol* 2000;84:572-8.
- 4 Holmes JM, Craker RT, Beck RW, Birch EE, Cotter SA, Everett DF, et al; Pediatric Eye Disease Investigator Group. A randomized trial of prescribed patching regimens for treatment of severe amblyopia in children. *Ophthalmology* 2003;110:2075-87.
- 5 Moseley MJ, Fielder AR, Irvin M, Jones JS, Auld RJ. Effectiveness of occlusion therapy in ametropic amblyopia: a pilot study. *Br J Ophthalmol* 1997;81:956-61.

Mild amblyopia should still be treated

EDITOR—In their randomised controlled trial Clarke et al showed directly for the first time that treatment improves visual acuity, at least for moderate amblyopia.¹

However, we take issue with the conclusion that vision of 6/12 in one eye and 6/9 in the other should not be treated. There are several possible reasons why no effect of treatment was found for this group.

6/9 is likely to be within the normal range of results for this age group for the crowded test used rather than for a single letter test, in which 6/9 would be more likely to be subnormal at this age.

Some of these children may well have had simple uncorrected refractive error rather than amblyopia because there was no refraction before randomisation and treatment.

Test-retest data for children show that visual acuity varies by at least one line between different occasions.^{2,3}

A ceiling effect at the mild end of the scale means that the effect size can be expected to be smaller in this group (less room for improvement). Given that the mild group probably contained several children

with normal sight owing to misclassification and that the effect size would be expected a priori to be smaller in the mild group, the study probably did not have sufficient power to show an effect of treatment in this group. The fact that a significant interaction term was found between mild and moderate groups would be expected with the presence of a ceiling restriction on the effect size in the milder group. Furthermore, no data are provided on binocular fusion or stereopsis, outcomes which might have improved in the absence of change in visual acuity.

This is a worthwhile study on which the research community can build, but it cannot be expected to answer all the relevant questions and it would be premature to draw sweeping conclusions based on its findings.

Richard Harrad *consultant ophthalmologist*
R.A.Harrad@bristol.ac.uk

Cathy Williams *consultant ophthalmologist*
John Sparrow *consultant ophthalmologist*
Bristol Eye Hospital, Bristol BS1 2LX

Competing interests: None declared.

- 1 Clarke MP, Wright CM, Hrisos S, Anderson JD, Henderson J, Richardson SR. Randomised controlled trial of treatment of unilateral visual impairment detected at preschool vision screening. *BMJ* 2003;327:251-4. (29 November.)
- 2 Kheterpal S, Jones HS, Auld R, Moseley MJ. Reliability of visual acuity in children with reduced vision. *Ophthalmic Physiol Opt* 1996;16:447-9.
- 3 Manny RE, Hussein M, Gwiazda J, Marsh-Tootle W. Repeatability of ETDRS visual acuity in children. *Invest Ophthalmol Vis Sci* 2003;44:3294-300.

Author's reply

EDITOR—O'Brien seems to attach greater weight to a study of apparently declining acuity in a non-compliant group than to our observation in a randomised controlled study of no deterioration in our control group, who showed the same eventual result from deferred treatment. This simply illustrates again the pitfalls inherent in natural history studies, which are plagued by unmeasurable biases that randomised controlled trials are designed to avoid.

Srinivas comments on patching dose. A recent study has shown that two hours' patching is as effective as six for children whose vision is in this range.¹

Harrad et al agree that an acuity of 6/9 may be within the normal range for age, and so it is hard to see why they do not endorse our conclusion that treatment of such children may be inappropriate. Screening does detect defective vision rather than amblyopia, but children found to have refractive errors without amblyopia are still currently treated.

We did have ample power to detect a useful effect in the mild subgroup (6/9 and 6/12 vision at recruitment). This contained 100 children and their maximum potential for acuity gain from treatment was 0.2-0.3 log units—for example, from 6/9.5 to 6/6. We had 80% power to detect a difference of the order of 0.1 log units and >99% power to detect a difference of 0.2 log units. However, the observed difference in this group between deferred and full treatment was only 0.04 log units (approximately half a Snellen line).

We believe that our study adds weight to the case for early school entry vision screening. However, in the current state of knowledge, treatment should be targeted only at children with poorer acuity to avoid inappropriate treatment of physiological refractive errors.

Michael Clarke *university reader in ophthalmology*
Claremont Wing, Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP
M.P.Clarke@ncl.ac.uk

Competing interests: None declared.

1 Pediatric Eye Disease Investigator Group. A randomised trial of patching regimens in the treatment of moderate amblyopia in children. *Arch Ophthalmol* 2003;121:603-11.

Volunteer counsellors for supporting breast feeding

Support for breast feeding must be proactive

EDITOR—Graffy et al's finding that offering voluntary support to breastfeeding mothers did not significantly affect breastfeeding rates is not surprising.¹

Many will know the chaos caused by a new baby's arrival. Mounds of paperwork, presents, and clothes build up in most homes as the usual order is abandoned. Contact numbers and advice leaflets are mislaid, sleep taking precedence over finding them. People do not like to admit defeat and feel foolish when phoning to ask for help. Often a mother new to breast feeding does not know what questions to ask.

Although volunteers are commendable for their efforts, they are simply volunteers. I found it difficult to contact my local counsellor of the National Childbirth Trust as she was often unavailable because of other commitments, including a full time job and the care of two young children. I then felt guilty for taking up her time.

Although breast feeding is best in most cases, it is not easy. Before giving birth women should be warned of potential problems alongside the benefits. When problems arise they will then at least not feel ashamed to admit difficulties.

Breastfeeding support must be proactive. It should be integrated into postnatal care delivered at home through midwives, health visitors, or dedicated counsellors. And it must be more than the cursory, "Breast feeding going OK?" on a list. Women should be offered practical help with feeds by someone sitting next to them. This is particularly important as pressure on hospital beds means that women are often discharged before their milk has come in and feeding is fully established.

Annette E Hay *senior house officer, medicine*
Aberdeen Royal Infirmary, Aberdeen AB25 2ZN
annettehaycrawford@doctors.net.uk

Competing interests: None declared.

1 Graffy J, Taylor J, Williams A, Eldridge S. Randomised controlled trial of support from volunteer counsellors for mothers considering breast feeding. *BMJ* 2004;328:26. (3 January.)

Peer support is worth further investigation

EDITOR—We need not be too pessimistic about the potential of volunteer counsellors in supporting breastfeeding mothers. Graffy et al examined only a fairly limited intervention, which relied heavily on the new mother to suspect a breastfeeding problem and initiate contact with a counsellor.¹

A woman is more likely to initiate and maintain breast feeding if she has a relative or friend who has breast fed successfully,² and it is at least plausible that support provided by a trained volunteer peer counsellor might have a similar effect. However, many women, particularly those who are less motivated and less socially skilled, find it difficult to initiate contact with a comparative stranger and wait for help to be offered.³

Furthermore, many central questions about peer support have yet to be addressed in methodologically adequate studies.⁴ There is little objective information on the qualities of a successful volunteer supporter, the experience and training she should have, how peer support could be most effectively offered, and the comparative importance of befriending and imparting information during the peer support process.

Commentators are quick to blame society for breastfeeding failure. Volunteer peer supporters might be the interface between society and the new mother where change could be most readily instigated. We urgently need effective methods of supporting the maintenance of breast feeding; volunteer peer support is worth further investigation.

Carol M A Campbell *community paediatrician (staff grade)*
Community Paediatric Unit, Bridgeview House,
Londonderry BT47 1TG
ccampbell@foylebv.n-i.nhs.uk

Competing interests: None declared.

- 1 Graffy J, Taylor J, Williams A, Eldridge S. Randomised controlled trial of support from volunteer counsellors for mothers considering breast feeding. *BMJ* 2004;328:26. (3 January.)
- 2 Hoddinott P, Pill R. Qualitative study of decisions about infant feeding among women in east end of London. *BMJ* 1999;318:30-4.
- 3 Hoddinott P, Pill R. Nobody actually tells you: a study of infant feeding. *Br J Midwifery* 1999;7:558-65.
- 4 Protheroe L, Dyson L, Renfrew MJ. *The effectiveness of public health interventions to promote the initiation of breast feeding: evidence briefing*. London: NHS Health Development Agency, 2003. Available at: www.hda.nhs.uk/evidence

American medicine meets the American dream

Moral dimension is important

EDITOR—Upshur points out that the distribution of medical care throughout the world is unequal and that this inequality is likely to worsen.¹ This is of course true. However, it is inappropriate to consider this problem of distribution of medical resources in isolation.

The sale of weapons and their development consume much of the world's "sur-

plus" resources, and, unlike frivolous health care, this industry is directly injurious. Similarly the widespread adoption of large four wheel drive vehicles increases pollution and injures those unfortunate enough to run into them.

Free markets and for profit companies exist to make money, not to provide the greatest good to the greatest number.

We must consider not just how medical services are developed and distributed, but all resources. For those fortunate enough to live in affluence all choices about resource consumption and personal expenditure contain an important moral dimension.

Stephen Workman *assistant professor*
Dalhousie University Department of Medicine,
Halifax, NS, Canada B3H 2Y9
sworkman@dal.ca

Competing interests: None declared.

1 Upshur R. Better than Well: American medicine meets the American dream. *BMJ* 2003;327:567. (6 September.)

Much inequality is self perpetuated

EDITOR—Upshur says that the problems associated with enhancement technologies are deeply related to affluence, which may seem unjust.¹ Yet we see self administration of unnecessary enhancements by poor people, often. Example: inner city people who sport sculpted hair styles weekly but won't get free health maintenance, even vaccinations.

In our compassionate zeal to flatten inequalities, we forget that much inequality is self perpetuated, preferring to think, "They can't help it." Conversely, sometimes enhancements are life saving. When is teenage acne—rejection, loss of self esteem, depression, suicide—a cosmetic illness?

The slippery slope of judgment requires educated, ethical, free doctors. This implies a medical system supplying the needy with basic care, not entitlements, paid by the group (government or insurance), and with enhancements paid by the individual unless modified by a doctor's considered opinion.

If enhancements require affluence, so be it. Those funding the first wave are the guinea pigs. With competition, prices fall and all can afford. Soon it's mainstream.

Pepi Granat *doctor*
7800 Red Road, Suite 202, South Miami, FL 33143,
USA
pgranat@pol.net

Competing interests: None declared.

1 Upshur R. Better than Well: American medicine meets the American dream. *BMJ* 2003;327:567. (6 September.)

Low prices for HIV drugs are only one step forward

EDITOR—Anything that cuts the price of drugs for HIV treatment is potentially beneficial to people with HIV infection in the developing world, as, for example, the recent deal from the World Health Organization.¹ But it remains only one step on a long path.

In many developing countries, the health systems have broken down. Infrastructure and skills through which to deliver complex health care are lacking. Education levels frustrate the most effective use of complex treatments, owing to inappropriate sharing of medication or lack of compliance, phenomena also to be found in Western populations. The risk is that, by focusing so much attention on the price of complex medicines, we will overlook the massive investment needed in developing countries fighting HIV.

However, without massive state aid, international stabilisation, and reduced weapons sales, better access to medicines, which may be unaffordable for many even at their lowest future price, is not a panacea.

Peter A West *director*
York Health Economics Consortium, University of York YO10 5NH
paw11@york.ac.uk

Competing interests: York Health Economics Consortium is a contract research company owned by the University of York. Clients include pharmaceutical companies, governments, and the NHS. The consortium has undertaken no work on patent and related issues for pharmaceutical companies or governments linked to current negotiations of the World Trade Organisation. York Health Economics Consortium has estimated savings from parallel trade in medicines in Europe for parallel traders. PAW has researched the cost effectiveness of HIV prevention strategies for non-governmental organisations and the UK government in many developing countries.

1 Pollock M, Price D. New deal from the World Health Organization. *BMJ* 29003;327:571. (13 September.)

Older people and the "patient centred" NHS

Choice for mentally ill patients will be even less with new legislation

EDITOR—Rowland and Pollock highlight another of the inequities of the NHS plan, with its heavy emphasis on acute hospital waiting and admission times, and the consequent impact on frail and chronically ill elderly people.¹

Elderly mental health beds will not be included in the delayed discharge legislation because they are not considered to be acute, though how beds that are needed to admit patients sectioned under the Mental Health Act are not considered acute beggars belief.

This exemption could be seen as recognition of the need for careful and often complex community care arrangements for these patients. However, social workers, who are already unable to provide the care packages they need to discharge these patients from acute hospitals, once subject to fines for the delays will inevitably see patients with depression or dementia, for whom they will not be fined, with even lower priority than they do now. The current staggering delays will lengthen further with their attendant effects on acute services.

Patients are already inappropriately admitted to and languish in acute wards, unable to be discharged or transferred to suitable therapeutic environments because of lack of investment, some of which is spent on devising ever more inventive ways of manipulating waiting lists.

The choice for these patients is already lamentable, but it will become even more restricted once this legislation is enacted.

David G Wilkinson *consultant in old age psychiatry*
Moorgreen Hospital, Southampton SO30 3JB

Competing interests: None declared

1 Rowland DR, Pollock AM. Choice and responsiveness for older people in the "patient centred" NHS. *BMJ* 2004;328:4-5. (3 January.)

Prioritisation will be for financial reasons rather than by need

EDITOR—Rowland and Pollock discuss many of the problems and inequities of the Community Care (Delayed Discharges) Act.¹ They do not mention that inpatients under the care of a psychiatrist have been specifically excluded, although most psychiatric inpatients also occupy acute beds which in many places are in short supply.

My service has always had an excellent working relationship with the local social services department, and I have found that delayed discharges are usually due to shortages in staffing and resources and not laziness or incompetence.

It seems now that people will have to be prioritised for services for financial reasons and not on the basis of need. People living in the community will become a lower priority, with psychiatric inpatients the lowest priority of all. The government does not seem to regard the blocking of psychiatric acute beds as a problem.

This legislation may also be counter-productive to multiagency working because inevitably resentment will build up when mental health teams see their patients marginalised in favour of general hospital inpatients.

Old age psychiatrists may think twice before transferring patients from medical or surgical beds to psychiatric assessment beds knowing that their social care will be prioritised if they stay where they are. This is likely to lead to admissions to residential or nursing home care for people who might have been rehabilitated at home, given more time and appropriate multidisciplinary and multiagency assessment and treatment.

Gillian E Moss *consultant in old age psychiatry*
Meadowbrook, Salford M6 8DD
gmoss@meadowbrook.bstnhp.nhs.uk

Competing interests: None declared.

1 Rowland DR, Pollock AM. Choice and responsiveness for older people in the "patient centred" NHS. *BMJ* 2004;328:4-5. (3 January.)

Proactive support in primary care can avert admissions

EDITOR—Rowland and Pollock's editorial induces reflection on the consequences to

the NHS and social services of patients admitted into secondary care in less than ideal circumstances.¹ In primary care such patients are recognised as on the brink of coping with chronic illness, increasing frailty, and confusion. In an increasingly mobile society many have no one close to them to offer support in precarious circumstances. Choice for such patients is indeed presented as an obstacle to the efficient functioning of the system that purports to protect them from destitution.

Our team recently reported a study seeking to identify such patients in primary care and offer them additional support, particularly over winter.² The data suggest that such patients can be identified and worked with proactively to reduce potential admission to hospital and an uncertain future. However, the effort of the primary health-care team must be concerted. It requires visits, calls, and interventions, which are welcomed even if not explicitly sought.

Despite our busy lives, few of us want to see our older relatives become a burden to the state. Our experience is that the involvement of an interested primary care team expressing concern is often enough to move relatives to act when action is required. In New Zealand substantial family input is reported to lighten general practitioners' load by reducing the need to resolve social issues.³ In the United Kingdom healthcare professionals must take the initiative, but when they do we have found that the family is not far behind.

Moyez Jiwa *lead research fellow*
University of Sheffield, Institute of General Practice, Community Sciences Centre, Northern General Hospital, Sheffield S5 7AU
m.jiwa@sheffield.ac.uk

Competing interests: None declared.

1 Rowland DR, Pollock AM. Choice and responsiveness for older people in the "patient centred" NHS. *BMJ* 2004;328:4-5. (3 January.)

2 Jiwa M, Gerrish K, Gibson A, Scott H. Preventing avoidable hospital admission of older people. *Br J Community Nurs* 2002;7:426-31.

3 Jones P. Elderly at home—lessons from New Zealand. *Health and Ageing* Dec 2003;22.

Cure is better than prevention

EDITOR—Prevention may not be better than cure.¹

Karl Popper, having fled from philistine philosophy, has already said: "In the realm of errors, cure is better than prevention."²

Overenthusiastic scientists, politicians, and so on, tend to forget that the realm of error in which we live is large. Prevention is permissible only in well understood and controllable conditions.

Jules Schagen van Leeuwen *gynaecologist*
St Antonius Hospital, 3435 EM Nieuwegein, Netherlands
jsvleeuw@knmg.nl

Competing interests: None declared.

1 Loeffler I. Prevention is better than cure. *BMJ* 2004;328:115. (10 January.)

2 Miller D, ed. Popper selections. Princeton, NJ: Princeton University Press, 1985:10.