Published in final edited form as:

J Am Chem Soc. 2012 April 11; 134(14): 6296–6308. doi:10.1021/ja211970p.

Experimental and Theoretical Studies on the Nazarov Cyclization/Wagner-Meerwein Rearrangement Sequence

David Lebœuf^a, Vincent Gandon^b, Jennifer Ciesielski^a, and Alison J. Frontier^a

Vincent Gandon: vincent.gandon@u-psud.fr; Alison J. Frontier: frontier@chem.rochester.edu

^aDepartment of Chemistry, University of Rochester, Rochester, NY 14627

^bICMMO, UMR CNRS 8182, Université Paris-Sud 11, 91405 Orsay cedex, France

Abstract

Highly functionalized cyclopentenones can be generated stereospecifically by a chemoselective copper(II)-mediated Nazarov/Wagner-Meerwein rearrangement sequence of divinyl ketones. A detailed investigation of this sequence is described including a study of substrate scope and limitations. After the initial 4π electrocyclization, this reaction proceeds *via* two different sequential [1,2]-shifts, with selectivity that depends upon either migratory ability or the steric bulkiness of the substituents at C1 and C5. This methodology allows the creation of vicinal stereogenic centers, including adjacent quaternary centers. This sequence can also be achieved by using a catalytic amount of copper(II) in combination with NaBAR_4^f , a weak Lewis acid. During the study of the scope of the reaction, a partial or complete *E/Z* isomerization of the enone moiety was observed in some cases prior to the cyclization, which resulted in a mixture of diastereomeric products. Use of a Cu(II)-bisoxazoline complex prevented the isomerization, allowing high diastereoselectivity to be obtained in all substrate types. In addition, the reaction sequence was studied by DFT computations at the UB3LYP/6-31G(d,p) level, which are consistent with the proposed sequences observed, including *E/Z* isomerizations and chemoselective Wagner-Meerwein shifts.

Introduction

Electrocyclizations are powerful pericyclic reactions that occur via simple orbital reorganization.¹ They are one of the few methods available to create carbon-carbon bonds stereospecifically. Cationic electrocyclizations, such as the Nazarov (4π cationic) cyclization, are initiated by formation of an extended cationic π -system, which also reacts with conservation of orbital symmetry.² In the past few years numerous examples of catalytic Nazarov cyclizations have been reported,^{3–11} and asymmetric induction in the Nazarov cyclization has been achieved using chiral auxiliaries,¹² chiral Lewis acids¹³ or organocatalysts,¹⁴ but the substrate scope is narrow in each case. The Nazarov cyclization can also serve as a cationic initiation step, generating an oxyallyl cation intermediate that can be intercepted by a suitable trapping agent.¹⁵ This oxyallyl cation can also be the starting point for rearrangement sequences.¹⁶ In a general sense, these reactions fit within a larger class of cationic cascades that involve activation of a carbonyl or alkene functionality to generate "stabilized" carbocation intermediates, which can undergo a variety of

Correspondence to: Vincent Gandon, vincent.gandon@u-psud.fr; Alison J. Frontier, frontier@chem.rochester.edu.

Supporting Information Available.

Complete reference 23, experimental procedures and characterization data for all new compounds are available free of charge via the Internet <http://pubs.acs.org>.

subsequent transformations. Cation-initiated C-C bond-forming reactions have great synthetic potential, if their reactivity and selectivity can be controlled, especially in the area of terpene and steroid biosynthesis.¹⁷

In our studies on the Nazarov cyclization, we found an interesting and unusual electrocyclization/rearrangement sequence that could efficiently compete with the traditional Nazarov cyclization when stoichiometric amounts of Ligand-Cu(SbF₆)₂ were employed.¹⁸ Experiments suggest that these reaction conditions suppress proton transfer, extending the lifetime of the oxyallyl cation and allowing rearrangement pathways to compete with elimination (Scheme 1). It is possible to achieve the stereospecific synthesis of unusual spirocyclic compounds with adjacent stereogenic centers, including adjacent quaternary centers. However, this initial study revealed the following limitations: (*i*) one full equivalent of Lewis acid promoter was required for rearrangement; (*ii*) the reaction scope was limited to cyclic substrates.

Recently, we have shown that this reaction does not require the use of a sophisticated ligand and can even be carried out with a catalytic amount of a copper (II) complex.¹⁹ Highly functionalized cyclopentenones can be prepared stereospecifically via conrotatory electrocyclization, with high chemoselectivity of subsequent Wagner-Meerwein shifts. In many cases, it is possible to install adjacent quaternary centers with complete diastereocontrol. The present article is devoted to the investigations conducted by our laboratory to evaluate the scope and the limitations of this copper(II)-mediated Nazarov cyclization/Wagner-Meerwein rearrangement sequence of acyclic substrates. To gain insight into the mechanism of the title reaction, the key steps have also been analyzed by means of DFT calculations.

Results and Discussion

Initial experiments were focused on substrate **1** bearing a 2,4,6-trimethoxyphenyl (TMP) group at C5 (Scheme 2), because the bulk and strong electron-donating character of the TMP group increases reaction rates.^{4b} The reaction was carried out in dichloromethane at room temperature in the presence of 100 mol% of (MeCN)₅Cu(SbF₆)₂. Complete conversion was obtained within 5 minutes to provide cyclopentenone **2** in 92% yield as a single diastereomer. Relative stereochemistry was established by nOe analysis.

The oxyallyl cation intermediate **3** is generated by a 4 π conrotatory electrocyclization. We previously reported that the cyclization of mixtures of *E* and *Z* isomers of alkylidene β -ketoesters was stereoconvergent and occurred *via* an efficient isomerization process to give the product with the stereochemistry corresponding to the *Z* isomer cyclization, which explains the exclusive formation of intermediate **3**.^{4b} It is well-known that the migratory aptitude of aryl groups is significantly greater than that of alkyl groups,²⁰ so it is not surprising to observe a chemoselective [1,2]-phenyl shift to give intermediate **4**. After this step, either [1,2]-hydride or carbon shift can occur. An electron-donating substituent on the aryl ring enhances its migratory ability. However, the propensity for migration can also be affected by steric factors. In the case of intermediate **4**, the sp² carbon shift is presumably interrupted by the bulkiness of the 2,4,6-trimethoxyphenyl, thus, cyclopentenone **2**, resulting from [1,2]-hydride shift, is the only product observed. The diastereoselectivity is dictated by both the conrotatory electrocyclization and the suprafacial nature of the subsequent Wagner-Meerwein migrations. Product **6**, the product expected from the conventional Nazarov cyclization/elimination sequence, is not observed.

Next, we conducted experiments to test the effect of the different substituents at C1 and C2, with a 2,4,6-trimethoxyphenyl substituent at C5 (Table 1). Consistent with Scheme 2, in

each case, the reaction is completely chemoselective with the exclusive migration of the R¹ group and [1,2]-hydride shift from C5 to give rise to the corresponding cyclopentenones in good yields (Table 1, Entries 1–4). We noticed that the rearrangement of substrate **11** (Table 1, Entry 3) led to the formation of two products **12** and **13** in a 1:1 ratio and a combined yield of 90%, the latter corresponding to the loss of the silyl group. This deprotection could result from the presence of the hexafluoroantimonate counterion in solution.²¹

We also examined the cyclization of 1,4-dien-3-ones with a phenyl group at C1 and various groups at C5, including 4-methoxyphenyl, phenyl, heteroaromatic, cinnamyl, and alkyl substituents. Almost all the reactions were carried out under refluxing conditions except for the substrates having an electron-donating 4-methoxyphenyl substituent at C5 (Table 2, Entries 4 and 5). In each case, cyclopentenones were obtained in high yields (up to 95%). In substrates **1**, **16**, **18**, and **34**, we observed the chemoselective [1,2]-shift of the phenyl from C1 to C2, and [1,2]-hydride shift from C5 to C1, consistent with the mechanism shown in Scheme 2 (Table 2, entries 1–3 and 11). However, a different reaction sequence was observed in substrates without a bulky TMP or isopropyl group at C5. In these cases, the second migration was a [1,2]-aryl or -alkenyl shift rather than a [1,2]-hydride shift (Table 2, Entries 4–10). Many of the products were obtained as mixtures of diastereomers, in ratios ranging from 4:1 to 20:1. The structures of the major products were ascertained by nOe analysis, which clearly indicated a *cis*-relationship between the phenyl and the alkyl moieties.

In the next set of experiments, the rearrangement of substrates with opposite C1–C2 double bond geometry (relative to the C1 Ph) was studied (Table 3). Interestingly, the major products obtained in most cases were the same as the ones previously observed with similar ratios (compare Table 2, Entries 4–7 and 9 with Table 3, Entries 3–7). These findings suggest that partial or complete isomerization of the C1–C2 bond occurs prior to cyclization in the presence of the copper(II) complex. Contrary to expectations, reduced chemoselectivity was observed in the rearrangement of substrate **37**. Rather than exclusive migration of the phenyl group as usual (Tables 2 and 3), a 3:1 ratio of phenyl vs. ethyl migration products was observed (Table 3, Entry 2).

Changes in chemoselectivity were also seen in the reactions of substrates possessing steric bulk at both C1 (R¹ = *i*Pr or Ph and R² = Ph) and C5 (R³ = aryl) (Table 4). In these substrates, the substituent with the lower migratory aptitude shifted in both the first and the second [1,2]-Wagner-Meerwein shift, compared to the corresponding substrates with less hindered pentadienyl cation termini (i.e. C1 and C5; see Tables 1–3). With R¹ = Me and R² = Me, cyclopentenone **45** was obtained in 66% yield after the expected migration of the electron-donating 4-methoxyphenyl group from C5 (Equation 1).

(1)

In contrast, with R¹ = *i*Pr or Ph and R² = Ph (Table 4, Entries 1 and 3), the rearrangement led to the formation of cyclopentenones **47** and **52** after hydride migration from C5. Similarly, hydride migration was observed in hindered substrates with a thienyl or a phenyl substituent at C5 (Table 4, Entries 4 and 5). When R¹ = *i*Pr, migration of the isopropyl group rather than the phenyl group was observed from C1 (Table 4, Entries 2–5).

It is clear from the results in Tables 1–4 that different substrates follow different reaction pathways during the cyclization/rearrangement sequence. From the substrates shown in Tables 2 and 3, we know that reaction of both *E* and *Z* isomers of a substrate give the same product mixture, which demonstrates that *E/Z* isomerization is facile under the reaction conditions. Consistent with these findings, *E/Z* isomerization could be observed by ^1H NMR prior to cyclization in representative cases. In contrast, *E/Z* isomerization was not observed in substrates with $\text{R}^1 = i\text{Pr}$ (cf. Table 4, Entries 2–5 to Table 3, Entries 1, 3, 5 and 7). The steric hindrance exerted by this group seems to prevent the σ -bond rotation of C1–C2 required for the *E/Z* isomerization. Additional study using DFT calculations allowed further elucidation of the mechanism (vide infra.)

Unfortunately, this isomerization leads to the isolation of a mixture of two diastereomeric products in a number of cases. If the fidelity of the double bond geometry can be preserved during the reaction, a highly diastereoselective reaction should occur. With this goal in mind, we anticipated that placement of a bulky ligand on copper might slow down the isomerization of the double bond. To test this hypothesis, we focused on substrate **57**, for which the use of $(\text{MeCN})_5\text{Cu}(\text{SbF}_6)_2$ gave rise to a mixture of the three products **58a**, **58b**, and **59** resulting from the isomerization of C1–C2 bond (Equation 2). In contrast, use of a bulky bidentate *t*BuBOX ligand allowed selective cyclization/rearrangement of **57**, producing cycloadduct **58a** as the sole product (Equation 3).

(2)

(3)

Cyclization/rearrangement of substrate **37** (with *Z* geometry) was examined next. Without the bulky ligand, *E/Z* isomerization was observed and the chemoselectivity of the first [1,2]-shift favored the phenyl shift (Table 3, Entry 2; 3:1 ratio of **19/38**). Promotion of the same reaction with *t*BuBox-ligated catalyst not only suppressed *E/Z* isomerization, but gave exclusively minor product **38** in a high yield of 91%.

(4)

Thus, using the bulky *t*BuBox ligand prevents the isomerization of both *Z* and *E* substrates, and is also able to reverse the chemoselectivity of the [1,2]-migration.

As noted above, while the chemoselectivity of [1,2]-shifts is governed by relative migratory aptitudes in most cases (Tables 1–3), in substrates with three bulky substituents at C1 and C5 (the reacting termini of the pentadienyl cation), reaction outcomes are counter to predictions (Table 4). In these cases, other effects overrule the chemoselectivity dictated by migratory aptitude predictions. In the case of the observed migration of isopropyl rather than phenyl (Table 4, entries 2–5), we propose that the reaction is dominated by thermodynamics: equilibration of cationic intermediates **60**, **61**, and **62** occurs to favor formation of **62** (see computational data). In cases when a hydride migration is observed rather than an aryl migration in the second [1,2]-shift (Table 4, entries 1–5), inspection of intermediate **62** is helpful. Migration of the expected aryl group must proceed through phenonium intermediate **64**, which would experience significant steric strain from both the presence of the phenonium bridge and the *syn* relationship of the phenyl group and the *isopropyl* group (Scheme 3). In comparison, the electronically disfavored hydride shift leads to **63**, which lacks the steric congestion of **64**.

Originally, a limitation to the method was the requirement for one equivalent of a Lewis acidic promoter for efficient cyclization/ rearrangement.^{18, 19} With catalytic asymmetric reaction as the ultimate goal, we sought to develop a transition-metal catalyzed version of the reaction. Our observations of both cyclization and rearrangement sequences suggested that Brønsted basic species in the reaction mixture facilitate the elimination of a proton, leading to formation of the normal Nazarov cycloadduct (see Scheme 1), while coordination of the basic carbonyl lone pairs with one equivalent of the copper complex slows elimination, allowing rearrangement pathway to compete.^{18b} Based on that model, we wanted to identify an additive that could bind to the carbonyl lone pairs of both substrate and product, could readily exchange with a reaction promoter, but would not promote the cyclization itself. Such an additive should work in combination with catalytic amount of a chiral Lewis acid to coordinate the lone pairs of the carbonyls, favoring rearrangement over elimination. In our initial attempts to achieve a catalytic rearrangement, we used a combination of 10 mol% of (MeCN)₅Cu(SbF₆)₂ and 90 mol% of several additives such as Zn(OAc)₂, Mn(acac)₂, Fe(acac)₂, LiClO₄, Mg(OTf)₂, NaSbF₆, NaPF₆, or NaBPh₄, but these reactions were sluggish and led to mixtures of products. Furthermore, all these Lewis acids displayed poor solubility in dichloromethane. BAr₄^f is a highly soluble, non-coordinating counterion, and we found that LiPF₆ itself promotes the cyclization, so it was not a suitable additive. Fortunately, experiments with NaBAr₄^f were ultimately successful.^{5b, 19} When dienone **66** was treated with (MeCN)₅Cu(SbF₆)₂ (10 mol%) and NaBAr₄^f (90 mol%) at room temperature for 10 minutes, cyclopentenone **67** was formed in 90% yield (Equation 5). It was surprising that a sodium ion, rather than a chelating additive was sufficient for suppression of the elimination pathway. The results suggest that only one carbonyl group plays a role in the proton capture.

(5)

Application of these optimal reaction conditions to a representative set of substrates demonstrated that the catalytic protocol was comparable to the stoichiometric reaction (Chart 1).

Study of Reaction Pathways by Density Functional Theory

To rationalize some of the mechanistic questions raised during this study, DFT computations were carried out using dienones **a1** as model substrates and Cu²⁺ as promoter (See Supporting Information for details).^{23, 24, 25} Dienones **a1** are *s-trans/s-trans* conformers with ideal alignment for Nazarov cyclization (Scheme 4). They are taken as reference compounds (free energy of 0.0 kcal/mol). The facile *E/Z* isomerization of the C1–C2 double bond, observed during some of the cyclization studies (Tables 2 and 3), served as the starting point of the simulation. The calculations suggest the following sequence for the isomerization: (i) σ -bond rotation from **a1** leads to *s-cis/s-trans* conformer **a2**, (ii) diastereomutation of **a2** gives **a3**, (iii) coordination of the phenyl group of **a3** to the metal center leads to **a4**, and (iv) a final σ -bond rotation of **a4** leads to the corresponding *s-trans/s-trans* isomer **a5**, aligned for Nazarov cyclization.²⁶

Then, conrotatory 4 π electrocyclicization could be computed from **a1** and **a5**, leading to oxallyl cations **b1** and **b2** respectively. From these two complexes, all possible Wagner-Meerwein rearrangements were modeled (suprafacial 1,2-shifts), leading to **c1–3**, and then to **d1–6**. The latter six complexes correspond to the six possible enones **e1–6** after copper dissociation.

The results are summarized in Tables 5–7, subdivided into the three main mechanistic categories (isomerization, electrocyclicization, and [1,2]-shifts) for clarity. With respect to isomerization, the overall transformation of pentadienyl cation **a1** into isomeric **a5** is either slightly exergonic (Y = Ph, 2-thienyl, 4-MeO-C₆H₄) or slightly endergonic (Y = (*E*)-cinnamyl or TMP). In contrast, the Ph-coordinated intermediates **a4** are always significantly more stable than **a1**, but **a4** does not represent a deep potential well. In each Y series, the rate-determining step is the diastereomutation itself (**a2**→**a3**), as one would expect, rather than any conformational interconversion. Those high-lying transition states show a virtually perpendicular arrangement of the PhC1R and the MeC2C planes (see Figure 1 with Y = 4-MeO-C₆H₄ and R = Me, MeC1C2Me = 79°).

Electron-withdrawing Y groups are expected to assist the isomerization process by lowering the C1–C2 bond order, as in resonance form **a2'**. On the other hand, electron-donating groups should favor **a2''** and make the isomerization more difficult (Equation 6).

(6)

In the phenyl series, the substrate with Y = TMP has a high inversion barrier, calculated at 19.1 kcal/mol (Table 5, Entry 5). The relationship between the inversion barrier and the electronic nature of the Y group is displayed in Table 6. As the energy required to reach TS_{a2-a3} increases, the C1–C2 bond order increases while the C1–C2 bond distance and the natural charge at C1 both decrease. These values indicate that the most important resonance form of **a2** should be **a2'** when Y = TMP.

Notably, the steric hindrance at C1 also plays a dramatic influence on the isomerization rate. For example, when R = Me is changed to R = Et, the free energy of activation of the **a2**→**a3** step increases by more than 4 kcal/mol to reach 24.6 kcal/mol (Table 5, Entries 5 and 6). When R = *i*Pr (Table 5, Entry 7), it was no longer possible to find the isomerization transition state, presumably because it is too high in energy. The absence of an isomerization

transition state on the potential energy surface when $R = iPr$ is consistent with the 100% stereoselective transformation of **49** into **50** (Table 4, Entry 3).

The rest of the mechanistic scenario depicted in Scheme 4 was computed for $R = Me$ and $Y = Ph$, (*E*)-cinnamyl, 4-MeOC₆H₄, and TMP (Table 7). The 4π electrocyclizations were all found to be endergonic. Nevertheless, those leading to the less sterically congested isomer **b1** are always kinetically and thermodynamically favored over those leading to **b2** (Table 7, Entries 1 and 2). Again, the magnitude of the activation barriers is critically dependent on the nature of Y . Electron-donating groups appear to raise the barrier for electrocyclization process (*e.g.* 4.1 for $Y = Ph$ vs 15.5 kcal/mol for $Y = 4-MeO-C_6H_4$), probably because these groups stabilize the pentadienyl cation.²⁸

Recently, the decomplexation of copper from the carbonyl oxygens of alkylidene β -ketoesters and oxazolidinones, resulting in an out-of-plane orientation of the C4 carbonyl, has been proposed to explain the stereochemical outcome of some copper(II)-catalyzed Nazarov cyclizations.^{3m} To examine this proposal in the context of our system, **TS_{a1-b1}** and **TS_{a5-b2}** were recomputed in the 4-MeO-C₆H₄ series after decomplexation of copper from the ester functionality. In the new transition states **TS_{a1'-b1'}** and **TS_{a5'-b2'}**, the metal establishes an agostic interaction with the methyl group at C2 (see Figure 2). However, the energies of these species relative to **a1** are much higher than those of **TS_{a1-b1}** and **TS_{a5-b2}** (56.4 and 58.3 kcal/mol respectively). An η^3 -dioxyallyl copper complex was also envisaged, yet we could not locate any electrocyclization transition state. Instead, a species corresponding to the rotation of the ester moiety converged. Again, this transformation requires a prohibitive free energy of activation of 50.3 kcal/mol. Thus, in both of the cases we modelled, dechelation is unlikely. Additional calculations using ligated copper (II) ((CH₃CN)₂Cu²⁺) instead of (Cu (II)²⁺) did not allow optimization of these high-lying transition states either (*vide infra*).

With respect to the [1,2]-shifts, the computations are in good agreement with the expected migratory aptitudes of methyl, hydride, phenyl or Y (Figure 3). For instance, the initial methyl shifts leading from **b1/b2** to **c1** (Table 7, Entries 3 and 4) require a considerably higher free energy of activation than the corresponding phenyl shifts (**b1** to **c2** or **b2** to **c3**; Table 8, Entries 5 and 6). For the subsequent step (from **c2/c3** to intermediates **d**) the [1,2] shifts of Y groups are kinetically favored over [1,2]-hydride shifts (Table 7, Entries 10 and 11 vs. Entries 9 and 12).

The geometries of the cyclopentenyl cations of type **d** also warrant comment. Indeed, the electron-deficient center C5 may receive electron density from the phenyl or the Y groups. This interaction is particularly obvious when $Y = (E)$ -cinnamyl, 4-MeO-C₆H₄, and TMP, which converge as bicyclo[3.1.0]hexane derivatives (see **d2'**, **d3'**, and **d5'**).²⁹ For example, as shown in Scheme 4, when $Y = Ph$ the carbocationic center is clearly located on C5, but delocalized throughout the π -system of the other electron-rich groups, including the (*E*)-cinnamyl group. In spite of our efforts, it was not possible to find minima representing both **d** and **d'** within the same series (Figure 4). This suggests that the bicyclic structure is always more stable than the localized secondary carbocation. Although much weaker, this neighbouring stabilization effect is also present in other intermediates, such as **b2** (tertiary carbocation at C2), **c2** and **c3** (tertiary carbocation at C1) (see Supporting Information).

To save computer time, all the calculations presented above involve unligated copper (II). It is quite clear that some of the computational results may change with the introduction of ligands, notably regarding the π -complexes **a4** and the agostic complexes shown in Figure 2. To address this issue, the computations related to the 4-MeO-C₆H₄ series were repeated with two acetonitrile units on Cu (Scheme 5). As expected, intermediate **a4** is no longer

observed in the **a1**→**a5** sequence. However, the isomerization transition state connecting **a2** and **a3** remains the highest-lying (25.7 kcal/mol), a value that is actually higher than those of the electrocyclization steps leading to **b1** and **b2** (18.3 and 20.3 kcal/mol respectively). Complex **b1** remains kinetically and thermodynamically favored over **b2**, since the electrocyclization steps are still endergonic, and the methyl shifts leading to **c1** remain slower than the phenyl shifts. In contrast with the unligated series, **c2** and **c3** do not converge. They collapse directly to **d2** and **d5**, with the migration of the phenyl group immediately followed by migration of the PMP. This finding is not unique to the calculations with ligated copper, since **c2** is also absent from the potential energy surface in the (*E*)-cinnamyl series (see Table 7). With acetonitrile ligands, the structures of **d2**, **d3**, and **d5** are quite distinct from the corresponding ones in the unligated series (phenoniums **d2'**, **d3'**, **d5'**; Scheme 4). This may be because the [(MeCN)₂Cu] moiety is more electron-rich, so the backdonation towards the organic backbone becomes more efficient. The stabilization of the charge at C5 (see **d2**, Scheme 5) by the adjacent PMP (i.e. the phenonium) is no longer required since the metal can now accommodate two formal charges, leading to Lewis structures that are much more like the final products. Despite these geometrical differences, the shifts leading to **d2** remain the most kinetically favorable, and **d2** represents the framework of the product experimentally obtained. Thus, both sets of calculations, using either (MeCN)₂Cu]²⁺ or the simpler unligated copper(II), predict the same experimental outcome. It is nonetheless interesting to note the favorable effect the ligands have on the migration at C5.

Summary, Conclusions and Perspectives

In summary, we have developed a diastereoselective method for preparation of highly functionalized cyclopentenones based on a copper(II)-promoted Nazarov/Wagner-Meerwein sequence. The scope of the reaction is broad, allowing preparation of products with adjacent stereogenic centers, and in some cases, adjacent quaternary centers. The reaction was always efficient using a stoichiometric amount of a copper-(II) complex, and in many cases it was possible to lower the amount of copper to 10 mol% by adding a sodium salt to attenuate the basicity of the carbonyl oxygen. Although chemoselectivity of the sequential [1,2] shifts is typically high, predicting the structures of the products is not as simple as one might expect. Sometimes the chemoselectivity is governed by relative migratory aptitude, and sometimes we find that other factors overwhelm these propensities, leading to the selective formation of an isomer with unexpected connectivity. Isomerization of the Nazarov cyclization substrate is another complication in some cases, which also makes prediction of product stereochemistry difficult. DFT calculations have helped us to gain mechanistic insight into the substrate isomerization and to understand the factors dictating selectivity in the cation rearrangements. Furthermore, we found it was possible to prevent the isomerization with the use of a bisoxazoline ligand on the copper (II) complex, allowing controlled, predictable electrocyclization of enone substrates with the expected stereospecificity. Ongoing research in the laboratory is focused upon the development of an asymmetric version of this reaction and application of this strategy towards the synthesis of natural products.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Science Foundation (Grant CHE-0847851, supporting D. L.) and the National Institutes of Health (NIGMS R01 GM079364, supporting J. C.) for funding this work. We used the computing facility of the CRIHAN (project 2006-013). We thank Prof R. Eisenberg (University of Rochester) for helpful discussions.

References

- (a) Woodward RB, Hoffmann R. *Angew. Chem. Int. Ed. Engl.* 1969; 8:781.(b) Woodward, RB.; Hoffmann, R. *The Conservation of Orbital Symmetry*. Verlag Chemie: Weinheim; 1970.
- (a) Santelli-Rouvier C, Santelli M. *Synthesis*. 1983;429.(b) Habermas, KL.; Denmark, SE.; Jones, TK. *Organic Reactions*. Paquette, L.A., editor. Vol. Vol. 45. New York: John Wiley & Sons, Inc.; 1994. p. 1-158.(c) Harmata M. *Chemtracts*. 2004; 17:416.(d) Pellissier H. *Tetrahedron*. 2005; 61:6479.(e) Frontier AJ, Collison C. *Tetrahedron*. 2005; 61:7577.(f) Tius MA. *Eur. J. Org. Chem.* 2005; 11:2193.(g) Nakanishi N, West FG. *Curr. Opin. Drug Discov. Dev.* 2009; 12:732.(h) Shimada N, Stewart C, Tius MA. *Tetrahedron*. 2011; 67:5881.(i) Vaidya T, Eisenberg R, Frontier AJ. *Chem Cat Chem*. 2011; 3:1531.
- For recent examples of Nazarov cyclizations, see: Williams DR, Robinson LA, Nevill CR, Reddy JP. *Angew. Chem. Int. Ed.* 2007; 46:915. He W, Huang J, Sun X, Frontier AJ. *J. Am. Chem. Soc.* 2008; 130:300. [PubMed: 18067294] Marcus AP, Lee AS, Davis RL, Tantillo DJ, Sarpong R. *Angew. Chem. Int. Ed.* 2008; 47:6379. Amere M, Blanchet J, Lasne MC, Rouden J. *Tetrahedron Lett.* 2008; 49:2541. Kokubo M, Kobayashi S. *Chem. Asian J.* 2009; 4:526. [PubMed: 19212964] Rieder CJ, Winberg KJ, West FG. *J. Am. Chem. Soc.* 2009; 131:7504. [PubMed: 19435345] Malona JA, Cariou K, Frontier AJ. *J. Am. Chem. Soc.* 2009; 131:7560. [PubMed: 19445456] Singh R, Parai MK, Panda G. *Org. Biomol. Chem.* 2009; 7:1858. [PubMed: 19590781] Cordier P, Aubert C, Malacria M, Lacôte E, Gandon V. *Angew. Chem. Int. Ed.* 2009; 48:8757. Wang M, Han F, Yuan H, Liu Q. *Chem. Commun.* 2010; 46:2247. Wender PA, Stemmler RT, Sirois LE. *J. Am. Chem. Soc.* 2010; 132:2532. [PubMed: 20141136] Murugan K, Srimurugan S, Chen C. *Chem. Commun.* 2010; 46:1127. Kerr DJ, White JM, Flynn BL. *J. Org. Chem.* 2010; 75:7073. [PubMed: 20873806] Marcus AP, Sarpong R. *Org. Lett.* 2010; 12:4560. [PubMed: 20843020] Cai Z, Harmata M. *Org. Lett.* 2010; 12:5668. [PubMed: 21080691] Hastings CJ, Pluth MD, Bergman RG, Raymond KN. *J. Am. Chem. Soc.* 2010; 132:6938. [PubMed: 20443566] Rieder CJ, Winberg KJ, West FG. *J. Org. Chem.* 2011; 76:50. [PubMed: 21105713] Spencer WT III, Levin MD, Frontier AJ. *Org. Lett.* 2011; 13:414. [PubMed: 21155592] Li W-DZ, Duo WG, Zhang CH. *Org. Lett.* 2011; 13:3538. [PubMed: 21650169] Brooks JL, Caruana PA, Frontier AJ. *J. Am. Chem. Soc.* 2011; 133:12454. [PubMed: 21780785]
- Cu(II): He W, Sun X, Frontier AJ. *J. Am. Chem. Soc.* 2003; 125:14278. [PubMed: 14624567] He W, Herrick IR, Atesin TA, Caruana PA, Kellenberger CA, Frontier AJ. *J. Am. Chem. Soc.* 2008; 130:1003. [PubMed: 18161974]
- Pd(II): Bee C, Leclerc E, Tius MA. *Org. Lett.* 2003; 5:4927. [PubMed: 14682731] Zhang J, Vaidya T, Brennessel WW, Frontier AJ, Eisenberg R. *Organometallics*. 2010; 29:3541. Subramaniam SS, Handa S, Miranda AJ, Slaughter LM. *ACS Catal.* 2011; 1:1371.
- Sc(III): Larini P, Guarna A, Occhiato EG. *Org. Lett.* 2006; 8:781. [PubMed: 16468766] Malona JA, Colbourne JM, Frontier AJ. *Org. Lett.* 2006; 8:5661. [PubMed: 17107097] Bitar AY, Frontier AJ. *Org. Lett.* 2009; 11:49. [PubMed: 19053717]
- IR(III): Janka M, He W, Frontier AJ, Eisenberg R. *J. Am. Chem. Soc.* 2004; 126:6864. [PubMed: 15174845] Janka M, He W, Frontier AJ, Flaschenriem C, Eisenberg R. *Tetrahedron*. 2005; 61:6193. Vaidya T, Atesin AC, Herrick IR, Frontier AJ, Eisenberg R. *Angew. Chem. Int. Ed.* 2010; 49:3363. Atesin AC, Zhang J, Vaidya T, Brennessel WW, Frontier AJ, Eisenberg R. *Inorg. Chem.* 2010; 49:4331. [PubMed: 20361755]
- V(IV): Walz I, Bertogg A, Togni A. *Eur. J. Org. Chem.* 2007:2650.
- Au(I) or Au(III): Zhang L, Wang S. *J. Am. Chem. Soc.* 2006; 128:1442. [PubMed: 16448102] Jin T, Yamamoto Y. *Org. Lett.* 2008; 10:3137. [PubMed: 18572946] Lemièrre G, Gandon V, Cariou K, Hours A, Fukuyama T, Dhiman A-L, Fensterbank L, Malacria M. *J. Am. Chem. Soc.* 2009; 131:2993. [PubMed: 19209868] Krafft ME, Vidhani DV, Cran JW, Manoharan M. *Chem. Commun.* 2011; 47:6707.
- Fe(II) or Co(I): Kawatsura M, Higuchi Y, Hayase S, Nanjo M, Itoh T. *Synlett*. 2008:1009. Kawatsura M, Kajita K, Hayase S, Itoh T. *Synlett*. 2010:1243.
- Pt(II): Zheng H, Xie X, Yang J, Zhao C, Jing P, Fang B, She X. *Org. Biomol. Chem.* 2011; 9:7555.
- (a) Pridgen LN, Huang K, Shilcrat S, Tickner-Eldridge A, DeBrosse C, Haltiwanger RC. *Synlett*. 1999:1612.(b) Tius MA, Harrington PE. *Org. Lett.* 2000; 2:2447. [PubMed: 10956518] (c)

- Harrington PE, Murai T, Chu C, Tius MA. *J. Am. Chem. Soc.* 2002; 124:10091. [PubMed: 12188673] (d) Kerr DJ, Metje C, Flynn BL. *Chem. Commun.* 2003:1380.(e) Banaag AR, Tius MA. *J. Org. Chem.* 2008; 73:8133. [PubMed: 18808186]
13. (a) Aggarwal VK, Belfield AJ. *Org. Lett.* 2003; 5:5075. [PubMed: 14682768] (b) Liang G, Trauner D. *J. Am. Chem. Soc.* 2004; 126:9544. [PubMed: 15291550] (c) Walz I, Togni A. *Chem. Commun.* 2008:4315.(d) Rueping M, Ieawsuwan W. *Adv. Synth. Catal.* 2009; 351:78.(e) Yaji K, Shindo M. *Synlett.* 2009:2524.(f) Cao P, Deng C, Zhou Y-Y, Sun XL, Zheng J-C, Xie W, Tang Y. *Angew. Chem. Int. Ed.* 2010; 49:4463.
14. (a) Rueping M, Ieawsuwan W, Antonchick AP, Nachtsheim BJ. *Angew. Chem. Int. Ed.* 2007; 46:2097.(b) Bow WF, Basak AK, Jolit A, Vicic DA, Tius MA. *Org. Lett.* 2010; 12:440. [PubMed: 20030316] (c) Basak AK, Shimada N, Bow WF, Vicic DA, Tius MA. *J. Am. Chem. Soc.* 2010; 132:8266. [PubMed: 20507121]
15. For a review, see: Grant TN, Rieder CJ, West FG. *Chem. Commun.* 2009:5676. and references cited therein. For other relevant examples, see: Janka M, He W, Haedicke IE, Fronczek FR, Frontier AJ, Eisenberg R. *J. Am. Chem. Soc.* 2006; 128:5312. [PubMed: 16620081] Dhoro F, Kristensen TE, Stockmann V, Yap GPA, Tius MA. *J. Am. Chem. Soc.* 2007; 129:7256. [PubMed: 17508753] Nie J, Zhu HW, Cui HF, Hua MQ, Ma JA. *Org. Lett.* 2007; 9:3053. [PubMed: 17629286] Fujiwara M, Kawatsura M, Hayase S, Nanjo M, Itoh T. *Adv. Synth. Catal.* 2009; 351:123. Marx VM, Cameron TS, Burnell DJ. *Tetrahedron Lett.* 2009; 50:7213. Marx VM, Burnell DJ. *J. Am. Chem. Soc.* 2010; 11:1685. [PubMed: 20078119] Rieder CJ, Fradette RJ, West FG. *Heterocycles.* 2010; 80:1413. Scadeng O, West FG. *Org. Lett.* 2011; 13:114. [PubMed: 21126097] Marx VM, LeFort FM, Burnell DJ. *Adv. Synth. Catal.* 2011; 353:64. Wu Y-K, McDonald R, West FG. *Org. Lett.* 2011; 13:3584. [PubMed: 21699172] Marx VM, Stoddard RL, Heverly-Coulson GS, Burnell SJ. *Chem. Eur. J.* 2011; 17:8098. [PubMed: 21656587] Rueping M, Ieawsuwan W. *Chem. Commun.* 2011; 47:11450.
16. Ohloff G, Schulte-Elte KH, Demole E. *Helv. Chim. Acta.* 1971; 54:2913.(b) Paquette LA, Fristad WE, Dime DS, Bailey TR. *J. Org. Chem.* 1980; 45:3017.(c) Denmark SE, Hite GAG. *Helv. Chim. Acta.* 1988; 71:195.(d) Motoyoshiya J, Yazaki T, Hayashi S. *J. Org. Chem.* 1991; 56:735.(e) Gruhn AG, Reusch W. *Tetrahedron.* 1993; 49:8159.(f) Kuroda C, Koshio H, Koito A, Sumiya H, Murase A, Hirono Y. *Tetrahedron.* 2000; 56:6441.(g) Chiu P, Li SL. *Org. Lett.* 2004; 6:613. [PubMed: 14961636]
17. For recent reviews, see: Wendt KU, Schulz GE, Corey EJ, Liu DR. *Angew. Chem. Int. Ed.* 2000; 39:2812. Yoder RA, Johnston JN. *Chem. Rev.* 2005; 105:4730. [PubMed: 16351060] Christianson DW. *Chem. Rev.* 2006; 106:3412. [PubMed: 16895335] Tantillo DJ. *Chem. Soc. Rev.* 2010; 39:2847. [PubMed: 20442917] Tantillo DJ. *Nat. Prod. Rep.* 2011; 28:1035. and references cited therein. [PubMed: 21541432]
18. (a) Huang J, Frontier AJ. *J. Am. Chem. Soc.* 2007; 129:8060. [PubMed: 17555319] (b) Huang J, Leboeuf D, Frontier AJ. *J. Am. Chem. Soc.* 2011; 133:6307. [PubMed: 21466152]
19. Leboeuf D, Huang J, Gandon V, Frontier AJ. *Angew. Chem. Int. Ed.* 2011; 50:10981.
20. (a) Bachmann WF, Ferguson JW. *J. Am. Chem. Soc.* 1934; 56:2081.(b) Collins CJ. *J. Am. Chem. Soc.* 1955; 77:5517.(c) Owen JR, Saunders WH. *J. Am. Chem. Soc.* 1966; 88:5809.(d) Heidke RL, Saunders WH. *J. Am. Chem. Soc.* 1966; 88:5816.
21. Brimble MA, Crimmins D, Trzoss M. *Arkivoc.* 2005; i:39.
22. The following minor product was obtained with a yield of 13%:

23. Frisch, MJ., et al. Gaussian 03 (v C02). Wallingford CT: Gaussian, Inc.; 2004.
24. The geometry optimizations and thermodynamic corrections were performed with hybrid density functional theory (UB3LYP, doublet spin state for Cu, +2 overall charge) with the 6-31G(d,p) basis set. The choice of this functional and basis set is justified in the Computational Methods section of ref 25a (see references therein). The B3LYP functional was also used in ref 25b. The spin contamination was checked from the final value of $\langle S^2 \rangle$ for all structures and found to be very small (< 0.76). Intrinsic reaction coordinate (IRC) calculations utilized the default implementation. They were carried out in the Y = Ph and Y = PMP series. Solvation corrections for dichloromethane were computed using the polarizable continuum model (PCM) with UFF radii using the gas phase optimized structures. All relative energies presented in this manuscript are free energies (ΔG) in kilocalories per mole. Solvation free energies were calculated by adding the solvation energies to the computed gas phase relative free energies (ΔG_{298}). Wiberg bond indices were calculated from NBO theory as implemented in Gaussian.
25. For rare theoretical reports on Cu^{2+} -mediated Nazarov reactions, see: Mayoral JA, Rodríguez-Rodríguez S, Salvatella L. Chem. Eur. J. 2008; 14:9274. [PubMed: 18756568] Cao P, Deng C, Zhou Y-Y, Sun X-L, Zheng J-C, Xie Z, Tang Y. Angew. Chem. Int. Ed. 2010; 49:4463. For selected theoretical reports on H^+ -mediated Nazarov reactions, see: (c) Nieto Faza O, Silva López C, Álvarez R, de Lera AR. Chem. Eur. J. 2004; 10:4324. [PubMed: 15352115] Cavalli A, Masetti M, Recanatini M, Prandi C, Guarna A, Occhiato EG. Chem. Eur. J. 2006; 12:2836. [PubMed: 16429476] Polo V, Andrés J. J. Chem. Theory Comput. 2007; 3:816. Cavalli A, Pacetti A, Recanatini M, Prandi C, Scarpi D, Occhiato EG. Chem. Eur. J. 2008; 14:9292. [PubMed: 18752233] Davis RL, Tantillo DJ. Curr. Org. Chem. 2010; 14:1561. For a theoretical report on the retro-Nazarov reaction, see: (h) Harmata M, Schreiner PR, Lee DR, Kirchhoefer PL. J. Am. Chem. Soc. 2004; 126:10954. [PubMed: 15339180]
26. Some minor conformational rearrangements with very low free energy of activations have been omitted for clarity.
27. (a) Wiberg KB. Tetrahedron. 1968; 24:1083.(b) Foster JP, Weinhold F. J. Am. Chem. Soc. 1980; 102:7211.(c) Reed AE, Weinstock RB, Weinhold F. J. Chem. Phys. 1985; 83:735.(d) Weinhold, F.; Carpenter, JE. The Structure of Small Molecules and Ions. Naaman, R.; Vager, Z., editors. Plenum; 1988. p. 227
28. For the influence of electronic effect on the Nazarov cyclization, see: Jones TK, Denmark SE. Helv. Chim. Acta. 1983; 66:2377. Jones TK, Denmark SE. Helv. Chim. Acta. 1983; 66:2397. Denmark SE, Habernas KL, Hite GA. Helv. Chim. Acta. 1988; 71:168. Denmark SE, Klix RC. Tetrahedron. 1988; 44:4043. Denmark SE, Wallace MA, Walker CB. J. Org. Chem. 1990; 55:5543.
29. (a) Cram DJ. J. Am. Chem. Soc. 1949; 71:3863.(b) Cram DJ, Davis R. J. Am. Chem. Soc. 1949; 71:3871.(c) Cram DJ. J. Am. Chem. Soc. 1949; 71:3875.(d) Cram DJ. J. Am. Chem. Soc. 1964; 86:3767.(e) Brown HC, Morgan KJ, Chloupek FJ. J. Am. Chem. Soc. 1965; 87:2137.(f) Vogel, P. Carbocation Chemistry. Amsterdam: Elsevier; 1985. Chapter 7. (g) Olah GA, Porter RD. J. Am.

Chem. Soc. 1970; 92:7627.(h) Del Rio E, Menendez MI, Lopez R, Sordo TL. J. Phys. Chem. A. 2000; 104:5568.

Figure 1.
Geometries and energies of the species involved in the **a1**→**a5** transformation with Y = 4-MeO-C₆H₄ and R = Me (UB3LYP/6-31G**//PCM correction, kcal/mol)

Figure 2.
Electrocyclization and ester rotation transition states (UB3LYP/6-31G**//PCM correction, kcal/mol) for Y = 4-MeO-C₆H₄ and R = Me (distances in Å)

TS_{b1-c1} (27.5, Me-shift) **TS_{c2-d1}** (15.2, H-shift) **TS_{b1-c2}** (13.7, Ph-shift) **TS_{c2-d2'}** (9.7, 4-MeO-C₆H₄-shift)

Figure 3.
Geometry of selected [1,2]-shift transition states (UB3LYP/6-31G**//PCM correction, kcal/mol) for R = Me (distances in Å)

Figure 4.
Geometry of **d2** and **d2'** cations (UB3LYP/6-31G**//PCM correction, distances in Å).

Scheme 1.
Spirocycle synthesis via Nazarov/ Wagner-Meerwein rearrangement sequence.

Scheme 2.
Copper-mediated cyclization of 1,4-dien-3-one **1** (TMP = 2,4,6-trimethoxyphenyl).

Scheme 3.

Scheme 4.
Computed intermediates for the Cu^{2+} -mediated cyclization of dienones

Scheme 5.

Computed intermediates for the Cu^{2+} -mediated cyclization of dienones with $Y = 4\text{-MeO-C}_6\text{H}_4$ (PMP) and acetonitrile on Cu (UB3LYP/6-31G**//PCM correction, kcal/mol)

Chart 1.

Copper-catalyzed cyclization of substituted 1,4-dien-3-ones.

[a] *Reaction conditions*: substrate in CH₂Cl₂ (0.03 M) in the presence of NaBAR₄^f (90 mol %) and (MeCN)₅Cu(SbF₆)₂ (10 mol%) at rt for 0.1–0.5 h. [b] Reaction carried out at 45 °C. BAR₄^f = 3,5-bis(trifluoromethyl)phenylborate. TMP = 2,4,6-trimethoxyphenyl. PMP = 4-methoxyphenyl.

Table 1

Entry	Substrate	Temp (°C)	Time (h)	Product	Yield (%)
1		25	0.1		79
2		25	0.1		51
3		25	0.5		45
4		25	1		82

Reaction conditions: substrate in CH₂Cl₂ (0.03 M) in the presence of one equivalent of (MeCN)₅Cu(SbF₆)₂ at the indicated temperature. PMP = 4-methoxyphenyl; TMP = 2,4,6-trimethoxyphenyl.

Table 2

Entry	Substrate	Temp (°C)	Time (h)	Product	Yield (%)
1		25	0.3		92
2		25	0.3		95
3		40	0.5		95
4		25	0.2		86
5		25	0.2		90
6		40	0.5		85
7		40	0.5		80
8		40	1		80

Entry	Substrate	Temp (°C)	Time (h)	Product	Yield (%)
9		40	1		95
10		40	1		86
11		40	3		68

Reaction conditions: substrate in CH₂Cl₂ (0.03 M) in the presence of one equivalent of (MeCN)₅Cu(SbF₆)₂ at the indicated temperature. TMP = 2,4,6-trimethoxyphenyl. PMP = 4-methoxyphenyl.

Table 3

Entry	Substrate	Temp (°C)	Time (h)	Product	Yield (%)
1	 36	25	0.2	 2	95
2	 37	25	0.2	 19:38 (3:1)	92
3	 39 R ¹ = Me	25	0.5	 21a/21b (5:1)	88
4	 40 R ¹ = Et	25	0.5	 23a/23b (7:1)	88
5	 41 R ¹ = Me	40	1	 25a/25b (9:1)	86
6	 42 R ¹ = Et	40	1.5	 27a/27b (8:1)	88

Entry	Substrate	Temp (°C)	Time (h)	Product	Yield (%)
7	43	40	1	31a/31b (4:1)	89

Reaction conditions: substrate in CH₂Cl₂ (0.03 M) in the presence of one equivalent of (MeCN)₅Cu(SbF₆)₂ at the indicated temperature. TMP = 2,4,6-trimethoxyphenyl. PMP = 4-methoxyphenyl.

Table 4

Entry	Substrate	Temp (°C)	Time (h)	Product	Yield (%)
1		40	1		68 ^[a]
2		40	0.5		98
3		40	1		70
4		40	1.5		56
5		40	2		89

Reaction conditions substrate in CH_2Cl_2 (0.03 M) in the presence of one equivalent of $(\text{MeCN})_5\text{Cu}(\text{SbF}_6)_2$ at the indicated temperature. [a]: reaction in 1,2-dichloroethane. TMP = 2,4,6-trimethoxyphenyl. PMP = 4-methoxyphenyl.

[a] The other diastereomer has also been isolated in 13% yield.²²

Table 5

Computed free energies for the isomerization process at 298K (UB3LYP/6-31G**//PCM correction, kcal/mol) relative to **a1** for the **a1**→**a5** transformation (free energies of activation are displayed in italics).

Isomerization									
Entry		a1→a2		a2→a3		a3→a4		a4→a5	
		$\Delta G^\ddagger$	ΔG	$\Delta G^\ddagger$	ΔG	$\Delta G^\ddagger$	ΔG	$\Delta G^\ddagger$	ΔG
1	Y = Ph, R = Me	8.8	1.4	12.1	[a]	-	-8.4	3.5	-2.7
2	Y = 2-thienyl, R = Me	5.6	1.4	15.4	0.6	3.6	-7.8	1.5	-0.6
3	Y = (<i>E</i>)-cinnamyl, R = Me	0.1	2.3	17.8	[a]	-	-5.5	2.4	0.2
4	Y = 4-MeO-C ₆ H ₄ , R = Me	10.3	2.6	18.6	-0.9	4.0	-7.0	4.7	-1.4
5	Y = TMP, R = Me	9.0	-0.6	19.1	1.0	4.8	-6.4	1.4	2.0
6	Y = TMP, R = Et	9.3	-6.1	24.6	0.8	2.4	-5.4	6.8	3.0
7	Y = TMP, R = <i>i</i> Pr	7.9	5.1	[b]	-0.6	1.8	-7.6	8.9	-0.8

[a] No convergence of this intermediate, direct collapse to the next one.

[b] Could not be located on the PES.

Relationship between the **a2**→**a3** free energies of activation (UB3LYP/6-31G**//PCM correction, kcal/mol) and the polarization of the C1–C2 bond in **a2**.

Table 6

Entry	Y	$\Delta G^\ddagger$	C1C2 (Å)	Natural charge at C1[a]	Wiberg bond index[a]
1	C ₆ H ₅	12.1	1.429	0.200	1.35
2	4-MeO-C ₆ H ₄	18.6	1.402	0.176	1.49
3	TMP	19.1	1.373	0.07	1.62

[a] Natural charges and Wiberg bond indices were calculated from NBO theory.²⁷

Table 7

Computed free energies for the electrocyclization and the [1,2] shifts at 298 K (UB3LYP/6-31G**//PCM correction, kcal/mol), relative to a1 with R = Me (free energies of activation are displayed in italics).

Entry		Y = Ph		Y = (E)-cinnamyl		Y = 4-MeO-C ₆ H ₄		Y = TMP	
		$\Delta G^\ddagger$	ΔG	$\Delta G^\ddagger$	ΔG	$\Delta G^\ddagger$	ΔG	$\Delta G^\ddagger$	ΔG
<i>Nazarov</i>									
1	a1→b1	4.1	3.7	21.0	18.4	15.5	12.6	19.0	8.7
2	a5→b2	13.0	5.7	23.3	20.8	18.7	13.3	25.7	19.7
<i>Shifts</i>									
3	b1→c1 (1,2-Me shift)	10.6	-8.1	27.7	5.7	18.6	0.4	27.5	3.0
4	b2→c1 (1,2-Me shift)	7.9	-8.1	33.6	5.7	22.2	0.4	25.9	3.0
5	b1→c2 (1,2-Ph shift)	5.4	2.2	18.6	[a]	13.7	11.3	19.1	8.5
6	b2→c3 (1,2-Ph shift)	3.5	3.6	19.3	18.4	15.0	5.0	23.7	12.3
7	c1→d3 (1,2-Y shift)	2.3	-3.7	7.8	10.8	3.9	1.7	8.9	2.7
8	c1→d4 (1,2-H shift)	6.6	-20.9	19.0	-14.0	2.7	-18.9	17.1	-14.0
9	c2→d1 (1,2-H shift)	8.9	-15.6	-	-14.8	15.2	-11.3	17.6	-11.3
10	c2→d2 (1,2-Y shift)	7.8	-4.2	-	-0.3	9.7	6.3	14.2	9.4
11	c3→d5 (1,2-Y shift)	7.8	-2.2	17.4	5.7	9.5	7.4	15.4	7.3
12	c3→d6 (1,2-H shift)	8.6	-14.4	21.3	-17.7	15.1	-10.7	17.4	-6.0

[a] No convergence of this intermediate, direct collapse to d2.