Published in final edited form as:

Tetrahedron Lett. 2012 April 4; 53(14): 1701–1704. doi:10.1016/j.tetlet.2012.01.049.

Utilization of the hydroxyalkylation reaction to prepare bis(benzocrown ethers)

Matthew Zielinski, Adam F. Tracy, and Douglas A. Klumpp*

Department of Chemistry and Biochemistry, Northern Illinois University, DeKalb, IL 60115

Abstract

The hydroxyalkylation reaction has been used to condense benzocrown ethers with various aldehydes and ketones. The condensation reactions are catalyzed by triflic or sulfuric acid. The products from the reactions are bis(benzocrown ethers) and they are formed in good yields (42–98%, 13 examples).

Keywords

crown compounds; condensation; heterocycles; electrophilic aromatic substitution; superacid

Since their discovery by Pederson in 1967, crown ethers and related substances have been useful in a variety of applications.¹ Their abilities to complex metal cations have made them useful as ionophores, phase transfer catalysts, and components in ionselective electrodes.² Because of the many applications of crown ethers and functionalized derivatives, there continues to be a need for new synthetic methods leading to crown ether products. Benzocrown ethers may be considered activated, electron rich arenes. As such, functionalized benzocrown ethers may be readily prepared using Friedel-Crafts type chemistry.³ We and others have recently demonstrated that the hydroxyalkylation condensation is an outstanding synthetic method leading triarylmethanes,⁴ functionalized heterocycles,⁵ and condensation polymers.⁶ The hydroxyalkylation reaction typically involves the condensation of an aldehyde or ketone with activated aromatic compounds, such as aryl ethers or phenols.⁷ Less reactive arenes may also form condensation products if the aldehyde or ketone generates highly electrophilic intermediates. In the following manuscript, we describe our studies involving the hydroxyalkylation reaction with benzocrown ethers leading to the preparation of bis(benzocrown ether) products.

Our initial studies involved reacting a series of aldehydes and ketones with benzo-18-crown-6 in the presence of the Brønsted superacid, CF₃SO₃H (eq 1 and Table 1). For example, 5-chloroisatin (**1**) is reacted with benzo-18-crown-6 (**2**) in CF₃SO₃H and the condensation product **3** is formed in good yield (eq 1). The reaction is best done with an excess of CF₃SO₃H (8 equivalents). Analysis of the crude mixture by NMR reveals very high regioselectivity in the reaction with benzo-18-crown-6 (**2**). Other isatins (**4–6**) likewise

© 2012 Elsevier Ltd. All rights reserved.

*Corresponding author, dklumpp@niu.edu; Tel.: 815-753-1959; fax: 815-753-4802.

Supplementary Material

Supplementary data (NMR spectra of products) associated with this article can be found, in the online version, at doi:

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

give the condensation products (**10-12**) in reactions with benzo-18-crown-6 (**2**). Ninhydrin (**7**) reacts efficiently with arenes at the 2-position, so condensation with compound **2** gives product **13** in good yield. In this case, H_2SO_4 is superior to $\text{CF}_3\text{SO}_3\text{H}$ in promoting the condensation. The quinone **8** also provides a bis(benzocrown ether) product (**14**), as does the heterocyclic aldehyde (**9**). Condensation reactions were also accomplished with benzo-15-crown-5 (**22**) and benzo-12-crown-4 (**23**, Table 2).

(1)

Superacid-promoted hydroxyalkylations have been utilized in several types of condensations leading to macromolecules.⁶ Recently, Zolotukhin and co-workers described the hydroxyalkylation reaction of isatin with a series of dibenzocrown ethers using methanesulfonic acid and P_2O_5 as an acid catalyst.⁸ By utilization of dibenzo-18-crown-6 and triflic acid, we have accomplished similar condensations to prepare crown-containing macromolecules (Scheme 1). Reaction of isatin **5** leads to the A_2B_2 -type condensation polymer **24**, a hard off-white solid which softens at about 170 °C. A similar condensation was accomplished with 2-pyridinecarboxaldehyde (**9**) to give the macromolecule **25**. Poor solubility in organic solvents prevented GPC analysis of polymers **24** and **25**, however NMR analysis could be accomplished by dissolving the polymers in $\text{CF}_3\text{CO}_2\text{D}$.¹¹ For structure **24**, the ^1H NMR spectrum is consistent with the presence of both oxyindole and crown ether structures. Further analysis was done by infrared spectroscopy,¹¹ comparing the carbonyl absorption bands of compound **5** and the polymeric product. *N*-Methylisatin exhibits strong absorption bands at 1727 and 1604 cm^{-1} , respectively, for the ketone and amide- type carbonyl groups. Analysis of the polymer **24** shows a single carbonyl absorption at 1684 cm^{-1} . This is consistent with the formation of a condensation polymer, because entrapped or unreacted isatin **5** would exhibit two carbonyl absorptions. Likewise, polymer **25** shows no absorptions in the carbonyl region of the infrared spectrum.⁹ As noted previously by Zolotukhin, the condensation reactions with dibenzo-18-crown-6 likely produces the pair of regioisomeric repeat units (electrophilic attack at the 4,4' positions and at the 4,5' positions of the dibenzo-18-crown-6).

In summary, we have found that hydroxyalkylation reactions can lead to products having two benzocrown units.¹⁰ These conversions are effectively promoted by triflic acid and in some cases sulfuric acid. With a variety of electrophilic systems, good regioselectivities are observed in these conversions. Bis(benzocrown ethers) are useful in a variety of applications and this chemistry provides ready access to these systems. In addition to the bis(benzocrown ether) products, we have found that macromolecules can be produced by hydroxyalkylation with dibenzo-18-crown-6 as a substrate.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the support of the NIH-National Institute of General Medical Sciences (GM085736-01A1).

References and notes

1. (a) Pederson CJ. *J Am Chem Soc.* 1967; 89:2495.(b) Pederson CJ. *J Am Chem Soc.* 1967; 89:7017.
2. (a) Hiraoka, M., editor. *Crown Ethers and Analogous Compounds.* Elsevier; Amsterdam: 1992. (b) Kralj M, Tusek-Bozic L, Frkanec L. *Chem Med Chem.* 2008; 3(10):1478. [PubMed: 18683175] (c) Rusalov MV, Uzhinov BM, Alifimov MV, Gromov SP. *Russ Chem Rev.* 2010; 79:1099.(d) Fery-Forgues S, Al-Ali F. *J Photochem Photobiol C Photochem Rev.* 2004; 5:139.
3. (a) Sharghi H, Jokar M, Doroodmand MM, Khalifeh R. *Adv Synth Cat.* 2010; 352(17):3031.(b) O'Connor MJ, Boblak KN, Spitzer AD, Gucciardo PA, Baumann AM, Peter JW, Chen CY, Peter R, Mitton AA, Klumpp DA. *Tetrahedron Lett.* 2010; 51:4984.(c) Kotha S, Brahmachary E. *Indian J Chem Sect B: Org Chem.* 2001; 40B(1):1.(d) Barboiu M, Supuran CT, Luca C, Popescu G, Barboiu C. *Liebigs Ann.* 1996; 6:959.(e) Akkus H, Gul A. *Org Prep Proc Int.* 1995; 27(6):668.(f) Warshawsky A, Kahana N. *J Am Chem Soc.* 1982; 104:2663.
4. (a) Prakash GKS, Panja C, Shakhmin A, Shah E, Mathew T, Olah GA. *J Org Chem.* 2009; 74:8659. [PubMed: 19863094] (b) Alvaro M, Garcia H, Sanjuan A, Espla M. *Appl Catal A.* 1998; 175:105. (c) Koltunov K, Yu, Walspurger S, Sommer J. *Catal Lett.* 2004; 98:89.
5. (a) Prakash GKS, Paknia F, Chacko S, Mathew T, Olah GA. *Heterocycles.* 2008; 76:783.(b) Sheets MA, Li A, Bower EA, Weigel AR, Abbott MP, Gallo RM, Mitton AA, Klumpp DA. *J Org Chem.* 2009; 73:2502. [PubMed: 19215114] (c) Klumpp DA, Zhang Y, Kindelin PJ, Lau S. *Tetrahedron.* 2006; 62:5915.(d) Zhang Y, Klumpp DA. *Tetrahedron Lett.* 2002; 43:6841.(e) Klumpp DA, Jones A, Lau S, DeLeon S, Garza M. *Synthesis.* 2000; 11:17.(f) Klumpp DA, Yeung KY, Prakash GKS, Olah GA. *J Org Chem.* 1998; 63:4481.
6. Gozman-Gutierrez MT, Nieto DR, Fomine S, Morales SL, Zolotukhin MG, Hernandez MCG, Kricheldorf H, Wilks ES. *Macromolecules.* 2011; 44:194., and references cited therein.
7. (a) Hofmann, JE.; Schriesheim, A. *Friedel-Crafts and Related Reaction.* Olah, GA., editor. Vol. 2. Wiley; New York, NY: 1964. p. 597-640.(b) March, J. *Advanced Organic Chemistry.* 4. Wiley; New York, NY: 1992. p. 548-549.
8. Zolotukhin MG, Hernandez MCG, Lopez AM, Fomina L, Cedillo G, Nogales A, Ezquerro T, Rueda D, Colquhoun HM, Fromm KM, Ruiz-Trevino A, Ree M. *Macromolecules.* 2006; 39:4696.
9. Data for polymeric products. **24**: ^1H NMR ($\text{CF}_3\text{CO}_2\text{D}$, 300 MHz): δ = 3.15-3.22, 3.60-3.90, 3.95-4.05, 6.50-6.64, 6.67-6.75, 6.75-6.88, 7.00-7.14, 7.24-7.38. IR: 2914, 2871, 1719, 1520, 1255, 1123, 750, 639 cm^{-1} . **25**: ^1H NMR ($\text{CF}_3\text{CO}_2\text{D}$, 300 MHz): δ = 3.4-3.8, 5.44 and 5.46, 6.10-6.12, 6.36-6.43, 7.15-7.20, 7.34-7.37, 7.93-7.95, 8.05-8.09. IR: 2855, 1506, 1249, 1118, 1052, 944, 751 cm^{-1} .
10. General procedure: The carbonyl substrate (0.1 g) is dissolved in 1-2 mL of anhydrous CHCl_3 and 2.0 equivalents of a benzocrown ether is added to the solution. To this mixture, $\text{CF}_3\text{SO}_3\text{H}$ (8.0 equivalents; H_2SO_4 may be used in some cases) is added dropwise with stirring. The reaction is stirred at room temperature for at least 2 hrs, after which, the mixture is poured over several grams of ice. The resulting solution is extracted three times with CHCl_3 . The organic phase is subsequently washed three times with water and dried over MgSO_4 solution. Removal of the solvent provides the product.

Scheme 1.

Acid-catalyzed reactions of carbonyl compounds (**4-9**) with benzo-18-crown-6 (**2**).

^a Reaction with 2.0 equivalents (**2**), 8 eq TfOH, CHCl₃, 25°C.
^b Reaction with 2.0 equivalents (**2**), 8 eq H₂SO₄, CHCl₃, 25°C.

^a Reaction with 2.0 equivalents (**2**), 8 eq TfOH, CHCl₃, 25°C.

^b Reaction with 2.0 equivalents (2), 8 eq H₂SO₄, CHCl₃, 25°C.

Table 2

Condensations with benzo-15-crown-5 and benzo-12-crown-4.

Starting material	Product	Yield
6	16, R = Ar ₁ 95% 19, R = Ar ₂ 92%
7	17, R = Ar ₁ 77% 20, R = Ar ₂ 65%
9	18, R = Ar ₁ 49% 21, R = Ar ₂ 94%	
<div style="border: 1px solid black; padding: 10px; display: flex; justify-content: space-around; align-items: center;"> <div> r </div> <div> Ar₂ </div> </div>		