Restriction endonuclease *RsrI* from *Rhodobacter sphaeroides*, an isoschizomer of *EcoRI*: purification and properties

Christopher Aiken and Richard I. Gumpert*

Department of Biochemistry, College of Medicine and School of Chemical Sciences, University of Illinois, Urbana, IL 61801, USA

Received May 16, 1988; Accepted July 11, 1988

ABSTRACT

We have purified *RsrI* endonuclease (R-*RsrI*), an isoschizomer of *EcoRI*, from *Rhodobacter sphaeroides* strain 630. The enzyme is homogeneous as judged by polyacrylamide gel electrophoresis and size-exclusion high-performance liquid chromatography. *RsrI* endonuclease is a dimer over the concentration range of 0.05 to 1.4 mg/ml. The reduced and denatured molecular weight of the enzyme is 30,000 Da. R-*RsrI*, like R-*EcoRI*, catalyzes the cleavage of duplex DNA and oligodeoxyribonucleotides between the first two residues of the sequence GAATTC. R-*RsrI* exhibits a K_M of 14 nM and a k_{cat} of 6.5 min^{-1} when reacting with pBR322 DNA at 25°C. R-*RsrI* differs from R-*EcoRI* in its N-terminal amino acid sequence, susceptibility to inhibition by antibodies, sensitivity to N-ethylmaleimide, isoelectric point, state of aggregation at high concentrations, temperature lability, and conditions for optimal reaction. R-*RsrI* displays a reduction of specificity ("star activity") under conditions that also relax the specificity of R-*EcoRI*.

INTRODUCTION

Type II restriction endonucleases are convenient and useful for studying site-specific protein-DNA interactions (see reference 1 for a review). These enzymes are small endodeoxyribonucleases that recognize and cleave DNA at defined palindromic sequences 4 to 8 base pairs long (1-3). Several such enzymes have been purified to homogeneity (1,2), and crystallized (4,5). *EcoRI* endonuclease (R-*EcoRI*) has been extensively characterized, and its structure, when complexed with a synthetic tridecadeoxyribonucleotide containing its recognition sequence, has been determined at high resolution (4,6). *EcoRI*, *BamHI*, *HpaI*, and *HhaII* obey Michaelis-Menten kinetics (7-12), and studies using synthetic oligodeoxyribonucleotides containing base-analogue substitutions have been carried out with *EcoRI* (11-13), *EcoRII* (14), *HpaI* (9), and *EcoRV* (15). *RsrI* endonuclease (R-*RsrI*)² is an isoschizomer of *EcoRI* (16). Here we report the purification of R-*RsrI* from *Rhodobacter sphaeroides* to homogeneity and present an initial characterization of its physical and catalytic properties. The enzyme and its cognate methylase will be useful for comparative studies of the mechanisms of site-specific DNA recognition and catalysis. A preliminary report of some of this work has appeared (17), and Greene and coworkers have reported a purification of the enzyme (18).

MATERIALS

Bacterial Strains

Rhodobacter sphaeroides strain 630 was provided by S. Kaplan (University of Illinois). *E. coli* strains HN493[*thy recA* (pBR322)] and MO were provided by J. Gardner (University of Illinois).

Enzymes and proteins

EcoRI endonuclease and methylase were gifts of P. Modrich, as was rabbit antiserum specific for *EcoRI* endonuclease. T4 polynucleotide kinase was a gift from D. Soltis and O. Uhlenbeck (University of Colorado at Boulder). ATP-dependent DNase (*Micrococcus lysodeikticus* Exo V) and HPLC gel-filtration markers were from United States Biochemical Corporation. Protein standards for SDS-polyacrylamide gel electrophoresis were from Bio-Rad. Isoelectric focusing protein standards were from Pharmacia.

DNA

Salmon sperm and calf thymus DNAs were from Calbiochem. Bacteriophage lambda was purified from phage-infected *E. coli* MO by a modification of the method of Gardner and Reznikoff (19), and the DNA purified using the method of Maniatis et. al. (20). ³H-labeled pBR322 DNA (10⁴-10⁵ cpm/μg) was isolated from *E. coli* HN493 after chloramphenicol amplification of the plasmid. Cells were grown at 37°C in 800 ml of M9CA medium (21) supplemented with 0.1 μg/ml of thiamine and 5 μg/ml of thymidine. At mid-log phase the culture was centrifuged at 12,000 xg for 20 min and the cells resuspended in fresh medium. After shaking for 10 min at 37°C, chloramphenicol was added to a final concentration of 150 μg/ml, followed by the addition of [³H]-thymidine (5.5 Ci/ml, 50 Ci/mmol, ICN Biomedicals, Inc.). Following overnight growth at 37°C, the plasmid DNA was purified by the alkaline lysis method of Maniatis et. al. (22) followed by isopycnic centrifugation in CsCl containing ethidium bromide. Ethidium bromide was removed by extraction with isoamyl alcohol. The isolated DNA was at least 95% in the covalently closed-circular form and was free of contaminating RNA. Oligonucleotides were synthesized on an Applied Biosystems 380A DNA synthesizer and were purified by reverse phase HPLC (23) or polyacrylamide gel electrophoresis. The self-complementary decamer d(CTGAATTCAG) was a gift of L. McLaughlin (Boston University). Oligonucleotides were labeled using polynucleotide kinase and [γ-³²P]ATP (>7000 Ci/mmol, New England Nuclear).

Other Materials

DEAE-cellulose (Whatman DE52) was equilibrated according to the manufacturer's instructions. Cellulose phosphate (Sigma C2383) was precycled and washed using the method of Greene et. al. (24). DNA-cellulose was prepared from Whatman CF-11 cellulose and native calf thymus DNA by the method of Alberts and Herrick (25). Agarose was from International Biotechnologies, Inc., and Polymin P (50% solution) was purchased from Miles Laboratories. Other chemicals were reagent grades.

METHODS

Growth of cells

R. sphaeroides 630 cells were grown aerobically at 30°C in 200 liters of Siström's minimal medium (26) containing 0.2% casamino acids in a Braun model 300D fermentor. The culture pH was maintained at 7.2 using phosphoric acid. At late-log to early-stationary phase, the culture was chilled to 10°C and harvested using a refrigerated Sharples centrifuge. The paste (800-1400 g) was frozen at -70°C and showed no loss of *RsrI* activity after 8 months.

Gel Electrophoresis

Enzyme purity and the reduced and denatured molecular weight were determined by polyacrylamide gel electrophoresis in the presence of SDS. Slab gels were prepared by the method of Hames (27), and the proteins were visualized by silver-staining (28) or by staining with Coomassie blue. Column fractions were qualitatively assayed for *RsrI* endonuclease activity with lambda DNA using agarose gel electrophoresis. Lambda DNA (0.5 µg) was incubated with enzyme in 20 µl of 50 mM Tris-HCl (pH 7.8), 10 mM MgCl₂, and 1 mM dithiothreitol (DTT). After 10 minutes at 23°C, 4 µl of gel-loading dye (0.1% bromophenol blue, 0.1% xylene cyanol, 40% sucrose) were added and the reactions heated to 65°C for 5 min prior to loading on a 1% agarose gel. Electrophoresis was performed in a buffer of 40 mM Tris-acetate (pH 8.3), 1 mM EDTA. The gel was stained with ethidium bromide and photographed using a Fotodyne transilluminator and Polaroid MP-4 camera with type 667 film using Fotodyne filters 3 and CC20Y. Isoelectric focusing in thin-layer polyacrylamide gels was carried out at 15°C using Ampholine PAG 5.5-8.5 plates with a Multiphor electrophoresis apparatus from LKB. The enzyme was desalted and concentrated using a Centricon 10 microconcentrator (Amicon), and samples of approximately 10 µg were loaded onto the gel with marker proteins. The samples were focused at 12 W constant power for 2.5 hours, the final voltage being 1600 V. Slices (1 cm) were cut from the gel and placed at 4°C in 2 ml of distilled water overnight, and the pH values measured to determine the pH gradient. One lane containing *RsrI* endonuclease was removed, and slices (1 cm) from this lane were soaked in *RsrI* dilution buffer overnight. These samples were subsequently assayed for endonuclease activity. The remainder of the gel was fixed with trichloroacetic acid (TCA) and stained with Coomassie blue. Electrophoresis under non-denaturing conditions was performed using a modification of the methods of Bryan (29) and Davis (30) as described in the Sigma Technical Bulletin No. MKR-137. Protein standards were purchased from Sigma, and polyacrylamide gels of 7%, 8%, 9%, and 10% acrylamide were used.

RsrI Endonuclease Reactions

The quantitative *RsrI* endonuclease assay is a modification of the method of Modrich and Zabel (31) for measuring *EcoRI* endonuclease activity. Reactions (50 µl) contained 50 mM Tris-HCl (pH 7.8), 10 mM NaCl, 10 mM MgCl₂, 10 µg/ml BSA, 1 mM DTT, and 1 pmol [³H]pBR322 (2.88 µg, 25,000 cpm/µg), and were initiated by the addition of *RsrI* endonuclease (5 µl)

diluted as necessary with 20 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 0.1 M NaCl, 10% glycerol, 0.1 mg/ml BSA, and 0.1 mM DTT (*RsrI* dilution buffer). Following incubation for 10 min at 23°C, 50 μ l of ice-cold twice-concentrated (2X) ATP-dependent DNase assay buffer [133 mM glycine-NaOH (pH 9.4), 60 mM MgCl₂, 16.6 mM 2-mercaptoethanol, 1 mM ATP] containing 0.5 unit of ATP-dependent DNase were added. Following incubation for 30 min at 37°C, 25 μ l of 1 mg/ml salmon sperm DNA and 25 μ l of 30% TCA were added. After 10 min on ice, the precipitate was removed by centrifugation at 14,000 xg for 10 min and an aliquot of the supernatant counted for ³H in Liquiscint (National Diagnostics). One unit of *RsrI* endonuclease converts 1 pmol of pBR322 DNA per minute at 23°C to a form which is sensitive to ATP-dependent DNase, and the assay was linear for 10 min with up to 0.050 unit of endonuclease. Because ATP-dependent DNase cleaves only at the termini of linear duplexes (32), the assay does not detect nicked DNA.

Other Methods

Size-exclusion chromatography was carried out at room temperature on a Beckman model 420 HPLC system using a 0.75 x 30 cm TSK G3000SW column (Altex). Proteins were eluted with Buffer A (as described below, but prepared at room temperature) and detected by their absorption at 280 nm. Radioisotopes were quantified by liquid scintillation counting using a Nuclear Chicago Isocap 300.

Protein was determined by the method of Bradford (33) using a kit from Bio-Rad. Lysozyme was used as the standard and its concentration was determined spectrophotometrically using $E^{0.1\%}_{281.5} = 2.64$ in 0.1 M KCl (34).

All steps in the purification of *RsrI* endonuclease, except for HPLC fractionations, were carried out at 0-4°C, and centrifugations were at 12,000 xg for 30 min.

Buffers

All buffer pH values were determined at the temperatures and concentrations at which they were used. Buffer A (4°C) was 20 mM Tris-HCl (pH 7.5), 10% (w/v) glycerol, 1 mM EDTA, and 10 mM 2-mercaptoethanol. Buffer B (4°C) was 20 mM potassium phosphate (KPi) pH 7.4, 10% (w/v) glycerol, 1 mM EDTA, and 10 mM 2-mercaptoethanol. Buffer C was identical to buffer B, but was prepared at 23°C.

RESULTS AND DISCUSSION

Purification of *RsrI* Endonuclease

Crude Extract and Polymin P Fractionation. *R. sphaeroides* 630 cell paste (100 g) was thawed overnight at 4°C and suspended in 300 ml of buffer A containing 0.5 M KCl and 0.1 mM phenylmethylsulfonylfluoride. Cells were lysed by three passes through a precooled French pressure cell at 16,000-20,000 lb/in², with the temperature of the cell suspension maintained below 30°C. The lysate was centrifuged, and to the supernatant (Fraction I, 370 ml) were added, dropwise with stirring over 10 min, 26 ml of an aqueous solution of 10% Polymin P (titrated to

pH 7.5 with HCl). The extract was stirred an additional 20 min and centrifuged. To the supernatant (Fraction II, 340 ml), ammonium sulfate (175 g) was added over 20 min to give an 80% saturated solution. The mixture was stirred an additional 20 min and centrifuged. The *RsrI* endonuclease activity in the pellet was stable at 4°C for at least 3 months.

DEAE-Cellulose Chromatography. The ammonium sulfate pellet was redissolved in 200 ml of buffer A and dialyzed against 1 liter of the same buffer using an Amicon CH2 spiral cartridge apparatus in the diafiltration mode. The dialysate (215 ml) was applied at a flow rate of 4 ml/min to a 240 ml column (3.9 cm x 20.4 cm) of DE52 previously equilibrated in buffer A. After washing with 600 ml of buffer A, the enzyme was eluted with a 2-liter linear gradient of 0 to 0.5 M KCl in buffer A. Fractions containing *RsrI* endonuclease activity, which eluted between 0.05 and 0.15 M KCl, were pooled (Fraction III, 192 ml). *RsrII* endonuclease (35) and *RsrI* methylase coeluted with *RsrI* endonuclease from this column. *RsrI* endonuclease in Fraction III was stable for at least two weeks when stored at 4°C.

Phosphocellulose Chromatography. Fraction III was dialyzed against 1 liter of buffer B using the spiral cartridge system. The dialysate was applied at a flow rate of 1 ml/min to a 45 ml column (1.9 cm x 15.4 cm) of phosphocellulose equilibrated in buffer B. The column was washed with 180 ml of buffer B and subsequently eluted with a 400 ml linear gradient of 0 to 0.8 M KCl in buffer B. Fractions containing *RsrI* endonuclease activity, which eluted between 0.15 and 0.27 M KCl, were pooled (Fraction IV, 42 ml). This step resolves *RsrI* endonuclease from *RsrII* endonuclease and *RsrI* methylase. In some purifications, *RsrII* endonuclease was also resolved from *RsrI* methylase, and can be concentrated and stored in 50% glycerol at -20°C. The *RsrII* endonuclease at this level of purity seems suitable for producing restriction digests because of the absence of apparent nonspecific nuclease activity when *RsrII* digests of lambda DNA were separated and visualized.

DNA-Cellulose Chromatography. Fraction IV was equilibrated twice, using conventional dialysis, with 500 ml of buffer B and applied at a flow rate of 0.5 ml/min to a 10 ml (1.5 cm x 6.2 cm) double-strand DNA-cellulose column equilibrated in buffer B. After washing with 40 ml of buffer B, the enzyme was eluted with a 100 ml linear gradient of 0 to 0.5 M NaCl in buffer B. Fractions containing *RsrI* endonuclease activity, which eluted between 0.18 and 0.28 M NaCl, were pooled (Fraction V, 9.6 ml).

Hydroxylapatite Chromatography. Fraction V was concentrated and desalted using a Centriprep-10 concentrator (Amicon). The sample (0.65 ml) was applied at a flow rate of 1 ml/min to a Regis KB hydroxylapatite HPLC column (0.6 cm x 10 cm) equilibrated in buffer C. The column was washed for 7 min, and eluted with a 40 ml linear gradient of 20 to 300 mM KP_i in buffer C. *RsrI* endonuclease eluted at 0.1 M KP_i . Fractions which were >95% pure as judged by SDS-polyacrylamide gel electrophoresis followed by silver staining were pooled, concentrated, and made 50% (v/v) in glycerol. The purified enzyme (Fraction V, 2 ml) was stored at -20°C. A summary of the purification is presented in Table I. This purification scheme for *RsrI*

Table I. Purification of *RsrI* Endonuclease from 100 g of *R. sphaeroides*

Fraction	Total Protein (mg)	Total Activity (U x 10 ⁻³)	Specific Activity (units/mg)	Yield (%)	Fold Purification
I. French Press	3,850	294	76	(100)	--
II. Polymyxin P	2,580	311	120	106	1.6
III. DEAE-cellulose	161	152	945	52	12
IV. Phosphocellulose	16.2	108	6,710	37	88
V. DNA-cellulose	1.0	72	71,400	24	940
VI. Hydroxylapatite HPLC	0.17	19	113,000	6.5	1,490

endonuclease is reproducible and yields homogeneous enzyme in modest amounts. After several attempts using conventional chromatography, we found that hydroxylapatite HPLC gave the additional purification needed to achieve homogeneity. The recovery from this column in the purification we reported was low (26%), but in another purification better yields were obtained (400 μ g, 67% recovery). It should be possible to scale up this purification, since preparative HPLC hydroxylapatite columns are available. In some cases, we have successfully employed chromatography on a Waters Protein-Pak DEAE 5PW column as an additional purification step. The amount of *RsrI* endonuclease in *R. sphaeroides* 630 depends on the growth conditions. O'Connor found that *RsrI* activity was maximal in cells grown in rich medium and harvested in late-log to early-stationary phase (36). We used Siström's medium to reduce the amount of foaming in the fermentor and to facilitate growing starter cultures of *R. sphaeroides* 630. We found that adding casamino acids (0.2%) to Siström's medium enhanced the yield of cells, and doubled the amount of *RsrI* endonuclease and methylase per gram of cell paste. Assuming a molecular weight of 30,000 (see below) and a specific activity for the homogeneous enzyme of 113,000 units/mg, we estimate that there are approximately 500-2000 monomer equivalents of *RsrI* endonuclease per cell. This is comparable to the value of 2000 estimated for *EcoRI* endonuclease in *E. coli* RY13 (31), a strain which carries the *EcoRI* endonuclease and methylase genes on a derepressed R factor (24). An attractive feature of the purification described here is that *RsrI* methylase copurifies with the endonuclease through the first chromatographic step of the procedure and is resolved from it at the following step. We have purified the *RsrI* methylase from this starting material (Kaszubska and Gumpert, unpublished results) and are examining its properties.

Physical Properties

Molecular Weight Determinations. The subunit molecular weight of the purified enzyme was determined by electrophoresis on 10% and 12.5% SDS-polyacrylamide slab gels, and the

Figure 1. Electrophoresis of purified *RsrI* endonuclease under denaturing conditions. Upper panel, 1.7 μ g of *RsrI* endonuclease (Fraction VI) was subjected to electrophoresis on a gel of 15% polyacrylamide containing SDS. The gel was stained with Coomassie brilliant blue, and scanned using an LKB scanning laser densitometer. Marks on the abscissa represent 1-cm distances along the gel. Lower panel, determination of reduced and denatured molecular weight of *RsrI* endonuclease by electrophoresis on calibrated SDS-polyacrylamide gels. Gels of 10% and 12% acrylamide were used.

proteins visualized by staining with Coomassie blue. A single band of $M_r = 30,000 \pm 2000$ was observed (Figure 1). *R-RsrI* comigrates with *R-EcoRI* on these gels. Subunit molecular weights of 30,000 Da (24) and 28,500 Da (31) have been reported for *R-EcoRI*; calculation of the molecular weight from the deduced amino acid sequence gave a value of 31,065 (37,38). Thus, *RsrI* and *EcoRI* endonucleases are indistinguishable on the basis of their reduced and denatured molecular weights. The molecular weight of the purified enzyme under non-denaturing conditions was determined by high performance size-exclusion chromatography. *RsrI* endonuclease eluted from the column as a single peak in the volume expected for a globular protein with a molecular weight of $59,600 \pm 2000$ Da (Figure 2). The endonuclease activity profile coincided with that of the protein. The fractions containing activity were examined by SDS-polyacrylamide gel electrophoresis and each displayed a single band migrating at the position of a protein with an $M_r = 30,000$. Three repetitions of this chromatography yielded

Figure 2. Determination of the molecular weight of native *RsrI* endonuclease by size-exclusion HPLC. Samples of *RsrI* endonuclease (7 to 28 μ g) were applied to a TSK G3000SW HPLC column as described in Methods. Protein standards were analyzed under identical conditions. The void volume of the column was determined using blue dextran. The elution time of the endonuclease was identical when it was co-injected with the standard proteins. The parameter K, as plotted on the ordinate, is a function of the elution volume, and was calculated as described (47).

essentially identical results. Using a different set of gel filtration standards (Bio-Rad), we obtained a value of $51,700 \pm 2000$ Da for the molecular weight (data not shown). Electrophoresis of the native enzyme on a series of polyacrylamide slab gels of differing acrylamide concentrations under nondenaturing conditions yielded a value of $77,000 \pm 4000$ Da for the molecular weight of *RsrI* endonuclease (Figure 3). A single band was consistently observed on these gels.

Collectively, these results suggest that R-*RsrI* exists as a dimer in solution over the range of concentrations from 0.05 to 1.4 mg/ml, and that the dimer may be the active form of the enzyme. R-*EcoRI* exists as an equilibrium mixture of dimers and tetramers, with the tetrameric form predominating at concentrations greater than 25 μ g/ml (31). We have not tested the possibility that R-*RsrI* also forms tetramers at higher concentrations than those examined. If so, the dissociation constant for the tetramer-dimer equilibrium for *RsrI* would be greater than the value of 1×10^{-7} M reported for R-*EcoRI* (31).

Isoelectric Point. The isoelectric pH of *RsrI* endonuclease was determined by focusing on thin-layer polyacrylamide gels. A single major protein and two closely associated bands of lesser intensity were observed. Each of these bands contained *RsrI* endonuclease activity, in approximately the same proportion as its relative intensity (data not shown). The pI value of the major protein band was 7.0 ± 0.1 (Figure 4), and of the minor protein bands 6.8 and 6.6.

Figure 3. Determination of the molecular weight of *Rsr*I endonuclease by electrophoresis on polyacrylamide gels under nondenaturing conditions. The slope value on the ordinate is that of the curve of the electrophoretic mobility versus acrylamide gel concentration plotted for each standard protein.

*Eco*RI endonuclease, focused on the same gel, was a single band with a pI of 6.4 ± 0.1 . This is in agreement with the value of 6.3 reported by Rubin and Modrich (39).

Amino Acid Sequence. Determination of the sequence of the amino-terminus of R-*Rsr*I was performed on an Applied Biosystems 470A Protein Sequenator. Phenylthiohydantoin(PTH)-amino acids were identified using an on-line Applied Biosystems 120A PTH Analyzer. Tryptophan and cysteine residues were not determined. A second determination of the first 29 amino acids using a different preparation of R-*Rsr*I confirmed the results. As shown in Table II, the sequence of the first 40 amino acids of R-*Rsr*I is considerably different from that of R-*Eco*RI. Greene and coworkers have shown that appropriate alignment of these two sequences

Figure 4. Isoelectric focusing of *Rsr*I endonuclease. The closed circles represent the pI values of the protein standards; open circles correspond to the pH values of the isolated gel slices, and are plotted as occurring at the center of each 1-cm segment.

Table II. N-Terminal Amino Acid Sequences^a

<i>RsrI</i> Endonuclease
ALA-GLY-GLU-VAL-GLU-PHE-LYS-GLY-LYS-GLY-GLN-ALA-LEU-ARG-LEU-GLY- ILE-GLN-GLN-(N.D.) ^b -LEU-GLY-GLY-GLY-PRO-LEU-SER-ILE-PHE-GLY-ALA-LYS- ALA-GLN-LYS-(N.D.) ^b -ALA-LEU-PRO-ALA
<i>EcoRI</i> Endonuclease ^c
(fMET)-SER-ASN-LYS-LYS-GLN-SER-ASN-ARG-LEU-THR-GLU-GLN-HIS-LYS-LEU- SER-GLN-GLY-VAL-ILE-GLY-ILE-PHE-GLY-ASP-TYR-ALA-LYS-ALA-HIS-ASP-LEU- ALA-VAL-GLY-GLU-VAL-SER-LYS-LEU

^aFirst 40 amino acids of amino terminus of each protein.

^bThe amino acid at this position was not identified.

^cAs deduced from the sequence of the *R.EcoRI* gene (37,38). The fMet is not present in the purified protein.

reveals 11 amino acid identities in 28 residues (17). We also tested Southern blots of *R. sphaeroides* 630 plasmid and chromosomal DNA for cross-hybridization using a labeled DNA restriction fragment containing the *EcoRI* endonuclease and methylase genes as a probe. Hybridization and washing was performed at 42°C according to the procedure of Meinkoth and Wahl (40). No hybridization was observed, suggesting that the DNA sequences of the *RsrI* and *EcoRI* endonuclease genes are different. Rabbit antiserum to *R.EcoRI* was used to test for cross-reactivity with *RsrI* endonuclease. By incubating dilutions of the antiserum with nearly equal activities of *R.RsrI* and *R.EcoRI*, we found that the antiserum was 1000 times less reactive toward *R.RsrI* than toward *R.EcoRI* as judged by inhibition of endonuclease activity (data not shown).

Taken together, these results demonstrate that although *RsrI* and *EcoRI* endonucleases have identical apparent subunit molecular weights and both recognize and cleave the same sequence (see below), they differ considerably in the sequences of their first 40 amino acids and show differences at the DNA sequence level. In addition, they exhibit different isoelectric points, and polyclonal antibodies distinguish between the two endonucleases.

Catalytic Properties

Determination of Recognition Site. *RsrI* and *EcoRI* endonucleases were incubated with the self-complementary dodecadeoxyribonucleotide [5'-³²P]d(pGTAGAATTCTAC), and the products separated by one-dimensional homochromatography (41). Hydrolysis with either enzyme resulted in the formation of the labeled tetranucleotide [5'-³²P]d(pGTAG) (Figure 5). A similar experiment using [5'-³²P]d(pCTGAATTCAG) resulted in the formation of the trinucleotide [5'-³²P]d(pCTG). No other labeled products were formed with these substrates, indicating the

Figure 5. Cleavage of oligonucleotide [5'-³²P]d(pGTAGAATTCTAC). The labeled dodecamer (0.2 μ M) was incubated with R-*RsrI* (5 nM) or R-*EcoRI* (5 nM) for 5 min at 20°, and samples were analyzed by homochromatography as described (41). Lane 1, no enzyme; lane 2, R-*EcoRI*; lane 3, R-*RsrI*.

absence of phosphatases and nonspecific endo- and exonucleases. Incubation of lambda DNA with the endonuclease followed by agarose gel electrophoresis resulted in a distinct DNA fragment pattern identical to that obtained by digestion with *EcoRI* endonuclease. A restriction digest in which both enzymes were present yielded no additional product bands. R-*RsrI* cleaved pBR322 at a single site. In addition, electrophoresis of a 5'-³²P-labeled DNA restriction fragment cleaved by R-*RsrI* alongside the same fragment subjected to Maxam-Gilbert sequencing reactions showed that the R-*RsrI*-cleaved fragment had been cut between the G and A residues at the *EcoRI* site (S. Lynn and J. Gardner, personal communication). Taken together, these data demonstrate that R-*RsrI* recognizes and cleaves the hexanucleotide site GAATTC in duplex DNA between the G and A residues.

Lambda DNA that had been methylated *in vivo* or *in vitro* by *EcoRI* methylase or *in vitro* by *RsrI* methylase was resistant to cleavage by both *RsrI* and *EcoRI* endonucleases (data not shown). Because both methylases catalyze the transfer of a methyl group to the central adenine residue of the sequence GAATTC (42-44; Kaszubska and Gumpert, unpublished results), methylation of this residue renders the site refractory to cleavage by R-*RsrI*.

Reaction Optimization. Using the quantitative assay (see Methods), R-*RsrI* activity was maximal in a solution containing 50 mM Tris-HCl (pH 7.8), 10 mM MgCl₂, 10 mM NaCl, and 1 mM DTT at a temperature of 23°C. A reaction containing 35 mM NaCl was 80% as active. Higher NaCl concentrations further inhibited the reaction. The pH optimum of the reaction was between 7.5 and 8.1. The reaction required Mg⁺² or Mn⁺². At 30°C, the reaction rate was 60% of the rate at 23°C. This was probably not due to a pH effect, since reaction buffers were prepared at

each temperature tested. The endonuclease was inactive in reactions at 37°C and 45°C. To test whether R-*RsrI* was being irreversibly inactivated at temperatures above 23°C, a series of assays was performed in which the endonuclease was incubated in the absence of DNA in reaction buffers at various temperatures for 10 min. An aliquot of each sample was then removed and assayed at 23°C. Pre-incubation at 23°C had no effect on the activity of the endonuclease, while pre-incubation at 37°C and 45°C for 10 min reduced the activity by 50% and 70%, respectively. Thus, under our assay conditions, the enzyme is unstable at temperatures above 30°C. R-*EcoRI* is routinely assayed at 37°C, and the reaction is stimulated two- to threefold by raising the NaCl concentration to 50 to 100 mM (24). Thus, R-*RsrI* is less stable than R-*EcoRI*, and requires lower ionic strength buffers for optimal activity. Incubation of R-*RsrI* in dilution buffer containing 10 mM N-ethylmaleimide at 15°C for 15 min completely eliminated activity, while identical treatment of *EcoRI* endonuclease resulted in less than 5% inhibition of activity. *EcoRI* methylase is sensitive to this reagent, while *EcoRI* endonuclease is not (38). Thus, unlike R-*EcoRI*, R-*RsrI* appears to contain an essential cysteine residue.

Kinetic Analysis. Steady-state kinetic constants for double-strand cleavage were determined at 25°C with ³H-labeled pBR322 DNA. The reaction shows Michaelis-Menten kinetics--a Lineweaver-Burk plot of the data was linear and yielded values of 14 nM and 6.5 min⁻¹ for K_M and k_{cat} , respectively (Figure 6). These values are similar to those reported by Modrich and Zabel (31) for R-*EcoRI* with ColE1 DNA at 37°C (8 nM and 3.8 min⁻¹). We have observed nicking of supercoiled pBR322 DNA in reactions analyzed by agarose gel electrophoresis. The

Figure 6. Steady-state kinetic analysis of *RsrI* endonuclease with pBR322 DNA at 25°C. Assays were performed under standard conditions in 250 μ l reactions containing 35 pM R-*RsrI* and 1.3 to 10 nM [³H]pBR322. Samples (50 μ l) were analyzed at 2-min intervals as described in Methods. The initial velocity was determined from the slope of the curve of a plot of product against time for each DNA concentration. Product was determined from the extent of cleavage obtained as the percent of a control reaction containing excess R-*RsrI*, which went to completion.

nicked product can be converted subsequently into a linear molecule by the enzyme. These results suggest a sequential strand cleavage mechanism analogous to that observed for R-*EcoRI* (44).

Star Activity. R-*RsrI* cleaves DNA at sites other than its canonical recognition sequence in a buffer of 12.5 mM Tris-HCl (pH 8.5), 15 mM NaCl, 2.5 mM MgCl₂, and 26% ethylene glycol at 25°C. Substitution of Mn²⁺ for Mg²⁺ in the same buffer resulted in a relaxation of specificity. Under these conditions, *EcoRI* endonuclease also exhibits a relaxation of specificity (45), which has been referred to as "star" activity (46). We observed star activity in reactions containing equal activities (defined for each enzyme using the quantitative assay for R-*RsrI* described in Methods) of either enzyme (Figure 7). Distinct and smaller bands were produced from the linear product arising from the cleavage of the unique canonical site in pBR322 DNA (Figure 7, lanes 2-4 and 5-7). Under these circumstances, R-*EcoRI* was more active at cleaving pBR322 DNA at noncanonical sites than was R-*RsrI*, and additional fragments were produced with R-*EcoRI*. The patterns of fragments produced by the enzymes consisted primarily of common fragments; however, there were some bands in the R-*EcoRI* digest of pBR322 DNA which never appeared in the R-*RsrI* digest. This suggests that the enzymes cleaved an overlapping set of noncanonical sites. The cleavage patterns using lambda DNA under these conditions appeared generally similar with both enzymes, but also contained bands unique to each enzyme. Thus, both

Figure 7. Cleavage of DNA by R-*RsrI* and R-*EcoRI* under conditions which produce star activity. Lane 1: R-*RsrI* incubated with pBR322 DNA under standard assay conditions. Lanes 2-4: 25, 12, and 6 units of R-*RsrI* incubated with 1 µg of pBR322 DNA for 10 h at 20°C under conditions of high pH, low ionic strength, and 26% ethylene glycol (see Results). Lanes 5-7: 25, 12, and 6 units of R-*EcoRI* incubated with 1 µg of pBR322 DNA for 10 h at 20°C under the same conditions as lanes 2-4. Lane 8: R-*RsrI* incubated with lambda DNA under standard assay conditions. Lanes 9-11: same as lanes 2-4, but with lambda DNA. Lanes 12-14: same as lanes 5-7, but with lambda DNA.

enzymes appear to cleave many of the same non-canonical sequences, but may also cleave some different sites or cleave these sites at different rates.

CONCLUSIONS

We have purified the *RsrI* endonuclease to apparent homogeneity and examined some of its properties. The purification is reproducible and provides from 0.15 to 0.40 mg of enzyme per 100 g of cell paste. The preparation is free of contaminating nuclease and phosphatase activities and is suitable for detailed mechanistic analysis. *R-RsrI* is an isoschizomer of *R-EcoRI*, and cleaves DNA at the same phosphodiester bond. *R-RsrI* also exhibits a reduction of specificity under conditions that relax the specificity of *R-EcoRI*. We have found that *RsrI* endonuclease differs from its *E. coli* counterpart in its N-terminal amino acid sequence, isoelectric point, inhibition by antibodies, sensitivity to N-ethylmaleimide, and optimal reaction components and temperature. The availability of pure *RsrI* and *EcoRI* endonucleases and methylases will allow comparison of the mechanisms by which these four enzymes interact with their common DNA recognition sequence. We are presently examining the steady-state kinetic behavior of the *RsrI* endonuclease with the same base-analogue substituted oligodeoxyribonucleotides that were used with *EcoRI* endonuclease (11,12).

ACKNOWLEDGEMENTS

We thank K. Milarski for assistance with the purification, Saw Kyin for protein sequencing, A. Jonas for the use of the TSK G3000SW column, J. Gardner and S. Kaplan for bacterial strains, and L. McLaughlin and P. Modrich for gifts and helpful discussions. We also thank P. Greene and coworkers for sharing their results with us prior to their publication. This work was supported in part by NIH Grant GM25621 and by a grant from the Earl M. Bane Charitable Trust of the University of Illinois College of Medicine. C. A. was supported in part by USPHS Training Grant GM07283-11.

*To whom correspondence should be addressed

Abbreviations: *R-RsrI*, *RsrI* endonuclease; *R-EcoRI*, *EcoRI* endonuclease; d(pNNN), d(pN-N-N); SDS, sodium dodecyl sulfate; DTT, dithiothreitol; TCA, trichloroacetic acid; HPLC, high performance liquid chromatography.

REFERENCES

1. Modrich, P. (1982) Crit. Rev. Biochem. **13**, 287-323.
2. Malcolm, A. D. B. and Snounou, G. (1987) In Neidle, S. (ed), Topics in Molecular and Structural Biology, Macmillan Press, London, Vol IX, pp. 193-222.
3. Blakesley, R. W. (1987) In Chirikjian, J. G. (ed), Gene Amplification and Analysis, Elsevier Science Publishing Co., New York, Vol. V, pp. 51-102.
4. McClarin, J. A., Frederick, C. A., Wang, B.-C., Greene, P., Boyer, H. W., Grable, J., and Rosenberg, J. M. (1986) Science **234**, 1526-1541.
5. D'Arcy, A., Brown, R. S., Zabeau, M., van Resandt, R. W., and Winkler, F. K. (1985) J. Biol. Chem. **260**, 1987-1990.

6. Frederick, C. A., Grable, J., Melia, M., Samudzi, C., Jen-Jacobsen, L., Lesser, D., Greene, P. J., Boyer, H. W., Itakura, K., and Rosenberg, J. M. (1984) *Nature* **309**, 327-331.
7. Greene, P. J., Poonian, M. S., Nussbaun, A. L., Tobias, L., Garfin, D. E., Boyer, H. W., and Goodman, H. M. (1975) *J. Mol. Biol.* **99**, 237-261.
8. Nardone, G. and Chirikjian, J. G. (1987) In Chirikjian, J. G. (ed), *Gene Amplification and Analysis*, Elsevier Science Publishing Co., New York, Vol. V., pp. 147-184.
9. Dwyer-Hallquist, P., Kezdy, Ferenc. J., and Agarwal, K. L. (1982) *Biochemistry* **21**, 4693-4700.
10. Kaddurah-Daouk, R., Cho, P., and Smith, H. O. (1985) *J. Biol. Chem.* **260**, 15345-15351.
11. Brennan, C. A., Van Cleve, M. D., and Gumpport, R. I. (1985) *J. Biol. Chem.* **261**, 7270-7278.
12. McLaughlin, L. W., Benseler, F., Graeser, E., Piel, N., and Scholtissek, S. (1987) *Biochemistry* **26**, 7238-7245.
13. Seela, F. and Kehne, A. (1987) *Biochemistry* **26**, 2232-2238.
14. Yolov, A. A., Vinogradova, M. N., Gromova, E. S., Rosenthal, A., Cech, D., Veiko, V. P., Metelev, V. G., Kosykh, V. D., Burynov, Y. I., Bayev, A. A., and Shabarova, Z. A. (1985) *Nucleic Acids Res.* **13**, 8983-8998.
15. Fliess, A., Wolfes, H., Rosenthal, A., Schwellnus, K., Blocker, H., Frank, R., and Pingoud, A. (1986) *Nucleic Acids Res.* **14**, 3643-3473.
16. Gardner, J. F., Cohen, L. K., Lynn, S. P., and Kaplan, S., personal communication.
17. Aiken, C., Milarski, K., and Gumpport, R. I. (1986) *Fed. Proc.* **45**, 1914.
18. Green, P. J., Ballard, B. T., Stephenson, F., Kohr, W. J., Rodriguez, H., Rosenberg, J. M., and Boyer, H. W. (1988) *Gene*, in press.
19. Gardner, J. F. and Reznikoff, W. S. (1978). *J. Mol. Biol.* **126**, 241-258.
20. Maniatis, T., Fritsch, E. F., and Sambrook, J. (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, p. 85.
21. Maniatis, T., Fritsch, E. F., and Sambrook, J. *ibid.*, pp. 440-441.
22. Maniatis, T., Fritsch, E. F., and Sambrook, J. *ibid.*, pp. 90-91.
23. Brennan, C. A. and Gumpport, R. I. (1984) *Nucleic Acids Res.* **13**, 8665-8684.
24. Greene, P. J., Betlach, M. C., and Boyer, H. W. (1974) In Wickner, R. B. (ed), *DNA Replication*, Marcel Dekker, Inc., New York, Vol. VII, 87-105.
25. Alberts, B. and Herrick, G. (1971) *Methods Enzymol.* **21**, 198-217.
26. Siström, W. R. (1982) *J. Gen. Microbiol.* **28**, 607-616.
27. Hames, B. D. (1981) In Hames, B. D. and Rickwood, D. (eds), *Gel Electrophoresis of Proteins: A Practical Approach*, IRL Press Limited, London and Washington DC, pp. 1-91.
28. Merrill, C. R., Goldman, D., Sedman, S. A., and Ebert, M. H. (1981) *Science* **211**, 1437-1438.
29. Bryan, J. K. (1977) *Anal. Biochem.* **78**, 513-519.
30. Davis, B. J. (1984) *Ann. NY Acad. Sci.* **121**, 404-427.
31. Modrich, P. and Zabel, D. (1976) *J. Biol. Chem.* **251**, 5866-5874.
32. Takagi, Y., Matsubara, K., and Anai, M. (1972) *Biochem. Biophys. Acta* **269**, 347-353.
33. Bradford, M. M. (1978) *Anal. Biochem.* **86**, 142-146.
34. Aune, K. C. and Tanford, C. (1969) *Biochemistry* **8**, 4579-4585.
35. O'Connor, C. D., Metcalf, E., Wrighton, C. J., Harris, T. J. R., and Saunders, J. R. (1984) *Nucleic Acids Res.* **12**, 6701-6708.
36. O'Connor, C. D., Walker, J. N. B., and Saunders, J. R. (1987) *Methods Enzymol.* **155**, 11-15.
37. Greene, P. J., Gupta, M., Boyer, H. W., Brown, W. E., and Rosenberg, J. M. (1981) *J. Biol. Chem.* **256**, 2143-2153.
38. Newman, A. K., Rubin, R. A., Kim, S.-H., and Modrich, P. (1981) *J. Biol. Chem.* **256**, 2131-2139.
39. Rubin, R. A. and Modrich, P. (1977) *J. Biol. Chem.* **252**, 7265-7272.
40. Meinkoth, J. and Wahl, G. (1984) *Anal. Biochem.* **138**, 267-284.

41. Jay, E., Bambara, R., Padmanabhan, R., and Wu, R. (1974) *Nucleic Acids Res.* **1**, 331-353.
42. Hedgepeth, J., Goodman, H. M., and Boyer, H. W. (1972) *Proc. Natl. Acad. Sci. U.S.A.* **69**, 3448-3452.
43. Dugaiczyk, A., Hedgepeth, J., Boyer, H. W., and Goodman, H. M. (1974) *Biochemistry* **13**, 503-512.
44. Rubin, R. A. and Modrich, P. (1978) *Nucleic Acids Res.* **5**, 2991-2997.
45. Jen-Jacobsen, L., Kurpiewski, M., Lesser, D., Grable, J., Boyer, H. W., Rosenberg, J. M., and Greene, P. J. (1983) *J. Biol. Chem.* **258**, 14638-14646.
46. Polisky, B., Greene, P., Garfin, D., McCarthy, B., Goodman, H., and Boyer, H. (1975) *Proc. Natl. Acad. Sci. U.S.A.* **72**, 3310-3314.
47. Reiland, J. (1971) *Methods Enzymol.* **22**, 287-321.