Published in final edited form as:

J Allergy Clin Immunol. 2012 July ; 130(1): 248–256.e9. doi:10.1016/j.jaci.2012.02.045.

Molecular mimicry between cockroach and helminth glutathione S-transferases promotes crossreactivity and cross-sensitization

Helton Santiago, MD, PhD^a, Elyse LeeVan, BA^a, Sasisekhar Bennuru, PhD^a, Flavia Ribeiro-Gomes, PhD^a, Ellen Mueller, BA^b, Mark Wilson, PhD^a, Thomas Wynn, PhD^a, David Garboczi, PhD^c, Joseph Urban, PhD^d, Edward Mitre, MD^b, and Thomas B. Nutman, MD

^aThe Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA

^bDepartment of Microbiology and Immunology, Uniformed Services University of the Health Sciences, Bethesda, Maryland, USA

^cLaboratory of Immunogenetics, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, Maryland, USA

^dDiet, Genomics, and Immunology Laboratory, Beltsville Human Nutrition Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland, USA

Abstract

Background—The extensive similarities between helminth proteins and allergens are thought to contribute to helminth-driven allergic sensitization.

Objective—To investigate the cross-reactivity between a major glutathione-S transferase (GST) allergen of cockroach (Bla g 5) and the GST of *Wuchereria bancrofti* (WbGST), a major lymphatic filarial pathogen of humans.

Methods—We compared the molecular and structural similarities between Bla g 5 and WbGST by in silico analysis and by linear epitope mapping. Levels of IgE, IgG and IgG4 antibodies were measured in filarial-infected and –uninfected patients. Mice were infected with *Heligmosomoides bakeri* (Hb) and skin tested for cross-reactive allergic responses.

Results—These two proteins are 30% identical at the amino acid level with remarkable similarity in the N-terminal region and overall structural conservation based on predicted three-dimensional models. Filarial infection was associated with IgE, IgG, and IgG4 anti-Bla g 5 Ab production, with a significant correlation between Abs (irrespective of isotype) to Bla g 5 and WbGST ($P < 0.0003$). Pre-incubation of sera from cockroach allergic subjects with WbGST partially depleted (by 50 to 70%) anti-Bla g 5 IgE, IgG, and IgG4 Abs. IgE epitope mapping of Bla g 5 revealed that two linear N-terminal epitopes are highly conserved in WbGST corresponding to Bla g 5 peptides partially involved in the inhibition of WbGST binding. Finally,

Reprint requests: Helton Santiago, Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, 4 Center Drive, Building 4, Room B1-05, NIH, Bethesda, MD 20892-0415. Telephone: 1-301-451-1889. FAX: 1-301-480-3757. helton.santiago@nih.gov.

Mark Wilson's present address is: National Institutes for Medical Research, London, England.

Disclosure of potential conflict of interest: The authors have declared that they have no conflicts of interest to disclose.

The online version of this article contains Supplemental Materials.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

mice infected with *Hb* developed anti-HbGST IgE and showed immediate type skin test reactivity to Bla g 5.

Conclusion—These data demonstrate that helminth GST and the aeroallergen Bla g 5 share epitopes that can induce allergic cross-sensitization.

Keywords

GST; cockroach; allergy; hygiene hypothesis; crossreactivity; filariasis

INTRODUCTION

Epidemiologic and experimental evidence suggests that helminth infections have a general ability to modulate allergic disease^{1–5} through both parasite-induced IL-10 and the expansion of natural regulatory T cells (nTregs)^{6–9}. Other studies, however, have shown that helminth infections may be associated with increased allergic manifestations^{10–12}, perhaps related to the Th2-dominated environment induced by the parasites¹³ and/or the potential crossreactivity among allergens and helminth proteins¹⁴. Indeed, crossreactivity between helminth tropomyosin and allergenic tropomyosin from mites and cockroaches has been described^{15–17}. Although crossreactivity among tropomyosins has dominated the discussion about helminth-aeroallergen crossreactivity¹⁸, several other allergens have orthologues in helminths¹⁹ that may be involved in cross sensitization.

In the present study, we investigated the crossreactivity among allergenic glutathione S-transferases (GST) and those in pathogenic helminths. GSTs play a variety of physiologic roles including detoxification of toxins, protection from oxidative stress, and eicosanoid metabolism²⁰. The cytosolic GSTs can be subdivided into different classes (alpha, mu, pi, sigma, delta, nu, and others) each of which contain two basic structural motifs: an N-terminal thioredoxin motif and a C-terminal GST motif that is involved in formation of the H-site (a hydrophobic pocket that binds substrate). The sequence variability of the C-terminal GST motif confers species specificity to the molecule, making it a target of drug development for allergic diseases, cancer treatment, and parasitic infections²⁰ including vaccines.

Although GSTs are important helminth vaccine candidates^{21–23}, GSTs are known to be strong inducers of IgE. For example, the GST of the cockroach *Blattella germanica* (Bla g 5) is a major cockroach allergen, being an allergen inducing IgE in 30% to 71% of those with cockroach allergies^{24–26}. Moreover, GST is a common allergen of house dust mite (HDM) — with IgE anti-GST being found in up to 96% of HDM allergic populations^{27, 28}.

The present study demonstrates marked similarities at the amino acid (aa) and structural level between filarial (and other helminth) GST and the major cockroach allergen Bla g 5. We have mapped the linear crossreactive epitopes in humans and have shown their involvement in the structural basis for this crossreactivity. Moreover, we have used mice infected with intestinal nematode *Heligmosomoides bakeri* (Hb) to demonstrate unequivocally that helminth infections induce parasite-specific IgE that sensitizes mice to Bla g 5. These data suggest that helminth infection promotes cross-sensitization to common allergens through molecular mimicry.

METHODS

Patients and sera

Sera from well characterized filaria-infected (Fil⁺) individuals were utilized in this study. All patients were seen by the Clinical Parasitology Section of the Laboratory of Parasitic

Diseases under protocols approved by the Institutional Review Board of the NIAID and registered (NCT00001230; NCT00001645). The Fil⁺ group in this study was composed of 47 patients with *Loa loa* (n=37), *O. volvulus* (n=6), or *W. bancrofti* (n=4). Among the 47, 39 were temporary residents of or travelers to filarial-endemic regions, while 8 were indigenous to these same regions. Sera from 29 filaria-uninfected (Fil⁻; normal) individuals were obtained from the Department of Transfusion Medicine, Clinical Center, NIH, under protocols approved by the Clinical Center, NIH IRB.

All sera were tested for IgE to common allergens using Phadiatop® technology (Phadia, Uppsala, Sweden). Serum samples with levels below 0.35 kUA/l were considered negative and categorized as non-atopic. Sera from Phadiatop®-positive subjects were further tested for cockroach-specific IgE using an Immunocap™ assay specific for *Blattella germanica* (Bla g) (Phadia). Individuals positive for Bla g (levels above 0.35 kUA/l) were considered atopic for cockroach. Based on these data, the 76 subjects were divided into four groups based on their cockroach allergy and filarial infection status: 1) Fil⁻ and non-atopic, Ni-NA; n = 15 individuals; 2) Fil⁻ and atopic, Ni-A; n = 14; 3) Fil⁺ and non-atopic, Fil+NA; n = 11; and 4) Fil⁺ and atopic, Fil+A; n = 36.

Antigens and peptides

cDNA encoding the GSTs of *B. germanica* (Bla g 5) or *W. bancrofti* (WbGST) were cloned into bacmids. Transformed baculoviruses were used to infect Hi5 cells for expression of Bla g 5 or WbGST. Cell lysates and supernatants were purified on glutathione columns. The purities of Bla g 5 and WbGST were assessed by SDS-PAGE. Some experiments utilized recombinant Bla g 5 purchased from Indoor Biotechnologies Inc. (Charlottesville, VA). Recombinant Bla g 4 was purchased from Indoor Biotechnologies Inc.

A library of 40 peptides with purities greater than 80% that spanned the entire length of Bla g 5 (Table E1) was synthesized by Mimotope. Each peptide was 15 aa residues in length and overlapped adjoining peptides by 10 residues. Peptide #31 could not be purified and was not used in the experiments. Peptides were dissolved in HPLC-grade DMSO (Fisher Scientific, Pittsburg, PA) at 10 mg/ml to produce stock solutions and were kept at -40°C until used.

Epitope mapping

An array of 39 overlapping peptides and the full-length Bla g 5 was blotted onto PVDF membranes using a 96-well mini-fold dot blotter (Schleicher & Schuell, Inc., Riviera Beach, FL). After vacuum aspiration, air drying and blocking, membranes were incubated with six positive and two negative sera, depleted of IgG by incubation with protein G beads (GE Biotechnologies, Piscataway, NJ), diluted 1:25 in PBS-milk 5%. Membranes were then incubated with goat anti-human IgE (R&D, Minneapolis, MN), followed by peroxidase-conjugated rabbit anti-goat IgG, or peroxidase-conjugated anti-human IgG (Jackson ImmunoResearch, West Grove, PA) and developed with enhanced chemiluminescence substrate (Amersham- GE Biotechnologies, Piscataway, NJ) according to the manufacture's recommendations.

ELISA for Bla g 5 and WbGST

Measurements of Bla g 5, Bla g 4 and WbGST Abs were performed by ELISA as previously described¹⁶. Geometric mean (GM) + 3 SD of the Ab levels of the Ni-NA group were used to set cut-off values to identify individuals positive and negative for Abs to Bla g 5, Bla g 4 and to WbGST.

Depletion ELISA

Depletion ELISA was performed as previously described¹⁶ using Bla g 5, WbGST, or a nonrelated recombinant antigen (OvTropomyosin) to capture specific antibodies and coated plates to measure the level of depletion.

Inhibition ELISA

Sera samples from infected and non-infected subjects that were IgE and IgG positive for Bla g 5 and WbGST were incubated overnight with 20 µg in 0.1 mL of selected peptides (Pep 2, Pep 7, Peps 12–13, Peps 22–23, and Peps 27–29) dissolved in DMSO (diluted in 1% BSA/PBS from 10 mg/mL stock). Full-length Bla g 5, or WbGST in DMSO at 2% v/v was used as positive controls and 2% DMSO was used as negative control. The following day sera were transferred to plates coated with Bla g 5 or WbGST and developed for IgE Ab reactivity. To increase the sensitivity of the IgE-based assays, these sera were depleted of IgG by protein-G incubation (GE Healthcare, Piscataway, NJ).

3-D modeling

3-D structural models of Bla g 5 and WbGST were constructed by homology modeling with GST sigma from *Drosophila melanogaster* (DmGST1-1; Acc. No P41043, PDB 1m0u)²⁹ and GST of *O. volvulus* (OvGST2; Acc. No P46427, PDB 1tu7)³⁰ using the Protein Model Portal (<http://www.proteinmodelportal.org/>)³¹. The 3-D crystal structure of *H. bakery* GST 2 (HbGST2; PDB 1tw9)³² was used to compare with the models predicted for Bla g 5 and WbGST (Fig E2 and Table E3). Molecular visualization of each model and 3-D alignment were performed using Swiss-pdb viewer (<http://www.expasy.org/spdbv/>).

Sequence alignment and phylogenetic tree

Lasergene MegAlign (DNASTar, Inc.) was used to compare the sequences of WbGST, Bla g 5, and HbGST2, and with themselves or with the protein sequences from other GST allergens of HDMs, cockroaches, and fungus (Table E4). Pair-wise or multiple-sequence alignments of the GSTs (Table E4) were performed using ClustalW (algorithmic Gonet series) and used to construct the phylogenetic tree. Bootstrap values were calculated using 1000 bootstrap repetition and 111 random seeds.

Mouse infection and skin testing

BALB/c female mice, 6–8 wk old, were purchased from Jackson Laboratories, housed at an Association for the Assessment and Accreditation of Laboratory Animal Care-approved facility at the National Institute of Allergy and Infectious Diseases (NIAID) and studied under an animal study proposal approved by the NIAID Animal Care and Use Committee. Mice were inoculated per os using a gavage tube with 200 infective third-stage larvae (L3) of Hb. Uninfected control mice (Ni) and mice infected with Hb for 30 d were skin tested. Mouse ear swelling assays³³ with modifications was performed to evaluate skin sensitivity. Mice were injected with 10 µg/10 µl of Bla g 4 (right ear) or Bla g 5 (left ear) and ear thickness measured with a caliper before and 15, 30, and 60 min after allergen injection. Alternatively, mice were injected i.v. with 200 µl of Evans' Blue solution (0.5% w/vol) 3 min after intradermal injection of the allergens and euthanized 15 min later. Ears were collected and the Evans' Blue dye extracted in formamide overnight at 63°C and quantified in a spectrophotometer at 620 nm.

Statistical analysis

GraphPad Prism v5.0 (GraphPad Software Inc., San Diego, California) was used for all of the statistical analyses (one-tailed Fisher's exact test, relative risks [RR] and odds ratios

[OR] with confidence intervals [CI], Spearman rank correlation and Wilcoxon matched-pairs signed rank test).

RESULTS

Bla g 5 and WbGST display related sequence and structure

We investigated the similarities between the cockroach GST allergen (Bla g 5) and a filarial GST from *W. bancrofti* (Fig 1). The sequences of Bla g 5 and WbGST were found to be closely related (30% identical, 47% similar) (Fig 1, A). Importantly, the N-terminal region of Bla g 5 and WbGST had a 41% identity over an 80-aa window, which classify them as “likely to crossreact” (WHO criteria³⁴). Phylogenetic analysis that included multiple allergenic GSTs demonstrated that the WbGST clusters closely with Bla g 5 (Fig 1, B). Moreover, the predicted 3-D structures of Bla g 5 and WbGST showed that both GSTs—although from different GST families (sigma and pi)—had well conserved structures, especially in the N-terminal domain (Fig 1, C).

Abs to Bla g 5 crossreact with WbGST

To address the crossreactivity between filarial GST and Bla g 5, sera from filaria-positive (Fil⁺) and filaria-negative (Fil⁻) individuals were subdivided into cockroach allergic or non-allergic based on specific reactivity to cockroach (*B. germanica*) extract giving four groups: 1) Fil⁻, non-atopic (Ni-NA); 2) Fil⁻, atopic (Ni-A); 3) Fil⁺, non-atopic (Fil+NA); and 4) Fil⁺, atopic (Fil+A). Levels of GST-specific IgE, IgG, and IgG4 were then assessed (Fig 2, A). As can be seen, filarial infection (irrespective of atopic status) was associated with increased levels of IgE, IgG, and IgG4 Abs to both WbGST and Bla g 5. When the prevalence of Bla g 5-specific Abs was assessed among the atopic individuals (Table I), the presence of filarial infection was associated with increased reactivity to Bla g 5. Indeed, the prevalence of GST-specific IgE was 28.6% (4/14) in the Ni-A group compared with 63.9% (23/36) in the Fil+A group ($P = 0.026$). For the anti-GST IgG, the prevalence increased from 14.3% (2/14) to 69.4% (25/36) ($P = 0.0005$), and for anti-GST IgG4, there was an increase in prevalence from 14.3% (2/14) to 45% (17/36) ($P = 0.0302$) when Ni-A and Fil+A were compared (Table I). Thus, the Fil⁺ individuals had significantly increased likelihoods of being positive for Bla g 5-specific IgE (RR of 2.9; OR = 4.4) and IgG (RR of 7.0; OR = 13.6) and IgG4 (OR = 5.3) (Table I). When similar analysis was performed for Bla g 4 (a different cockroach allergen that does not have a parasite orthologue) no significant association between infection status and antibody levels was found. For example: Ni-A and Fil+A showed 8.3% and 31% of anti-Bla g 4 IgE prevalence ($P = 0.102$), respectively, and 8.3% and 25% ($P = 0.199$) for IgG. Furthermore, a positive correlation was found between levels of the IgE, IgG, and IgG4 anti-WbGST Abs and the respective levels of anti-Bla g 5 (IgE, $r = 0.38$ and $P = 0.0003$; IgG, $r = 0.68$ and $P < 0.0001$; and IgG4, $r = 0.52$ and $P < 0.0001$) (Fig 2, B).

To test more specifically the crossreactivity between the two GST proteins, depletion analyses were performed (Fig 3). Pre-incubation of cockroach GST positive sera with WbGST was able to deplete 70% of Bla g 5-specific IgE, IgG, and IgG4, whereas pre-incubation with a recombinant, structurally unrelated filarial control protein known also to be allergenic (OvTropomyosin)¹⁶ failed to deplete Bla g 5-specific Abs. Incubation of sera with full-length Bla g 5 or WbGST was able to inhibit the IgE binding to both proteins, but the data suggested that the cross-reactivity was not complete (data not shown).

Involvement of two N-terminal linear epitopes in crossreactivity between Bla g 5 and WbGST

To investigate if linear epitopes were involved in the molecular basis of Bla g 5/WbGST crossreactivity, overlapping peptides (15 mers) covering the full length of Bla g 5 (Table E1) were used to map the epitopes involved in IgE and IgG binding (Fig 4) with sera of 4 filaria-infected (S#1,2,5 and 6) and 2 non-infected subjects (S#3 and 4). As seen, four linear regions of Bla g 5 were identified that were reactive to both IgG and IgE: two regions in the N-terminal domain (peptide 7, aa residues 31–45 and peptides 12–13, aa residues 56–75) and two regions in the C-terminal domain (peptides 22–23, aa residues 106–125 and peptides 27–29, aa residues 131–155). Three of the regions (Pep 7, Peps 12–13, and Peps 22–23) corresponded to exposed regions of Bla g 5 that were predicted to be antigenic (Fig 4, B). There was strong reactivity to Pep 7 in 3/6 for IgE and 4/6 positive for IgG (Fig 4, A). Four of six had a weak signal to Peps 12–13 for IgE, but for IgG there was strong reactivity. The region of the molecule corresponding to Peps 22–23, predicted to be highly antigenic (Fig 4, B), had weak but measurable reactivity for IgE (in 3/6) and IgG (in 4/6) (Fig 4, A). Surprisingly, there was consistent binding of IgE (2/6) and IgG (6/6) to Peps 27–29, a region predicted to be poorly antigenic. We next mapped the four antigenic regions to 3-D models of Bla g 5 (Fig E1 and Table E2) and compared them to the corresponding regions in the 3-D model of WbGST (Fig E2 and Table E3). Peptides 7 and 12–13 were found to be highly conserved between Bla g 5 and WbGST with 40% sequence identity and Carbon-alpha root square mean difference of 2.4 Å and 1.7 Å, respectively, at the structural level.

Each of the antigenic peptides was clearly able to partially inhibit the binding of IgE from infected individuals to Bla g 5 (Fig 5A). For example, peptide 7, peptides 12–13, peptides 22–23 and peptides 27–29 were able to inhibit the binding of IgE to plate immobilized Bla g 5 ($P < 0.04$). In contrast, peptide 2 that was not recognized by any of the subjects tested could not inhibit binding at all. The fact the peptides could not inhibit completely IgE binding to Bla g 5 suggests that both non-linear conformational and linear (possibly conformational) epitopes are involved in the IgE response to this allergen. In the present study, we examined the role played by these linear regions in the cross-reactivity to WbGST. When the peptides were used to inhibit the WbGST ELISA, only peptide 7 and peptides 12–13 could inhibit IgE binding suggesting that the other peptides are not relevant for cross-reactivity. Relevant non-linear conformational epitopes could not be evaluated in this study.

Conversely, using sera from filaria-infected patients (Fig E3) or Bla g sensitized mice (ip immunizations) (Fig E4) in an inhibition ELISA to verify the specificity of the cross-reactivity, we were able to show that peptides from WbGST and HpGST-2 corresponding to the Bla g 5 peptides 7 and 12–13 could inhibit anti-Bla g 5 IgE equally well (Fig E3 and E4). In addition, Bla g 5 and HpGST peptides could also inhibit the WbGST human IgE ELISA (Fig E3B).

Helminth infection can induce cross-sensitization in mice

To assess whether helminth infection could induce crossreactive IgE and cause cross-sensitization in vivo, we infected BALB/c mice with *Heligmosomoides bakeri* (Hb), a parasite that contains a GST (HpGST-2) that is 32% identical (50% similar) to Bla g 5 and contains conserved sequences to the linear epitopes mapped for Bla g 5 (Fig E2 and Table E3), and likely also shows conserved conformational epitopes. More importantly, mice infected with Hb developed IgE to Bla g 5 (and not to an unrelated cockroach antigen, Bla g 4) (Fig 6, A); further this IgE mediated immediate-type hypersensitivity responses as measured intradermally administered antigen (Bla g 5 in one ear or Bla g 4 in the contralateral ear). Hb-infected mice developed immediate hypersensitivity reactions to Bla g 5 but not to Bla g 4 (Fig 6, B). To confirm this finding, skin-tested animals were injected

with Evans' blue dye i.v. 3–5 min after intradermal injection, and dermal edema was assessed by dye leakage (Fig 6, C). As seen, Hb-infected mice challenged with Bla g 5 had clear evidence of vascular leakage, whereas Bla g 4 (in the context of Hb infection) failed to induce an immediate hypersensitivity reaction. This was not due to a failure of Bla g 4 to induce sensitization, as mice sensitized (by i.p. immunizations) to either Bla g 4 or Bla g 5 could mount an immediate hypersensitivity response to the homologous antigen challenge (data not shown). As can also be observed in Fig 6, C, uninfected (Ni) mice reacted to neither Bla g 5 nor Bla g 4.

DISCUSSION

Although the “Hygiene Hypothesis”³⁵ suggests that a decrease in helminth (and/or other) infections may be responsible for the global increase in allergic diseases, there are data to show helminth infection can actually increase the risk for atopic disease^{10–12} and asthma^{35–38}. One concept that has emerged—in an attempt to understand the allergy-promoting capacity of certain helminth infections—has been the induction of parasite-specific IgE responses that can crossreact with allergens (particularly aeroallergens), thereby inducing allergic effector responses^{14–16, 19, 39}.

In regions of the world that are highly endemic for parasitic worm infections, cockroach and HDM are responsible for the overwhelming majority of aeroallergen sensitization^{10, 11, 40, 41}. Among the 10 defined allergens of cockroach in the WHO/OIUIS Allergen Nomenclature Database (www.allergen.org), GSTs (especially Bla g 5) are among the important cockroach allergens^{24, 42}. Although the GSTs studied here belong to different families—Bla g 5 is a GST sigma, while WbGST and HbGST2 are GSTs pi and nu—they are evolutionarily related⁴³. Immunologic crossreactivity for GSTs of different families that are evolutionarily conserved, although considered to be rare^{43, 44}, has been demonstrated for Der p 8 (a HDM GST mu) and Bla g 5 (GST sigma)²⁷. We found by in silico analysis of Bla g 5 and WbGST that the N-terminal motif of WbGST displayed 41% identity over an 80-aa residue span, fulfilling the WHO paradigm used to predict crossreactivity³⁴. In addition, the 3-D modeling onto which the allergenic linear epitopes were mapped (Fig E2) provided further molecular and structural basis for the crossreaction seen.

Aeroallergens such as Bla g 5, in contrast with food allergens that tend to have mostly linear epitopes due to proteolytic digestion of the antigen, have primarily conformational epitopes displayed on the surface of the protein. Of note, some linear epitopes can also be conformational if that sequence is surface-exposed⁴⁵, and, perhaps, this is the case of the linear epitopes identified in this study for Bla g 5. Because the peptides identified as IgE epitopes were unable to completely inhibit IgE binding to Bla g or WbGST, our data suggests that conformational epitopes are quite important, as suggested by the difference in the potency of ELISA inhibition between the entire protein and the peptides. Nevertheless our results suggest that the N-terminal portion of Bla g 5, the most conserved region of the molecule, was likely to be the primary region involved in the crossreactivity to WbGST. These data agree with data from epitope mapping of the bovine filarial parasite *Setaria cervi* GST that has recently shown that the N-terminal end was highly immunogenic⁴⁶. Because our data suggest only partial crossreactivity (i.e., WbGST could not inhibit completely anti-Bla g 5 Abs), we conclude that Bla g 5-specific Abs are comprised of both crossreactive and non-crossreactive Abs. The non-crossreactive Abs may be directed to a less conserved portion of the molecules or may reflect small differences in the peptide structure or in non-linear conformational epitopes.

The most important finding in the present study was the demonstration that this induction of crossreactive, allergen-specific IgE could be recapitulated in vivo using a model of a rodent

gastrointestinal nematode in BALB/c mice. We were able to show that this helminth infection induced crossreactive anti-Bla g 5 IgE that mediated skin sensitivity (immediate hypersensitivity reactions) to Bla g 5, but not to Bla g 4 (a non-orthologous cockroach allergen). The use of Bla g 4 was important to exclude the possibility of helminth-induced polyclonal IgE activation through a Th2-skewed immune response that could induce indiscriminate allergic sensitization^{47, 48}. Our data suggest that cross-sensitization was based primarily on the identity/similarity of the proteins of the parasite to their aeroallergen orthologues.

Cross-sensitization may have important implications for understanding the hygiene hypothesis¹⁶ and to helminth-associated allergy¹⁴. We show that parasite can induce crossreactive IgE with possible implications of cross-sensitization in vivo that may have significant implications for vaccine development against helminth parasites as suggested by the allergic reactivity (related to the presence of pre-existing IgE to the vaccine protein) to a first generation hookworm vaccine²². Because GSTs are potential vaccine candidates for *S. mansoni*²¹, *N. americanus*^{22, 49} and *W. bancrofti*²³, not only is the presence of pre-existing vaccine-specific IgE of concern, but also is the potential for cross reactions to aeroallergen (e.g. cockroach) orthologues.

Together, our data demonstrating induction of crossreactive allergen-specific Abs by helminth infection in both humans and experimentally infected mice clearly help to understand the conflicting data that have emerged in the study of the helminth/allergy interface. Not only have we provided a structural basis for the induction of crossreactive Abs, but we have also identified concerns that must be addressed in terms of allergy epidemiology, vaccine development²² and therapeutic approaches to allergic diseases.

Acknowledgments

This work was supported by the Intramural Research Program of the Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health. We thank NIAID intramural editor Brenda Rae Marshall for assistance.

Abbreviations used

Bla g 5	GST of the cockroach <i>Blattella germanica</i>
CI	confidence interval
Bla g	<i>Blattella germanica</i>
ELISA	enzyme-linked immunosorbent assay
Fil⁺	filaria-infected
Fil⁻	filaria-uninfected
Fil-A	Fil ⁺ and atopic
Fil-NA	Fil ⁺ and non-atopic
GM	geometric mean
GST	glutathione-S transferase
Hb	<i>Heligmosomoides bakeri</i>
HDM	house dust mite
HbGST-2	GST of <i>H. bakeri</i>

IQR	interquartile range
Ni-A	Fil ⁻ and atopic
Ni-NA	Fil ⁻ and non-atopic
nTreg	natural regulatory T cell
OR	odds ratio
SmGST	GST of <i>Schistosoma mansoni</i>
WbGST	GST of <i>Wuchereria bancrofti</i>
RR	relative risk

References

- Cooper PJ, Chico ME, Rodrigues LC, Ordonez M, Strachan D, Griffin GE, et al. Reduced risk of atopy among school-age children infected with geohelminth parasites in a rural area of the tropics. *J Allergy Clin Immunol.* 2003; 111:995–1000. [PubMed: 12743563]
- Dittrich AM, Erbacher A, Specht S, Diesner F, Krokowski M, Avagyan A, et al. Helminth infection with *Litomosoides sigmodontis* induces regulatory T cells and inhibits allergic sensitization, airway inflammation, and hyperreactivity in a murine asthma model. *J Immunol.* 2008; 180:1792–9. [PubMed: 18209076]
- Flohr C, Tuyen LN, Lewis S, Quinnell R, Minh TT, Liem HT, et al. Poor sanitation and helminth infection protect against skin sensitization in Vietnamese children: A cross-sectional study. *J Allergy Clin Immunol.* 2006; 118:1305–11. [PubMed: 17157661]
- Hagel I, Lynch NR, Perez M, Di Prisco MC, Lopez R, Rojas E. Modulation of the allergic reactivity of slum children by helminthic infection. *Parasite Immunol.* 1993; 15:311–5. [PubMed: 8361773]
- Maizels RM, Yazdanbakhsh M. Immune regulation by helminth parasites: cellular and molecular mechanisms. *Nat Rev Immunol.* 2003; 3:733–44. [PubMed: 12949497]
- Metenou S, Dembele B, Konate S, Dolo H, Coulibaly SY, Coulibaly YI, et al. At homeostasis filarial infections have expanded adaptive T regulatory but not classical Th2 cells. *J Immunol.* 2010; 184:5375–82. [PubMed: 20357251]
- Mitre E, Chien D, Nutman TB. CD4(+) (and not CD25+) T cells are the predominant interleukin-10-producing cells in the circulation of filaria-infected patients. *J Infect Dis.* 2008; 197:94–101. [PubMed: 18171291]
- Wilson MS, Taylor MD, Balic A, Finney CA, Lamb JR, Maizels RM. Suppression of allergic airway inflammation by helminth-induced regulatory T cells. *J Exp Med.* 2005; 202:1199–212. [PubMed: 16275759]
- Yazdanbakhsh M, Wahyuni S. The role of helminth infections in protection from atopic disorders. *Curr Opin Allergy Clin Immunol.* 2005; 5:386–91. [PubMed: 16131911]
- Dold S, Heinrich J, Wichmann HE, Wjst M. *Ascaris*-specific IgE and allergic sensitization in a cohort of school children in the former East Germany. *J Allergy Clin Immunol.* 1998; 102:414–20. [PubMed: 9768582]
- Hunninghake GM, Soto-Quiros ME, Avila L, Ly NP, Liang C, Sylvia JS, et al. Sensitization to *Ascaris lumbricoides* and severity of childhood asthma in Costa Rica. *J Allergy Clin Immunol.* 2007; 119:654–61. [PubMed: 17336615]
- Palmer LJ, Celedon JC, Weiss ST, Wang B, Fang Z, Xu X. *Ascaris lumbricoides* infection is associated with increased risk of childhood asthma and atopy in rural China. *Am J Respir Crit Care Med.* 2002; 165:1489–93. [PubMed: 12045121]
- Anthony RM, Rutitzky LI, Urban JF Jr, Stadecker MJ, Gause WC. Protective immune mechanisms in helminth infection. *Nat Rev Immunol.* 2007; 7:975–87. [PubMed: 18007680]
- Caraballo L, Acevedo N. Allergy in the tropics: the impact of cross-reactivity between mites and *Ascaris*. *Front Biosci (Elite Ed).* 2011; 3:51–64. [PubMed: 21196284]

15. Acevedo N, Sanchez J, Erler A, Mercado D, Briza P, Kennedy M, et al. IgE cross-reactivity between *Ascaris* and domestic mite allergens: the role of tropomyosin and the nematode polyprotein ABA-1. *Allergy*. 2009; 64:1635–43. [PubMed: 19624559]
16. Santiago HC, Bennuru S, Boyd A, Eberhard M, Nutman TB. Structural and immunologic cross-reactivity among filarial and mite tropomyosin: Implications for the hygiene hypothesis. *J Allergy Clin Immunol*. 2011; 127:479–86. [PubMed: 21185070]
17. Santos AB, Rocha GM, Oliver C, Ferriani VP, Lima RC, Palma MS, et al. Cross-reactive IgE antibody responses to tropomyosins from *Ascaris lumbricoides* and cockroach. *J Allergy Clin Immunol*. 2008; 121:1040–6. e1. [PubMed: 18275995]
18. Sereda MJ, Hartmann S, Lucius R. Helminths and allergy: the example of tropomyosin. *Trends Parasitol*. 2008; 24:272–8. [PubMed: 18450511]
19. Fitzsimmons CM, Dunne DW. Survival of the fittest: allergology or parasitology? *Trends Parasitol*. 2009; 25:447–51. [PubMed: 19744885]
20. Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annu Rev Pharmacol Toxicol*. 2005; 45:51–88. [PubMed: 15822171]
21. Capron A, Capron M, Riveau G. Vaccine development against schistosomiasis from concepts to clinical trials. *Br Med Bull*. 2002; 62:139–48. [PubMed: 12176856]
22. Hotez PJ, Bethony JM, Diemert DJ, Pearson M, Loukas A. Developing vaccines to combat hookworm infection and intestinal schistosomiasis. *Nat Rev Microbiol*. 2010; 8:814–26. [PubMed: 20948553]
23. Veerapathran A, Dakshinamoorthy G, Gnanasekar M, Reddy MV, Kalyanasundaram R. Evaluation of *Wuchereria bancrofti* GST as a vaccine candidate for lymphatic filariasis. *PLoS Negl Trop Dis*. 2009; 3:e457. [PubMed: 19513102]
24. Arruda LK, Vailes LD, Platts-Mills TA, Hayden ML, Chapman MD. Induction of IgE antibody responses by glutathione S-transferase from the German cockroach (*Blattella germanica*). *J Biol Chem*. 1997; 272:20907–12. [PubMed: 9252418]
25. Jeong KY, Jeong KJ, Yi MH, Lee H, Hong CS, Yong TS. Allergenicity of sigma and delta class glutathione S-transferases from the German cockroach. *Int Arch Allergy Immunol*. 2009; 148:59–64. [PubMed: 18716404]
26. Satinover SM, Reefer AJ, Pomes A, Chapman MD, Platts-Mills TA, Woodfolk JA. Specific IgE and IgG antibody-binding patterns to recombinant cockroach allergens. *J Allergy Clin Immunol*. 2005; 115:803–9. [PubMed: 15806002]
27. Huang CH, Liew LM, Mah KW, Kuo LE, Lee BW, Chua KY. Characterization of glutathione S-transferase from dust mite, Der p 8 and its immunoglobulin E cross-reactivity with cockroach glutathione S-transferase. *Clin Exp Allergy*. 2006; 36:369–76. [PubMed: 16499649]
28. O'Neill GM, Donovan GR, Baldo BA. Glutathione S-transferase a major allergen of the house dust mite, *Dermatophagoides pteronyssinus*. *Immunol Lett*. 1995; 48:103–7. [PubMed: 8719107]
29. Agianian B, Tucker PA, Schouten A, Leonard K, Bullard B, Gros P. Structure of a *Drosophila* sigma class glutathione S-transferase reveals a novel active site topography suited for lipid peroxidation products. *J Mol Biol*. 2003; 326:151–65. [PubMed: 12547198]
30. Perbandt M, Hoppner J, Betzel C, Walter RD, Liebau E. Structure of the major cytosolic glutathione S-transferase from the parasitic nematode *Onchocerca volvulus*. *J Biol Chem*. 2005; 280:12630–6. [PubMed: 15640152]
31. Arnold K, Kiefer F, Kopp J, Battey JN, Podvinec M, Westbrook JD, et al. The Protein Model Portal. *J Struct Funct Genomics*. 2009; 10:1–8. [PubMed: 19037750]
32. Schuller DJ, Liu Q, Kriksunov IA, Campbell AM, Barrett J, Brophy PM, et al. Crystal structure of a new class of glutathione transferase from the model human hookworm nematode *Heligmosomoides polygyrus*. *Proteins*. 2005; 61:1024–31. [PubMed: 16189827]
33. Proust B, Astier C, Jacquenet S, Ogier V, Magueur E, Roitel O, et al. A single oral sensitization to peanut without adjuvant leads to anaphylaxis in mice. *Int Arch Allergy Immunol*. 2008; 146:212–8. [PubMed: 18268389]
34. Organization FaA. Evaluation of allergenicity of genetically modified foods: Report of a joint FAO/WHO expert consultation. Rome: 2001.

35. Benicio MH, Ferreira MU, Cardoso MR, Konno SC, Monteiro CA. Wheezing conditions in early childhood: prevalence and risk factors in the city of Sao Paulo, Brazil. *Bull World Health Organ.* 2004; 82:516–22. [PubMed: 15508196]
36. Hagel I, Cabrera M, Hurtado MA, Sanchez P, Puccio F, Di Prisco MC, et al. Infection by *Ascaris lumbricoides* and bronchial hyper reactivity: an outstanding association in Venezuelan school children from endemic areas. *Acta Trop.* 2007; 103:231–41. [PubMed: 17698018]
37. Lynch NR, Palenque M, Hagel I, DiPrisco MC. Clinical improvement of asthma after anthelmintic treatment in a tropical situation. *Am J Respir Crit Care Med.* 1997; 156:50–4. [PubMed: 9230725]
38. Takeuchi H, Zaman K, Takahashi J, Yunus M, Chowdhury HR, Arifeen SE, et al. High titre of anti-*Ascaris* immunoglobulin E associated with bronchial asthma symptoms in 5-year-old rural Bangladeshi children. *Clin Exp Allergy.* 2008; 38:276–82. [PubMed: 18070165]
39. Santos AB, Chapman MD, Aalberse RC, Vailes LD, Ferriani VP, Oliver C, et al. Cockroach allergens and asthma in Brazil: identification of tropomyosin as a major allergen with potential cross-reactivity with mite and shrimp allergens. *J Allergy Clin Immunol.* 1999; 104:329–37. [PubMed: 10452753]
40. Cooper PJ, Chico ME, Bland M, Griffin GE, Nutman TB. Allergic symptoms, atopy, and geohelminth infections in a rural area of Ecuador. *Am J Respir Crit Care Med.* 2003; 168:313–7. [PubMed: 12714349]
41. Kim JS, Ouyang F, Pongracic JA, Fang Y, Wang B, Liu X, et al. Dissociation between the prevalence of atopy and allergic disease in rural China among children and adults. *J Allergy Clin Immunol.* 2008; 122:929–35. e4. [PubMed: 18805578]
42. Gore JC, Schal C. Cockroach allergen biology and mitigation in the indoor environment. *Annu Rev Entomol.* 2007; 52:439–63. [PubMed: 17163801]
43. Sheehan D, Meade G, Foley VM, Dowd CA. Structure, function and evolution of glutathione transferases: implications for classification of non-mammalian members of an ancient enzyme superfamily. *Biochem J.* 2001; 360:1–16. [PubMed: 11695986]
44. Hayes JD, Mantle TJ. Use of immuno-blot techniques to discriminate between the glutathione S-transferase Yf, Yk, Ya, Yn/Yb and Yc subunits and to study their distribution in extrahepatic tissues. Evidence for three immunochemically distinct groups of transferase in the rat. *Biochem J.* 1986; 233:779–88. [PubMed: 3707525]
45. Pomés A. Relevant B Cell Epitopes in Allergic Disease. *International Archives of Allergy and Immunology.* 2010; 152:1–11.
46. Yadav M, Liebau E, Haldar C, Rathaur S. Identification of major antigenic peptide of filarial glutathione-S-transferase. *Vaccine.* 2011
47. Jarrett E, Bazin H. Elevation of total serum IgE in rats following helminth parasite infection. *Nature.* 1974; 251:613–4. [PubMed: 4547380]
48. Urban JF Jr. Cellular basis of the non-specific potentiation of the immunoglobulin E response after helminth parasite infection. *Vet Parasitol.* 1982; 10:131–40. [PubMed: 6982561]
49. Zhan B, Liu S, Perally S, Xue J, Fujiwara R, Brophy P, et al. Biochemical characterization and vaccine potential of a heme-binding glutathione transferase from the adult hookworm *Ancylostoma caninum*. *Infect Immun.* 2005; 73:6903–11. [PubMed: 16177370]

Key message

- Cockroach and helminth GST crossreact because of remarkable molecular and structural similarities
- Filarial infection is associated with increased prevalence of cross-reactive anti-Bla g 5 IgE with possible clinical implications
- Experimental infection with helminth induced cross-reactive allergy as measured by skin test
- Vaccine development for helminths should be concerned about the potential impact of crossreactivity with common allergens

FIG 1.

Sequence and structural similarities between Bla g 5 and WbGST. **A**, Alignment of Bla g 5 and WbGST sequences showing identical (shaded in black) and similar (shaded in gray) amino acids. **B**, Phylogenetic tree of allergenic GSTs. **C**, Predicted 3-D structure of Bla g 5 and WbGST color-coding basic structure for comparison.

FIG 2.

Correlation between WbGST and Bla g 5 Ab levels. (A) Level of anti-Bla g 5 and anti-WbGST IgE, IgG, and IgG4 Abs in non-infected and non-atopic (Ni-NA), non-infected and atopic (Ni-A), filaria-infected and non-atopic (Fil+NA) and filaria-infected and atopic (Fil+A) individuals. Individuals above the dotted lines (representing GM + 3 SD of Ni-NA) were considered positive for anti-Bla g 5 antibodies. B, Correlation between anti-WbGST and anti-Bla g 5 IgE, IgG, and IgG4; each point represents an individual. $n = 76$.

FIG 3.

WbGST-specific Abs crossreact with Bla g 5 due to conserved epitopes. Sera of patients positive for anti-Bla g 5 IgE, IgG, and IgG4 were incubated in plates coated with a control antigen (OvTropomyosin), WbGST or Bla g 5. Following incubation, sera were transferred to Bla g 5-coated plates and developed for IgE, IgG, and IgG4. Each dot represents one individual.

FIG 4.

Linear epitope mapping of Bla g 5. **(A)** Thirty-nine overlapping peptides spanning the entire sequence of Bla g 5 was used to perform western dot blots for IgE and IgG binding epitopes (numbers reflect peptide identity, S stands for serum and P for recombinant Bla g 5). **(B)**, The results were compared to in silico epitope prediction based on the Bla g 5 sequence.

FIG 5.

Antigenic peptides can inhibit IgE binding to Bla g 5 and WbGST. Sera from infected patients positive for anti-Bla g 5 and anti-WbGST Abs were incubated with DMSO (open squares) or specific peptides or recombinant proteins (closed circles) and then transferred to Bla g 5- or WbGST-coated multiwell plates. Optical densities (OD) are shown for all sera tested. Differences were compared using the Wilcoxon matched-signed rank test.

FIG 6. Helminth infection can cause cross-sensitization to Bla g 5. Mice infected with *Heligmosomoides bakeri* (Hb) developed IgE to Bla g 5 but not to Bla g 4 as measured by ELISA (**A**) and skin test reactivity by ear thickness (**B**) or by Evans' blue dye extravasation (**C**). Data represent one of four experiments performed ($n = 4-5$ per group per experiment). Bars (in A and B) represent means $\pm$ SE. In Panel C, each dot represents one animal and the dotted lines represent the GM of the Ni group + 3 SD.

TABLE I

Filarial infection is associated with higher prevalences of anti-Bla g 5 antibodies

	Prevalence (%) [*]		RR (CI)	Odds ratio (CI)	P value [†]
	Ni-A [‡]	Fil+A [§]			
IgE	28.6	63.9	2.9 (1.06–8.12)	4.4 (1.15–17.0)	0.0262
IgG	14.3	69.4	7.0 (1.75–28.3)	13.6 (2.60–71.5)	0.0005
IgG4	14.3	47.2	3.6 (0.92–14.7)	5.3 (1.05–27.5)	0.0302

^{*} Prevalence was calculated using cut-off values as explained in Fig 2.

[†] P value was calculated using one-sided Fisher's exact test.

[‡] Ni-A, non-infected and atopic; $n = 14$.

[§] Fil+A, filaria-infected and atopic; $n = 36$.