RFLPs for the human pepsinogen A haplotypes (PGA)R.Thomas Taggart^{1,2*}, Farzin B.Boudi² and Graeme I.Bell³

¹Department of Molecular Biology and Genetics, Wayne State University School of Medicine, 540 East Canfield, Detroit, MI 48201, ²Human Genetics Research, Veterans Administration Medical Center, Sepulveda, CA 91343, and ³The Howard Hughes Medical Institute, University of Chicago, IL 60637, USA

SOURCE/DESCRIPTION: PGA 101 is a 1340 bp cDNA clone containing exons 1-9 of the predicted human pepsinogen A coding sequence (1).

POLYMORPHISM: Two distinct polymorphisms are detected with EcoRI and Bgl II. Analysis with these enzymes provides for discrimination of the PGA haplotypes A, B, and C containing three, two and one PGA genes respectively (see below).

		PGA PHENOTYPES					PGA HAPLOTYPES		
		AA	AB	BB	AC	BC	A	B	C
EcoRI	22 kb	-	-	-	+	+	-	-	+
	17.8 kb	+	+	+	+	+	+	+	-
	17.2 kb	+	+	+	+	+	+	+	-
	13.5 kb	+	+	-	+	-	+	-	-
	3.9 kb	+	+	+	+	+	+	+	-
Bgl II	6.3 kb	-	+	+	+	+	-	+	+
	1.6 kb	+	+	-	+	-	+	-	-

FREQUENCY: Estimated from 55 unrelated Caucasians.

A haplotype .54

B haplotype .37

C haplotype .06

NOT POLYMORPHIC FOR: AvaI, MspI, PvuII, RsaI, Sau3AI, and TaqI

CHROMOSOMAL LOCALIZATION: The PGA complex is located at 11q13 (2).

MENDELIAN INHERITANCE: Demonstrated in 20 families.

PROBE AVAILABILITY: Contact R. T. Taggart.

OTHER COMMENTS: Low background at high stringency wash conditions (0.1 X SSC, 0.05% SDS).

REFERENCES:

1. Taggart, R. T., Boudi, F.B., Lothe, R.A., White, R., and Bell, G. I. Cytogenet. Cell Genet. 46:701, 1987.
2. Nakai, H., Byers, M.G., Shows, T.B., and Taggart, R.T. Cytogenet. Cell Genet. 43:215-217, 1986.

ACKNOWLEDGEMENT: This work was supported by grant AM32015 from the National Institutes of Health.

*To whom correspondence should be addressed