Published in final edited form as:

Org Biomol Chem. 2012 July 21; 10(27): 5194–5196. doi:10.1039/c2ob25821h.

Total Synthesis of Indole-3-Acetonitrile-4-Methoxy-2-*C*- β -D-Glucopyranoside. Proposal for Structural Revision of the Natural Product

Akop Yepremyan^a and Thomas G. Minehan^a

Thomas G. Minehan: thomas.minehan@csun.edu

^aDepartment of Chemistry and Biochemistry, California State University, Northridge, 18111 Nordhoff Street, Northridge, CA 91330, U.S.A. Fax: 1 818 677 4062; Tel: 1 818 677 3315

Abstract

Indole-3-acetonitrile-4-methoxy-2-*C*- β -D-glucopyranoside (**1**), a novel *C*-glycoside from *Isatis indigotica* with important cytotoxic activity, has been prepared in ten steps from ethynyl- β -*C*-glycoside **3** and 2-iodo-3-nitrophenyl acetate **6**. Key steps in the synthesis include a Sonogashira coupling and a CuI-mediated indole formation. NMR spectroscopic data for synthetic **1** differs from that reported for the natural product. A revised structure for the natural product, containing an alternate carbohydrate substituent, is proposed.

C-aryl glycosides are a class of natural products that exhibit a range of important biological properties.¹ Numerous members of this family display potent antitumor, antiviral, and antibiotic activities,² and there is ample experimental evidence that *C*-aryl glycosides bind duplex DNA.³

Two novel alkaloids recently isolated from the roots of the plant *Isatis indigotica* possess an indole-*C*-glycoside core.⁴ Indole-3-acetonitrile-4-methoxy-2-*C*- β -D-glucopyranoside (**1**, Figure 1) displays cytotoxic activity against human myeloid leukemia HL60 cells (IC₅₀=1.3 mM) and human liver cancer HepG2 cells (IC₅₀=2.1 mM). The structural isomer of **1**, *N*-methoxy-indole-3-acetonitrile-2-*C*- β -D-glucopyranoside (**2**), shows cytotoxic activity against both HL60 cells (IC₅₀=5.1 mM) and human myeloid leukemia Mata cells (IC₅₀=12.1 mM). In view of its promising biological profile, and with the ultimate aim of exploring the DNA-binding properties of indole-*C*-glycosides, we decided to undertake a total synthesis of **1**.

We envisioned that the crucial linkage between the indole and glycoside moieties could be fashioned from protected alkynyl-*C*-glycoside **3** and 2-iodo-3-nitrophenol **6** through a Sonogashira coupling, nitro group reduction, and intramolecular amine-alkyne cyclization sequence (Figure 2). Subsequent installation of the acetonitrile moiety under standard conditions and deprotection was envisaged to provide the natural product.

The assembly of the alkynyl-*C*-glycoside coupling partner **3**, possessing the requisite β -stereochemistry at the anomeric carbon, required efficient access to 2,3,4,6-tetra-*O*-benzyl gluconolactone **5** (Scheme 1).

Correspondence to: Thomas G. Minehan, thomas.minehan@csun.edu.

[†]Electronic Supplementary Information (ESI) available: full experimental details and characterization data for all compounds reported. See DOI: 10.1039/b000000x/

Allylation of dextrose under acidic conditions,⁵ followed by exhaustive benzylation provided allyl glycoside **4** as a mixture of C.1 anomers in 75% yield. Removal of the allyl ether by standard base-mediated olefin isomerization and enol ether hydrolysis⁶ furnished 2,3,4,6-tetra-*O*-benzyl glucose, which upon Swern oxidation⁷ provided lactone **5**. Following literature precedent,^{8,9} reaction of **5** with lithium(trimethylsilyl) acetylide in the presence of CeCl₃ led to an intermediate lactol which was immediately reduced with BF₃•OEt₂/Et₃SiH to provide the silyl-protected alkynyl glycoside. Subsequent treatment with aqueous NaOH gave rise to alkyne **3**.⁹

The synthesis of the aryl iodide coupling partner commenced from commercially available 2-amino-3-nitrophenol; diazotization¹⁰ in the presence of NaI furnished iodide **6a**, which could be methylated (K₂CO₃, CH₃I, DMF) or acylated (AcCl, Et₃N, DCM, 0°C) to give **6b** or **6c**, respectively.

Initial attempts at coupling methyl ether **6b** with alkyne **3** under standard Sonogashira conditions¹¹ met with limited success (Scheme 2). The major product isolated in these reactions was invariably symmetrical diyne **3d**. We hypothesized that, due to the electron-donating character of the methoxy substituent, oxidative addition of the palladium catalyst to **6b** was slow relative to alkyne dimerization. Gratifyingly, employing acetoxymethyl aryl iodide **6c** in the cross-coupling reaction cleanly gave rise to alkyne **7** in 66% yield with minimal formation of dimer **3d**. Aminolysis of the acetate ester, followed by hydroxyl methylation and reduction of the nitro group provided aniline **8b** in 61% yield over three steps.

Next, a base-mediated indolization was attempted using Knochel's *t*-BuOK/NMP system (Scheme 3).¹² Treatment of **8b** with potassium tert-butoxide in NMP for 15 minutes furnished variable yields of indole *C*-glycoside **9** in the range of 33–55%. Prolonged exposure to the reaction conditions resulted in significant substrate decomposition. A superior protocol for producing **9** reproducibly involved exposure of **8b** to excess CuI (2–5 equivalents) at 145°C in DMF for 2 hours.¹³ These conditions allowed indole **9** to be secured in 65% yield.

With **9** in hand, we next tested installation of the acetonitrile moiety via a three-step protocol.¹⁴ Treatment of **9** with formalin, diethylamine, and acetic acid overnight, isolation and subjection of the crude amine to methyl iodide in CH₂Cl₂, followed by refluxing the resulting quaternary ammonium salt with sodium cyanide in ethanol (80°C), gave rise to nitrile **10** in 85% overall yield. However, attempted deprotection of the benzyl ether protecting groups by hydrogenolysis over Pearlman's catalyst¹⁵ led to the formation of significant quantities of amine **11** in addition to desired nitrile **1**. Hydrogenolysis of **9** instead, followed by alcohol acylation provided peracetate **12**, the three-dimensional structure of which was confirmed by NOESY spectroscopy.

An alternative, higher-yielding route to **1** was secured via persilyl derivative **13** (Scheme 4), prepared in 67% yield from **9** (H₂, Pd(OH)₂; TBS-Cl, imid, DMF, 60°C). Acetonitrile installation under the aforementioned conditions gave rise to nitrile **14** in 60% overall yield. Removal of the silyl ether protecting groups was accomplished in 72% yield by exposure of **14** to TBAF (5 equiv) in THF for 9 hours, providing synthetic **1** as an amorphous white solid. The ¹H and ¹³C NMR data for synthetic **1** (recorded in acetone-*d*₆) differed from those reported by Hu et al.⁴ for natural **1** (Figure 3). Proton and carbon assignments for synthetic **1** were obtained from COSY, HSQC, and HMBC experiments, which also provided support for the hydrogen-carbon and carbon-carbon connectivity of the molecule. To further confirm the structure of our synthetic material, compound **1** was converted to peracetate **15** (Ac₂O, Pyr, 95%) and COSY and NOESY experiments were performed. Analysis of the crosspeaks in the NOESY spectrum of **15** revealed a connectivity pattern

consistent with the findings for compound **12**: a glucopyranosyl carbohydrate moiety is attached at the C.2 carbon atom of the indole ring.

Careful study of the spectroscopic data in figure 3 indicates that major differences in chemical shift between synthetic and natural **1** occur for protons on the carbohydrate substituent and for protons and carbon atoms near the site of attachment of the sugar moiety to the indole ring. The 9.1 Hz coupling constant^{16a} reported for H1' of the natural product suggests that the carbohydrate moiety is indeed a hexopyranose, and likely a diastereomer of synthetic **1** such as allose (the C.3 epimer) or galactose (the C.4 epimer).^{16b}

In summary, we have developed a concise route to indole-3-acetonitrile-4-methoxy-2-*C*-β-D-glucopyranoside, the proposed structure of a natural indole *C*-glycoside from *Isatis indigotica*. Comparison of spectroscopic data for synthetic and natural **1** indicate that the natural product likely contains a diastereomeric hexopyranose moiety. Preparation of the galactopyranose- and allopypyranose-containing indole *C*-glycosides is underway and comparisons of their spectroscopic data with that of the natural material will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Institutes of Health (SC3 GM 096899-01) and the CSUN Office of Research and Sponsored Projects for their generous support of our research program.

Notes and references

1. Hansen MR, Hurley LH. *Acc Chem Res.* 1996; 29:249.
2. Hacksell U, Daves GD. *Prog Med Chem.* 1985; 22:1. [PubMed: 3915364]
3. (a) Cairns MJ, Murray V. *Biochem Mol Biol Int.* 1998; 46:267. [PubMed: 9801795] (b) Hardman LC, Murray V. *Biochem Mol Biol Int.* 1997; 42:349. [PubMed: 9238534] (c) Murray V, Moore AG, Matias C, Wickham G. *BBA-Gene Struct Expr.* 1995; 1261:195. (d) Prakash AS, Moore AG, Murray V, Matias C, McFadyen WD, Wickham G. *Chem-Biol Interact.* 1995; 95:17. [PubMed: 7697748] (e) Kamei N, Sekiguchi T, Tsuboi M, Ishii S, Ohtsubo E, Maeda Y. *Nucl Acid S.* 1994; 31:87. (f) Sun D, Hansen M, Hurley L. *J Am Chem Soc.* 1995; 117:2430.
4. Wu Y, Zhang ZX, Hu H, Li D, Qiu G, Hu X, He X. *Fitoterapia.* 2011; 82:288. [PubMed: 20971167]
5. Schindler CS, Bertschi L, Carreira EM. *Angew Chem Int Ed.* 2010; 49:9229.
6. (a) Gent PA, Gigg R. *Carbohydr Res.* 1976; 49:325. [PubMed: 963695] (b) Rollin P, Sinay P. *J Chem Soc, Perkin Trans.* 1977; 1:2513.
7. Mancuso AJ, Huang S-L, Swern D. *J Org Chem.* 1978; 43:2480.
8. (a) Imamoto T. *Pure Appl Chem.* 1990; 62:747. (b) Alzeer J, Vasella A. *Helv Chim Acta.* 1995; 78:177.
9. Lowary T, Meldal M, Helmboldt A, Vasella A, Block K. *J Org Chem.* 1998; 63:9657.
10. Banwell MG, Jones MT, Loong DT, Lupton DW, Pinkerton DM, Ray JK, Willis AC. *Tetrahedron.* 2010; 66:9252.
11. Sonogashira K. *J Organomet Chem.* 2002; 653:46.
12. Rodriguez A, Koradin C, Dohle W, Knochel P. *Angew Chem Int Ed.* 2000; 39:2488.
13. Ezquerro J, Pedregal C, Lamas C, Barluenga J, Perez M, Garcia-Martin MA, Gonzalez JM. *J Org Chem.* 1996; 61:5804.
14. Perez-Alvarez V, Morales-Rios MS, Hong E, Joseph-Nathan P. *J Pharm Pharmacol.* 1997; 49:246. [PubMed: 9231339]

15. Pearlman WM. Tetrahedron Lett. 1967:1663.
16. Hu, et al. report two separate coupling constants for H1' in ref. 2: 9.1 Hz and 9.5 Hz. (b) The >9 Hz coupling constant for H1' in the natural product is indicative of an equatorial hydroxyl group at C.2 of the carbohydrate; thus a mannopyranose unit is not a likely candidate for the carbohydrate substituent of the natural product.

Figure 1.
Indole C-glycosides from *Isatis Indigotica*

Figure 2.
Retrosynthetic Analysis of **1**.

Position	1 natural		1 synthetic	
	δ_C	$\delta_H(J \text{ in Hz})$	δ_C	$\delta_H(J \text{ in Hz})$
1		11.26 (s)		10.23 (s)
2	121.8		133.1	
3	110.8		102.2	
4	153.2		154.2	
4 α	116.0		118.0	
5	99.7	6.54 (d, 7.8)	99.5	6.52 (d, 7.6)
6	123.9	7.09 (dd, 8.2, 7.8)	122.7	7.01 (dd, 8.0, 7.6)
7	104.7	6.95 (d, 8.2)	104.7	6.94 (d, 8.4)
7 α	137.9		137.2	
α	14.9	4.10 (s)	13.9	4.05 (s)
β	119.5		119.2	
1'	87.5	4.33 (d, 9.1)	74.5	4.64 (d, 8.8)
2'	72.2	2.76 (m)	74.5	3.60-3.49 (m)
3'	77.8	3.13 (m)	78.4	3.60-3.49 (m)
4'	69.5	2.96 (m)	70.2	3.60-3.49 (m)
5'	81.0	3.13 (m)	80.7	3.60-3.49 (m)
6'	61.0	3.67 (m)	61.8	3.85 (d, 13.3)
		3.43 (m)		3.73 (dd, 12.0, 4.0)
OCH ₃	69.6	3.88 (s)	54.7	3.91 (s)

Fig. 3.
Comparison of NMR chemical shift data for natural (ref. 4) and synthetic **1** recorded in acetone-d₆.

Scheme 1.
Preparation of coupling fragments **3** and **6**

Scheme 2.
Sonogashira fragment coupling

Scheme 3.
C-glycosyl indole synthesis and attempted deprotection; NOESY correlations for **12**.

Scheme 4.

An alternative route to **1** and structure assignment of **15**.