Published in final edited form as:

Placenta. 2012 August ; 33(8): 591–598. doi:10.1016/j.placenta.2012.04.011.

Survival by self-destruction: A role for autophagy in the placenta?

I Bildirici^a, MS Longtine^b, B Chen^b, and DM Nelson^b

^aDepartment of Obstetrics and Gynecology, Acibadem University School of Medicine, Istanbul, Turkey

^bDepartment of Obstetrics and Gynecology, Washington University School of Medicine, St Louis, MO 63110, USA

Abstract

Autophagy is a burgeoning area of research from yeast to humans. Although previously described as a death pathway, autophagy is now considered an important survival phenomenon in response to environmental stressors to which most organs are exposed.

Despite an ever expanding literature in non-placental cells, studies of autophagy in the placenta are lagging. We review the regulation of autophagy, summarize available placental studies of autophagy, and highlight potential areas for future research. We believe that such studies will yield novel insights into how placentas protect the survival of the species by “self-eating”.

Keywords

Placenta; trophoblast; autophagy; apoptosis; mTOR; ATG genes

Overview

Autophagy is Latin for “self eating”, a term that now describes a vital biological phenomenon. Autophagy is a highly regulated process by which damaged proteins, injured organelles, and invading organisms are contained within double membrane-bound vesicles called autophagosomes that fuse with lysosomes, which leads to degradation of cargo in the resulting autolysosome to recycle biomolecules. A cassette of autophagy (Atg) genes were initially described in the yeast, *Saccharomyces cerevisiae*, and to date, 35 such genes are identified, with many conserved in species from slime mould to humans [1–6]. Apoptosis, autophagy, and necrosis were originally designated as cell death types I, II, and III, respectively [7]. However, this nomenclature is no longer pertinent, as autophagy is recognized to promote cell survival and rarely, does this cause cell death [8]. Moreover, autophagy declines in animal cells during aging *in vivo* and *in vitro*, linking the process to longevity [9–10]. Indeed, caloric restriction is both an inducer of autophagy and the only intervention known to promote longevity [11]. We below review the stressors for the

© 2012 Elsevier Ltd. All rights reserved.

Corresponding author: Ibrahim Bildirici, MD, Department of Obstetrics and Gynecology, Acibadem University School of Medicine, Gulsuyu Mah. Fevzi Cakmak Cad. Divan Sok. No:1, Maltepe, Istanbul. bildirici@superonline.com, Phone: +90 2123044042, Fax: +90 2122854584.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

induction, and the mechanisms for execution, of autophagy in mammalian cells. We next describe what little is known about autophagy in the placenta, we lastly highlight key areas for research as we opine that such research will shed light on the placental dysfunction associated with pregnancy maladies.

Induction and execution of autophagy

There are three types of autophagy. Microautophagy involves engulfment of cytoplasmic components by invaginations or protrusions of the lysosomal membrane [12]. In chaperone-mediated autophagy (CMA), a KFERQ-like pentapeptide in the target proteins is bound by a cytosolic chaperone, which results in translocation of the substrate to the lysosome for degradation [13]. Macroautophagy, designated hereafter as autophagy, is the focus of this review and involves delivery of cytosolic contents to lysosomes by autophagosomes. Table 1 summarizes mediators of the autophagy pathway.

Autophagy induction and cargo recognition leads to formation of the phagophore, a C-shaped double membrane organelle that sequesters cytoplasm or a specific organelle (Fig 1.). Closure of the phagophore around the cargo results in a double-membrane organelle, the autophagosome. The autophagosome then fuses with lysosomes via the help of the cytoskeleton. Fusion of the outer membrane of the autophagosome with the lysosomal membrane results in the entry of the inner membrane of the autophagosome into the membrane of the lysosome. Lysosomal enzymes degrade the inner membrane of the autophagosome and the cargo therein contained, releasing free fatty acids, monosaccharides, and amino acids into the cytoplasm for re-use.

Induction of autophagy requires an Unc-51-like kinase (ULK) complex that contains two related kinases, ULK1 and ULK2, a scaffold protein, FIP200 (the focal adhesion kinase family interacting protein of 200 kD), and mAtg13 [14-16]. Inactivation of mTOR yields dephosphorylation and activation of ULK1 and ULK2 and allows phosphorylation of FIP200 and mAtg13. The recently identified Atg101 binds and stabilizes Atg13 in the ULK-Atg13-FIP200 complex [17]. The nucleation and assembly of the initial phagophore membrane involves a class III phosphatidylinositol 3-complex (PI3K) consisting of the lipid kinase, vacuolar protein sorting 34 (VPS34), VPS15, Beclin 1, Atg14L and the activating molecule in Beclin 1-regulated autophagy (AMBRA 1) [18]. Phosphorylation of AMBRA 1 by ULK and association of Beclin 1 increase the PI3 kinase activity of VPS34. The activated PI3K complex results in the generation of phosphatidylinositol triphosphate (PI3P), which is essential for phagophore elongation and for the recruitment of heterodimers of Atg5 and Atg12, in combination with a homodimer of Atg16, to phagophore assembly sites. Membranes for the phagophore are initially derived from the ER, trans-Golgi network and late endosomes [19], but *de novo* synthesis of membranes is likely involved in the expansion of the phagophore. An ubiquitin-like system modulates the evolution of the autophagosomal membranes and yields the processing of microtubule-associated protein light chain 3-I (LC3-I) into the LC3-II isoform, which covalently attaches phosphatidylethanolamine. LC3 II is incorporated into phosphatidylethanolamine-rich lipid membranes during autophagosome genesis and thus, is commonly used as a specific marker for autophagosome formation [20]. Atg9 is an integral membrane protein, which can form multimers and which cycles between the autophagosomal membrane and other membranes. This may enhance phagophore expansion and phagophore tethering to the lysosome [21]. After phagophore expansion, closure and capture of cargo occurs, forming a mature autophagosome. Autophagosomes are trafficked to lysosomes upon microtubules by dynein motors, and en route, these may fuse with endosomes to form amphisomes. Fusion with lysosomes occurs, and the degradation of the lysosomal contents by hydrolases releases monosaccharides, amino acids, and free fatty acids into the cytoplasm for recycling.

Regulation of autophagy

Nutrient deprivation is a potent inducer of autophagy, and mTOR is at the center of nutrient sensing in cells. MTOR provides the catalytic subunit for two distinct multiprotein complexes, mTORC1 and mTORC2 (Fig 2.), which share some proteins, e.g., GβL and Deptor, but have others, such as raptor in mTORC1 and rictor in mTORC2, that reflect the unique functions of each complex [22-23]. For example, raptor binds to mTOR signaling motifs on downstream targets, notably on ribosomal S6 protein kinase 1 (S6K1) and eukaryotic initiation factor 4-binding protein 1 (4EBP1). Proline-rich Akt substrate 40 kDa (PRAS40) associates with raptor to mediate the TOR signaling motif and this functions as a bridge downstream of mTORC1 and upstream of the effectors S6K1 and 4EBP1 [24-30]. Importantly, mTORC1 coordinates lipid and protein synthesis to enhance cell growth. Conversely, mTORC2 is implicated in the control of the actin cytoskeleton [31]. Inhibition of mTORC1 by rapamycin, for example, inhibits translation and activates ULK1/2 complex, which yields autophagy. An important regulator of mTOR is a small GTP-ase module consisting of Rheb, a small G-protein that activates mTORC1, and of the TSC1 (Hamartin)/TSC2 (Tuberin), which can stimulate Rheb GTPase activity [32].

Sufficiency of amino acids, energy, and growth factors promote mTORC1, whereas regulation of mTORC2 remains mostly unknown [33]. In HeLa cells, bidirectional transport of glutamine out of the cell and leucine into the cell by large neutral amino acids transporter small subunit 1 (SLC7A5 or LAT-1) induce phosphorylation of mTORC1 and inhibit autophagy. In skeletal muscle, increased availability of essential amino acids is shown to both stimulate mTORC1 signaling and increase amino acid transporter expression [34-35]. In the presence of amino acids, Rag proteins, the Ras-related small GTPases, also activate mTOR and inhibit autophagy [36]. Conversely, limited availability of these amino acids inhibits mTORC1 and induces autophagy.

In yeast, Ras/cAMP-dependent protein kinase A (PKA) signaling pathway plays an important role in glucose sensing and negatively regulates autophagy in parallel with the mTOR pathway [37-38]. In nutrient-rich conditions, the small GTPases, Ras 1 and Ras 2 are active and enhance cAMP generation, resulting in activation of protein kinase A (PKA), with downstream suppression of TSC and activation of mTORC1.

Growth factors suppress TSC function by both PI3K-dependent and independent pathways. The latter occurs via Ras and MEK downstream pathways with activation of the mTORC1 signaling [32]. Insulin and insulin-like growth factors (IGF) regulate mTOR through the Class I PI3K. Insulin binding to the insulin receptor induces autophosphorylation, and recruitment of insulin receptor substrates (IRS) 1 and 2, where PI3K binds to yield phosphatidylinositol (3,4,5)-triphosphate (PIP₃), increases membrane recruitment of protein kinase B (Akt) and phosphorylation of phosphoinositide-dependent protein kinase 1 (PDK1). PDK1 phosphorylates Akt which in turn phosphorylates TSC2, blocking its interaction with TSC1 [39] and thereby allowing GTP-Rheb to bind and maintain active mTORC1, inhibiting autophagy [40-42]. Conversely, the 3'-phosphoinositide phosphatase PTEN reverses PIP₃ production, decreasing Akt signaling and activating autophagy [43].

Phosphorylation and deacetylation are among important post-translational modifications for the regulation of autophagy. Recently, specific mTOR phosphorylation sites such as Ser¹²⁶¹, Ser²¹⁵⁹ and Thr 2164 that regulate mTORC1 function were identified [44-45]. The NAD-dependent deacetylase sirtuin-1 forms a complex with several proteins of the autophagy machinery and deacetylates them to allow autophagy to proceed [46].

Energy levels are detected through the 5'-AMP-activated protein kinase (AMPK), which senses reduced cellular ATP levels. A decreased ATP/AMP ratio activates AMPK through

the upstream LKB1 kinase. AMPK then phosphorylates the TSC1/2 complex, which promotes GTP-Rheb activity, inhibiting mTOR activity [47]. AMPK also inhibits mTOR by directly phosphorylating raptor [48]. Unlike necrosis or apoptosis, cellular ATP production increases throughout the autophagic process that recycles nutrients. Moreover, the LKB1-AMPK pathway activates the cyclin-dependent kinase inhibitor, P27^{kip1}, leading to cell-cycle arrest [49]. The end result of the induced autophagy is survival through partial self-destruction, cell cycle arrest, and avoidance of apoptotic cell death, despite the presence of nutrient deprivation.

Hypoxia induces autophagy by at least two mechanisms. First, prolonged hypoxia reduces ATP levels and thus activates AMPK. Secondly, hypoxia stabilizes hypoxia-inducible factor 1 (Hif 1), a transcription factor that is a major regulator of the cellular response to hypoxia [50]. Increased expression of Redd1 in hypoxia which is in part due to transcription activity by Hif 1, activates TSC1/2, suppresses mTORC1 activity and results in autophagy [51]. Hif 1 also induces transcription of Bcl-2/adenovirus E1a 19-kda interacting protein 3 (BNIP3), and the related BNIP3L [52-54]. Bnp3/Bnip3L binds Bcl-2 and Bcl-XL, disrupting the interaction with Beclin, allowing Beclin to associate with the class I PI3K complex and activate autophagy. Notably, low oxygen levels play a vital role in early placental development, as invasive trophoblast normally differentiate in a pO₂ < 20 mm Hg [55]. However, by 10-12 weeks of gestation, oxygen delivery to the intervillous space by maternal spiral arterioles raises the pO₂ to about 60 mm Hg, and the developing fetus is dependent upon adequate placental oxygen delivery for the remainder of pregnancy.

Oxidative stress plays a major role in the induction of autophagy, especially by generation of superoxide anions [56-57]. The electron transport system that provides energy from mitochondria generates multiple reactive oxygen species (ROS) and hydrogen peroxide, all of which must be eliminated by cell anti-oxidant defenses to avoid cellular damage. When cell oxidant levels exceed anti-oxidant defenses, lipids, proteins, DNA, mitochondria, and other organelles are damaged. Induction of autophagy can eliminate these damaged components. Induced mitophagy eliminates damaged mitochondria, reducing ROS generation and contributing to cell survival [58-59]. The cysteine protease Atg4 cleaves Atg8/LC3 from the autophagosome outer membrane. ROS inhibit Atg4 protease activity and enhance lipidation of Atg8/LC3, which is essential for autophagy [60]. Atg4 and ROS through the activity of poly (ADP-ribose) polymerase-1 (PARP-1) stimulates the LKB1-AMPK pathway to induce autophagy [61].

Crosstalk between apoptosis and autophagy

Autophagy and apoptosis are both important responses to cellular stress [62-64]. Apoptosis and autophagy can act cooperatively to yield cell death, but recent data show that autophagy often promotes cell survival by antagonizing apoptosis. Not surprisingly, cross talk between the two phenomena is extensive, as recently reviewed [65-67]. There has been significant progress in the identification of mediators that participate in both autophagy and apoptosis.

Notably, Beclin 1, identified as Atg6, was found to be a binding partner for Bcl-2, the first identified member of a family of proteins that controls apoptosis [68]. Bcl-2 family members play opposing roles in the regulation of apoptosis and autophagy [69], while Beclin 1 is a key regulator of autophagy. The anti-apoptotic proteins Bcl-2 and Bcl-XL can each bind to a BH3 domain in Beclin 1 to inhibit autophagy [70], especially when nutrients are plentiful [71]. When nutrients are depleted, JNK1 phosphorylates Bcl-2, Death Associated Protein Kinase (DAPK) phosphorylates Beclin 1 and both phosphorylations facilitate Bcl-2 separation from Beclin 1, the latter activating autophagy through interaction with AMBRA 1, VPS 15 and 34 [72-73]. During prolonged starvation, the phosphorylated

Bcl-2 binds to the proapoptotic protein Bax to inhibit apoptosis, but when survival is no longer possible, JNK1 mediates hyperphosphorylation of Bcl-2, Bax detaches, and apoptosis ensues [74-75]. Atg5 is cleaved by calpains during apoptosis and cleaved Atg5 interacts with Bcl-XL in mitochondria to control apoptosis. The absence of mitochondrial Atg5 yields autophagy [76].

Flc-inhibitory protein (FLIP) plays a double role, as a negative regulator of the extrinsic pathway to apoptosis and as an inhibitor of autophagy by blocking Atg3 conjugation to LC3 [77]. If caspases are activated, they not only degrade apoptotic substrates but also degrade autophagic proteins such as Beclin 1 to inhibit autophagy and potentiate apoptosis [78-80]. Conversely, cell survival is promoted through autophagy when Beclin 1 physically interacts with survivin to prolong the function of this inhibitor of apoptosis [81].

Sequestosome (p62) is a multifunctional protein that binds to Atg8/LC3 yielding proteasome activity and autophagy [82]. Importantly, p62 is also required for formation and full activation of the caspase-8 complex, important as an initiator caspase in the extrinsic pathway [83]. Moreover, LC3-II binds p62 to deliver misfolded proteins and injured organelles to the autophagosome for degradation.

P53 is a pivotal protein in effecting apoptosis and this protein regulates autophagy in a bidirectional manner, depending upon the sub-cellular location of p53 action. Cytoplasmic p53 blocks autophagy while nuclear p53 functions in transcriptional regulation of sestrins (SES) 1 and 2 and provides these proteins to phosphorylate AMP-responsive protein kinase (AMPK), which in turn phosphorylates TSC2 and activates autophagy. Other nuclear effects of p53 include increased IGFBP3, PTEN, and ARF. Interestingly, damage-regulated autophagy modulator (DRAM) is also essential in p53-mediated apoptosis further illustrating the intertwining of autophagy and apoptosis [84-85]. P53 promotes phosphorylation of Bcl-2, which dissociates the latter protein from Beclin 1 to enhance autophagy. P53 regulates DRAM expression and this lysosomal protein also enhances autophagy. Conversely, the p53-induced glycolysis and apoptosis regulator (TIGAR) inhibits glycolysis and decreases cellular ROS levels to inhibit autophagy [86]. The complexity of p53 regulation of autophagy is intimately related to the activity of p53 in apoptosis.

Autophagy in placenta

A role for autophagy during the trophoblast interactions with the endometrium in early implantation was implicated more than 35 years ago [87]. However, only a few studies to date focus on autophagy in placental function. Approximately 15 years ago, APG9L2, a mammalian homologue of the first identified autophagy protein (Atg9p) in yeast, was shown to be differentially expressed in the placenta and pituitary gland but not other tissues, while another homologue APG9L1 was ubiquitously expressed in adult human tissues [88]. Notably, APG9L2 showed a four fold decrease in expression in syncytiotrophoblast compared to cytotrophoblast, suggesting a developmental regulation during trophoblastic differentiation.

Autophagy occurs in trophoblast from term placental villi, with a higher level of autophagy in villi from cesarean section placentas reflected by accumulation of more LC3-II, compared to placentas from spontaneous vaginal delivery [89]. The lower cord blood glucose levels of infants delivered by cesarean section, from maternal fasting prior to the operation, is suggested to be the etiology of this counterintuitive proposition. The result may reflect the fact that autophagy is activated as a survival mechanism during nutrient deprivation [89]. Roh and colleagues [90] showed increased expression of LC3-II, but not Beclin 1, in placentas of women with severe preeclampsia, compared to controls. In the JEG-3

choriocarcinoma cell line, hypoxia resulted in increased LC3-II and diminished Beclin 1. Interestingly, TNF- α is also a known inducer of apoptosis in primary human trophoblasts [91] and this cytokine increases LC3-II expression in the JEG cells [90].

The mTOR system is a pivotal regulator of autophagy, as described above, and this nutrient sensing system has recently been studied in primary human trophoblasts [92]. Glucose deprivation down-regulates system L activity in an mTOR dependent manner in cultures of trophoblast [92]. Conversely, IGF-1 increases system A and insulin increases system L amino acid transport activity, both in an mTOR dependent manner [92]. mTOR signaling regulates the activity of key amino acid transporters [93] and both mTOR signaling and placental amino acid transporters are down regulated by maternal protein deprivation [94]. Interestingly, placentas from pregnancies complicated by IUGR have reduced mTOR activity [95-96].

A recent study found that, in addition to mTOR, human placental villous trophoblasts express Deptor, Raptor, and Rictor, other components of mTORC1. Interestingly, Deptor expression is reduced in placental villi exposed to maternal stress *in vivo* and in trophoblast cell lines exposed to the stress hormone cortisol. Thus, maternal stress may influence autophagy in placenta [97].

An interesting study by Hung and colleagues [98] noted concomitant exposure to vitamins C and E differentially affected trophoblast apoptosis and autophagy in villous explants under conditions simulating normoxia and hypoxia with re oxygenation *in vivo*. Specifically, vitamins C and E reduced the levels of apoptotic markers under normoxic conditions but increased apoptosis when hypoxia re oxygenation occurred. Moreover, vitamin treatment led to reduced LC3 immunostaining and reduced LC3B expression on immunoblot analysis, in explants exposed to 8% oxygen to simulate normoxic conditions. Conversely, there was increased expression of LC3B-II in both trophoblast and stromal cells in hypoxia re oxygenation conditions when vitamin C and E were present, but not in the 8% oxygen paradigm. Notably, cultured cytotrophoblasts treated with vitamins C and E under hypoxia re-oxygenation conditions showed decreased expression of the anti-apoptotic proteins, Bcl-2 and Bcl-XL, increased expression of pro-apoptotic Bak, enhanced cytosolic LC3-II and decreased interaction of Beclin 1 with Bcl-2 and the Bcl-XL/Beclin 1 complex. These results are consistent with enhanced autophagy under conditions of re-oxygenation [98]. These studies suggest that insults to placental villi and trophoblast cells yield condition-specific responses in the balance between trophoblast autophagy and apoptosis.

The importance of placental autophagy in fetal development and survival is underscored by effects of maternal starvation in a mouse model of the imprinted gene PEG 3 (paternally expressed gene 3) [99]. Undernutrition in mid-gestation desynchronizes the expression of PEG 3, increasing PEG 3 expression in fetal hypothalamus and decreasing PEG 3 expression in placenta. This desynchronization results in enhanced placental autophagy, specifically ribophagy, which produces nutrients to sustain energy demands of the hypothalamus. These data suggest that the fetus in general and the fetal brain in particular is spared during short-term acute maternal starvation, at the expense of the placenta.

Future research on autophagy and the placenta

The regulation of autophagy and the mechanisms involved with induction in non-placental tissues have made substantial strides in the last decade. The role mTOR in nutrient sensing of the placenta is established, but what role this protein plays in the balance of trophoblast, and thus fetal survival and demise is unknown. Moreover, autophagy has only recently been identified in human placentas. Several key questions must be addressed. First, what role does autophagy play in trophoblast homeostasis? Does autophagy contribute to placental

dysfunction in the maladies of pregnancy? Alternatively, does autophagy provide protection for placentas exposed to exogenous stressors during pregnancy, perhaps by inhibiting apoptosis and aiding in nutrient recycling? What is the interplay of autophagy and apoptosis in the trophoblast layer, particularly in complicated pregnancies? We speculate that future research will identify key roles for autophagy in placental function of women with preeclampsia, IUGR, or both. We also predict that autophagy will be pivotal in the response of trophoblasts to stressors present in normal pregnancies, allowing the trophoblast, in particular, and placental villi in general, to adapt to the variety of insults normally encountered in pregnant women. We believe that self-destruction by autophagy has a unique mission to protect placental function during stress and thereby, to optimize fetal development. We propose that studies evaluating the regulation of autophagy in the placenta will offer new insights into placental function under homeostatic conditions and the placental dysfunction that accompanies exposure to exogenous insults.

Acknowledgments

This review was supported in part by NIH HD 29190 (DMN).

References

- [1]. Tsukada M, Ohsumi Y. Isolation and characterization of autophagy-defective mutants of *Saccharomyces cerevisiae*. FEBS Lett. 1993; 333:169–74. [PubMed: 8224160]
- [2]. Thumm M, Egner R, Koch B, Schlumberger M, Straub M, Veenhuis M, et al. Isolation of autophagocytosis mutants of *Saccharomyces cerevisiae*. FEBS Lett. 1994; 349:275–80. [PubMed: 8050581]
- [3]. Harding TM, Morano KA, Scott SV, Klionsky DJ. Isolation and characterization of yeast mutants in the cytoplasm to vacuole protein targeting pathway. J Cell Biol. 1995; 131:591–602. [PubMed: 7593182]
- [4]. Matsuura A, Tsukada M, Wada Y, Ohsumi Y. Apg1p, a novel protein kinase required for the autophagic process in *Saccharomyces cerevisiae*. Gene. 1997; 192:245–50. [PubMed: 9224897]
- [5]. Klionsky DJ, Ohsumi Y. Vacuolar import of proteins and organelles from the cytoplasm. Annu Rev Cell Dev Biol. 1999; 15:1–32. [PubMed: 10611955]
- [6]. Klionsky DJ, Codogno P, Cuervo AM, Deretic V, Elazar Z, Fueyo-Margareto J, et al. A comprehensive glossary of autophagy-related molecules and processes. Autophagy. 2010; 6:438–48.
- [7]. Lockshin RA, Zakeri Z. Apoptosis, autophagy, and more. Int J Biochem Cell Biol. 2004; 36:2405–19. [PubMed: 15325581]
- [8]. Kroemer G, Levine B. Autophagic cell death: the story of a misnomer. Nat Rev Mol Cell Biol. 2008; 9:1004–10. [PubMed: 18971948]
- [9]. Del Roso A, Vittorini S, Cavallini G, Donati A, Gori Z, Masini M, et al. Ageing-related changes in the in vivo function of rat liver macroautophagy and proteolysis. Exp Gerontol. 2003; 38:519–27. [PubMed: 12742529]
- [10]. Donati A, Cavallini G, Paradiso C, Vittorini S, Pollera M, Gori Z, et al. Age-related changes in the regulation of autophagic proteolysis in rat isolated hepatocytes. J Gerontol A Biol Sci Med Sci. 2001; 56:B288–93. [PubMed: 11445593]
- [11]. Cavallini G, Donati A, Gori Z, Bergamini E. Towards an understanding of the anti-aging mechanism of caloric restriction. Curr Aging Sci. 2008; 1:4–9. [PubMed: 20021367]
- [12]. Mijaljica D, Prescott M, Devenish RJ. Microautophagy in mammalian cells: revisiting a 40-year-old conundrum. Autophagy. 2011; 7:673–82. [PubMed: 21646866]
- [13]. Orenstein SJ, Cuervo AM. Chaperone-mediated autophagy: molecular mechanisms and physiological relevance. Semin Cell Dev Biol. 2010; 21:719–26. [PubMed: 20176123]
- [14]. Mizushima N. The role of the Atg1/ULK1 complex in autophagy regulation. Curr Opin Cell Biol. 2010; 22:132–9. [PubMed: 20056399]

- [15]. Neufeld TP. TOR-dependent control of autophagy: biting the hand that feeds. *Curr Opin Cell Biol.* 2010; 22:157–68. [PubMed: 20006481]
- [16]. Wang RC, Levine B. Autophagy in cellular growth control. *FEBS Lett.* 2010; 584:1417–26. [PubMed: 20096689]
- [17]. Mercer CA, Kaliappan A, Dennis PB. A novel, human Atg13 binding protein, Atg101, interacts with ULK1 and is essential for macroautophagy. *Autophagy.* 2009; 5:649–62. [PubMed: 19287211]
- [18]. Funderburk SF, Wang QJ, Yue Z. The Beclin 1-VPS34 complex--at the crossroads of autophagy and beyond. *Trends Cell Biol.* 2010; 20:355–62. [PubMed: 20356743]
- [19]. Rubinsztein DC, Shpilka T, Elazar Z. Mechanisms of autophagosome biogenesis. *Curr Biol.* 2012; 22:R29–34. [PubMed: 22240478]
- [20]. Glick D, Barth S, Macleod KF. Autophagy: cellular and molecular mechanisms. *J Pathol.* 2010; 221:3–12. [PubMed: 20225336]
- [21]. He C, Baba M, Cao Y, Klionsky DJ. Self-interaction is critical for Atg9 transport and function at the phagophore assembly site during autophagy. *Mol Biol Cell.* 2008; 19:5506–16. [PubMed: 18829864]
- [22]. Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, et al. Ablation in mice of the mTORC components raptor, rictor, or mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKCalpha, but not S6K1. *Dev Cell.* 2006; 11:859–71. [PubMed: 17141160]
- [23]. Gao D, Inuzuka H, Tan MK, Fukushima H, Locasale JW, Liu P, et al. mTOR drives its own activation via SCF(β TrCP)-dependent degradation of the mTOR inhibitor DEPTOR. *Mol Cell.* 2011; 44:290–303. [PubMed: 22017875]
- [24]. Brown EJ, Albers MW, Shin TB, Ichikawa K, Keith CT, Lane WS, et al. A mammalian protein targeted by G1-arresting rapamycin-receptor complex. *Nature.* 1994; 369:756–8. [PubMed: 8008069]
- [25]. Sarbassov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, et al. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Mol Cell.* 2006; 22:159–68. [PubMed: 16603397]
- [26]. Hara K, Maruki Y, Long X, Yoshino K, Oshiro N, Hidayat S, et al. Raptor, a binding partner of target of rapamycin (TOR) mediates TOR action. *Cell.* 2002; 110:177–89. [PubMed: 12150926]
- [27]. Nojima H, Tokunaga C, Eguchi S, Oshiro N, Hidayat S, Yoshino K, et al. The mammalian target of rapamycin (mTOR) partner, raptor, binds the mTOR substrates p70 S6 kinase and 4E-BP1 through their TOR signaling (TOS) motif. *J Biol Chem.* 2003; 278:15461–4. [PubMed: 12604610]
- [28]. Kim DH, Sarbassov DD, Ali SM, Latek RR, Guntur KV, Erdjument-Bromage H, et al. GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol Cell.* 2003; 11:895–904. [PubMed: 12718876]
- [29]. Fonseca DB, Ewan MS, Lee VH, MacKintosh C, Proud CG. PRAS40 is a target for mammalian target of rapamycin complex 1 and is required for signaling downstream of this complex. *J Biol Chem.* 2007; 282:24514–24. [PubMed: 17604271]
- [30]. Nyfeler B, Bergman P, Triantafellow E, Wilson CJ, Zhu Y, Radetich B, et al. Relieving autophagy and 4EBP1 from rapamycin resistance. *Mol Cell Biol.* 2011; 31:2867–76. [PubMed: 21576371]
- [31]. Jacinto E, Loewith R, Schmidt A, Lin S, Ruegg MA, Hall A, et al. Mammalian TOR complex 2 controls the actin cytoskeleton and is rapamycin insensitive. *Nat Cell Biol.* 2004; 6:1122–8. [PubMed: 15467718]
- [32]. Huang J, Manning BD. The TSC1-TSC2 complex: a molecular switchboard controlling cell growth. *Biochem J.* 2008; 412:179–90. [PubMed: 18466115]
- [33]. Foster KG, Fingar DC. Mammalian target of rapamycin (mTOR): conducting the cellular signaling symphony. *J Biol Chem.* 2010; 285:14071–7. [PubMed: 20231296]
- [34]. Nicklin P, Bergman P, Zhang B, Triantafellow E, Wang H, Nyfeler B, et al. Bidirectional transport of amino acids regulates mTOR and autophagy. *Cell.* 2009; 136:521–34. [PubMed: 19203585]

- [35]. Drummond MJ, Glynn EL, Fry CS, Timmerman KL, Volpi E, Rasmussen BB. An increase in essential aminoacid availability upregulates aminoacid transporter expression in human skeletal muscle. *Am J Physiol Endocrinol Metab.* 2010; 298:E1011–8. [PubMed: 20304764]
- [36]. Kim E, Goraksha-Hicks P, Li L, Neufeld TP, Guan KL. Regulation of TORC1 by Rag GTPases in nutrient response. *Nat. Cell Biol.* 2008; 10:935–45. [PubMed: 18604198]
- [37]. Budovskaya YV, Stephan JS, Reggiori F, Klionsky DJ, Herman PK. The Ras/cAMP-dependent protein kinase signaling pathway regulates an early step of the autophagy process in *Saccharomyces cerevisiae*. *J Biol Chem.* 2004; 279:20663–71. [PubMed: 15016820]
- [38]. Schmelzle T, Beck T, Martin DE, Hall MN. Activation of the RAS/cyclicAMP pathway suppresses a TOR deficiency in yeast. *Mol Cell Biol.* 2004; 24:338–51. [PubMed: 14673167]
- [39]. Manning BD, Tee AR, Logsdon MN, Blenis J, Cantley LC. Identification of the tuberous sclerosis complex-2 tumor suppressor gene product tuberlin as a target of the phosphoinositide 3-kinase/Akt pathway. *Mol Cell.* 2002; 10:151–62. [PubMed: 12150915]
- [40]. Long X, Lin Y, Ortiz-Vega S, Yonezawa K, Avruch J. Rheb binds and regulates the mTOR kinase. *Curr Biol.* 2005; 15:702–13. [PubMed: 15854902]
- [41]. Alessi DR, James SR, Downes CP, Holmes AB, Gaffney PR, Reese CB, et al. Characterization of α 3-phosphoinositide-dependent protein kinase which phosphorylates and activates protein kinase B. *Curr Biol.* 1997; 7:261–9. [PubMed: 9094314]
- [42]. Stokoe D, Stephens LR, Copeland T, Gaffney PR, Reese CB, Painter GF, et al. Dual role of phosphatidylinositol-3,4,5-trisphosphate in the activation of protein kinase B. *Science.* 1997; 277:567–70. [PubMed: 9228007]
- [43]. Arico S, Petiot A, Bauvy C, Dubbelhuis PF, Meijer AJ, Codogno P, et al. The tumor suppressor PTEN positively regulates macroautophagy by inhibiting the phosphatidylinositol 3-kinase/protein kinase B pathway. *J Biol Chem.* 2001; 276:35243–6. [PubMed: 11477064]
- [44]. Acosta-Jaquez HA, Keller JA, Foster KG, Ekim B, Soliman GA, Feener EP, et al. Site-specific mTOR phosphorylation promotes mTORC1 mediated signaling and cell growth. *Mol Cell Biol.* 2009; 29:4308–24. [PubMed: 19487463]
- [45]. Ekim B, Magnuson B, Acosta-Jaquez HA, Keller JA, Feener EP, Fingar DC. mTOR kinase domain phosphorylation promotes mTORC1 signaling, cell growth, and cell cycle progression. *Mol Cell Biol.* 2011; 31:2787–801. [PubMed: 21576368]
- [46]. Lee IH, Cao L, Mostoslavsky R, Lombard DB, Liu J, Bruns NE, et al. A role for the NAD-dependent deacetylase Sirt1 in the regulation of autophagy. *Proc Natl Acad Sci USA.* 2008; 105:3374–9. [PubMed: 18296641]
- [47]. Inoki K, Zhu T, Guan KL. TSC2 mediates cellular energy response to control cell growth and survival. *Cell.* 2003; 115:577–90. [PubMed: 14651849]
- [48]. Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, et al. AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol Cell.* 2008; 30:214–26. [PubMed: 18439900]
- [49]. Liang J, Shao SH, Xu ZX, Hennessy B, Ding Z, Larrea M, et al. The energy sensing LKB1-AMPK pathway regulates p27(kip1) phosphorylation mediating the decision to enter autophagy or apoptosis. *Nat Cell Biol.* 2007; 9:218–24. [PubMed: 17237771]
- [50]. Majmundar AJ, Wong WJ, Simon MC. Hypoxia-inducible factors and the response to hypoxic stress. *Mol Cell.* 2010; 40:294–309. [PubMed: 20965423]
- [51]. DeYoung MP, Horak P, Sofer A, Sgroi D, Ellisen LW. Hypoxia regulates TSC1/2-mTOR signaling and tumor suppression through REDD1-mediated 14-3-3 shuttling. *Genes Dev.* 2008; 22:239–51. [PubMed: 18198340]
- [52]. Azad MB, Chen Y, Henson ES, Cizeau J, McMillan-Ward E, Israels SJ, et al. Hypoxia induces autophagic cell death in apoptosis-competent cells through a mechanism involving BNIP3. *Autophagy.* 2008; 4:195–204. [PubMed: 18059169]
- [53]. Zhang H, Bosch-Marce M, Shimoda LA, Tan YS, Baek JH, Wesley JB, et al. Mitochondrial autophagy is an HIF-1-dependent adaptive metabolic response to hypoxia. *J Biol Chem.* 2008; 283:10892–903. [PubMed: 18281291]

- [54]. Bellot G, Garcia-Medina R, Gounon P, Chiche J, Roux D, Pouyssegur J, et al. Hypoxia-induced autophagy is mediated through hypoxia-inducible factor induction of BNIP3 and BNIP3L via their BH3 domains. *Mol Cell Biol*. 2009; 29:2570–81. [PubMed: 19273585]
- [55]. Genbacev O, Zhou Y, Ludlow JW, Fisher SJ. Regulation of human placental development by oxygen tension. *Science*. 1997; 277:1669–72. [PubMed: 9287221]
- [56]. Kiffin R, Bandyopadhyay U, Cuervo AM. Oxidative stress and autophagy. *Antioxid Redox Signal*. 2006; 8:152–62. [PubMed: 16487049]
- [57]. Chen Y, Azad MD, Gibson SB. Superoxide is the major reactive oxygen species regulating autophagy. *Cell Death Differ*. 2009; 16:1040–52. [PubMed: 19407826]
- [58]. Narendra D, Tanaka A, Suen DF, Youle RJ. Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *J Cell Biol*. 2008; 183:795–803. [PubMed: 19029340]
- [59]. Rodriguez-Hernandez A, Cordero MD, Salviati L, Artuch R, Pineda M, Briones P, et al. Coenzyme Q deficiency triggers mitochondria degradation by mitophagy. *Autophagy*. 2009; 5:19–32. [PubMed: 19115482]
- [60]. Scherz-Shouval R, Shvets E, Fass E, Shorer H, Gil L, Elazar Z. Reactive oxygen species are essential for autophagy and specifically regulate the activity of Atg4. *EMBO J*. 2007; 26:1749–60. [PubMed: 17347651]
- [61]. Huang Q, Wu YT, Tan HL, Ong CN, Shen HM. A novel function of poly(ADP-ribose) polymerase-1 in modulation of autophagy and necrosis under oxidative stress. *Cell Death Differ*. 2009; 16:264–77. [PubMed: 18974775]
- [62]. Terman A, Brunk UT. Autophagy in cardiac myocyte homeostasis, aging, and pathology. *Cardiovasc Res*. 2005; 68:355–65. [PubMed: 16213475]
- [63]. Nishida K, Kyo S, Yamaguchi O, Sadoshima J, Otsu K. The role of autophagy in the heart. *Cell Death Differ*. 2009; 16:31–8. [PubMed: 19008922]
- [64]. Mizushima N, Levine B. Autophagy in mammalian development and differentiation. *Nat Cell Biol*. 2010; 12:823–30. [PubMed: 20811354]
- [65]. Giansanti V, Torriglia A, Scovassi AI. Conversation between apoptosis and autophagy: “Is it your turn or mine?”. *Apoptosis*. 2011; 16:321–33. [PubMed: 21404107]
- [66]. Gump JM, Thorburn A. Autophagy and apoptosis: what is the connection? *Trends Cell Biol*. 2011; 21:387–92. [PubMed: 21561772]
- [67]. Maiuri MC, Zalckvar E, Kimchi A, Kroemer G. Self-eating and self-killing: crosstalk between autophagy and apoptosis. *Nat Rev Mol Cell Biol*. 2007; 8:741–52. [PubMed: 17717517]
- [68]. Liang XH, Kleeman LK, Jiang HH, Gordon G, Goldman JE, Berry G, et al. Protection against fetal Sindbis virus encephalitis by beclin, a novel Bcl-2-interacting protein. *J Virol*. 1998; 72:8586–96. [PubMed: 9765397]
- [69]. Zhou F, Yang Y, Xing D. Bcl-2 and Bcl-xL play important roles in the crosstalk between autophagy and apoptosis. *FEBS J*. 2011; 278:403–13. [PubMed: 21182587]
- [70]. Maiuri MC, Le Toumelin G, Criollo A, Rain JC, Gautier F, Julin P, et al. Functional and physical interaction between Bcl-X(L) and a BH3-like domain in Beclin-1. *EMBO J*. 2007; 26:2527–39. [PubMed: 17446862]
- [71]. Pattingre S, Tassa A, Qu X, Garuti R, Liang XH, Mizushima N, et al. Bcl-2 antiapoptotic proteins inhibit Beclin 1-dependent autophagy. *Cell*. 2005; 122:927–39. [PubMed: 16179260]
- [72]. Zalckvar E, Berissi H, Mizrachy L, Idelchuk Y, Koren I, Eisenstein M, et al. DAP-kinase-mediated phosphorylation on the BH3 domain of beclin 1 promotes dissociation of beclin 1 from Bcl-XL and induction of autophagy. *EMBO Rep*. 2009; 10:285–92. [PubMed: 19180116]
- [73]. Bovellan M, Fritzsche M, Stevens C, Charras G. Death associated protein kinase (DAPK) and signal transduction: blebbing in programmed cell death. *FEBS J*. 2010; 277:58–65. [PubMed: 19878312]
- [74]. Wei Y, Pattingre S, Sinha S, Bassik M, Levine B. JNK1-mediated phosphorylation of Bcl-2 regulates starvation-induced autophagy. *Mol Cell*. 2008; 30:678–88. [PubMed: 18570871]
- [75]. Wei Y, Sinha S, Levine B. Dual role of JNK1-mediated phosphorylation of Bcl-2 in autophagy and apoptosis regulation. *Autophagy*. 2008; 4:949–51. [PubMed: 18769111]

- [76]. Yousefi S, Perozzo R, Schmid I, Ziemiecki A, Schaffner T, Scapozza L, et al. Calpain-mediated cleavage of Atg5 switches autophagy to apoptosis. *Nat Cell Biol.* 2006; 8:1124–32. [PubMed: 16998475]
- [77]. Lee JS, Li Q, Lee JY, Lee SH, Jeong JH, Lee HR, et al. FLIP-mediated autophagy regulation in cell death control. *Nat Cell Biol.* 2009; 11:1355–62. [PubMed: 19838173]
- [78]. Luo S, Rubinsztein DC. Apoptosis blocks Beclin 1-dependent autophagosome synthesis: an effect rescued by Bcl-xL. *Cell Death Differ.* 2010; 17:268–77. [PubMed: 19713971]
- [79]. Cho DH, Jo YK, Hwang JJ, Lee YM, Roh SA, Kim JC. Caspase-mediated cleavage of ATG6/Beclin-1 links apoptosis to autophagy in HeLa cells. *Cancer Lett.* 2009; 274:95–100. [PubMed: 18842334]
- [80]. Djavaheri-Mergny M, Maiuri MC, Kroemer G. Cross talk between apoptosis and autophagy by caspase-mediated cleavage of Beclin 1. *Oncogene.* 2010; 29:1717–1719. [PubMed: 20101204]
- [81]. Niu TK, Cheng Y, Ren X, Yang JM. Interaction of Beclin 1 with survivin regulates sensitivity of human glioma cells to TRAIL-induced apoptosis. *FEBS Lett.* 2010; 584:3519–24. [PubMed: 20638385]
- [82]. Pankiv S, Clausen TH, Lamark T, Brech A, Bruun JA, Outzen H, et al. p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem.* 2007; 282:24131–45. [PubMed: 17580304]
- [83]. Jin Z, Li Y, Pitti R, Lawrence D, Pham VC, Lill JR, et al. Cullin3-based polyubiquitination and p62-dependent aggregation of caspase-8 mediate extrinsic apoptosis signalling. *Cell.* 2009; 137:721–35. [PubMed: 19427028]
- [84]. Crighton D, Wilkinson S, O'Prey J, Syed N, Smith P, Harrison PR, et al. DRAM, a p53-induced modulator of autophagy, is critical for apoptosis. *Cell.* 2006; 126:121–34. [PubMed: 16839881]
- [85]. Crighton D, Wilkinson S, Ryan KM. DRAM links autophagy to p53 and programmed cell death. *Autophagy.* 2007; 3:72–4. [PubMed: 17102582]
- [86]. Bensaad K, Cheung EC, Vousden KH. Modulation of intracellular ROS levels by TIGAR controls autophagy. *EMBO J.* 2009; 28:3015–26. [PubMed: 19713938]
- [87]. Leiser R, Wille KH. Cytochemical establishment of acid phosphatase in the bovine endometrium and trophoblast during implantation. *Anat Embryol (Berl).* 1975; 148:159–73. [PubMed: 1211660]
- [88]. Yamada T, Carson AR, Caniggia I, Umebayashi K, Yoshimori T, Nakabayashi K, et al. Endothelial nitric-oxide synthase antisense (NOS3AS) gene encodes an autophagy-related protein (APG9-like2) highly expressed in trophoblast. *J Biol Chem.* 2005; 280:18283–90. [PubMed: 15755735]
- [89]. Signorelli P, Avagliano L, Virgili E, Gagliostro V, Doi P, Braidotti P, et al. Autophagy in term normal human placentas. *Placenta.* 2011; 32:482–5. [PubMed: 21459442]
- [90]. Oh SY, Choi SJ, Kim KH, Cho EY, Kim JH, Roh CR. Autophagy-related proteins, LC3 and Beclin-1, in placentas from pregnancies complicated by preeclampsia. *Reprod Sci.* 2008; 15:912–20. [PubMed: 19050324]
- [91]. Garcia-Lloret MI, Yui J, Winkler-Lowen B, Guilbert LJ. Epidermal growth factor inhibits cytokine-induced apoptosis of primary human trophoblasts. *J Cell Physiol.* 1996; 167:324–32. [PubMed: 8613474]
- [92]. Roos S, Lagerlof O, Wennergren M, Powell TL, Jansson T. Regulation of amino acid transporters by glucose and growth factors in cultured primary human trophoblast cells is mediated by mTOR signaling. *Am J Physiol Cell Physiol.* 2009; 297:C723–31. [PubMed: 19587219]
- [93]. Roos S, Kanai Y, Prasad PD, Powell TL, Jansson T. Regulation of placental amino acid transporter activity by mammalian target of rapamycin. *Am J Physiol Cell Physiol.* 2009; 296:C142–50. [PubMed: 18987252]
- [94]. Rosario FJ, Jansson N, Kanai Y, Prasad PD, Powell TL, Jansson T. Maternal protein restriction in the rat inhibits placental insulin, mTOR, and STAT3 signaling and down-regulates placental amino acid transporters. *Endocrinology.* 2011; 152:1119–29. [PubMed: 21285325]

- [95]. Roos S, Jansson N, Palmberg I, Säljö K, Powell TL, Jansson T. Mammalian target of rapamycin in the human placenta regulates leucine transport and is down-regulated in restricted fetal growth. *J Physiol*. 2007; 582:449–59. [PubMed: 17463046]
- [96]. Yung HW, Calabrese S, Hynx D, Hemmings BA, Cetin I, Charnock-Jones DS, et al. Evidence of placental translation inhibition and endoplasmic reticulum stress in the etiology of human intrauterine growth restriction. *Am J Pathol*. 2008; 173:451–62. [PubMed: 18583310]
- [97]. Mpampakas D, Zachariades E, Goumenou A, Gidron Y, Karteris E. Placental DEPTOR as a stress sensor during pregnancy. *Clin Sci (Lond)*. 2012; 122:349–59. [PubMed: 21992080]
- [98]. Hung TH, Chen SF, Li MJ, Yeh YL, Hsieh TT. Differential effects of concomitant use of vitamins C and E on trophoblast apoptosis and autophagy between normoxia and hypoxia-reoxygenation. *PLoS One*. 2010; 5:e12202. [PubMed: 20808946]
- [99]. Broad KD, Keverne EB. Placental protection of the fetal brain during short-term food deprivation. *Proc Natl Acad Sci U S A*. 2011; 108:15237–41. [PubMed: 21810990]
- [100]. Kim E, Guan KL. RAG GTPases in nutrient-mediated TOR signaling pathway. *Cell Cycle*. 2009; 8:1014–8. [PubMed: 19270521]
- [101]. Hwang SO, Lee GM. Effect of Akt overexpression on programmed cell death in antibody-producing Chinese hamster ovary cells. *J Biotechnol*. 2009; 139:89–94. [PubMed: 18983878]
- [102]. Degtyarev M, De Mazière A, Orr C, Lin J, Lee BB, Tien JY, et al. Akt inhibition promotes autophagy and sensitizes PTEN-null tumors to lysosomotropic agents. *J Cell Biol*. 2008; 183:101–16. [PubMed: 18838554]
- [103]. Fimia GM, Stoykova A, Romagnoli A, Giunta L, Di Bartolomeo S, Nardacci R, et al. Ambra1 regulates autophagy and development of the nervous system. *Nature*. 2007; 447:1121–5. [PubMed: 17589504]
- [104]. Cecconi F, Di Bartolomeo S, Nardacci R, Fuoco C, Corazzari M, Giunta L, et al. A novel role for autophagy in neurodevelopment. *Autophagy*. 2007; 3:506–8. [PubMed: 17622796]
- [105]. Strapozzon F, Vietri-Rudan M, Campello S, Nazio F, Florenzano F, Fimia GM, et al. Mitochondrial BCL-2 inhibits AMBRA1-induced autophagy. *EMBO J*. 2011; 30:1195–208. [PubMed: 21358617]
- [106]. Hardie DG. AMP-activated protein kinase: an energy sensor that regulates all aspects of cell function. *Genes Dev*. 2011; 25:1895–908. [PubMed: 21937710]
- [107]. Mihaylova MM, Shaw RJ. The AMPK signalling pathway coordinates cell growth, autophagy and metabolism. *Nat Cell Biol*. 2011; 13:1016–23. [PubMed: 21892142]
- [108]. Meley D, Bauvy C, Houben-Weerts JH, Dubbelhuis PF, Helmond MT, Codogno P, et al. AMP-activated protein kinase and the regulation of autophagic proteolysis. *J Biol Chem*. 2006; 281:34870–9. [PubMed: 16990266]
- [109]. Shimizu S, Kanaseki T, Mizushima N, Mizuta T, Arakawa-Kobayashi S, Thompson CB, et al. Role of Bcl-2 family proteins in a non-apoptotic programmed cell death dependent on autophagy genes. *Nat Cell Biol*. 2004; 6:1221–8. [PubMed: 15558033]
- [110]. Pattingre S, Levine B. Bcl-2 inhibition of autophagy: a new route to cancer? *Cancer Res*. 2006; 66:2885–8. [PubMed: 16540632]
- [111]. Levine B, Sinha S, Kroemer G. Bcl-2 family members: dual regulators of apoptosis and autophagy. *Autophagy*. 2008; 4:600–6. [PubMed: 18497563]
- [112]. Liang XH, Jackson S, Seaman M, Brown K, Kempkes B, Hibshoosh H, et al. Induction of autophagy and inhibition of tumorigenesis by beclin 1. *Nature*. 1999; 402:672–6. [PubMed: 10604474]
- [113]. Ku B, Woo JS, Liang C, Lee KH, Jung JU, Oh BH. An insight into the mechanistic role of Beclin 1 and its inhibition by prosurvival Bcl-2 family proteins. *Autophagy*. 2008; 4:519–20. [PubMed: 18334862]
- [114]. Park KJ, Lee SH, Lee CH, Jang JY, Chung J, Kwon MH, et al. Upregulation of Beclin-1 expression and phosphorylation of Bcl-2 and p53 are involved in the JNK-mediated autophagic cell death. *Biochem Biophys Res Commun*. 2009; 382:726–9. [PubMed: 19318089]
- [115]. Tracy K, Macleod KF. Regulation of mitochondrial integrity, autophagy and cell survival by BNIP3. *Autophagy*. 2007; 3:616–9. [PubMed: 17786027]

- [116]. Byun YJ, Kim SK, Kim YM, Chae GT, Jeong SW, Lee SB. Hydrogen peroxide induces autophagic cell death in C6 glioma cells via BNIP3-mediated suppression of the mTOR pathway. *Neurosci Lett*. 2009; 461:131–5. [PubMed: 19539716]
- [117]. Quinsay MN, Thomas RL, Lee Y, Gustafsson AB. Bnip3-mediated mitochondrial autophagy is independent of the mitochondrial permeability transition pore. *Autophagy*. 2010; 6:855–62. [PubMed: 20668412]
- [118]. Azad MB, Gibson SB. Role of BNIP3 in proliferation and hypoxia-induced autophagy: implications for personalized cancer therapies. *Ann N Y Acad Sci*. 2010; 1210:8–16. [PubMed: 20973794]
- [119]. Thomas RL, Kubli DA, Gustafsson AB. Bnip3-mediated defects in oxidative phosphorylation promote mitophagy. *Autophagy*. 2011; 7:775–7. [PubMed: 21460627]
- [120]. Stevens C, Lin Y, Harrison B, Burch L, Ridgway RA, Sansom O, et al. Peptide combinatorial libraries identify TSC2 as a death-associated protein kinase (DA PK) death domain-binding protein and reveal a stimulatory role for DAPK in mTORC1 signaling. *J Biol Chem*. 2009; 284:334–44. [PubMed: 18974095]
- [121]. Lorin S, Pierron G, Ryan KM, Codogno P, Djavaheri-Mergny M. Evidence for the interplay between JNK and p53-DRAM signalling pathways in the regulation of autophagy. *Autophagy*. 2010; 6:153–4. [PubMed: 19949306]
- [122]. O'Prey J, Skommer J, Wilkinson S, Ryan KM. Analysis of DRAM-related proteins reveals evolutionarily conserved and divergent roles in the control of autophagy. *Cell Cycle*. 2009; 8:2260–5. [PubMed: 19556885]
- [123]. Cagnol S, Chambard JC. ERK and cell death: mechanisms of ERK-induced cell death--apoptosis, autophagy and senescence. *FEBS J*. 2010; 277:2–21. [PubMed: 19843174]
- [124]. Ugland H, Naderi S, Brech A, Collas P, Blomhoff HK. cAMP induces autophagy via a novel pathway involving ERK, cyclin E and Beclin 1. *Autophagy*. 2011; 7:1199–211. [PubMed: 21750416]
- [125]. Wong CH, Iskandar KB, Yadav SK, Hirpara JL, Loh T, Pervaiz S. Simultaneous induction of non-canonical autophagy and apoptosis in cancer cells by ROS-dependent ERK and JNK activation. *PLoS One*. 2010; 5:e9996. [PubMed: 20368806]
- [126]. Wang J, Whiteman MW, Lian H, Wang G, Singh A, Huang D, et al. A non-canonical MEK/ERK signaling pathway regulates autophagy via regulating Beclin 1. *J Biol Chem*. 2009; 284:21412–24. [PubMed: 19520853]
- [127]. Corcelle E, Djerbi N, Mari M, Nebout M, Fiorini C, Fénichel P, et al. Control of the autophagy maturation step by the MAPK ERK and p38: lessons from environmental carcinogens. *Autophagy*. 2007; 3:57–9. [PubMed: 17102581]
- [128]. Sengupta A, Molkentin JD, Paik JH, DePinho RA, Yutzey KE. FoxO transcription factors promote cardiomyocyte survival upon induction of oxidative stress. *J Biol Chem*. 2011; 286:7468–78. [PubMed: 21159781]
- [129]. Chiacchiera F, Matrone A, Ferrari E, Ingravalle G, Lo Sasso G, Murzilli S, et al. p38alpha blockade inhibits colorectal cancer growth in vivo by inducing a switch from HIF1alpha- to FoxO-dependent transcription. *Cell Death Differ*. 2009; 16:1203–14. [PubMed: 19343039]
- [130]. Zhao J, Brault JJ, Schild A, Cao P, Sandri M, Schiaffino S, et al. FoxO3 coordinately activates protein degradation by the autophagic/lysosomal and proteasomal pathways in atrophying muscle cells. *Cell Metab*. 2007; 6:472–83. [PubMed: 18054316]
- [131]. Zhao J, Brault JJ, Schild A, Goldberg AL. Coordinate activation of autophagy and the proteasome pathway by FoxO transcription factor. *Autophagy*. 2008; 4:378–80. [PubMed: 18227643]
- [132]. Mazure NM, Pouyssegur J. Hypoxia-induced autophagy: cell death or cell survival? *Curr Opin Cell Biol*. 2010; 22:177–80. [PubMed: 20022734]
- [133]. Semenza GL. HIF-1: upstream and downstream of cancer metabolism. *Curr Opin Genet Dev*. 2010; 20:51–6. [PubMed: 19942427]
- [134]. Mellor HR, Harris AL. The role of the hypoxia-inducible BH3-only proteins BNIP3 and BNIP3L in cancer. *Cancer Metastasis Rev*. 2007; 26:553–66. [PubMed: 17805942]

- [135]. Shimizu S, Konishi A, Nishida Y, Mizuta T, Nishina H, Yamamoto A, et al. Involvement of JNK in the regulation of autophagic cell death. *Oncogene*. 2010; 29:2070–82. [PubMed: 20101227]
- [136]. Kabeya Y, Mizushima N, Ueno T, Yamamoto A, Kirisako T, Noda T, Kominami E, et al. LC3, a mammalian homologue of yeast Apg8p, is localized in autophagosome membranes after processing. *EMBO J*. 2000; 19:5720–8. [PubMed: 11060023] *EMBO J*. 2003; 22:4577. Erratum in:
- [137]. Tanida I, Ueno T, Kominami E. LC3 conjugation system in mammalian autophagy. *Int J Biochem Cell Biol*. 2004; 36:2503–18. [PubMed: 15325588]
- [138]. Tanida I, Ueno T, Kominami E. LC3 and Autophagy. *Methods Mol Biol*. 2008; 445:77–88. [PubMed: 18425443]
- [139]. Cherra SJ 3rd, Kulich SM, Uechi G, Balasubramani M, Mountzouris J, Day BW, et al. Regulation of the autophagy protein LC3 by phosphorylation. *J Cell Biol*. 2010; 190:533–9. [PubMed: 20713600]
- [140]. Germain M, Nguyen AP, Le Grand JN, Arbour N, Vanderluit JL, Park DS, et al. MCL-1 is a stress sensor that regulates autophagy in a developmentally regulated manner. *EMBO J*. 2011; 30:395–407. [PubMed: 21139567]
- [141]. Germain M, Slack RS. MCL-1 regulates the balance between autophagy and apoptosis. *Autophagy*. 2011; 7:549–51. [PubMed: 21412051]
- [142]. Kim J, Kundu M, Viollet B, Guan KL. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol*. 2011; 13:132–41. [PubMed: 21258367]
- [143]. Jung CH, Ro SH, Cao J, Otto NM, Kim DH. mTOR regulation of autophagy. *FEBS Lett*. 2010; 584:1287–95. [PubMed: 20083114]
- [144]. Ganley IG, Lam du H, Wang J, Ding X, Chen S, Jiang X. ULK1, ATG13, FIP 200 complex mediates mTOR signaling and is essential for autophagy. *J Biol Chem*. 2009; 284:12297–305. [PubMed: 19258318]
- [145]. Pattingre S, Espert L, Biard-Piechaczyk M, Codogno P. Regulation of macroautophagy by mTOR and Beclin 1 complexes. *Biochimie*. 2008; 90:313–23. [PubMed: 17928127]
- [146]. Feng Z, Zhang H, Levine AJ, Jin S. The coordinate regulation of the p53 and mTOR pathways in cells. *Proc Natl Acad Sci U S A*. 2005; 102:8204–9. [PubMed: 15928081]
- [147]. Muñoz-Gámez JA, Rodríguez-Vargas JM, Quiles Pérez R, Aguilar-Quesada R, Martín-Oliva D, de Murcia G, et al. PARP-1 is involved in autophagy induced by DNA damage. *Autophagy*. 2009; 5:61–74. [PubMed: 19001878]
- [148]. Huang Q, Shen HM. To die or to live: the dual role of poly(ADP-ribose) polymerase-1 in autophagy and necrosis under oxidative stress and DNA damage. *Autophagy*. 2009; 5:273–6. [PubMed: 19139632] *Autophagy*. 2010; 6:1232. Erratum in: [PubMed: 21079416]
- [149]. Petiot A, Ogier-Denis E, Blommaert EF, Meijer AJ, Codogno P. Distinct classes of phosphatidylinositol 3'-kinases are involved in signaling pathways that control macroautophagy in HT-29 cells. *J Biol Chem*. 2000; 275:992–8. [PubMed: 10625637] *J Biol Chem*. 2000; 275:12360. Erratum in:
- [150]. Backer JM. The regulation and function of Class III PI3Ks: novel roles for Vps34. *Biochem J*. 2008; 410:1–17. [PubMed: 18215151]
- [151]. Wu YT, Tan HL, Huang Q, Ong CN, Shen HM. Activation of the PI3K-Akt-mTOR signaling pathway promotes necrotic cell death via suppression of autophagy. *Autophagy*. 2009; 5:824–34. [PubMed: 19556857]
- [152]. Casas-Terradellas E, Tato I, Bartrons R, Ventura F, Rosa JL. ERK and p38 pathways regulate amino acid signalling. *Biochim Biophys Acta*. 2008; 1783:2241–54. [PubMed: 18809440]
- [153]. McClung JM, Judge AR, Powers SK, Yan Z. p38 MAPK links oxidative stress to autophagy-related gene expression in cachectic muscle wasting. *Am J Physiol Cell Physiol*. 2010; 298:C542–9. [PubMed: 19955483]
- [154]. Jin S. p53, Autophagy and tumor suppression. *Autophagy*. 2005; 1:171–3. [PubMed: 16874039]
- [155]. Levine B, Abrams J. p53: The Janus of autophagy? *Nat Cell Biol*. 2008; 10:637–9. [PubMed: 18521069]

- [156]. Maiuri MC, Galluzzi L, Morselli E, Kepp O, Malik SA, Kroemer G. Autophagy regulation by p53. *Curr Opin Cell Biol.* 2010; 22:181–5. [PubMed: 20044243]
- [157]. Parkhitko A, Myachina F, Morrison TA, Hindi KM, Auricchio N, Karbowiczek M, et al. Tumorigenesis in tuberous sclerosis complex is autophagy and p62/sequestosome 1 (SQSTM1)-dependent. *Proc Natl Acad Sci U S A.* 2011; 108:12455–60. [PubMed: 21746920]
- [158]. Ryan KM. p53 and autophagy in cancer: guardian of the genome meets guardian of the proteome. *Eur J Cancer.* 2011; 47:44–50. [PubMed: 21112207]
- [159]. Bjorkoy G, Lamark T, Johansen T. P62/SQSTM1: a missing link between protein aggregates and the autophagy machinery. *Autophagy.* 2006; 2:138–9. [PubMed: 16874037]
- [160]. Ueno T, Sato W, Horie Y, Komatsu M, Tanida I, Yoshida M, et al. Loss of Pten, a tumor suppressor, causes the strong inhibition of autophagy without affecting LC3 lipidation. *Autophagy.* 2008; 4:692–700. [PubMed: 18424911]
- [161]. Furuta S, Hidaka E, Ogata A, Yokota S, Kamata T. Ras is involved in the negative control of autophagy through the class I PI3-kinase. *Oncogene.* 2004; 23:3898–904. [PubMed: 15064741]
- [162]. Zhou X, Ikenoue T, Chen X, Li L, Inoki K, Guan KL. Rheb controls misfolded protein metabolism by inhibiting aggresome formation and autophagy. *Proc Natl Acad Sci U S A.* 2009; 106:8923–8. [PubMed: 19458266]
- [163]. Salminen A, Kaarniranta K. SIRT1: regulation of longevity via autophagy. *Cell Signal.* 2009; 21:1356–60. [PubMed: 19249351]
- [164]. Hariharan N, Maejima Y, Nakae J, Paik J, Depinho RA, Sadoshima J. Deacetylation of FoxO by Sirt1 Plays an Essential Role in Mediating Starvation-Induced Autophagy in Cardiac Myocytes. *Circ Res.* 2010; 107:1470–82. [PubMed: 20947830]
- [165]. Yan J, Kuroyanagi H, Kuroiwa A, Matsuda Y, Tokumitsu H, Tomoda T, et al. Identification of mouse ULK1, a novel protein kinase structurally related to *C. elegans* UNC-51. *Biochem Biophys Res Commun.* 1998; 246:222–7. [PubMed: 9600096]
- [166]. Jung CH, Jun CB, Ro SH, Kim YM, Otto NM, Cao J, et al. ULK-Atg13-FIP200 complexes mediate mTOR signalling to the autophagy machinery. *Mol Biol Cell.* 2009; 20:1992–2003. o. [PubMed: 19225151]

Fig. 1.
Features of autophagy execution.

Fig. 2.

Signaling pathways that regulate autophagy. Abbreviations not listed here are included in the text.

Leu: Leucine

mSin1: mammalian stress-activated protein kinase [SAPK]-interacting protein

RAF, MEK and ERK (mitogen-activated protein kinase): serine/threonine selective protein kinases in the MAPK/ERK pathway.

RSK: Ribosomal S6 kinase (activated by MAPK/ERK pathway)

SLC7A5: Large neutral amino acids transporter small subunit 1; L-type amino acid transporter 1 (LAT1)

Table 1**Mediators of Autophagy**

Mediator	Alternate names	Function	References
Akt	Protein kinase B	Key downstream effector of the PI3K signaling pathway; regulates cell survival and proliferation. Akt inhibition leads to activation of autophagy. PI3K-Akt-mTOR pathway suppresses autophagy.	101-102
AMBRA 1	Activating molecule in Beclin 1 regulated autophagy	Regulates autophagy and development of the nervous system. AMBRA 1 binds to Beclin 1 and favors Beclin 1/VPS34 interaction.	103-105
AMPK	AMP-activated protein kinase	Senses reduced cellular energy level and regulates many aspects of cell function. AMPK inhibits mTOR-dependent signaling. It is required for autophagy in both yeast and mammalian cells. Activated AMPK triggers destruction of defective mitochondria (mitophagy), biogenesis of new mitochondria, and restrains cell growth and proliferation.	106-108
Bcl-2	B-Cell lymphoma 2 family of regulator proteins encoded by the Bcl-2 gene	Dual regulator of apoptosis and autophagy; binds and sequesters Beclin 1. Dissociation of Beclin 1 from Bcl-2 is required for autophagy induction.	69, 109-111
Beclin 1	Mammalian orthologue of the yeast Atg6 gene	Encodes a Bcl-2 interacting coiled-coil protein with autophagy and tumor suppressor function. Regulates kinase activity of VPS34.	68, 112-114
BNIP3	Bcl-2/adenovirus E1b 19-kda interacting protein 3	Competitively binds to Bcl-2 and releases Beclin 1, a known inducer of autophagy; specific activator of mitochondrial autophagy; inhibits mTOR activity and induces autophagy; crucial for hypoxia-induced autophagy	54, 115-119
DAPK	Death-associated protein kinase	Phosphorylates Beclin 1 causing dissociation from Bcl-2 and Bcl-X _L , which in turn induces autophagy; phosphorylates TSC2 leading to TSC1-TSC2 dissociation to stimulate mTOR in response to growth factor activation; plays a role in membrane blebbing.	72-73, 120
DRAM	Damage-regulated autophagy modulator	This lysosomal protein is critical for p53 induction of autophagy and cell death, has a tumor-suppressive function.	84-85, 121-122
ERK	Extracellular signal-regulated kinase	Depending on the cell type and stimulus, ERK activity will mediate different antiproliferative events, such as apoptosis, autophagy and senescence, in vitro and in vivo.	123-127
FOXO3	Forkhead Box O3	FoxO transcription factors are critical mediators of oxidative stress resistance in multiple cell types. FoxO transcription factors promote autophagy in skeletal muscle and have additional roles in regulation of cell size, proliferation, and metabolism. Decreased IGF-1-PI3K-Akt signaling activates autophagy not only through mTOR, but also by a transcription-dependent mechanism involving FoxO3.	128-131
Hif 1	Hypoxia inducible factor 1	Induces a cell survival response engaging autophagy mediated by BNIP3/BNIP3L	50, 132-134
JNK	c-Jun N-terminal kinase	JNK activation modulates autophagy through two distinct mechanisms: 1. Promotes Bcl-2 phosphorylation resulting in the dissociation of the Beclin 1-Bcl-2 complex. 2. Upregulates DRAM, a p53 target gene.	74-75, 114, 121, 135
LC3	Microtubule-associated protein Light Chain 3; Mammalian homolog of Atg8	Cytosolic form of LC3 (LC3-I) is conjugated to phosphatidylethanolamine to form LC3-II, which is recruited to autophagosomal membranes.	136-139
MCL-1	Myeloid leukemia cell differentiation protein	Has a key role in controlling both autophagy and apoptosis processes in a developmentally regulated manner.	140-141
mTOR	Mammalian target of rapamycin	Key negative regulator of autophagy; stress conditions and starvation inactivate mTOR.	142-146
PARP1	Poly (ADP-ribose) polymerase-1	Plays a dual role in the modulation of autophagy, apoptosis, and necrosis under oxidative stress	61, 147-148

Mediator	Alternate names	Function	References
PI3K	Phosphatidylinositol 3-kinase	Class III PI3K regulates autophagy as well as protein synthesis through the mTOR pathway	149-151
p38	p38 mitogen-activated protein kinase	Plays a role in the control of autophagy at the sequestration and maturation step; regulates amino acid signaling, along with ERK 1/2 kinase	127, 152-153
p53	Protein 53	Modulates autophagy in a dual fashion, depending on the subcellular localization in the nucleus versus the cytoplasm. Activation of p53 in the nucleus inhibits mTOR activity. The mechanisms by which p53 regulates mTOR activity in the cytoplasm involves AMP kinase activation and requires the tuberous sclerosis (TSC)1/TSC2 complex, both of which respond to energy deprivation in cells.	84, 146, 154-158
p62	Sequestosome	Links polyubiquitinated protein aggregates to the autophagy machinery; required for formation and activation of the caspase-8 complex	82-83, 159
PTEN	Phosphatase and tensin homolog	Positively regulates autophagy by inhibiting the phosphatidylinositol 3-kinase/protein kinase B pathway; antagonizes insulin dependent cell signaling; acts as a tumor suppressor gene through the action of its phosphatase protein product	43, 160
Rag1 and 2	Ras-related GTPase	Small GTPases that regulate mTORC1 activity in response to amino acids, likely independently of TSC1/2 and Rheb.	100
Raptor	Regulatory associated protein of mTOR	mTOR partner in mTORC1. Binds substrates, including S6K1 and 4EBP1	26-27
Ras	Rat sarcoma protein	Small GTPase that acts as a negative regulator for nutrient-deprivation induced autophagy through the Class I PI3K signaling pathway; activation mutations lead to malignant cell transformation	37-38, 161
Rheb	Ras homologue enriched in brain	Small GTPase that regulates mTOR activity; regulates aggresome formation by inhibiting dynein dependent transportation of misfolded proteins	162
SIRT1	Sirtuin-1; NAD-dependent deacetylase sirtuin	SIRT1 can form a molecular complex with several essential components of the autophagy machinery, including autophagy genes Atg5, Atg7, and Atg8. SIRT1 can directly deacetylate these components, in an NAD-dependent fashion. Regulates longevity via autophagy.	46, 163-164
TSC1/2 complex	Tuberous sclerosis complex Hamartin (TSC1) Tuberin (TSC2)	TSC1/2-Rheb-mTOR pathways play a central role in modulating cellular protein synthesis and autophagy.	146, 157
ULK1 & 2	Unc-51-Like Kinase 1 & 2 Mammalian Atg1 homologues	Formation of ULK complex (ULKs-Atg13-FIP200) plays an essential role in the initiation of autophagy	14, 165-166