Design, synthesis and expression of a human interleukin-2 gene incorporating the codon usage bias found in highly expressed *Escherichia coli* genes

D.P.Williams¹, D.Regier, D.Akiyoshi⁺, F.Genbauffe⁺ and J.R.Murphy*

Evans Department of Clinical Research and Department of Medicine, The University Hospital and

¹Department of Microbiology, Boston University School of Medicine, Boston, MA 02118, USA

Received September 15, 1988; Accepted October 18, 1988

Accession no. X07256

ABSTRACT

A synthetic gene encoding human interleukin-2 (IL-2) was designed such that the codon usage bias resembled that found in highly expressed *Escherichia coli* genes. The percentage of preferred codons was increased from 43% in the native cDNA sequence to 85% in the synthetic sequence. The cDNA and synthetic IL-2 genes were placed under the control of the *trc* promoter and expressed in *E. coli* JM101. While Northern blot analysis of IL-2 mRNA from each genetic construct demonstrated equivalent message half-lives, immunoblot and bio-activity analyses showed the synthetic gene to direct the synthesis of up to 16 times more IL-2 than the native cDNA sequence.

INTRODUCTION

DNA sequence analysis of genes encoding either high abundance or low abundance proteins in *E. coli* has revealed a non-random usage of degenerate codons for most of the 20 amino acids (1-5). Highly expressed genes show the greatest degree of codon bias, and favored codons correspond to the most abundant tRNAs in the cell. It has been postulated that the preferred nucleotide triplets reflect the optimization of codon:anticodon interaction energy (3), the influence of tRNA availability on the rate of peptide chain elongation (4), or natural selective pressure for translational efficiency (5). Codon usage bias has also been observed in yeast, where clusters of rarely used codons at exon (protein domain) boundaries of genes are thought to comprise important ribosome pause sites (6). Despite the strong correlation between codon usage patterns and the level of expression of endogenous genes in *E. coli*, little has been published regarding the effects of altering the DNA sequence of heterologous genes such that their codon bias is consistent with that found in highly expressed *E. coli* genes.

In the present study, we examine the effect of codon usage on the level of expression of human interleukin-2 (IL-2) in *E. coli*. IL-2, a hormone-like polypeptide growth factor produced by T lymphocytes in response to antigen or mitogen, has been purified from several sources including peripheral blood

lymphocytes, splenic lymphocytes, and T-cell leukemia cultures (7-9). In addition, cloned IL-2 cDNAs have been expressed in E. coli and shown to produce biologically active lymphokine (10,11). However, codon usage in the human IL-2 cDNA sequence does not reflect the codon bias found in highly expressed E. coli genes (3). Fewer than 45% of the codons found in the cDNA sequence for mature IL-2 (10,12) are preferred E. coli codons. Thus, it was of interest to compare the level of recombinant protein produced from the native human IL-2 cDNA sequence with that produced from a synthetic version of the IL-2 gene having preferred codon usage.

Here we describe the design and construction of a synthetic IL-2 gene having 85% of its codons corresponding to those preferred for high level expression in E. coli. We have placed the synthetic and cDNA versions of the IL-2 gene under the control of the trc promoter and find that E.coli JM101 carrying the synthetic IL-2 gene construct produce more lymphokine than JM101 carrying the native cDNA sequence. Moreover, we show by Northern blot analysis that the increased level of IL-2 produced from the synthetic IL-2 construct does not result from an increase in the stability of the IL-2 specific messenger RNA.

MATERIALS AND METHODS

Oligonucleotide Synthesis and Purification

Oligonucleotides were synthesized using methoxy phosphoramidite chemistry on an Applied Biosystems Model 380A DNA synthesizer and purified by standard gel electrophoresis methods (13,14). Electroeluted material was precipitated with 5 volumes of ethanol or desalted and concentrated by passage over NENsorb columns (NEN/DuPont) as directed by the manufacturer. Purified oligonucleotides were resuspended in TE buffer (10 mM Tris pH 8.0, 1 mM EDTA). To anneal complementary strands, equimolar amounts were mixed in the presence of 100 mM NaCl, heated for 10 minutes at 90°C, and allowed to cool slowly to room temperature.

Bacterial Strains and DNA Manipulations

E. coli strain DH5 α (Bethesda Research Laboratories) was used for routine plasmid screening and plasmid DNA sequencing, and strain JM101 (New England Biolabs) was used for M13 sequencing, screening of IL-2 expression constructs, and expression of IL-2. Recombinant E. coli were grown in LB medium containing 100 μ g/ml ampicillin or in 2X YT medium (15). JM101 containing IL-2 expression plasmids grew poorly at 37°C and therefore were maintained at 30°C subsequent to the transformation heat shock step.

Sources of enzymes and plasmid DNA were as follows: restriction endonucleases, T4 DNA polymerase, T4 polynucleotide kinase, and M13 DNA sequencing pack from New England Biolabs; restriction endonucleases, calf intestine alkaline phosphatase, M13mp19, and pUC19 from Boehringer Mannheim Biochemicals; T4 DNA ligase from Bethesda Research Laboratories; GemSeq plasmid sequencing kit from Promega; and pKK233-2 from Pharmacia. All enzymes were used as recommended by the manufacturers.

Routine electrophoresis was carried out in 1% agarose/TBE gels and restriction fragments were purified by electroelution as described by Maniatis et al. (14). Conditions for ligation of DNA, *E. coli* transformation, and large-scale plasmid DNA purification have been described (14). "Mini-lysate" plasmid DNA and small-scale M13 phage RF DNA was prepared by the method of Birnboim and Doly (16), and M13 single-stranded template DNA was prepared as described by Williams et al. (17). DNA sequence analysis of M13 and pUC19 DNA was by the Sanger dideoxy chain termination method (18) using universal and other synthetic primers.

Expression of IL-2 in *E. coli*

E. coli JM101 harboring plasmid pNIL2-22, pNIL2-22', pSIL2-32, pSIL2-32', or pKK233-2 was grown at 30°C to an OD₅₉₀ of 0.2 and IL-2 expression was induced by the addition of isopropyl-β-D-thiogalacto-pyranoside (IPTG, Sigma Chemical Co.) to a final concentration of 2 mM. Following a 3 hour induction period, the optical density of each culture was determined and approximately 6×10^9 cells were harvested by centrifugation (19). *E. coli* extracts were prepared as follows: cultures were chilled on ice and all subsequent operations carried out at 4°C; cells were pelleted, drained, resuspended in 2 ml lysis buffer (7 M guanidine hydrochloride, 100 mM Tris-HCl, pH 8.0) and sonicated for 3 minutes at power level 5 and 50% duty cycle (Branson sonifier). Extracts were transferred to 3500 MWCO tubing (SpectraPor) and dialysed against phosphate-buffered saline overnight. Dialysed extracts were clarified by brief centrifugation and supernatants assayed for protein concentration using Pierce Protein Reagent (Pierce Chemical Co.).

SDS-polyacrylamide gel and Immunoblot Analysis

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis was performed according to the method of Laemmli (20) using 15% polyacrylamide gels and a Mini-Protean II gel apparatus (BioRad). Recombinant human IL-2 (rIL-2; Biogen, Inc.) was used as a reference and low molecular weight protein standards were from Pharmacia. Proteins were either silver-stained (BioRad silver stain reagent kit) or transferred to nitrocellulose paper (BA85, 0.45μ

Schleicher & Schuell) for immunoblot analysis as described by Towbin et al. (21). For immuno-dot blots, the protein concentration of E. coli extracts was adjusted to 400 µg/ml with phosphate-buffered saline and applied to a Bio-Dot microfiltration apparatus (BioRad) as recommended by the manufacturer. Monoclonal anti-recombinant human IL-2 (MAC-3; Celltech, Ltd.) and horseradish peroxidase-conjugated goat anti-mouse IgG (Cappel/Organon Teknika) were used to process all immunoblots. Non-specific cross-reactive antibodies were removed from the goat anti-mouse preparation by precipitation with E. coli lysate (BioRad). Incubation buffer contained 0.5 M NaCl and blots were developed in phosphate-buffered saline with 0.5 mg/ml 3',3'-diaminobenzidine tetra-hydrochloride (Sigma Chemical Co.).

Biological Assay for IL-2

Biologically active IL-2 was detected as the stimulation of the IL-2-dependent murine cytotoxic T-cell line, CTLL-2. Cell proliferation was quantitated by incorporation of ³H-thymidine as described by Gillis et al. (22). CTLL-2 cells were maintained in RPMI 1640 medium (Gibco) containing 10% fetal bovine serum (Hyclone), 2 mM glutamine (Gibco), and 100 µg/ml each of penicillin and streptomycin (Gibco). Extracts of recombinant E. coli were diluted with complete medium containing 10% serum and assayed for IL-2 activity.

RNA Isolation and Northern Blot Analysis

E. coli JM101(pKK233-2), JM101(pNIL2-22), and JM101(pSIL2-32) were grown at 30°C to an OD₅₉₀ of 0.4 prior to induction with 2.0 mM IPTG (final concentration). One hour after the addition of IPTG, transcription was blocked and total cellular RNA was prepared from each culture as follows: 15 ml aliquots of the induced culture were transferred to each of 3 pre-warmed flasks. At 2-minute intervals, rifampicin (Sigma) was added at 0.2 mg/ml to each of the 3 flasks. Immediately following the addition of rifampicin to the "0 minute" flask, all 3 flasks were plunged into an ice/water bath. Bacteria were harvested by centrifugation at 12,000 x g for 5 min at 4°C. Cell pellets were frozen in a dry ice/ethanol bath and thawed with vigorous mixing in the presence of 0.5 ml TE (10 mM Tris pH 7.5, 1 mM EDTA) containing 500 µg of proteinase K (Boehringer Mannheim Biochemicals) per ml (23). RNA was extracted according to an adaptation of the method of Chomczynski and Sacchi (24). Briefly, 5 ml of denaturing solution (4 M guanidinium thiocyanate, 25 mM sodium citrate, pH 7.0, 0.5% sarcosyl, 0.1 M 2-mercaptoethanol) was added to the thawed cell suspension and mixed vigorously. Sequentially, 0.5 ml of

2 M sodium acetate, pH 4.0, 5.0 ml of water-saturated phenol, and 1.0 ml chloroform-isoamyl alcohol mixture (49:1) were added to the lysate, with rapid and thorough mixing after the addition of each reagent. The final suspension was chilled on ice for 15 min, and the phases were separated by centrifugation at 10,000 x g for 20 min at 4°C. RNA was precipitated from the aqueous phase by addition of 5.0 ml of isopropanol. After 1 hr at -20°C, RNA was collected by centrifugation (10,000 x g, 20 min, 4°C). The resulting RNA pellets were drained, dissolved in 0.3 ml of denaturing solution, and transferred to 1.5 ml Eppendorf tubes. RNA was re-precipitated with 1 volume of isopropanol at -20°C for 1 hr. Following centrifugation, the RNA pellets were washed twice with 70% ethanol, dried, and resuspended in 100 µl of CE buffer (10 mM sodium citrate, pH 7.0, 1 mM EDTA). RNA was precipitated a third time by addition of 0.15 volume of 2 M sodium acetate, pH 5.2, and 2.5 volumes of 100% ethanol. The final RNA pellet was dissolved in CE containing 0.5% SDS (final concentration).

RNA samples were electrophoresed on 1.5% agarose-formaldehyde gels (14) and blotted onto nitrocellulose as recommended by Schleicher and Schuell (25). RNA molecular size ladder was from Bethesda Research Laboratories. Blots were probed with a 14-base oligonucleotide complementary to both IL-2 encoding genes as indicated in Figure 1. The probe was labeled with [³²P-ATP] using T₄ polynucleotide kinase and purified over NENsorb cartridges. Hybridizations were performed at 31°C under conditions recommended for oligonucleotide probes (25). Autoradiograms were analyzed by densitometry and mRNA half-lives were determined by least-squares analysis of semilogarithmic plots of band intensity as a function of time (26).

RESULTS

Design and Assembly of a Synthetic Human IL-2 Gene

A synthetic gene coding for the 133 amino acid mature form of IL-2 was designed in two steps. First, amino acid codons found in the human IL-2 cDNA (12) were compared to amino acid codons found in highly expressed E. coli genes (3,4). Only seven of 22 Leu codons were of the type found in highly expressed E. coli genes. Similarly, only one of six Phe codons was found to be of the preferred sequence, TTC. Overall, only 43% of the codons found in the native IL-2 cDNA were those favored for high level expression in E. coli. Therefore, a modified coding sequence was designed wherein nucleotide substitutions generated preferred E. coli codons while maintaining human IL-2

```

      1                               10
IL-2 amino acid:  Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln
cDNA sequence:   gca cct act tca agt tct aca aag aaa aca cag cta caa
codon modifications:      t c c c c g g
synthetic DNA sequence: GCA CCT ACT TCT AGC TCT ACC AAG AAA ACC CAG CTG CAG

      20                               30
Leu Glu His Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys Asn
ctg gag cat tta ctg ctg gat tta cag atg att ttg aat gga att aat aat tac aag aat
      c c g g c c c c t c c
CTCGAGCAC CTG CTG CTG GAT TTG CAG ATG ATC CTG AAC GGT ATC AAC AAT TAC AAG AAC
XhoI

      40                               50
Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro Lys Lys Ala Thr Glu Leu
ccc aaa ctc acc agg atg ctc aca ttt aag ttt tac atg ccc aag aag gcc aca gaa ctg
      g g c t g c c c g c c
CCG AAA CTG ACGGT ATG CTG ACC TTC AAG TTC TAC ATG CCG AAG AAG GCC ACC GAA CTG
MluI

      60                               70
Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys Pro Leu Glu Glu Val Leu Asn Leu Ala
aaa cat ctt cag tgt cta gaa gaa gaa ctc aaa cct ctg gag gaa gtg cta aat tta gct
      c g g g g g g t g c c g
AAA CAC CTG CAG TG TCT AGAA GAA GAA CTG AAA CCG CTG GAG GAA GTT CTG AAC CTG GCT
XbaI

      80                               90
Gln Ser Lys Asn Phe His Leu Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val
caa agc aaa aac ttt cac tta aga ccc agg gac tta atc agc aat atc aac gta ata gtt
      g tct c c g c t g c t c g tct c c
CAG TCT AAA AAC TTC CAC CT GCGGCCGC GT GAC CTG ATC TCT AAC ATC AAC GTA ATC GTT
NotI

      100                              110
Leu Glu Leu Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala Thr
ctg gaa cta aag gga tct gaa aca aca ttc atg tgt gaa tat gct gat gag aca gca acc
      g t c c c c c c c
CTG GAA CTG AAGGCTCTGAA ACC ACC TTC ATG TGT GAA TAC GCT GAT GAG ACC GCA ACC
BamHI

      120                              130
Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile Ile Ser Thr Leu Thr STOP
att gta gaa ttt ctg aac aga tgg att acc ttt tgt caa agc atc atc tca aca cta act tga
      c c c t c c g tct t c g c
ATC GTA GAATTC CTG AAC CGT TGG ATC ACC TTC TGT CAG TCT ATC ATC TCT ACC CTG ACC TGA
EcoRI

```

Figure 1. Amino acid and nucleic acid sequence of the mature form of human IL-2 (10). "Codon modifications" indicates base changes required to incorporate *E. coli* preferred codons. Bases changed from the preferred sequence to generate unique restriction sites are underlined. Bold underline indicates a region of IL-2 DNA sequence common to both the cDNA and synthetic genes and complementary to a 14-base oligonucleotide probe used for Northern blot analysis.

amino acid sequence. For amino acids such as Asp and Cys where no clear codon preference has been demonstrated, the original cDNA nucleotide triplet was retained.

Figure 2. Construction of the synthetic IL-2 gene. The 5' to 3' assembly of the synthetic IL-2 gene was accomplished in 7 vectorial cloning steps. Synthetic fragment 1 was ligated to BamHI / SalI digested pUC19. Putative clones were selected on the basis of acquisition of a unique XhoI restriction site, and inserts were confirmed by DNA sequence analysis. DNA from a fragment 1-positive clone was digested with XhoI and SalI and ligated to synthetic fragment 2. Clones having acquired a unique MluI site were subjected to DNA sequence analysis, etc. Following this scheme, synthetic fragments 3 through 7 were added to the 3' end of fragment 2 in successive cloning/screening steps. The complete synthetic IL-2 gene was recloned into the BamHI and SalI sites of M13mp19 and sequenced using the universal primer and a synthetic primer corresponding to nucleotides 165 to 182 of synthetic IL-2.

The second step was the incorporation of unique restriction sites to divide the synthetic gene into smaller segments amenable to *in vitro* synthesis and cloning. The modified IL-2 sequence was found to contain a complete EcoRI site and partial target sites for XhoI, MluI, XbaI, NotI, and BanII. These partial sites were converted to complete target sites by single base changes which did not alter the amino acid sequence, but which did change preferred codons to non-preferred codons. Figure 1 compares the original cDNA sequence for IL-2 with that of the synthetic IL-2 gene.

The overall strategy used for the molecular cloning of the synthetic IL-2 gene is depicted in Figure 2. Each of the seven restriction fragments comprising the complete gene was synthesized as a pair of complementary oligonucleotides. As diagrammed in Figure 2, the 5'-end of each fragment consisted of a $\frac{1}{2}$ restriction site homologous to a unique restriction site in the preceding fragment. In addition to the IL-2 encoding region, a 6 bp spacer and a $\frac{1}{2}$ SalI site were added to the 3'-end of each segment to facilitate vectorial cloning. The spacer region was necessary for efficient cleavage of adjacent restriction sites. The 5'-end of the synthetic IL-2 gene was modified by the addition of a 24 bp linker. This sequence was analogous to a linker that had been added to the IL-2 cDNA clone (pDW6, Figure 3) and includes a BamHI site allowing for in-frame fusion to the heat-inducible

Figure 3. Construction of IL-2 expression plasmids. The IL-2 cDNA gene was excised from pDW6 by digestion with *Sph*I and *Stu*I. The synthetic IL-2 gene was released from pDW15 by digestion with *Sph*I only. *Sph*I 3' extensions were removed with T4 DNA polymerase and blunt-ended fragments were ligated to phosphorylated, self-complementary 12 bp *Nco*I linkers (5' CCATGGCCATGG 3'). Following digestion with *Nco*I, IL-2 encoding inserts were electroeluted from agarose gels, ligated to *Nco*I-digested alkaline phosphatase-treated pKK233-2, and used to transform *E. coli* JM101. Resulting plasmids, pNIL2-22 and pSIL2-32, carry the native cDNA and synthetic alleles, respectively. The vector portions of pNIL2-22 and pSIL2-32 were exchanged by digestion with *Nco*I, isolation of the respective fragments, and religation (as shown) to generate a second set of IL-2 expression plasmids designated pNIL2-22' and pSIL2-32'. In each case, expression is directed from the *trc* promoter (arrowheads) and the translation initiation codon (ATG) is adjacent to the codon for Ala¹ of mature IL-2. (*Xho*I, *Xba*I, *Not*I, *Ban*II, and *Eco*RI sites were omitted from the restriction map of the synthetic IL-2 gene to reduce the complexity of the diagram. See Figure 1).

coliphage λ P_R (27). A unique *Sph*I site was also included in the linker to facilitate the genetic fusion of IL-2 to a truncated form of diphtheria toxin as previously described (28). The complete synthetic IL-2 gene was ligated to

BamHI / SalI digested pUC19 and the resulting plasmid, pDW15, is shown in Figure 3.

Construction of IL-2 Expression Plasmids

IL-2 expression plasmids were constructed using vector pKK233-2. This vector contains the trc promoter, the lacZ ribosome binding site, an ATG initiation codon contained within a unique NcoI site, and strong rrnB transcriptional terminators (29). Figure 3 shows the cloning strategy used to place the native and synthetic IL-2 genes under the control of the trc promoter. IL-2 encoding fragments from plasmids pDW6 (cDNA) and pDW15 (synthetic gene) were fused to the trc promoter such that the translational initiation codon (ATG) within the NcoI site was adjacent to the codon for Ala¹ of mature IL-2. Resulting plasmids were designated pNIL2-22 and pSIL2-32, denoting the native and synthetic gene constructs, respectively. Nucleotide sequence analysis showed the trc promoter regions (through Thr³ of IL-2) of both constructs to be identical (data not shown). To ensure that any differences in IL-2 expression between JM101(pNIL2-22) and JM101(pSIL2-32) were not a result of incidental vector mutations, pNIL2-22 and pSIL2-32 were digested with NcoI, and each insert was ligated to the reciprocal vector fragment to generate pNIL2-22' and pSIL2-32' (Figure 3).

Expression and Biological Activity

Extracts of IPTG-induced cultures of JM101(pNIL2-22, pNIL2-22', pSIL2-32, and pSIL2-32') were prepared as described in METHODS and found to contain an anti-IL-2 reactive protein that co-migrated with purified rIL-2 (Figure 4B). Furthermore, cells carrying the synthetic IL-2 gene appeared to produce more lymphokine than those carrying the native IL-2 cDNA as evidenced in silver-stained gels and immunoblots (Figures 4A & 4B). In order to quantitate the amount of biologically active lymphokine produced by JM101(pNIL2-22) and JM101(pSIL2-32), cell extracts were tested in a standard CTLL-2 growth assay (Figure 5). Using probit analysis (22), the concentration of biologically active IL-2 in JM101(pSIL2-32) extracts was determined to be 8 times greater than that in JM101(pNIL2-22) extracts.

Dot-Blot Quantitation

Previous studies have shown that the use of guanidine hydrochloride to solubilize and extract recombinant human IL-2 from E. coli must be followed by a renaturation period in order to recover bioactivity. Bailon et al.(30) reported recovery of 43% of total IL-2 activity after 24 hours of renaturation. However, recovery approached 100% when refolding was allowed to proceed for up to one week (30). These results suggest that the IL-2

Figure 4. SDS-polyacrylamide gel analysis and immunoblot detection of human IL-2. Whole cell extracts of *E. coli* JM101 carrying plasmids pKK233-2, pNIL2-22, pNIL2-22', pSIL2-32, and pSIL2-32' were prepared as described in Methods. 10 μ g total protein was loaded per lane. A) Silver stained SDS-polyacrylamide gel. LMWM, low molecular weight protein standards; rIL-2, 1 μ g pure rIL-2. Arrow indicates rIL-2. B) Immunoblot of proteins shown in Panel A probed with monoclonal antibody to IL-2. rIL-2, 150 ng pure rIL-2.

bioactivity assay may actually underestimate the levels of IL-2 present in guanidine extracts of recombinant *E. coli*. Therefore, extracts of JM101(pNIL2-22) and JM101(pSIL2-32) were assayed by immuno-dot blot analysis using monoclonal antibody to IL-2. As shown in Figure 6, extracts from JM101(pSIL2-32) cultures contained up to 16 times more immunoreactive IL-2 than did extracts of JM101(pNIL2-22). This result, although in close agreement with data from the bioactivity assay (above), indicates that a portion of the IL-2 present in the extracts may be biologically inactive and that longer renaturation periods may be required to recover full activity.

Messenger RNA Stability

From the results of the dot-blot quantitation studies it was clear that

Figure 5. IL-2 dependent [^3H]-thymidine incorporation by CTLL-2 cells. Cells were washed and seeded at 10^4 cells / well. Whole cell extracts of *E. coli* JM101 carrying plasmids pNIL2-22 ($\square$) and pSIL2-32 ($\circ$) were adjusted to a starting concentration of 80 $\mu\text{g/ml}$ total protein. Each point is the average of triplicate analysis. (----) represents background [^3H]-TdR incorporation of medium alone or JM101(pKK233-2) controls.

the codon changes incorporated into the synthetic IL-2 gene had, in some way, increased its efficiency of expression. Since altering the nucleotide sequence of the IL-2 gene may have altered the potential secondary structure and/or stability of the mRNA, we determined the *in vivo* decay rates of both the cDNA and synthetic gene transcripts. Total RNA from IPTG induced cultures of JM101(pNIL2-22) and JM101(pSIL2-32) was prepared at different time points after the addition of the transcription initiation inhibitor, rifampicin (31). Following electrophoresis and transfer of RNA to nitrocellulose sheets, blots were hybridized with an IL-2 specific DNA probe. The decay of intact IL-2 specific mRNA was monitored by laser densitometry of autoradiograms as described in Methods. As shown in Figure 7, the synthetic IL-2 gene transcript and the IL-2 cDNA transcript decayed at the same rate with an apparent half-life of approximately 1 minute.

DISCUSSION

In an effort to examine the effect of altering the DNA sequence of a eukaryotic gene such that its codon usage bias more closely resembles that

Figure 6. Dot-blot quantitation of IL-2 in crude extracts of recombinant *E. coli*. Serial 4-fold dilutions of *E. coli* extracts, and serial 2-fold dilutions of rIL-2 were applied from left to right. Initial protein concentrations were 400 µg/ml for *E. coli* extracts, and 4 µg/ml for rIL-2. All sample volumes were adjusted to 100 µl. Duplicate rows were as follows: rows 1 & 2, pKK233-2; rows 3 & 4, pNIL2-22; rows 5 & 6, pNIL2-22'; rows 7 & 8, pSIL2-32; rows 9 & 10, pSIL2-32'; and rows 11 & 12, pure rIL-2. The limit of detection for rIL-2 was 6.25 ng/well.

found in highly expressed *E. coli* genes, we have assembled a synthetic human IL-2 gene in which the proportion of *E. coli* preferred codons was increased from 43% to 85%. By direct comparison, we have shown that the synthetic IL-2 gene directs the production of 8 to 16 times more IL-2 than does the native cDNA sequence when expressed from the *trc* promoter in *E. coli* JM101. Furthermore, in order to rule out potential vector differences, IL-2-encoding *Nco*I fragments of the native cDNA and synthetic gene expression plasmids were exchanged, and no alteration in the relative amount of lymphokine produced by either gene was found. Since crude extracts were used for all assays of rIL-2, we have avoided bias that may have been introduced during purification steps. In addition, Northern blot analysis of IL-2 specific mRNA following rifampicin treatment demonstrated equivalent decay rates for the cDNA and synthetic IL-2 messages. Moreover, computer-assisted analysis of the two gene sequences showed no significant difference in predicted mRNA secondary

Figure 7. Northern blot analysis of the decay of IL-2 mRNA. (UPPER) Samples (20 μ g) of total cellular RNA isolated from JM101(pNIL2-22), lanes 3-5, and from JM101(pSIL2-32), lanes 6-8, at 0 min (lanes 3,6), 2 min (lanes 4,7), and 4 min (lanes 5,8) after rifampicin treatment were fractionated on 1.5% agarose-formaldehyde gels, blotted onto nitrocellulose and probed with an end-labeled oligonucleotide complementary to both forms of the IL-2 gene. A sample of purified pNIL2-22 plasmid DNA (lane 1) was included as a positive control, and 20 μ g of total cellular RNA isolated from JM101(pKK233-2) provided the negative control (lane 2). Molecular size standards are given in kilobases. (LOWER) Semilogarithmic plot of the decay of the NIL-2 (O) and SIL-2 (●) messages as determined by densitometric analysis of autoradiograms.

structures (PCGene; IntelliGenetics, Inc.). Thus, the increased level of rIL-2 found in *E. coli* cultures harboring the synthetic IL-2 gene is most likely a direct result of codon usage bias.

Robinson *et al.* (32) inserted synthetic oligonucleotides containing either four preferred ARG codons or four rarely used ARG codons into the structural gene for chloramphenicol acetyltransferase (CAT). The insertion of four rare codons into an otherwise efficiently translated sequence was shown to reduce CAT synthesis by as much as 1/3, but only at very high rates of

transcription. George et al. (33) found that changing 13 of the first 23 codons of mature bovine growth hormone (bGH) to reflect the codon bias of highly expressed E. coli genes increased hormone accumulation by 10- to 15-fold. In this case, however, analysis of mRNA secondary structure predicted configurations within the translational initiation region of the codon-modified sequence which differed substantially from those of the native gene sequence. Since mRNA configuration greatly influences the efficiency of translation (34) it was difficult to ascertain what role, if any, codon modification played in increasing bGH synthesis. More recently, Zurawski et al. (35) synthesized a gene for human interleukin-1 α (IL-1 α) in which 95% of the codons were preferred versus only 47% in the native sequence. These investigators, however, did not use parallel expression constructs to compare protein yields from their synthetic gene with those from the native gene, and therefore the high level of expression of IL-1 α from the synthetic gene could not be directly attributed to the increased proportion of preferred codons.

While we are not certain that all codon changes incorporated into the synthetic IL-2 gene are significant, and we have not attempted to further optimize IL-2 production, it is clear from results presented here that codon usage bias is an important consideration for maximizing human IL-2 gene expression in E. coli.

ACKNOWLEDGMENTS

We are grateful to Drs. Craig Morita, David Beller, Barbara Pugh, and Raymond Donnelly for their assistance with the CTLL-2 assay. We also thank Karen Parker for helpful discussions. This work was supported by a grant from Seragen Inc., Hopkinton, MA, and by Public Health Service grants CA-41746 from the National Cancer Institute and AI-21628 from the National Institute of Allergy and Infectious Diseases.

*To whom correspondence should be addressed

+Present address: Seragen Inc., Hopkinton, MA 01748, USA

REFERENCES

1. Ikemura, T. (1981) J. Mol. Biol. 146, 1-21.
2. Ikemura, T. (1981) J. Mol. Biol. 151, 389-409.
3. Grosjean, H. and Fiers, W. (1982) Gene 18, 199-209.
4. deBoer, H.A. and Kastelein, R.A. (1986) in *Maximizing Gene Expression*, Butterworths, Boston.
5. Sharp, P.M. and Li, W.-H. (1987) Nucleic Acids Res. 15, 1281-1295.
6. Swaffield, J.C., Purvis, I.J., and Brown, A.J.P. (1987) Protein Engineering 1, 228.
7. Smith, K.A. (1980) Immun. Rev. 51, 337-357.

8. Robb, R.J., Kutny, R.M., Panico, M., Morris, H.R., and Chowdry, V. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6486-6490.
9. Hansen, J.A., Martin, P.J., and Nowinski, R.C. (1980) *Immunogenetics* 10, 247-252.
10. Devos, R., Plaetinck, G., Cheroutre, H., Simons, G., Degrave, W., Tavernier, J., Remaut, E., and Fiers, W. (1983) *Nucleic Acids Res.* 11, 4307-4322.
11. Rosenberg, S.A., Grimm, E.A., McGrogan, M., Doyle, M., Kawasaki, E., Kothe, K., and Mark, D.F. (1984) *Science* 233, 1412-1415.
12. Taniguchi, T., Matsui, H., Fujita, T., Takaoka, C., Kashima, N., Yoshimoto, R., and Hamuro, J. (1983) *Nature* 302, 305-310.
13. Evaluation and Purification of Synthetic Oligonucleotides (1984) in *DNA Synthesizer User Bulletin*, issue no. 13, Applied Biosystems, Inc., Foster City, CA.
14. Maniatis, T., Fritsch, E.F., and Sambrook, J. (1982) *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor Laboratory, New York.
15. Miller, J. (1972) in *Experiments in Molecular Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
16. Birnboim, H.C. and Doly, J. (1979) *Nucleic Acids Res.* 7, 1513.
17. Williams, S.A., Slatko, B.E., Moran, L.S., and Desimone, S.M. (1986) *BioTechniques* 4, 138-147.
18. Sanger, F., Nicklen, S., and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
19. Silhavy, T.J., Berman, M.L., and Enquist, L.W. (1984) in *Experiments With Gene Fusions*. Cold Spring Harbor Laboratory, New York.
20. Laemmli, U.K. (1970) *Nature* 227, 680-685.
21. Towbin, H., Staehelin, T., and Gordon, J. (1979) *Proc. Natl. Acad. Sci. USA* 76, 4350-4354.
22. Gillis, S., Ferm, M.M., Ou, W., and Smith, K.A. (1978) *J. Immunol.* 120, 2027-2032.
23. Arico, B. and Rappuoli, R. (1987) *J. Bacteriol.* 169, 2847-2853.
24. Chomczynski, P. and Sacchi, N. (1987) *Analytical Biochemistry* 162, 156-159.
25. *Transfer and Immobilization of Nucleic Acids to S&S Solid Supports*. (1987) Schleicher & Schuell, Inc., Keene, NH.
26. von Gabain, A., Belasco, J.G., Schottel, J.L., Chang, C.Y., and Cohen, S.N. (1983) *Proc. Natl. Acad. Sci. USA* 80, 653-657.
27. Dente, L., Sollazzo, M., Baldari, C., Cesareni, G., and Cortese, R. (1985) in *DNA Cloning. A Practical Approach*. Glover, D.M., Ed. Vol 1, pp 101-107. IRL Press Limited, Oxford.
28. Williams, D.P., Parker, K., Bacha, P., Bishai, W., Borowski, M., Genbauffe, F., Strom, T.B., and Murphy, J.R. (1986) *Protein Engineering* 1, 493-498.
29. Amann, E., and Brosius, J. (1985) *Gene* 40, 183-190.
30. Bailon, P., Weber, D.V., Keeney, R.F., Fredericks, J.E., Smith, C., Familletti, P.C. and Smart, J.E. (1987) *Bio/Technology* 5, 1195-1198.
31. Hartmann, G., Honikel, K.O., Knusel, F., and Nuesch, J. (1967) *Biochim. Biophys. Acta* 145, 843-844.
32. Robinson, M., Lilley, R., Little, S., Emtage, J.S., Yarranton, G., Stephens, P., Millican, A., Eaton, M., and Humphreys, G. (1984) *Nucleic Acids Res.* 12, 6663-6671.
33. George, H.J., L'Italien, J.J., Pilacinski, W.P., Glassman, D.L., and Krzyzek, R.A. (1985) *DNA* 4, 273-281.
34. Gheysen, D., Iserentant, C., Derom, C., and Fiers, W. (1982) *Gene* 17, 55-63.
35. Zurawski, S.M., Pope, K., Cherwinski, H., and Zurawski, G. (1986) *Gene* 49, 61-68.