Isolation and mapping of a polymorphic DNA sequence (pEFD64.1) on chromosome 3 [D3S42]

Y.Nakamura, E.Fujimoto, P.O'Connell, M.Leppert, G.M.Lathrop, J.-M.Lalouel and R.White

---

The Howard Hughes Medical Institute, University of Utah Medical School, Salt Lake City, UT 84132, USA

SOURCE/DESCRIPTION: A 5.2 kb TaqI fragment of cosmid EFD64 isolated by a HBV-3 oligonucleotide (1) was subcloned into AccI site of pUC18.

POLYMORPHISM: HinfI identifies two systems of VNTR polymorphisms with bands between 0.5 and 1.5 kb (9 alleles - system 1), or 1.7 and 1.9 kb (3 alleles - system 2). TaqI, RsaI, BglII, EcoRI, BamHI, HindIII and PvuII identify the 1 system VNTR polymorphism.

FREQUENCY: With HinfI, 85 % (system 1) and 60 % (system 2) heterozygosity were observed in 84 unrelated Caucasians.

NOT POLYMORPHIC FOR: Not known

CHROMOSOMAL LOCALIZATION: pEFD64.1 has been assigned to chromosome 3 by linkage analysis (2) with the locus (APOD) known to span this chromosome(3).

MENDELIAN INHERITANCE: Co-dominant segregation of the HinfI RFLP was observed in 42 three generation families.

PROBE AVAILABILITY: Freely available (contact Y.N.) or will be available from ATCC.

OTHER COMMENTS: RFLPs were observed under the normal hybridization and washing conditions.

REFERENCES:

1. Y. Nakamura et al., Science **235**:1616-1622 (1987)
2. G.M. Lathrop et al., Am. J. Hum. Genet. **37**:482-498 (1985)
3. D. Drayna et al., DNA, **6**:194-204 (1987)