Isolation and mapping of a polymorphic DNA sequence (cTB14.36) on chromosome 10 [D10S35]

T.Bragg, Y.Nakamura, E.Fujimoto, C.Jones¹ and R.White

The Howard Hughes Medical Institute, University of Utah Medical School, Salt Lake City, UT 84132 and

¹Eleanor Roosevelt Institute for Cancer Research, Denver, CO 80262, USA

SOURCE/DESCRIPTION: This cosmid was isolated from a somatic hybrid cell line 762-8A which included chromosome 10 and Y as human chromosomes (1). The cosmid includes a nearly 40 kb of human DNA digested partially with Sau3AI at the BamHI site of pWE15.

POLYMORPHISM: Msp I identifies a two allele polymorphism with a band at 4.4 kb (M1) or 3.1 kb (M2).

FREQUENCY : Estimated from 64 unrelated Caucasians.

M1 : 0.51

M2 : 0.49

NOT POLYMORPHIC FOR: TaqI, RsaI, BglII, PstI and PvuII

CHROMOSOMAL LOCALIZATION: cTB14.36 has been assigned to chromosome 10 by multipoint linkage analysis (2) with loci mapped on chromosome 10 (3).

MENDELIAN INHERITANCE: Co-dominant segregation of RFLP was observed in 32 three generation families.

PROBE AVAILABILITY: Freely available (contact Y.N.).

OTHER COMMENTS: RFLPs were observed after competitive hybridization with total human DNA.

REFERENCES:

1. Fisher et al., Am. J. Hum. Genet. 40 :503-511 (1987)
2. G.M. Lathrop et al., Am. J. Hum. Genet. 37:482-498 (1985)
3. G.M. Lathrop et al., Genomics 2:157-164 (1988)