Published in final edited form as:

Neuroscience. 2011 October 13; 193: 399–410. doi:10.1016/j.neuroscience.2011.07.039.

Phosphodiesterase-5 (PDE5) is a Therapeutic Target for Peripheral Neuropathy in Diabetic Mice

Lei Wang¹, Michael Chopp^{1,2}, Alexandra Szalad¹, Zhongwu Liu¹, Marianne Bolz¹, Francisco Moniche Álvarez¹, Mei Lu¹, Li Zhang¹, Yisheng Cui¹, Rui Lan Zhang¹, and Zheng Gang Zhang¹

¹Department of Neurology, Henry Ford Health Sciences Center, 2799 W. Grand Boulevard, Detroit, Michigan 48202

²Department of Physics, Oakland University, Rochester, Michigan 48309

Abstract

Peripheral neuropathy is a common and major complication of diabetes, the underlying mechanisms of which are not fully understood. Using a mouse model of type II diabetes, the present study investigated the role of phosphodiesterase-5 (PDE5) in peripheral neuropathy. BKS.Cg-*m*+/*Lepr*^{db}/*J*(db/db) mice were treated with sildenafil, a specific inhibitor of PDE5, at doses of 2 and 10 mg/kg or saline. Levels of PDE5 and morphometric parameters in sciatic nerve tissue as well as the motor and sensory function were measured in these mice. In diabetic mice, PDE5 expression in sciatic nerve tissue was significantly upregulated, while the myelin sheath thickness, myelin basic protein (MBP), and subcutaneous nerve fibers were significantly reduced. Treatment with sildenafil, significantly improved neurological function, assayed by motor and sensory conducting velocities and thermal and mechanical noxious stimuli, concomitantly with increases in myelin sheath thickness, MBP levels, and subcutaneous nerve fibers. In vitro, hyperglycemia upregulated PDE5 in Schwann cells, and reduced Schwann cell proliferation, migration and expression of brain-derived neurotrophic factor (BDNF). Blockage of PDE5 with sildenafil, increased cGMP, and completely abolished the effect of hyperglycemia on Schwann cells. Sildenafil upregulated cGMP-dependent protein kinase G1 (PKG1), whereas inhibition of PKG1 with a PKG inhibitor, KT5823, suppressed the inhibitory effect of sildenafil on Schwann cells. These data indicate that hyperglycemia substantially upregulates PDE5 expression and that the cGMP/PKG signaling pathway activated by sildenafil mediates the beneficial effects of sildenafil on diabetic peripheral neuropathy.

Keywords

sildenafil; PDE5; peripheral neuropathy; function; diabetes; mice

Peripheral neuropathy is a major complication of diabetes, and a cause of considerable morbidity (Feldman et al., 1997). Diabetic peripheral neuropathy is characterized by the loss and/or degeneration of neurons, Schwann cells, and neuronal fibers and the slowing of nerve

© 2011 IBRO. Published by Elsevier Ltd. All rights reserved.

Please send all correspondence to: Dr. Zheng Gang Zhang, Department of Neurology, Henry Ford Hospital, 2799 West Grand Boulevard, Detroit, MI 48202, Tel: 313-916-5456, Fax 313-916-1318, zhazh@neuro.hfh.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

conduction velocities (Brownlee et al., 1986, Greene et al., 1992). The underlying mechanisms of which are not fully understood.

Sildenafil, a phosphodiesterase type 5 (PDE-5) antagonist, attenuates catabolism and causes intracellular accumulation of cyclic guanosine monophosphate (cGMP). cGMP is generated by cytoplasmic soluble guanylate cyclases and degraded by PDE5 (Ignarro, 1991, Corbin and Francis, 1999). cGMP effector molecules include cGMP-dependent protein kinases (PKGs) and cGMP-gated ion channels (Butt et al., 1993). PKGs are activated by cGMP and classified into two types, PKGI and PKGII (Lohmann et al., 1997, Pfeifer et al., 1999). cGMP/PKGI regulates cell proliferation, axon guidance and synaptic plasticity (Zhuo et al., 1994).

Treatment of diabetic peripheral neuropathy with sildenafil improves blood supply to the vasa nervorum and functional outcome through NO-cGMP in rodents (Patil et al., 2004, Neeraj Arora, 2007). Case report data show that patients with diabetic peripheral neuropathy treated with sildenafil improve their symptoms (Hackett, 2006). However, the molecular mechanisms by which PDE5 ameliorates diabetic peripheral neuropathy have not been extensively investigated.

Using a mouse model of type II diabetes which develops severe peripheral neuropathy, we investigated the role of PDE5 in diabetic peripheral neuropathy and report that blockage of PDE5 with sildenafil substantially improved neurological outcome and suppressed PDE5 expression upregulated by hyperglycemia in Schwann cells and that the cGMP/PKG signaling pathway activated by sildenafil in Schwann cells mediates the beneficial effects of sildenafil on diabetic peripheral neuropathy.

EXPERIMENTAL PROCEDURES

All experimental procedures were carried out in accordance with NIH Guide for the Care and Use of Laboratory Animals and approved by the institutional Animal Care and Use Committee of Henry Ford Hospital. Male BKS.Cg-*m*+/*+**Lepr*^{*db*}/*J* (db/db) mice (Jackson Laboratories, Bar Harbor, Maine) aged 16 weeks were used. Age-matched heterozygotes mice (db/m), a non-penetrant genotype (Jackson Laboratories), were used as the control animals. For isolation of Schwann cell, 4 day-old C57BL6 mice (Jackson Laboratories) were be used.

Sildenafil treatment

db/db mice at age 16 weeks were treated daily with sildenafil (2 mg/kg/d or 10 mg/kg/d, subcutaneous, n=10/each group, Viagra, Pfizer Inc.) for 28 days. For control, db/db mice and db/m mice were treated with the same volume of saline as a vehicle (n=10/each group), respectively. We defined the day just prior to the initiation of the treatment as day 0 and days after initial sildenafil treatment as days after treatment. All mice were sacrificed 56 days after initial treatment. Doses of 2 and 10 mg/kg/d of sildenafil were selected based on published studies showing that this therapeutic dosing level is comparable in human (Walker et al., 1999). Sildenafil powder was dissolved in saline.

Bromodeoxyuridine labeling

To examine proliferating cells, bromodeoxyuridine (BrdU, 100 mg/kg, i.p, Sigma-Aldrich, St.Louis, MO), the thymidine analog that is incorporated into the DNA of dividing cells during S-phase (Zhang et al., 2001), was injected (i.p) to the mice daily for 14 days starting at the initial day of sildenafil treatment.

Blood glucose levels were measured by using an instant check meter (Roche Diagnostics, Indianapolis, IN). Blood glucose levels and body weight were measured at day 0, and day 56 after the treatment. Electrophysiological measurements and functional tests were performed in all mice at baseline (day 0) and then at day 28 after the treatment, a time point just termination of the sildenafil treatment, and day 56 after the treatment. All procedures and analyses were performed by investigators who were blinded to the treatment administered.

Neurophysiological Measurements

Sciatic nerve conduction velocity was assessed with orthodromic recording techniques, as described by Li 2005 (Li et al., 2005). Briefly, trigger single square wave current pulses were delivered using an isolated pulse stimulator (Model 2100, A-M Systems, Everett, WA). The simultaneous electromyographies were recorded by two sterilized electrodes placed into the intrinsic foot muscles with a Grass Amplifier (Model P5, Grass Instruments, Quincy, MA). During the measurements, animal rectal temperature was kept at $37 \pm 1.0^{\circ}\text{C}$ using a feedback controlled water bath. Motor nerve conduction velocity (MCV) and sensory nerve conduction velocity (SCV) were calculated according to a published study (Li et al., 2005).

Tail-flick and hot plate tests

To examine thermal hyperalgesia, tail-flick and hot plate tests were employed according to published methods (South and Smith, 1998, Vanderah et al., 2001). Briefly, for tail-flick test, a mouse was restrained in a conical polypropylene tube with an opening through which its tail was exposed. Approximately 2 cm of the mouse's tail was immersed into a $52^{\circ}\text{C} \pm 0.2$ water bath and the time until the rodent flicks or removes its tail was recorded (Vanderah et al., 2001). For hot plate test, a mouse was placed within a plexiglass chamber on a transparent glass surface and allowed to acclimate for at least 20 min. A thermal stimulation meter (IITC Model 39 Hot Plate Analgesia Meter, IITC Life Science) was used with floor temperature at 55°C (manufacturer's setup). The latency of paw withdrawal in response to the radiant heat was recorded (South and Smith, 1998). Cut-off periods of 10 and 15s were employed to avoid tissue damage for the tail-flick and hot plate tests, respectively. In both tests, at least three readings per animal were taken at 15 min intervals, and the average was calculated.

Tactile allodynia test

To examine tactile allodynia, we employed Von Frey filaments (Stoelting, Wood Dale, IL) to stimulate paw withdraw according to published protocols (Chaplan et al., 1994). Briefly, a series of filaments with force ranged from 0.02 to 1.4 g were applied to the plantar surface of the left hindpaw with a pressure causing the filament to buckle. A paw withdraw in response to each stimulus was recorded and a 50% paw withdrawal threshold was calculated according to a published formula (Dixon, 1980, Chaplan et al., 1994).

Tissue preparation

The sciatic nerves were harvested at the mid-thigh level according to published protocol (Mousa et al., 2001). The left and right side sciatic nerves were fixed in 4% paraformaldehyde for immunohistochemistry (Zhang et al., 2001) and 2.5% purified glutaraldehyde buffer (Electron Microscopy Sciences, PA) for morphometric studies (Di Scipio et al., 2008), respectively. Epidermal foot pads from left hind feet were fixed in 4% paraformaldehyde for immunohistochemistry.

Staining myelin sheets

We adapted a protocol to stain myelin sheets in paraffin-embedded tissue (Di Scipio et al., 2008). Briefly, the sciatic nerves were fixed in the 2.5% glutaraldehyde and 0.5% saccharose

(Sigma) on PBS buffer for 6–8 h, and then immersed in 2% osmium tetroxide (Sigma) for 2h. The specimens were then dehydrated with numerous alcohol passages and embedded in paraffin (Di Scipio et al., 2008). Semi-thin transverse sections (2- μ m thick) were cut and stained with 1% toluidine blue and three semi-thin sections per mouse were analyzed.

Immunohistochemistry

Sciatic nerve tissues and epidermal foot pads were isolated and embedded in paraffin. Three cross sections (6- μ m-thick) or three longitudinal sections (6- μ m-thick) in the one in ten series (60 μ m apart) for each animal were used for immunostaining according to our published protocols (Zhang et al., 2001). The following primary antibodies were used: polyclonal rabbit anti-myelin basic protein (MBP, 1:400, Dako Denmark), polyclonal anti myelin associated glycoprotein (MAG, 1:200 R&D system), polyclonal anti-neurofilament, heavy chain (NF-H, 1:10000, Thermo Scientific), polyclonal rabbit anti brain-derived neurotrophic factor (BDNF, 1:300, Santa Cruz Biotechnology), polyclonal goat anti-cGKI (1:300, Santa Cruz Biotechnology), monoclonal mouse anti-bromodeoxyuridine (BrdU, 1:100, Dako), and polyclonal rabbit anti-protein gene product 9.5 (PGP9.5, 1:1,000; MILLIPORE). Normal rabbit or goat IgG was used as a negative control. Sections were counterstained with 4',6-diamidino-2-phenylindole (DAPI, 1:5000, Pierce Biotechnology, Rockford, IL).

Image Analysis and Quantification

Image acquisition and analysis were performed using a computer imaging analysis system (MicroComputer Imaging Device, MCID, Imaging Research, Linton, Cambridge, UK) (Zhang et al., 2001). For morphometric analysis of sciatic nerves, three fields of view per section were randomly acquired under a 100 $\times$ oil immersion objective (BX40; Olympus Optical, Tokyo, Japan). Myelinated fiber diameter, axon diameter, and myelin sheath thickness were measured. The g-ratio (the quotient axon diameter/fiber diameter) was calculated to measure the degree of myelination. At least 200 myelinated fibers were measured per animal (Sameni et al., 2008).

For immunostaining, three sections spaced as 60 μ m interval for each staining were used for analysis from each mouse and three fields of the view per section were randomly imaged under a 40 $\times$ objective. For quantization of MBP, NF-H, MAG and BDNF immunoreactive, cross sections or longitudinal sections were digitized. Data are presented as the percentage of immunoreactive area within the total imaged area. For measurements of proliferative Schwann cells, S100 and BrdU-positive cells were counted in longitudinal sections. Data are presented as the percentage of S100/BrdU-positive cells.

For subcutaneous innervations, every PGP9.5-positive nerve fiber in dermis within a 50- μ m radius from the epidermal/dermal junction was measured by means of MCID and the number of PGP9.5-immunoreactive nerve fibers with a cross-sectional area of 20 μ m² was counted. Data are presented as nerve fiber density calculated based a published formula (Kakinoki et al., 2006).

Real-time RT-PCR

Total RNA samples from cells were isolated using the Stratagene Absolutely RNA MicroRNA isolation kit (Stratagene, La Jolla, CA), according to the manufacturer's instructions. The complementary DNA (cDNA) was reversely transcribed from the same concentrations of total RNA products using random hexamers and M-MLV reverse-transcriptase (Invitrogen). Using the SYBR Green real-time PCR method (Wang et al., 2005), quantitative PCR was performed on an ABI 7000 PCR instrument (Applied Biosystems, Foster City, CA) by means of three-stage program parameters provided by the

manufacturer, as follows; 2 min at 50 °C, 10 min at 95 °C, and then 40 cycles of 15 s at 95 °C and 1 min at 60 °C. Specificity of the produced amplification product was confirmed by examination of dissociation reaction plots. Each sample was tested in triplicate, and samples obtained from three independent experiments were used for analysis of relative gene expression using the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). The following primers for real-time PCR were designed using Primer Express software (ABI): Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (FWD, AGA ACA TCA TCC CTG CAT CC; REV, CAC ATT GGG GGT AGG AAC AC), BDNF (FWD, TAC TTC GGT TGC ATG AAG GCG; REV, GTC AGA CCT CTC GAA CCT GCC), mPDE5 (FED, TGC ATT GGA AGT TTG CTT TG; REV, GCT TCC AGG AAC TGC TCA TC) and hPDE5 (FED, ACT TGC CAT TGC TGA TTG CTG; REV, AGT AAA GCT GGG CAA GTG GA).

Western blot analysis

Western blot was performed according to published methods (Wang et al., 2005). Briefly, equal amounts of proteins were loaded on 10% SDS-polyacrylamide gel. After electrophoresis, the proteins were transferred to nitrocellulose membranes, and the blots were subsequently probed with the following antibodies: PDE5 (1:1000; Cell Signaling Technology), cGKI (1:1000; Santa Cruz Biotechnology) and BDNF (N-19) (1:1000; Santa Cruz Biotechnology). For detection, horseradish peroxidase-conjugated secondary antibodies were used (1:2000) followed by enhanced chemiluminescence development (Pierce Biotechnology). Normalization of results was ensured by running parallel Western blot with β -actin antibody. The optical density was quantified using an image processing and analysis program (Scion Image, Ederick, MA).

Cell culture

Primary mouse Schwann cell culture was performed according to published studies (Paivalainen et al., 2008). Briefly, Schwann cells were harvested from 4 day old C57BL/6 mouse sciatic nerves and seeded on 60 cm plastic culture dishes coated with Poly D-line (Sigma) in primary Schwann cell growth medium which contains Dulbecco's Modified Eagle's Medium (DMEM), heregulin- β 1 (2 ng/ml, Sigma), L-glutamine (2 mM, R&D system), forskolin (0.5 μ M, Sigma), bovine pituitary extract (20 μ g/ml, Sigma), basic fibroblast growth factor (bFGF, 10 ng/ml, R&D system) and 10% house serum (R&D system) (Paivalainen et al., 2008). The purity of Schwann cells was approximately 95% based on cellular morphology and immunocytochemical staining with antibodies against S100 protein and glial fibrillary acidic protein.

Rat immortalized Schwann cells line (American Type Culture Collection, ATCC, Manassas, VA) was maintained in DMEM with 10% fetal bovine serum (GIBCO). For optimally growing primary mouse Schwann cells, DMEM containing 25 mmol/L glucose was used, this concentration of glucose has been used by many others for studying primary mouse Schwann cell biological function (Honkanen et al., 2007, Paivalainen et al., 2008). However, to mimic physiological condition, we also employed immortalized rat Schwann cells that were cultured DMEM containing 5.5 mmol/L glucose (Li et al., 2005;). Therefore, we defined 5.5 and 25 mmol/L glucose concentrations as the normal glucose (NG) for immortalized and primary Schwann cells, respectively, while glucose levels at 25.5 and 45 mmol/L were defined as high glucose (HG) for immortalized and primary Schwann cells, respectively. These glucose concentrations for the in vitro hyperglycemia experiments have been used by others (Russell et al., 2002, Li et al., 2005, Vincent et al., 2005).

Human Schwann cells (ScienCell Research Laboratories, Carlsbad, CA) derived from primary culture were maintained in manufacture's Schwann cell medium (ScienCell Research Laboratories).

To examine the effect of high glucose on human Schwann cells, human Schwann cells were cultured in Schwann cell medium containing 5.5 and 25.5 mmol/L glucose.

Transfection of primary Schwann cells with small interfering RNA against PDE5 (siRNA-PDE5)

primary mouse Schwann cells were transfected with siRNA against mouse PDE5 (siRNA-PDE5, Dharmacon, Inc. Chicago, IL) using lipofectamine 2000 (Invitrogen) following the manufacturer's instructions. Briefly, primary mouse Schwann cells were plated out in 6 well plates and cultured until they were 80% confluent. They were then transfected with 100 pmol siRNA-PDE5 in Opti-MEM® I Reduced-Serum Medium. Levels of PDE5 mRNA and protein were measured 48 and 72h after transfection.

To investigate the effect of sildenafil on Schwann cell proliferation, Schwann cells were seeded in a 24-well dish at a density of 2.5×10^4 cells per well and incubated in normal and high glucose levels with sildenafil at concentrations from 0 to 300 nM for 72 hours. BrdU (10 μ mol/L) was added 4 and 24 hours in immortalized and primary Schwann cells, respectively, prior to termination of the experiments. The cells were fixed for BrdU immunostaining. The number of BrdU positive cells was counted in 5 fields of view under a 20X objective. To further examine the effect of inhibition of PDE5 on cell proliferation, primary mouse Schwann cells were transfected with siRNA-PDE5. To examine whether the PKG pathway is involved in the effect of sildenafil on cell proliferation, a specific PKG inhibitor, KT5823 (1 μ M, CALBIOCHEM) was used in presence or absence of sildenafil.

To examine the effect of sildenafil on migration of primary mouse Schwann cells, a modified Boyden's chamber assay was employed, as described previously (Wang et al., 2006). Briefly, the polycarbonate filter (8- μ m pore size) (Neuro Probe, Inc. Gaithersburg, MD) was coated with 50 μ g/ml fibronectin (Chemicon) and 0.1% gelatin (Sigma) and placed between upper and lower chambers. The cells were pre-incubated in normal and high glucose levels with sildenafil at concentrations of 0 to 300 nM in the presence or absence of the specific PKG inhibitor, KT5823 (1 μ M) for 72 hours. Cell suspensions (5×10^4 cells per well) were placed in the upper chamber, and the lower chamber was filled with medium containing human recombinant nerve growth factor (100 ng/ml, R&D Systems). The chamber was incubated for 5 hours at 37°C and 5% CO₂. Migrating cells caught in the membrane were then stained using hematoxylin and eosin (Anatech LTD). The number of cells that migrated through the filter was counted in 5 fields of view under a 40X objective.

To examine the effect of hyperglycemia on cGMP and PKG, we measured cGMP levels and PKG activity using a cGMP ELISA Kit (R&D system) or a PKG Assay Kit (MBL International Corporation), respectively, according to the manufacturer's instructions. Briefly, primary mouse Schwann cells were cultured on 6-well plates at 70% to 80% confluence under normal and high glucose conditions with or without sildenafil (150 nM) for 72 hours (Wang et al., 2005). Culture medium or cells were collected for the measurement of cGMP and activation of PKG, respectively. The concentration of cGMP or PKG was measured as an OD value at 540 nm or 450 nm.

To investigate the effect of sildenafil on gene expression in Schwann cells, Schwann cells were incubated in normal and high glucose levels with sildenafil at concentrations of 0 to 300 nM in the presence or absence of KT5823 (1 μ M) for 24 or 72 hours, respectively. Levels of mRNAs and proteins were measured with real-time RT-PCR or Western blots, respectively.

Statistical analysis

Data were evaluated for normality. Data transformation was considered if data were not normal. As the result, ranked data were used for the analysis. The global test using Generalized Estimating Equation (GEE) were implanted to test the group difference on functional recovery measured. The analysis started testing the treatment overall group effect, followed by a subgroup analysis at the 0.05 level, if the overall group effect was detected at the 0.05 level. The global test on multiple outcomes is more efficient than a single outcome, when the dose effects are consistent on all the outcomes (e.g., the positive correlation). A significant dose effect on the functional recovery ($p < 0.05$ based on the global test) would be further tested on individual outcome at 0.05. One-way ANOVA was used to test the overall group effect. The data were presented as mean $\pm$ SE. A value of $P < 0.05$ was taken as significant.

RESULTS

Sildenafil improves neurological function in diabetic mice

Male BKS.Cg-*m*^{+/+}*Lepr*^{db}/*J* (db/db) mice at age 4 weeks developed diabetes, as indicated by elevation of blood glucose levels. To examine whether diabetic mice acquire peripheral neuropathy, we measured the motor and sensory conducting velocity (MCV and SCV) in sciatic nerves of these mice. Electrophysiological recordings showed that diabetic mice had significant decreases of the MCV and SCV compared with the age matched non-diabetic mice (Fig. 1A and 1B). In addition, thermal and mechanical noxious stimuli measured by tail flick test, hot plate test and tactile allodynia test, respectively, revealed that diabetic mice exhibited a significant increase in response threshold to hot and mechanical stimuli compared with non-diabetic mice (Fig. 1C, 1D and 1E). Collectively, these data indicate that diabetic mice develop severe peripheral neuropathy. Western blot analysis revealed that PDE5 levels in sciatic nerve tissue from diabetic mice significantly increased compared with the tissue from non-diabetic mice (Fig. 1F). Treatment of the diabetic mice with sildenafil at doses of 2 and 10 mg/kg significantly attenuated the reduced MCV and SCV starting 28 days after treatment. This increase in MCV and SCV persisted at least for 56 days after treatment, and was associated with substantial improvements in response to thermal and mechanical stimuli compared with mice treated with saline during 28 to 56 days after treatment (Fig. 1), suggesting that inhibition of PDE5 by sildenafil reduces peripheral neuropathy. Chronic treatment with sildenafil significantly reduces animal body weight (Ayala et al., 2007). Interestingly, we found that diabetic mice treated with saline increased their mean body weight by approximately 4 g during 56 days, while diabetic mice treated with sildenafil at 2 and 10 mg/kg did not gain any body weight, although differences between the saline and sildenafil groups did not reach statistical significance (Table 1). Treatment with sildenafil did not significantly affect blood glucose levels of the diabetic mice (Table 1).

The effect of sildenafil on myelinated neuronal fibers and Schwann cells in diabetic mice

Degeneration of neuronal fibers and Schwann cells leads to decreases in motor and sensory conducting velocity (Foster et al., 1980). To investigate effects of sildenafil on subcutaneous innervation and Schwann cells, we examined subcutaneous nerve fibers, sciatic nerves and Schwann cells in diabetic mice. Immunostaining of mouse plantar skin with an antibody against PGP9.5, which is a cytoplasmic ubiquitin C-terminal hydrolase expressed in all types of efferent and afferent peripheral nerve fibers (Bianchi et al., 2004), revealed that nerve fiber density was 40.6 ± 2.6 counts/mm in non-diabetic mice, which is comparable to published data (Kakinoki et al., 2006). Diabetic mice exhibited significant reduction of the number of nerve fibers, indicating that hyperglycemia induces loss of subcutaneous

innervation (Fig. 2). However, diabetic mice treated with sildenafil showed a marked increase in nerve fiber density (Fig. 2).

Using a myelin sheet staining protocol developed by Di Scipio et al (Di Scipio et al., 2008), we then measured myelinated sciatic nerves. We found substantial reductions in sciatic nerve fiber diameter and myelin sheath thickness and significant increases in g ratio (axon diameter/fiber diameter) in the diabetic mice treated with saline compared with non-diabetic mice (Table 2). Treatment with sildenafil significantly reversed mean myelin sheath thickness reduction and the g ratio increase (Table 2). In addition, the myelinated fiber and axon diameters were displaced toward larger sizes compared with those of saline treated diabetic mice.

To confirm morphological findings on the sciatic nerves, we performed double immunofluorescent staining with antibodies against myelin basic protein (MBP) and neurofilament heavy chain (NF-H) to detect myelin and neurofilaments, respectively. Consistent with morphological data, double immunostaining revealed that diabetic mice treated with saline had substantial reduction of MBP and NF-H immunoreactivity and that many NF-H immunoreactive neurofilaments were not surrounded by MBP (Fig. 3). In contrast, treatment with sildenafil reversed MBP and NF-H immunoreactive levels approximately back to levels found in non-diabetic mice (Fig. 3). Moreover, treatment of diabetic mice with sildenafil significantly increased myelin associated glycoprotein (MAG) immunoreactivity ($8.4 \pm 0.8\%$, $n=10$) compared with diabetic mice treated with saline ($4.19 \pm 0.7\%$, $n=10$, $p<0.05$). Collectively, these data suggest that in addition to subcutaneous nerve fibers, sildenafil may prevent loss of sciatic nerves and enhance myelinating Schwann cells. Indeed, double immunofluorescent staining at 56 days after treatment showed that sildenafil significantly increased proliferating Schwann cells, measured by the number of bromodeoxyuridine (BrdU) and S100, a marker of Schwann cells, compared with the saline treatment (Fig. 4).

The effect of sildenafil on Schwann cell biological function

Aforementioned data that diabetic mice had elevated PDE5 levels and sildenafil may augment myelinating Schwann cells in diabetic mice prompted us to investigate the direct effect of sildenafil on Schwann cells. Using primary mouse Schwann cells harvested from sciatic nerve tissue, we examined whether glucose levels affect PDE5 expression in Schwann cells. Primary mouse Schwann cells were cultured with normal and high glucose. We found that primary mouse Schwann cells expressed PDE5 and that the high glucose level substantially increased PDE5 levels measured by Western blot (Fig. 5). Concurrently, ELISA revealed that the high glucose significantly ($p<0.05$) reduced concentrations of cGMP in primary mouse Schwann cells (7.8 ± 0.6 pmol/mg, $n=3$) compared with the normal glucose group (14.4 ± 1.4 pmol/mg, $n=3$). Furthermore, the high glucose significantly decreased the number of BrdU positive primary mouse Schwann cells (Fig. 5), but did not affect the number of TUNEL positive cells ($4.9 \pm 0.5\%$ in the control vs. $5.9 \pm 0.6\%$ in high glucose, $n=3$ /group), indicating that the high glucose decreases primary mouse Schwann cell proliferation. The high glucose also substantially suppressed primary mouse Schwann cell migration measured using a Boyden chamber assay (Fig. 5). To examine whether inhibition of PDE5 by sildenafil could attenuate the effect of elevated glucose on Schwann cell biological function, primary mouse Schwann cells were treated with sildenafil at various concentrations. We found that sildenafil at a dose of 150 nM completely rescued Schwann cell proliferation and migration when cultured with the high glucose (Fig. 5). Sildenafil elevated cGMP levels in Schwann cells (11.8 ± 1.03 pmol/mg, $n=3$) compared with the high glucose group (7.8 ± 0.6 pmol/mg, $n=3$, $p<0.05$). To verify the specific effect of cGMP on Schwann cell proliferation and migration, primary mouse Schwann cells were treated with 8-Br-cGMP (10 μ mol/L), a stable analog of cGMP. Br-cGMP significantly increased the

proliferation and migration of primary mouse Schwann cells when cultured with the high glucose (Fig. 5). To further confirm the effects of high glucose and sildenafil on Schwann cell proliferation, we employed immortalized Schwann cells that permit us to use glucose concentrations at clinical relevant levels. Immortalized Schwann cells cultured in a high glucose level exhibited significant ($p<0.05$) reduction of cell proliferation ($17.3\pm2.0\%$, $n=3$) compared the cells cultured at normal glucose level ($33.0\pm3.0\%$, $n=3$). Sildenafil (150 nM) suppressed the inhibitory effect of high glucose on the cell proliferation ($31.8\pm3.0\%$, $n=3$, $p<0.05$). These data indicate that high glucose reduces Schwann cell proliferation, although primary Schwann cells have different characteristics compared with immortalized Schwann cells. To further examine the specific effect of PDE5 on Schwann cells, we attenuated endogenous PDE5 with siRNA-PDE5. Transfection of siRNA-PDE5 in primary mouse Schwann cells substantially reduced high glucose-upregulated PDE5 levels compared to the cells transfected with scrambled sequences (Fig. 5) and suppressed high glucose-reduced Schwann cell proliferation ($23\pm2\%$ BrdU⁺ cells vs. $17\pm2\%$ BrdU⁺ cells in high glucose and $15\pm2\%$ BrdU⁺ cells in high glucose and scrambled sequences, $n=3$, $p<0.05$), indicating that PDE5 mediates Schwann cell proliferation under high glucose condition. In addition, high glucose upregulated PDE5 expression in human Schwann cells (Fig. 5). Collectively, these data indicate that PDE5 upregulated by hyperglycemia in Schwann cells causes reduction of cGMP, which leads to decreasing proliferation and migration of Schwann cells.

The effect of sildenafil on PKG

A primary action of elevated cGMP levels is the stimulation of PKG (Wang and Robinson, 1997). To examine whether PKG is involved in the effect of cGMP on Schwann cells, expression of PKG was measured by Western blot. Levels of PKGI significantly decreased in primary mouse Schwann cells when cultured with high glucose compared with that cultured with normal glucose (Fig. 5). High glucose also reduced PKG activity (1 ± 0.01 , $n=3$) compared with normal glucose (2.25 ± 0.02 , $n=3$, $p<0.05$). Treatment of primary mouse Schwann cells with sildenafil under a high glucose condition increased PKGI protein levels (Fig. 5) and PKG activity (2.0 ± 0.01 vs. 1 ± 0.01 , $p<0.05$, $n=3$ /group). Furthermore, blockage of PKG with a PKG inhibitor, KT5823, abolished sildenafil-augmented primary mouse Schwann cell proliferation and migration (Fig. 5). These findings indicate that PKGI mediates the effect of cGMP on primary mouse Schwann cells.

To examine whether Schwann cells express PKGI *in vivo*, double immunofluorescent staining was performed. We found that S100 positive Schwann cells were PKGI immunoreactive (Fig. 6).

The effect of sildenafil on BDNF

BDNF, a neurotrophic factor, enhances peripheral nervous system myelination by promoting proliferation and migration of Schwann cells (Tolwani et al., 2004). We thus examined the effect of sildenafil on BDNF expression. Immunohistological analysis of sciatic nerve tissue revealed that the diabetic mice treated with saline exhibited a significantly reduction of BDNF immunoreactivity compared with the non-diabetic mice (Fig. 6), whereas the diabetic mice treated with sildenafil at dose 2 mg/kg and 10 mg/kg did not show decreases in BDNF immunoreactivity compared with non-diabetic mice 56 days after treatment (Fig. 6). The effect of glucose and sildenafil on BDNF was also examined in immortalized Schwann cells. The high glucose level downregulated BDNF expression in immortalized Schwann cells (Fig. 6). Sildenafil and Br-cGMP abolished the effect of the high glucose on BDNF expression (Fig. 6), whereas blockage of PKG by KT5823 suppressed sildenafil-increased BDNF (Fig. 6).

DISCUSSION

The present study provides novel insights into mechanisms of PDE5 in mediating diabetic peripheral neuropathy by demonstrating that hyperglycemia substantially upregulated PDE5 in Schwann cells and that the cGMP/PKG signaling pathway activated by sildenafil promotes the beneficial effects of sildenafil on diabetic peripheral neuropathy.

Cyclic GMP is involved in the regulation of many neural functions, including neurotransmission, long term potentiation, gene expression, and even neurotoxicity and neurodegeneration (Wang and Robinson, 1997). The PDE5 enzyme is highly specific for hydrolysis of cGMP (Corbin and Francis, 1999). Severe peripheral neuropathy develops in type II diabetic mice. Studies of PDE5 primarily focus on vasculature (Jackson et al., 1999). The role of PDE5 in diabetic peripheral neuropathy has not been fully investigated. Experimental data show that treatment of diabetic peripheral neuropathy with sildenafil, a novel inhibitor of PDE5, improves blood supply to the vasa nervorum (Neeraj Arora, 2007). Analysis of human data from case reports show that patients with erectile dysfunction treated with sildenafil improved their diabetic peripheral neuropathy symptoms, which was attributed to improvement of blood supply (Hackett, 2006). The present study demonstrates that Schwann cells expressed PDE5 and that hyperglycemia upregulated PDE5 expression. More importantly, our data indicate that the beneficial effect of sildenafil on diabetic peripheral neuropathy results from targeting PDE5 in Schwann cells.

Schwann cells play an important role in regenerating peripheral nerve system by their capacity to proliferate, migrate, and secrete numerous factors that control nerve regeneration (Yamauchi et al., 2004, Bhatheja and Field, 2006). Schwann cells migrate along axons, ensheath individual axons, and eventually form the myelin sheath (Yamauchi et al., 2004). Decreased proliferation of Schwann cells has also been demonstrated in both human and experimental models of diabetes (Ballin and Thomas, 1968, Gurny et al., 2008). The present study demonstrates that reduction of cGMP levels by hyperglycemia suppresses Schwann cell proliferation and migration, and downregulates BDNF expression, whereas elevation of cGMP either by blockage of PDE5 with sildenafil or by the cGMP analog, Br-cGMP, reverses the effect of hyperglycemia on Schwann cells. Thus, these data indicate that cGMP levels mediate Schwann cell biological function and thereby affect the development of diabetic peripheral neuropathy.

The development of diabetic peripheral neuropathy is correlated with segmental demyelination and axonal degeneration of the sciatic nerve in diabetic mice and is characterized by severe nerve conduction velocity deficits and thermal hypoalgesia (Shirasaki and Rosenbluth, 1991). Myelinating Schwann cells express MBP and MAG, which are required for peripheral nerve myelin formation and involved in both experimental and human demyelinating disease (Ginalski-Winkelmann et al., 1983, Farhadi et al., 2003). Neurofilaments are critical components of axons and reduction of neurofilaments in diabetic neuropathy have been reported (Yagihashi et al., 1990, Scott et al., 1999). Using morphology and immunohistochemistry, the present study shows that reduction of myelin sheets, MBP, MAG, and neurofilaments was associated with impairments of nerve conduction velocities in diabetic mice, which is consistent with previous reports that reduction of myelin proteins such as MBP and MAG decreases MCV in peripheral neuropathy (Trojaborg et al., 1995). Based on the present data together with published studies, we speculate that augmentation of myelinating proteins by sildenafil contributes to improvement of neurological function in diabetic mice, as observed in the present study.

In addition to impairment of nerve conduction velocity that is an indicator of large fiber function, diabetic mice exhibited substantial reduction of intraepidermal nerve fiber density,

which is consistent with common findings in diabetic neuropathy patients (Pittenger GL, 2004, Smith et al., 2005, Kakinoki et al., 2006). The effect of sildenafil on increasing subcutaneous nerve fiber density may contribute to observed improvement of thermal and mechanical stimuli, suggesting that sildenafil also prevents the loss of the cutaneous innervation.

Previous studies demonstrated that cGMP induces Schwann cells to synthesize nerve growth factor (NGF) and neurotrophin 3 (NT3) (Thippeswamy et al., 2005). Our data extend the previous findings by demonstrating that PDE5/cGMP regulates BDNF expression in Schwann cells of diabetic mice. BDNF promotes the recovery of neural function and subcutaneous injections of BDNF enhance myelin formation in the sciatic nerve, indicating that BDNF is required for myelination and regeneration of injured sciatic nerve (Zhang et al., 2000). Together with these prior findings, our data suggest that upregulated BDNF in Schwann cells by cGMP may reduce sciatic nerve injury caused by diabetes.

PKGs are major effectors for cGMP and mediate many of the effects of cGMP in target tissues (Butt et al., 1993). The current study shows that Schwann cells expressed PKG1 and suggests that PKG1 is likely a downstream target of cGMP, because inhibition of PKG1 abolished the effect of cGMP on Schwann cells. However, additional experiments by using different ways to block PKG1 and PKG2 are warranted.

There are some limitations that need to be considered in the present study. Data on development of abnormal behavioral response to mechanical and thermal stimuli are inconsistent in the db/db mouse that shares a number of features with human diabetic neuropathy and is widely used mouse model of type II diabetes (Ii et al., 2005, Kakinoki et al., 2006, Leonelli et al., 2007, Sullivan et al., 2007, Cheng et al., 2009). Our observations that the db/db mouse at age of 16 weeks developed abnormal behavioral response to mechanical and thermal stimuli in the mouse are consistent with published studies (Ii et al., 2005, Kakinoki et al., 2006, Sullivan et al., 2007, Wright et al., 2007), but contrast with others (Walwyn et al., 2006, Wright et al., 2007). We therefore employed several different methods to verify presence of abnormal behavioral changes. A glucose concentration used for culturing primary Schwann cells is another issue. In the present study, glucose at 25 mmol/L was employed to optimally grow primary mouse Schwann cells, which has been used by many others to study Schwann cell biological functions (Honkanen et al., 2007, Paivalainen et al., 2008). However, in vivo a normal glucose level equals to approximately 5.5 mmol/L and a glucose concentration above 10 mmol/L is considered to be diabetic (Acheampong et al., 2009). We therefore employed immortalized Schwann cells, which permitted us to used glucose levels close to in vivo condition. The results of the sildenafil on both Schwann cells are comparable when the cells cultured in normal and high glucose concentrations. Furthermore, a high glucose level close to in vivo condition upregulated PED5 expression in human Schwann cells. We, therefore, believe that PDE5 plays an important role in mediating diabetic peripheral neuropathy.

In summary, the present study reveals a role for PDE5 in Schwann cells in mediating diabetic peripheral neuropathy by showing that inhibition of PDE5 in Schwann cells by sildenafil augments cGMP levels that lead to reduction of the detrimental effect of diabetics on Schwann cells. Our results indicate that sildenafil may have a beneficial effect on the clinical treatment of diabetic peripheral.

Research Highlights

Diabetics upregulated PDE5 expression in sciatic nerve tissue.> Inhibition of PDE5 with sildenafil substantially improved neurological outcome.> cGMP levels augmented by

sildenafil lead to improve Schwann cell biological function. > sildenafil may have a beneficial effect on the treatment of diabetic peripheral neuropathy.

Acknowledgments

This work was supported by the Dykstra Foundation, and NINDS grants PO1 NS23393, and RO1HL 64766. The authors acknowledge Dr. Douglas W. Losordo and Dr. Haruki Sekiguchi's assistance in setting up neurophysiological measurements. The authors thank Cynthia Roberts, and Qing-e Lu for their technical assistance and Deborah Jewell for secretarial support.

LIST OF ABBREVIATIONS

BDNF	brain-derived neurotrophic factor
BrdU	bromodeoxyuridine
bFGF	basic fibroblast growth factor
cGMP	cyclic guanosine monophosphate
db/db	BKS.Cg- <i>m</i> +/ <i>+</i> <i>Lepr</i> ^{<i>db</i>} / <i>J</i> mice
db/m	heterozygotes mice
DAPI	4',6'-diamidino-2-phenylindole
DMEM	Dulbecco's Modified Eagle's Medium
ELISA	enzyme-linked immunosorbent assay
GAPDH	Glyceraldehyde-3-phosphate dehydrogenase
RT-PCR	reverse-transcription polymerase chain reaction
NF-H	neurofilament, heavy chain
MCID	MicroComputer Imaging Device
MCV	Motor nerve conduction velocity
MBP	myelin basic protein
MAG	myelin associated glycoprotein
NO	Nitric Oxide
PDE5	phosphodiesterase type 5
PKG	cGMP-dependent protein kinase G
PGP 9.5	protein gene product 9.5
SCV	sensory nerve conduction velocity

REFERENCES

- Acheampong EA, Roschel C, Mukhtar M, Srinivasan A, Rafi M, Pomerantz RJ, Parveen Z. Combined effects of hyperglycemic conditions and HIV-1 Nef: a potential model for induced HIV neuropathogenesis. *Virology*. 2009; 6:183. [PubMed: 19878567]
- Ayala JE, Bracy DP, Julien BM, Rottman JN, Fueger PT, Wasserman DH. Chronic treatment with sildenafil improves energy balance and insulin action in high fat-fed conscious mice. *Diabetes*. 2007; 56:1025–1033. [PubMed: 17229936]
- Ballin RH, Thomas PK. Hypertrophic changes in diabetic neuropathy. *Acta Neuropathol*. 1968; 11:93–102. [PubMed: 5697605]

- Bhatheja K, Field J. Schwann cells: origins and role in axonal maintenance and regeneration. *Int J Biochem Cell Biol.* 2006; 38:1995–1999. [PubMed: 16807057]
- Bianchi R, Buyukakilli B, Brines M, Savino C, Cavaletti G, Oggioni N, Lauria G, Borgna M, Lombardi R, Cimen B, Comelekoglu U, Kanik A, Tataroglu C, Cerami A, Ghezzi P. Erythropoietin both protects from and reverses experimental diabetic neuropathy. *Proc Natl Acad Sci U S A.* 2004; 101:823–828. [PubMed: 14718663]
- Brownlee M, Vlassara H, Cerami A. Trapped immunoglobulins on peripheral nerve myelin from patients with diabetes mellitus. *Diabetes.* 1986; 35:999–1003. [PubMed: 3743908]
- Butt E, Geiger J, Jarchau T, Lohmann SM, Walter U. The cGMP-dependent protein kinase--gene, protein, and function. *Neurochem Res.* 1993; 18:27–42. [PubMed: 8385276]
- Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile allodynia in the rat paw. *J Neurosci Methods.* 1994; 53:55–63. [PubMed: 7990513]
- Cheng HT, Dauch JR, Hayes JM, Hong Y, Feldman EL. Nerve growth factor mediates mechanical allodynia in a mouse model of type 2 diabetes. *J Neuropathol Exp Neurol.* 2009; 68:1229–1243. [PubMed: 19816194]
- Corbin JD, Francis SH. Cyclic GMP phosphodiesterase-5: target of sildenafil. *J Biol Chem.* 1999; 274:13729–13732. [PubMed: 10318772]
- Di Scipio F, Raimondo S, Tos P, Geuna S. A simple protocol for paraffin-embedded myelin sheath staining with osmium tetroxide for light microscope observation. *Microsc Res Tech.* 2008; 71:497–502. [PubMed: 18320578]
- Dixon WJ. Efficient analysis of experimental observations. *Annu Rev Pharmacol Toxicol.* 1980; 20:441–462. [PubMed: 7387124]
- Farhadi HF, Lepage P, Forghani R, Friedman HC, Orfali W, Jasmin L, Miller W, Hudson TJ, Peterson AC. A combinatorial network of evolutionarily conserved myelin basic protein regulatory sequences confers distinct glial-specific phenotypes. *J Neurosci.* 2003; 23:10214–10223. [PubMed: 14614079]
- Feldman EL, Stevens MJ, Greene DA. Pathogenesis of diabetic neuropathy. *Clin Neurosci.* 1997; 4:365–370. [PubMed: 9358981]
- Foster RE, Whalen CC, Waxman SG. Reorganization of the axon membrane in demyelinated peripheral nerve fibers: morphological evidence. *Science.* 1980; 210:661–663. [PubMed: 6159685]
- Ginalski-Winkelmann H, Omlin FX, Matthieu JM. Myelin basic protein deficit in the PNS of mld mutant mice recovers during development. *Dev Neurosci.* 1983; 6:118–126. [PubMed: 6205845]
- Greene DA, Sima AA, Stevens MJ, Feldman EL, Lattimer SA. Complications: neuropathy, pathogenetic considerations. *Diabetes Care.* 1992; 15:1902–1925. [PubMed: 1464245]
- Gumy LF, Bampton ET, Tolkovsky AM. Hyperglycaemia inhibits Schwann cell proliferation and migration and restricts regeneration of axons and Schwann cells from adult murine DRG. *Mol Cell Neurosci.* 2008; 37:298–311. [PubMed: 18024075]
- Hackett G. PDE5 inhibitors in diabetic peripheral neuropathy. *Int J Clin Pract.* 2006; 60:1123–1126. [PubMed: 16939556]
- Honkanen H, Lahti O, Nissinen M, Myllyla RM, Kangas S, Paivalainen S, Alanne MH, Peltonen S, Peltonen J, Heape AM. Isolation, purification and expansion of myelination-competent, neonatal mouse Schwann cells. *Eur J Neurosci.* 2007; 26:953–964. [PubMed: 17714189]
- Ignarro LJ. Heme-dependent activation of guanylate cyclase by nitric oxide: a novel signal transduction mechanism. *Blood Vessels.* 1991; 28:67–73. [PubMed: 1672101]
- Li M, Nishimura H, Kusano KF, Qin G, Yoon YS, Wecker A, Asahara T, Losordo DW. Neuronal nitric oxide synthase mediates statin-induced restoration of vasa nervorum and reversal of diabetic neuropathy. *Circulation.* 2005; 112:93–102. [PubMed: 15983249]
- Jackson G, Benjamin N, Jackson N, Allen MJ. Effects of sildenafil citrate on human hemodynamics. *Am J Cardiol.* 1999; 83:13C–20C.
- Kakinoki B, Sekimoto S, Yuki S, Ohgami T, Sejima M, Yamagami K, Saito K. Orally active neurotrophin-enhancing agent protects against dysfunctions of the peripheral nerves in hyperglycemic animals. *Diabetes.* 2006; 55:616–621. [PubMed: 16505223]

- Leonelli E, Bianchi R, Cavaletti G, Caruso D, Crippa D, Garcia-Segura LM, Lauria G, Magnaghi V, Roglio I, Melcangi RC. Progesterone and its derivatives are neuroprotective agents in experimental diabetic neuropathy: a multimodal analysis. *Neuroscience*. 2007; 144:1293–1304. [PubMed: 17187935]
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 2001; 25:402–408. [PubMed: 11846609]
- Lohmann SM, Vaandrager AB, Smolenski A, Walter U, De Jonge HR. Distinct and specific functions of cGMP-dependent protein kinases. *Trends Biochem Sci*. 1997; 22:307–312. [PubMed: 9270304]
- Mendelsohn ME. Viagra: now mending hearts. *Nat Med*. 2005; 11:115–116. [PubMed: 15692588]
- Mousa SA, Zhang Q, Sitte N, Ji R, Stein C. beta-Endorphin-containing memory-cells and mu-opioid receptors undergo transport to peripheral inflamed tissue. *J Neuroimmunol*. 2001; 115:71–78. [PubMed: 11282156]
- Neeraj Arora DK, Veerareddy Srikar, Langston Will, Patel Rakesh, Kevil Christopher. Sildenafil enhances angiogenesis and blood flow in response to ischemia in Diabetic mice. *The FASEB Journal*. 2007; 21:7066. 2007.
- Paivalainen S, Nissinen M, Honkanen H, Lahti O, Kangas SM, Peltonen J, Peltonen S, Heape AM. Myelination in mouse dorsal root ganglion/Schwann cell cocultures. *Mol Cell Neurosci*. 2008; 37:568–578. [PubMed: 18206387]
- Patil CS, Singh VP, Singh S, Kulkarni SK. Modulatory effect of the PDE-5 inhibitor sildenafil in diabetic neuropathy. *Pharmacology*. 2004; 72:190–195. [PubMed: 15452368]
- Pfeifer A, Ruth P, Dostmann W, Sausbier M, Klatt P, Hofmann F. Structure and function of cGMP-dependent protein kinases. *Rev Physiol Biochem Pharmacol*. 1999; 135:105–149. [PubMed: 9932482]
- Pittenger GLRM, Burcus NI, McNulty P, Basta B, Vinik AI. Intraepidermal nerve fibers are indicators of small-fiber neuropathy in both diabetic and nondiabetic patients. *Diabetes Care*. 2004 Aug; 27(8):1974–1979. 2004. [PubMed: 15277426]
- Russell JW, Golovoy D, Vincent AM, Mahendru P, Olzmann JA, Mentzer A, Feldman EL. High glucose-induced oxidative stress and mitochondrial dysfunction in neurons. *Faseb J*. 2002; 16:1738–1748. [PubMed: 12409316]
- Sameni HR, Panahi M, Sarkaki A, Saki GH, Makvandi M. The neuroprotective effects of progesterone on experimental diabetic neuropathy in rats. *Pak J Biol Sci*. 2008; 11:1994–2000. [PubMed: 19266905]
- Scott JN, Clark AW, Zochodne DW. Neurofilament and tubulin gene expression in progressive experimental diabetes: failure of synthesis and export by sensory neurons. *Brain*. 1999; 122(Pt 11): 2109–2118. [PubMed: 10545396]
- Shirasaki N, Rosenbluth J. Structural abnormalities in freeze-fractured sciatic nerve fibres of diabetic mice. *J Neurocytol*. 1991; 20:573–584. [PubMed: 1919605]
- Smith AG, Howard JR, Kroll R, Ramachandran P, Hauer P, Singleton JR, McArthur J. The reliability of skin biopsy with measurement of intraepidermal nerve fiber density. *J Neurol Sci*. 2005; 228:65–69. [PubMed: 15607212]
- South SM, Smith MT. Apparent insensitivity of the hotplate latency test for detection of antinociception following intraperitoneal, intravenous or intracerebroventricular M6G administration to rats. *J Pharmacol Exp Ther*. 1998; 286:1326–1332. [PubMed: 9732396]
- Sullivan KA, Hayes JM, Wiggan TD, Backus C, Su Oh S, Lentz SI, Brosius F 3rd, Feldman EL. Mouse models of diabetic neuropathy. *Neurobiol Dis*. 2007; 28:276–285. [PubMed: 17804249]
- Thippeswamy T, McKay JS, Morris R, Quinn J, Wong LF, Murphy D. Glial-mediated neuroprotection: evidence for the protective role of the NO-cGMP pathway via neuron-glia communication in the peripheral nervous system. *Glia*. 2005; 49:197–210. [PubMed: 15390094]
- Tolwani RJ, Cosgaya JM, Varma S, Jacob R, Kuo LE, Shooter EM. BDNF overexpression produces a long-term increase in myelin formation in the peripheral nervous system. *J Neurosci Res*. 2004; 77:662–669. [PubMed: 15352212]
- Trojaborg W, Hays AP, van den Berg L, Younger DS, Latov N. Motor conduction parameters in neuropathies associated with anti-MAG antibodies and other types of demyelinating and axonal neuropathies. *Muscle Nerve*. 1995; 18:730–735. [PubMed: 7540258]

- Vanderah TW, Suenaga NM, Ossipov MH, Malan TP Jr, Lai J, Porreca F. Tonic descending facilitation from the rostral ventromedial medulla mediates opioid-induced abnormal pain and antinociceptive tolerance. *J Neurosci.* 2001; 21:279–286. [PubMed: 11150345]
- Vincent AM, McLean LL, Backus C, Feldman EL. Short-term hyperglycemia produces oxidative damage and apoptosis in neurons. *FASEB J.* 2005; 19:638–640. [PubMed: 15677696]
- Walker DK, Ackland MJ, James GC, Muirhead GJ, Rance DJ, Wastall P, Wright PA. Pharmacokinetics and metabolism of sildenafil in mouse, rat, rabbit, dog and man. *Xenobiotica.* 1999; 29:297–310. [PubMed: 10219969]
- Walwyn WM, Matsuka Y, Arai D, Bloom DC, Lam H, Tran C, Spigelman I, Maidment NT. HSV-1-mediated NGF delivery delays nociceptive deficits in a genetic model of diabetic neuropathy. *Exp Neurol.* 2006; 198:260–270. [PubMed: 16427624]
- Wang L, Gang Zhang Z, Lan Zhang R, Chopp M. Activation of the PI3-K/Akt pathway mediates cGMP enhanced-neurogenesis in the adult progenitor cells derived from the subventricular zone. *J Cereb Blood Flow Metab.* 2005; 25:1150–1158. [PubMed: 15815584]
- Wang L, Zhang ZG, Zhang RL, Gregg SR, Hozeska-Solgot A, LeTourneau Y, Wang Y, Chopp M. Matrix metalloproteinase 2 (MMP2) and MMP9 secreted by erythropoietin-activated endothelial cells promote neural progenitor cell migration. *J Neurosci.* 2006; 26:5996–6003. [PubMed: 16738242]
- Wang X, Robinson PJ. Cyclic GMP-dependent protein kinase and cellular signaling in the nervous system. *J Neurochem.* 1997; 68:443–456. [PubMed: 9003029]
- Wright DE, Johnson MS, Arnett MG, Smittkamp SE, Ryals JM. Selective changes in nociceptive behavior despite normal cutaneous axon innervation in leptin receptor-null mutant (db/db) mice. *J Peripher Nerv Syst.* 2007; 12:250–261. [PubMed: 18042135]
- Yagihashi S, Kamijo M, Watanabe K. Reduced myelinated fiber size correlates with loss of axonal neurofilaments in peripheral nerve of chronically streptozotocin diabetic rats. *Am J Pathol.* 1990; 136:1365–1373. [PubMed: 2141449]
- Yamauchi J, Chan JR, Shooter EM. Neurotrophins regulate Schwann cell migration by activating divergent signaling pathways dependent on Rho GTPases. *Proc Natl Acad Sci U S A.* 2004; 101:8774–8779. [PubMed: 15161978]
- Zhang JY, Luo XG, Xian CJ, Liu ZH, Zhou XF. Endogenous BDNF is required for myelination and regeneration of injured sciatic nerve in rodents. *Eur J Neurosci.* 2000; 12:4171–4180. [PubMed: 11122329]
- Zhang R, Zhang L, Zhang Z, Wang Y, Lu M, Lapointe M, Chopp M. A nitric oxide donor induces neurogenesis and reduces functional deficits after stroke in rats. *Ann Neurol.* 2001; 50:602–611. [PubMed: 11706966]
- Zhuo M, Hu Y, Schultz C, Kandel ER, Hawkins RD. Role of guanylyl cyclase and cGMP-dependent protein kinase in long-term potentiation. *Nature.* 1994; 368:635–639. [PubMed: 7908417]

Figure 1. Effect of sildenafil on neurological function in diabetic mice

Treatment of diabetic mice with sildenafil improves neurological function measured by MCV (A), SCV (B), Tail-flick testing (C), hot plate test (D) and tactile von Frey filaments test (E) at 28 and 56 days after treatment. * $P < 0.01$, ** $P < 0.001$ versus the non diabetic group. # $P < 0.05$, ## $P < 0.01$, ### $P < 0.001$ versus the diabetic group treated with saline. $n = 10$ /group. Western blot analysis of PDE5 levels in sciatic nerve tissue and β -actin as an internal control (F). * $P < 0.05$ versus the non-diabetic group. $n = 6$ /group.

Figure 2. Effect of sildenafil on the subcutaneous innervation in diabetic mice

Panels A to C show PGP 9.5 immunoreactive fibers (arrows) in the plantar skin from a representative non-diabetic mouse (A), diabetic mouse treated with saline (B) and diabetic mouse treated with sildenafil (10mg/kg, C) 56 days after treatment. Panel D is quantitative data. Bar = 50 μ m. * P < 0.05 versus the non-diabetic group, # P < 0.05 versus diabetic mice treated with Saline. n = 10/group.

Figure 3. Effect of sildenafil on MBP, NF-H and morphomeric change of myelinated sciatic nerves

Double immunofluorescent staining (A to I) shows MBP (A, C, D, F, G, and I, red) and NFH (B, C, E, F, H, and I, green) immunoreactive sciatic nerves of a representative non-diabetic mouse (A to C), diabetic mouse treated with saline (D to F), and diabetic mouse treated with sildenafil (10 mg/kg, G to I) 56 days after treatment. Panels J to L show semi-thin toluidine blue-stained cross sections of sciatic nerves from a representative non-diabetic mouse (J), diabetic mouse treated with saline (K), and diabetic mouse treated with sildenafil (10 mg/kg, L) 56 days after treatment. Panels M and N show quantitative data of the percentage of MBP (M) and NF-H (N) immunoreactive area. Bars in I = 50μm, L = 25μm.

* $P < 0.05$ versus the non-diabetic group, # $P < 0.05$ versus diabetic mice treated with Saline.
n=10/group.

Figure 4. Effect of sildenafil on Schwann cell proliferation in vivo

Double immunostaining (A to C) shows BrdU (green, arrow) and S100 (red) positive cells in sciatic nerve tissue on the longitudinal sections from a representative non-diabetic mouse (A), diabetic mouse treated with saline (B), and diabetic mouse treated with sildenafil (10 mg/kg, C) 56 days after treatment. Panel D shows quantitative data of the percentage of BrdU and S100 positive cells. Bar in C = 50 μ m. * $P < 0.05$ versus the non-diabetic group, # $P < 0.05$ versus diabetic mice treated with Saline. $n = 10$ /group.

Figure 5. Effect of sildenafil on primary Schwann cell proliferation and migration

Panels A through D show BrdU immunoreactive cells (red) in primary mouse Schwann cells cultured in normal glucose (NG, A), high glucose (HG, B), sildenafil (sil, 150 nM) (C) and sildenafil (150 nM) with KT5823 (KT, D). Panel E shows quantitative data of the percentage of BrdU immunoreactive cells. Panel F shows quantitative data of the percentage of migration cells assayed by a modified Boyden chamber. Bar in D = 100 μ m. *P < 0.05 and #P < 0.05 versus the normal glucose (NG) and sildenafil (150 nM) group, respectively. n = 6/group. Panels G and H show Western blot analysis of PDE5 and PKGI in primary mouse Schwann cells cultured in normal glucose (NG), high glucose (HG) and sildenafil (+sil, 150 nM). Panels I and J show real-time RT-PCR and Western blot, respectively, analyses of

PDE5 in primary mouse Schwann cells cultured in high glucose (HG), Schwann cells transfected with siRNA-PDE5 (+si) or scrambled sequences (+ss) in high glucose condition. Panel K and L show real-time RT-PCR and Western blot, respectively, analysis of PDE5 in primary human Schwann cell cultured in normal glucose (NG) and high glucose (HG) conditions. GAPDH and β -actin were used as internal controls for mRNA and proteins, respectively. * $P < 0.05$ and # $P < 0.05$ versus the normal glucose (NG) and high glucose (HG) group, respectively. $n = 6/\text{group}$.

Figure 6. Effect of sildenafil on PKG1 and BDNF

Double immunostaining (A to D) shows PKGI (A and D, green, arrow) and S100 (B and D, red, arrow) immunoreactive cells in sciatic nerve tissue on the longitudinal section. Panels E to G show BDNF immunoreactivity in the cross sections of sciatic nerves of a representative non-diabetic mouse (E), diabetic mouse treated with saline (F), and diabetic mouse treated with sildenafil (10 mg/kg, G) 56 days after treatment. Panel H shows quantitative data of the percentage of BDNF immunoreactive area within the cross sections. Bars in D and G = 50µm. *P<0.05 and #P<0.05 versus the non-diabetic and diabetic mice treated with saline, respectively. n=10/group. Panels I and K show quantitative data of BDNF mRNA levels (I) and Western blot analysis (J and K) of BDNF in immortalized Schwann cells cultured in normal glucose (NG), high glucose (HG), high glucose with sildenafil, sildenafil (150 nM) with KT5823 (+KT) and Br-cGMP (cGMP, 10 µmol/L). GAPDH and β-actin were used as internal control for real-time RT-PCR and Western blot analysis, respectively. *P<0.05 and #P<0.05 versus the normal glucose (NG) and sildenafil (150 nM) group, respectively. n=6/group.

Table 1

Effect of sildenafil on blood glucose and bodyweight

Groups	Glucose, mg/dl		Weight, g	
	0d	56d	0d	56d
Non-diabetic	139.4±10.3 *	164.2±7.9 *	29.5±0.4 *	32.3±0.3 *
Diabetic+saline	513.8±13.9	538.4±22.6	55.7±1.7	59.4±3.1
+sildenafil (2mg/kg)	521.9±19.3	542.4±19.9	54.3±1.0	54.7±2.2
+sildenafil(10mg/kg)	539.5±17.1	566.2±15.2	58.1±0.6	57.4±0.9

Values are mean±SE.

* P<0.01 versus Diabetic+Saline group. n=10/group. 0d and 56d indicate that the day before and after the treatment, respectively.

Table 2

Effect of sildenafil on histomorphometric parameter of sciatic nerves

Property	Non-diabetic	Diabetic		
		+Saline	+sildenafil (2mg/kg)	+sildenafil (10mg/kg)
Fiber diameter(μm)	8.76 $\pm$ 0.08 *	7.21 $\pm$ 0.03	8.30 $\pm$ 0.08 *	8.25 $\pm$ 0.05 *
Axon diameter(μm)	6.09 $\pm$ 0.05 *	5.44 $\pm$ 0.01	5.95 $\pm$ 0.05 *	6.02 $\pm$ 0.03 *
Myelin thickness(μm)	1.32 $\pm$ 0.04 *	0.88 $\pm$ 0.01	1.18 $\pm$ 0.03 *	1.12 $\pm$ 0.02 *
Fiber area(μm^2)	350.1 $\pm$ 10.1 *	247.1 $\pm$ 8.46	344.4 $\pm$ 12.1 *	357.21 $\pm$ 7.5 *
Axon area(μm^2)	269.4 $\pm$ 6.97 *	183.1 $\pm$ 6.61	251.2 $\pm$ 7.96 *	246.8 $\pm$ 6.44 *
G ratio	0.69 $\pm$ 0.006 *	0.76 $\pm$ 0.002	0.71 $\pm$ 0.005 *	0.72 $\pm$ 0.005 *

Values are mean $\pm$ SE.*
p<0.01 vs. Diabetic+Saline group. n=10/group