Published in final edited form as:

J Psychiatr Res. 2012 August ; 46(8): 1036–1044. doi:10.1016/j.jpsychires.2012.04.015.

Sleep Promotes Consolidation and Generalization of Extinction Learning in Simulated Exposure Therapy for Spider Fear

Edward F. Pace-Schott, Ph.D.^{1,2}, Patrick Verga¹, Tobias Bennet¹, and Rebecca M.C. Spencer, Ph.D.¹

¹Department of Psychology and Neuroscience, University of Massachusetts, Amherst, MA

²Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, MA

Abstract

Simulated exposure therapy for spider phobia served as a clinically naturalistic model to study effects of sleep on extinction. Spider-fearing, young adult women (N=66), instrumented for skin conductance response (SCR), heart rate acceleration (HRA) and corrugator electromyography (EMG), viewed 14 identical 1-min videos of a behaving spider before a 12-hr delay containing a normal night's Sleep (N=20) or continuous daytime Wake (N=23), or a 2-hr delay of continuous wake in the Morning (N=11) or Evening (N=12). Following the delay, all groups viewed this same video 6 times followed by six 1-min videos of a novel spider. After each video, participants rated disgust, fearfulness and unpleasantness. In all 4 groups, all measures except corrugator EMG diminished across Session 1 (extinction learning) and, excepting SCR to a sudden noise, increased from the old to novel spider in Session 2. In Wake only, summed subjective ratings and SCR to the old spider significantly increased across the delay (extinction loss) and were greater for the novel vs. the old spider when it was equally novel at the beginning of Session 1 (sensitization). In Sleep only, SCR to a sudden noise decreased across the inter-session delay (extinction augmentation) and, along with HRA, was lower to the novel spider than initially to the old spider in Session 1 (extinction generalization). None of the above differentiated Morning and Evening groups suggesting that intervening sleep, rather than time-of-testing, produced differences between Sleep and Wake. Thus, sleep following exposure therapy may promote retention and generalization of extinction learning.

Keywords

Sleep; Extinction; Spider Phobia; Fear Conditioning; Exposure Therapy

Introduction

Abnormal expression of fear, as occurs in anxiety disorders such as post-traumatic stress disorder (PTSD) and specific phobia, may result from abnormally strong fear conditioning

© 2012 Elsevier Ltd. All rights reserved

Corresponding Author: Edward F. Pace-Schott, Ph.D., Department of Psychology, University of Massachusetts, Amherst, Tobin Hall, Rm. 522, 135 Hicks Way, Amherst, MA 01003, Phone: 508-523-4288, Fax: 413-545-0996, epacesch@psych.umass.edu.

Financial Disclosure: None of the authors report any potential conflicts of interest

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

(Armfield, 2006; Lissek et al., 2005; Mineka & Oehlberg, 2008; Orr et al., 2000), deficiency of inhibitory mechanisms that normally moderate fear expression (Craske et al., 2008; Hofmann, 2008; Milad, Rauch, Pitman, & Quirk, 2006), or both. Key among such inhibitory processes is extinction — learning that a once-feared object or event is no longer dangerous (Milad et al., 2006). Rather than erasing a fearful memory, extinction forms a new “safety memory” that competes with the fear memory when the once-feared object or event is re-encountered (Hermans, Craske, Mineka, & Lovibond, 2006; Quirk & Mueller, 2008).

Formation of such therapeutic extinction memories is the neurocognitive basis for the efficacy of exposure therapy, a first-line behavioral treatment for anxiety disorders (Craske et al., 2008; McNally, 2007). In order for exposure therapy to be successful, consolidation and retention of extinction learning acquired during therapy is essential (Craske et al., 2008). In addition, such learning must generalize in order to ensure that the reduction of fearful responding to specific cues in treatment will extend to stimuli encountered outside the therapist’s office (Rowe & Craske, 1998; Vansteenwegen et al., 2007).

Using an experimental fear-conditioning paradigm (Milad et al., 2007), normal sleep has been shown to promote the generalization of extinction memories (Pace-Schott et al., 2009). However, unlike such experimentally induced *de-novo* fears, anxiety disorders are associated with long-standing fears that have complex, multi-factorial origins and perpetuating factors (Armfield, 2006). Specific phobias, such as spider phobia, are highly prevalent, mild anxiety disorders (LeBeau et al., 2010; Lissek et al., 2007) in which treatment strategies, such as exposure therapy, can be studied in a non-clinical setting (e.g., Vansteenwegen et al., 2007).

Sleep enhances consolidation of emotional memory (reviewed in Walker, 2009). Here we characterize the effect of sleep on the retention and generalization of a specific emotional memory-- the extinction of spider fear produced by simulated exposure therapy. We hypothesized that sleep following simulated exposure therapy in spider-phobic subjects would lead to greater retention of fear extinction for the spider to which they were repeatedly exposed. In addition, we predicted that sleep would enhance generalization of this extinction memory to a novel spider.

Methods and Materials

Participants

Participants were 66 females (18–28 yrs, mean=19.9) with significant fear of spiders operationally defined using thresholds of 80 on the Fear of Spiders Questionnaire [FSQ (Szymanski & O’Donohue, 1995)] and 15 on the Spider Phobia Questionnaire [SPQ (Klorman, Hastings, Weerts, Melamed, & Lang, 1974)]. Previous research has determined that these thresholds reflect significant fear of spiders (Muris and Merckelbach, 1996, Rodriguez et al., 1999, Guastella et al., 2007, Vansteenwegen et al., 2007). The FSQ was included in a large pre-screening for research participation by students at the University of Massachusetts, Amherst. Starting with the highest FSQ scores and working downward, such individuals were offered the opportunity to earn academic credit for participation if they confirmed eligibility via the SPQ.

Qualified respondents were pseudo-randomly assigned to Sleep (N=20) and Wake (N=23) experimental groups as well as Morning (N=11) and Evening (N=12) control groups. Invitations specified that participants must be non-smoking, without psychiatric, sleep, medical or neurological disorders and not using psychiatric or sleep-affecting drugs. A 23-item screening questionnaire administered at the first session queried these criteria. Only 12 individuals had one or more deviations from the invitation criteria (Table 1). However,

because deviations were distributed between the groups and because co-morbidities are common in specific phobias (American Psychiatric Association, 2000), these individuals were included in analyses (see Supplementary Methods for additional details). Physiological data could not be analyzed in 1 Wake, 2 Sleep and 1 Evening participants leaving N=18 (Sleep), 22 (Wake), 11 (Morning) and 11 (Evening) for the physiological measures. This study was approved by the University of Massachusetts, Amherst, IRB and all participants provided written informed consent.

Procedure

Participants completed two sessions (Figure 1) from approximately 8:00–9:00PM and 8:00–9:00AM the following morning (Sleep), 8:00–9:00AM and 8:00–9:00PM on a single day (Wake), 7:00–8:00 and 10:00–11:00AM (Morning) or 7:00–8:00 and 10:00–11:00PM (Evening). Procedures for the experimental groups (Sleep and Wake) and those for the control groups (Morning and Evening) were identical, except for the duration of the inter-session interval (12 hr experimental vs. 2 hours control). All participants were instructed to abstain from alcohol, recreational drugs and daytime napping from the day before Session 1 (S1) until completing Session 2 (S2). At S1 (Sleep and Wake) or immediately following S1 (Morning and Evening), participants completed a sleep diary that retrospectively queried sleep duration and quality on the 2 preceding nights. The Sleep group also completed this diary for the night between sessions. On the night before S1 (all groups) and between S1 and S2 (Sleep group), participants were instructed to allow themselves the opportunity for at least 7 hours sleep and to have no caffeine after arising until the end of S2. Wake, Morning and Evening groups were specifically instructed to remain continuously awake between sessions. Between S1 and S2, all participants completed the Epworth Sleepiness Scale [ESS (Johns, 1994)], the Pittsburgh Sleep Quality Index [PSQI (Buysse, Reynolds, Monk, Berman, & Kupfer, 1989)], the Morningness-Eveningness questionnaire [MEQ (Horne & Ostberg, 1976)], the Revised NEO Personality Inventory [NEO-PI-R (Costa & McCrae, 1992)], the Disgust Propensity and Sensitivity Scale [DPSS-R (van Overveld, de Jong, Peters, Cavanagh, & Davey, 2006)] and the STAI-Trait version [STAI-T (Spielberger, Gorsuch, & Lushene, 1990)].

Participants completed the STAI State version (STAI-S) and the Stanford Sleepiness Scale [SSS (Hoddes, Zarcone, Smythe, Phillips, & Dement, 1973)] at the beginning and end of both sessions. They were instructed to imagine themselves “being there with the spider” in the video and told that they “should not resist their fear” and that the “fear will go away by itself” (Vansteenwegen et al., 2007). Participants sat approximately 2 feet from a 17-inch computer screen. Following each video, a screen instructed participants to rate how disgusting, fearful and unpleasant they found the spider using three 11-point rating scales with maxima (+10), minima (–10) and midpoint (0) indicated and intervening increments of 2 indicated by bullets. The order of scales was counterbalanced across subjects. After 20 sec, another screen briefly appeared instructing participants to await the next video followed by a 10 sec inter-stimulus interval (ISI) after which the next video began. During S1, participants viewed 14 identical videos of a spider and, during S2, participants viewed six videos of this same, “old” spider followed by six videos of a novel spider.

Stimuli

Three *YouTube* videos of individual spiders (see Supplementary Methods) were edited to remove sound, achieve a standard 60-sec length and optimize homogeneity of background. Each subject viewed two of these videos, one as the old spider and a second as the novel spider. The three videos were counterbalanced across participants. Three videos were used in order to reduce the effect of systematic differences in fearfulness between any specific pair of videos thereby increasing the generalizability of findings. In other words, to increase

importance of the experimental assignment (old vs. novel) over any inherent differences between any specific pair.

To probe evoked sympathetic reactivity, during some videos and ISIs, a 10-msec, 83dB white noise stimulus was delivered through headphones. During S1, this noise stimulus occurred during 9 of 14 videos and 5 of 14 ISI's. During S2, the noise stimulus occurred during 4 of 6 videos and 4 of 6 ISI's each for the old and novel spider. For habituation, each session began with 10 unpredictable repetitions of this noise stimulus. The noise stimulus occurred only during the last 30 sec of a video with its latency pseudo-randomly varying between 5–26 sec into these final 30 sec. Similarly, the noise stimulus occurred with a latency of 2–8 sec into the 10-sec ISI. Noise latencies were identical for all subjects.

Psychophysiological measurements

Skin conductance response (SCR), electrocardiography (ECG) and electromyography (EMG) were recorded using the MP150 data acquisition unit (BIOPAC Systems, Inc., Goleta, CA) and BIOPAC AcqKnowledge 3.9.2 software for the Macintosh (see Supplementary Methods for hardware details). One event marker indicated the beginning of each video and another the onset of the noise stimulus allowing precise synchronization of each stimulus onset with ongoing physiological recording. Sampling rate was 2000 Hz.

Skin conductance level (SCL), a reliable index of sympathetic activation (Dawson, Schell, & Filion, 2007), was recorded using disposable adhesive sensors, separated by 10 mm, attached to the hypothenar surface of the non-dominant hand. Two measures of SCR were computed. First, video-related skin conductance change (Video SCR) was computed by subtracting the mean SCL in microSiemens (μ S) during the first 1 sec of the video, the typical SCR latency (Dawson et al., 2007), from the maximum SCL during the subsequent 29 sec (i.e., before any of the noise-stimuli occurred). Second, noise-related SCR (Noise SCR) was computed by subtracting the mean SCL during the 2 sec preceding the noise-stimulus onset from the maximum SCL during sec 2–6 following stimulus onset during videos (Noise-SCR-in-Video) or during ISIs (Noise-SCR-in-ISI). SCRs were square-root transformed and, if the untransformed SCR was negative, the negative sign was retained after calculating the square root of the SCR's absolute value (Orr et al., 2000).

ECG electrodes were attached to the torso over the first intercostal space (right) and below the lowest rib (left). Heart-Rate acceleration (HRA) was defined as the maximum beats per minute (bpm) achieved during the first 30 sec of each video (i.e., before any noise stimulus) minus the mean of the initial 2 seconds of the preceding 10-sec ISI (again before any noise stimulus). A correction for extreme outlying values was applied (see Supplementary Methods).

Corrugator supercilli EMG is a reliable index of both overt and covert negative emotion (Bradley & Lang, 2007). Electrode attachment followed Fridlund & Cacioppo (1986). In the majority of participants, impedance was confirmed to be below 10 KOhm (see Supplementary Methods). EMG was integrated over a 250-msec time constant and response to each video was calculated by subtracting the mean signal amplitude during the initial 2 seconds of the 10-sec ISI from the maximum during the first 30 sec of the succeeding video.

Statistical Analyses

A single self-report outcome measure, Composite Negative Ratings, was calculated by summing each participants' –10 to +10 ratings of disgust, fearfulness and unpleasantness for each video.

Data were analyzed in four phases (Figure 1). “Early Exposure” contained responses during videos (“Trials”) 1–6 of S1; “Late Exposure,” Trials 7–12 of S1; “Old Spider,” Trials 1–6 of S2; and “Novel Spider,” Trials 7–12 of S2. Although 14 trials were presented in S1 in order to maximally promote extinction learning (Vansteenwegen et al., 2007), videos 13 and 14 were excluded from analyses to balance the nested analysis of variance (ANOVA) design described below.

Composite Negative Ratings, corrugator EMG, HRA and Video SCR were analyzed using 3-factor mixed ANOVA with one between-subjects variable, Group, and 2 within-subject variables, 6 Trials nested in 4 Phases (fewer trials for Noise SCR described in Supplementary Methods). Because the inter-session duration differed between experimental and control groups, the Sleep vs. Wake groups and Morning vs. Evening groups were separately compared, however, their phase definitions and analyses were identical.

Figure 1 illustrates operational definitions of emotional memory processes that were measured for each outcome variable. First, “Extinction Learning” was analyzed in S1 data by 2 separate methods using mixed ANOVA and the between-subjects variable, Group (see Supplementary Methods). Next, each outcome variable was analyzed across all 4 phases using 3-factor mixed ANOVA (see above). Each group was then analyzed separately by repeated measures ANOVA and, when there was a significant main effect of Phase, the remaining 3 emotional memory processes were evaluated using post-hoc means comparisons between the 4 Phases. “Extinction Retention” was defined as the maintenance of lowered responding at Old Spider following the inter-session delay. “Response Return” was defined as the increase in responding at Novel Spider relative to Old Spider. “Sensitization” was defined as the degree to which responses at Novel Spider exceeded those to the old spider at Early Exposure during S1 when it was equally “novel.” For both Response Return and Sensitization, change in the opposite direction represented “Generalization” of extinction learning.

Of specific interest were instances in which a significant Group x Phase interaction indicated group differences between one or more of the above processes. When such an interaction occurred, the above, within-group post-hoc comparisons of Phase were used to determine which specific processes differed between groups (e.g., occurred in one group but not the other). Such Group x Phase interactions, however, also reflected group differences in response magnitude at specific phases, differences that, in turn, were determined, in part, by individual differences in baseline reactivity. Additionally, the absolute magnitude of responses at baseline (S1) could potentially influence the ability to retain extinction across the delay. Therefore, the above 3-factor mixed ANOVA for each outcome measure was repeated with the inclusion of each subject’s maximum S1 response as a covariate. Similarly, S1 differences in Extinction Learning might influence subsequent extinction memory processes. Therefore, when Group main effects or Group x Phase interactions occurred at S1, ANOVA was repeated across all 4 phases using only those individuals showing Absolute Extinction Learning (i.e., Early Extinction - Late Extinction > 0).

The Greenhouse-Geisser correction was applied to all within-subject main effects and their interactions. To examine outcome variables without individuals deviating from advertised inclusion criteria (see Table 1), analyses were repeated excluding these individuals in a stepwise manner (see Supplementary Methods).

One-way ANOVA, followed, when significant, by Bonferroni-Dunn post-hoc tests, compared groups’ questionnaire (e.g., FSC) results. Mixed ANOVA compared repeated measure questionnaires (e.g., SSS) between groups.

Results

Comparison of group characteristics

As shown in Table 1 and Supplementary Results, the four groups showed highly similar spider fear, habitual sleep and psychological trait measures. All groups averaged above 100 on the FSQ and 20 on the SPQ (Table 1) indicative of a highly spider-fearful sample (Muris and Merckelbach, 1996). Notably, when Group main effects were present (NEO-PI-R Extraversion, sleep duration on night before S1), this reflected one control group differing from the remaining groups rather than differences between Sleep and Wake groups (Table 1, Supplementary Results). Group similarity also extended to sleepiness (SSS) and state anxiety (STAI-S) measured at the beginning and end of each session (Supplementary Results). Collapsing across groups, significantly greater STAI-S at the end vs. beginning of sessions [$F(1,61)=74.59$, $p<.0001$] confirmed the impact of videos on participants' state anxiety.

Extinction learning, retention and generalization

Comparisons across both S1 and S2 (e.g., Extinction Retention) are detailed below for each outcome measure. However, Extinction Learning within S1 itself is summarized below with details provided in Supplementary Results.

Rating scales

Session 1: Across S1, significant Extinction Learning for Composite Negative Ratings was evident in both the experimental (Sleep and Wake) and control (Morning and Evening) groups. However, there was no S1 Group main effect or Group interactions with Phase or Trial (Fig. 2A, Supplementary Results).

Experimental groups: Across both Sessions, comparing Sleep and Wake groups for Composite Negative Ratings, there was a significant Group x Phase interaction [$F(3,123)=3.43$, $p=.026$] that remained significant when maximum S1 Composite Negative Rating was included as a covariate [$F(3,120)=2.99$, $p=.044$]. Main effects of Phase occurred in both Sleep [$F(3,57)=6.23$, $p=.004$] and Wake [$F(3,66)=14.45$, $p=.0001$] groups. Both groups showed significant Response Return during S2 with ratings during Novel Spider significantly greater than during Old Spider (Sleep: $F=8.47$, $p=.012$; Wake: $F=18.38$, $p=.0004$, Fig. 2A). However, only in the Wake group was Old Spider at S2 rated more negatively than at Late Exposure during S1 ($F=4.35$, $p=.055$, Fig. 2A). Similarly, only in the Wake group was Novel Spider in S2 rated more negatively than was the old spider at Early Exposure during S1 when it was equally novel ($F=7.23$, $p=.017$, Fig. 2A). Therefore, for Composite Negative Ratings, only in the Wake group was there incomplete Extinction Retention at Old Spider and significant Sensitization at Novel Spider.

Control group: Comparing Composite Negative Ratings in the Morning and Evening (control) groups across both sessions, the Group x Phase interaction that was observed when comparing Sleep and Wake groups was absent [$F(3,63)=0.38$, $p=.74$]. Both control groups showed significant Response Return as well as full Extinction Retention and no Sensitization (Fig. 2B).

Physiological measures

Video SCR

Session 1: Across S1, significant Extinction Learning in Video SCR was evident in both Sleep and Wake groups (Fig. 3A, Supplementary Results). However, Extinction Learning

was more pronounced in the Sleep relative to the Wake group as evidenced by a Group x Phase interaction trend [$F(1,38)=3.48$, $p=.07$].

Experimental groups: Across both Sessions, comparing Sleep and Wake groups for Video SCR (Fig. 3A), there was a significant Group x Phase interaction [$F(3,114)=3.16$, $p=.035$] that remained significant when maximum S1 Video SCR was included as a covariate [$F(3,111)=3.20$, $p=.034$]. (Further analyses of the effects of the S1 Group x Phase interaction on S2 performance for Video SCR are provided in Supplementary Results.) Main effects of Phase occurred in both Sleep [$F(3,51)=12.57$, $p=.0001$] and Wake [$F(3,63)=6.68$, $p=.002$] groups. Response Return during S2 was significant in the Sleep group ($F=12.53$, $p=.001$) and a trend in the Wake group ($F=3.03$, $p=.095$). However, only in the Wake group was Video SCR to Old Spider during S2 greater than at Late Exposure during S1 ($F=6.97$, $p=.016$). This loss of extinction memory for the old spider across the inter-session interval in the Wake group may account for this group's lesser Response Return (i.e., the S2 old vs. novel comparison) because responding was already elevated to the Old Spider when Novel Spider was introduced. Notably, despite a lesser increase relative to Old Spider, only in the Wake group was Video SCR to Novel Spider in S2 greater than it was to the old spider during S1 at Early Exposure when it was equally novel ($F=7.08$, $p=.015$). Therefore, for Video SCR, only in the Wake group was there incomplete Extinction Retention and significant Sensitization.

Control groups: In Morning and Evening groups (Fig. 3B), there was no Group x Phase interaction for Video SCR across both sessions ($p=.52$).

Noise SCR

Session 1: Across S1, Extinction Learning, as measured by Noise-SCR-in-Video, occurred in all 4 groups (Fig. 4A, C). In the experimental groups, S1 Group main effect and Group x Phase interaction trends also occurred (Supplementary Results).

Experimental groups: Across both sessions, comparing Noise-SCR-in-Video for Sleep and Wake groups, there was a significant Group x Phase interaction ($F[3,114]=6.00$, $p=.002$). Although this interaction was due, in part, to the above baseline Group differences, when each subject's maximum S1 Noise-SCR-in-Video was added as a covariate to the mixed ANOVA model, the Group x Phase interaction remained significant ($F[3,114]=3.87$, $p=.017$). (Two further analyses of the effects of these S1 group differences on S2 Noise-SCR-in-Video are provided in Supplementary Results.) Decomposing this interaction, the Sleep group showed a main effect of Phase [$F(3,51)=18.73$, $p<.0001$] but this was absent in the Wake group ($p=.38$). In the Sleep group, there was a trend for Noise-SCR-in-Video at Old Spider to further decrease relative to Late Exposure ($F=4.11$, $p=.056$, Fig. 4A). Additionally, at Novel Spider, Noise-SCR-in-Video was significantly lower than at Early Exposure when the old spider was equally novel ($F=46.65$, $p<.0001$, Fig. 4A).

Therefore, among experimental participants, only the Sleep group showed further augmentation of Extinction Retention for Noise-SCR-in-Video across the inter-session delay as well as "negative" Sensitization (i.e., Generalization) (Fig. 4A). That observed differences between Sleep and Wake groups were specific to noises presented during spider videos was indicated by the lack of such group effects for Noise-SCR-in-ISI (Fig. 4B, see Supplementary Results).

Control groups: Analysis of Morning and Evening groups across both sessions showed no Group x Phase interaction for Noise-SCR-in-Video ($p=.72$, Fig. 4C) or Noise-SCR-in-ISI ($p=.49$, Fig. 4D).

Heart Rate Acceleration

Session 1: Across S1, significant Extinction Learning for HRA was evident in experimental and control groups (Fig. 5A,B, Supplementary Materials). However, there was no S1 Group main effect or Group interactions with Phase or Trial.

Experimental groups: Across both sessions, comparing HRA in Sleep and Wake groups, there was a significant Group x Phase interaction [$F(3,111)=3.42$, $p=.025$] that remained significant when maximum S1 Video SCR was included as a covariate [$F(3,108)=2.96$, $p=.042$]. There was a main effect of Phase in the Sleep [$F(3,51)=4.89$, $p=.007$] but not the Wake ($p=.39$) group. In the Sleep group, HRA at Novel Spider was significantly lower than at Early Exposure when the old spider was equally novel ($F=7.74$, $p=.01$, Fig 5A). Therefore, for HRA in the Sleep group, there was “negative” Sensitization (i.e., Generalization) whereas, in the Wake group, phases did not significantly differ.

Control groups: In Morning and Evening groups, there was no Group x Phase interaction for HRA across both sessions ($p=.78$; Fig 5B).

Corrugator EMG—Across S1, no significant Extinction Learning for corrugator EMG was evident in experimental or control groups (Figure 6A,B, Supplementary Results). Across both sessions, neither the experimental nor control groups showed a Group x Phase interaction ($p=.75$ and $.57$ respectively) or a Group main effect ($p=.39$ and $.22$). However, both the experimental and control groups showed Response Return and Sensitization (Fig. 6A,B, Supplementary Results).

Discussion

In spider-fearing young adult women, simulated exposure therapy acutely diminished both subjective and physiological measures of negative affect. Following a 12-hr delay, those exposed in the evening, who then slept, showed better extinction retention and generalization when tested in the morning than those exposed in the morning who remained awake until tested in the evening. The majority of findings remained unchanged after exclusion of participants with potentially confounding factors (see Supplementary Results) and results from control groups are inconsistent with a circadian explanation of experimental group differences (see below).

Circadian vs. sleep-dependent processes

Because, among control groups, those both exposed and tested in the morning did not differ from those exposed and tested in the evening, intervening sleep rather than circadian effects better explains the differences between Sleep and Wake groups seen in 4 of 5 outcome variables. Nonetheless, certain circadian effects were evident. Most notably, at S1, the evening-exposed groups displayed greater Noise-SCR-in-Video than the morning-exposed groups (see Supplementary Results)—a diurnal pattern of electrodermal reactivity that has previously been described for SCR to emotional images (Hot, Leconte, & Sequeira, 2005).

We chose Morning and Evening groups as circadian controls for two main reasons. First, when measuring physiological parameters as expression of emotional memory, major concerns are possible circadian effects upon these same parameters independent of memory processes (as noted for SCR). Because we define emotional memory processes by intra-group changes in each outcome variable (see Methods), the control groups tested whether time-of-day might influence changes in outcome variables from one phase to another despite having maintained the same behavioral state (wake) across the delay. Second, we ruled out use of an alternative 24-hr circadian control (i.e., AM to AM and/or PM to PM) because

work in animals has shown that extinction memory consolidation, pharmacologically blocked for 24 hours, can consolidate across the subsequent 24 hours (Santini et al. 2001). Therefore, in the current study, 24-hour controls would confound the duration of wake prior to sleep with the effects of sleep itself.

Rather than circadian effects, we hypothesize that sleep-dependent processes allow new emotional learning, such as extinction, to influence pre-existing memories. A network of limbic and paralimbic, fear-related anatomic regions have repeatedly been shown to be activated by *de-novo* fear conditioning (Alvarez, Biggs, Chen, Pine, & Grillon, 2008; LaBar, Gatenby, Gore, LeDoux, & Phelps, 1998), extinction learning for such conditioning (Knight, Smith, Cheng, Stein, & Helmstetter, 2004; Phelps, Delgado, Nearing, & LeDoux, 2004) and recall of memory for that extinction (Kalisch et al., 2006; Milad et al., 2007). A recent meta-analysis has shown greater amygdala and insula activity during emotional processing tasks in individuals with PTSD, Social Phobia and Specific Phobia compared with controls, suggesting underlying similarity in neural circuits affected by these anxiety disorders (Etkin & Wager, 2007). The amygdala and insula are components of an “anterior paralimbic REM activation area” that selectively reactivates during REM (Nofzinger et al., 2004) providing opportunity during sleep for new emotional learning, such as extinction, to be integrated with preexisting fear-related memory.

Extinction and habituation

Habituation and extinction, are overlapping forms of learning (McSweeney & Swindell, 2002) and its underlying neuroplasticity (Storvse, McNally, & Richardson 2010). However, whereas habituation is a non-associative learning process whereby behavioral and physiological responses during initial exposure to a stimulus diminish with its repeated presentation (Grissom and Bhatnagar, 2009; Leussis and Bolivar, 2006), extinction constitutes new associative learning whereby a stimulus previously associated with danger also becomes associated with the absence of danger (Hermans et al. 2006). During exposure therapy, habituation learning occurs concurrently with extinction. For example, in the current study, *intra*-session habituation undoubtedly contributed to diminishing responses across S1 as well as, possibly, *inter*-session habituation to such diminution in SCR across the delay (see Pace-Schott et al., 2011). However, extinction learning is a key contributor to reduction of fear in exposure therapy (Hermans et al., 2006; McNally, 2007; Milad et al., 2006) and may be more important than habituation for the therapeutic efficacy of this treatment (Craske et al., 2008). Similarly, although declarative and vicarious learning may have contributed to participants’ original spider fear, we assume that fear conditioning also played a role in its origin (Armfield, 2006; Fyer, 1998; Mineka et al., 2008; Rachman, 2002). Other emotional processes that may have contributed to inter-individual variability in subjective and objective responses are considered in Supplementary Discussion.

Clinical significance

Sleep-dependent generalization of extinction memories (Pace-Schott et al., 2009) suggests that conducting exposure therapy sessions in temporal proximity to sleep may enhance its efficacy. The current findings suggest an additional mechanism by which sleep might enhance exposure treatment, protection from sensitization. Sensitization-like processes play a role in the perpetuation of anxiety disorders. One such process is fear generalization, in which fear of a conditioned stimulus specifically associated with an inherently aversive stimulus comes to be elicited by stimuli that resemble but are not identical to the original conditioned stimulus (Lissek et al., 2008). A second such process is secondary conditioning whereby fear of a previously neutral stimulus is established by its association with a stimulus that was itself associated with the original, inherently aversive stimulus or event.

Increased secondary conditioning has been demonstrated both in PTSD (Wessa & Flor, 2007) and in spider phobia (Schweckendiek et al., 2011).

Conclusions

A period of sleep following simulated exposure therapy resulted in better retention and generalization of subjective and physiological responses to phobic stimuli. Sleep may confer protection from sensitization to such stimuli. Findings encourage further clinical investigation into the timing of exposure therapy in relation to sleep.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Scott Orr, Ph.D. scientific and technical advice, Philip Cusimano for technical advice, and Shanzay Haider, Shannon McKeon, Joanna Hong, Jessica Green and Lana Kim for research assistance. Research supported by NIA R00AG029710 and NIMH R21MH090357.

References

- Alvarez RP, Biggs A, Chen G, Pine DS, Grillon C. Contextual fear conditioning in humans: cortical-hippocampal and amygdala contributions. *Journal of Neuroscience*. 2008; 28:6211–6219. [PubMed: 18550763]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR*. 4. Washington, DC: American Psychiatric Association; 2000. (Text Revision)
- Armfield JM. Cognitive vulnerability: a model of the etiology of fear. *Clinical Psychology Review*. 2006; 26:746–768. [PubMed: 16806621]
- Bradley, MM.; Lang, PJ. Emotion and motivation. In: Cacioppo, JT.; Tassinary, LG.; Berntson, GG., editors. *Handbook of Psychophysiology*. 3. Cambridge, UK: Cambridge University Press; 2007.
- Buyse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research*. 1989; 28:193–213. [PubMed: 2748771]
- Costa, PT.; McCrae, RR. *Revised NEO Personality Inventory (NEO-PI-R) and NEO five-factor inventory (NEO-FFI) Professional Manual*. Odessa FL: Psychological Assessment Resources, Inc; 1992.
- Craske MG, Kircanski K, Zelikowsky M, Mystkowski J, Chowdhury N, Baker A. Optimizing inhibitory learning during exposure therapy. *Behaviour Research and Therapy*. 2008; 46:5–27. [PubMed: 18005936]
- Dawson, ME.; Schell, AM.; Fillion, DL. The electrodermal system. In: Cacioppo, JT.; Tassinary, LG.; Berntson, GG., editors. *Handbook of Psychophysiology*. 3. New York: Cambridge University Press; 2007. p. 159-181.
- Etkin A, Wager TD. Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*. 2007; 164:1476–1488. [PubMed: 17898336]
- Fridlund AJ, Cacioppo JT. Guidelines for human electromyographic research. *Psychophysiology*. 1986; 23:567–589. [PubMed: 3809364]
- Fyer AJ. Current approaches to etiology and pathophysiology of specific phobia. *Biological Psychiatry*. 1998; 44:1295–1304. [PubMed: 9861472]
- Grissom N, Bhatnagar S. Habituation to repeated stress: get used to it. *Neurobiology of Learning and Memory*. 92:215–224. [PubMed: 18667167]
- Guastella AJ, Dadds MR, Lovibond PF, Mitchell P, Richardson R. A randomized controlled trial of the effect of D-cycloserine on exposure therapy for spider fear. *J Psychiatr Res*. 2007; 41:466–471. [PubMed: 16828803]

- Hermans D, Craske MG, Mineka S, Lovibond PF. Extinction in human fear conditioning. *Biological Psychiatry*. 2006; 60:361–368. [PubMed: 16503330]
- Hoddes E, Zarcone V, Smythe H, Phillips R, Dement WC. Quantification of sleepiness: a new approach. *Psychophysiology*. 1973; 10:431–436. [PubMed: 4719486]
- Hofmann SG. Cognitive processes during fear acquisition and extinction in animals and humans: implications for exposure therapy of anxiety disorders. *Clinical Psychology Review*. 2008; 28:199–210. [PubMed: 17532105]
- Horne JA, Ostberg O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International Journal of Chronobiology*. 1976; 4:97–110. [PubMed: 1027738]
- Hot P, Leconte P, Sequeira H. Diurnal autonomic variations and emotional reactivity. *Biological Psychology*. 2005; 69:261–270. [PubMed: 15925029]
- Johns MW. Sleepiness in different situations measured by the Epworth Sleepiness Scale. *Sleep*. 1994; 17:703–710. [PubMed: 7701181]
- Kalisch R, Korenfeld E, Stephan KE, Weiskopf N, Seymour B, Dolan RJ. Context-dependent human extinction memory is mediated by a ventromedial prefrontal and hippocampal network. *Journal of Neuroscience*. 2006; 26:9503–9511. [PubMed: 16971534]
- Klorman R, Hastings J, Weerts T, Melamed B, Lang PJ. Psychometric description of some specific fear questionnaires. *Behav Ther*. 1974; 5:401–409.
- Knight DC, Smith CN, Cheng DT, Stein EA, Helmstetter FJ. Amygdala and hippocampal activity during acquisition and extinction of human fear conditioning. *Cogn Affect Behav Neurosci*. 2004; 4:317–325. [PubMed: 15535167]
- LaBar KS, Gatenby JC, Gore JC, LeDoux JE, Phelps EA. Human amygdala activation during conditioned fear acquisition and extinction: a mixed-trial fMRI study. *Neuron*. 1998; 20:937–945. [PubMed: 9620698]
- LeBeau RT, Glenn D, Liao B, Wittchen HU, Beesdo-Baum K, Ollendick T, Craske MG. Specific phobia: a review of DSM-IV specific phobia and preliminary recommendations for DSM-V. *Depression and Anxiety*. 2010; 27:148–167. [PubMed: 20099272]
- Lissek S, Biggs AL, Rabin SJ, Cornwell BR, Alvarez RP, Pine DS, Grillon C. Generalization of conditioned fear-potentiated startle in humans: experimental validation and clinical relevance. *Behaviour Research and Therapy*. 2008; 46:678–687. [PubMed: 18394587]
- Lissek S, Orme K, McDowell DJ, Johnson LL, Luckenbaugh DA, Baas JM, Cornwell BR, Grillon C. Emotion regulation and potentiated startle across affective picture and threat-of-shock paradigms. *Biological Psychology*. 2007; 76:124–133. [PubMed: 17692453]
- Lissek S, Powers AS, McClure EB, Phelps EA, Woldehawariat G, Grillon C, Pine DS. Classical fear conditioning in the anxiety disorders: a meta-analysis. *Behaviour Research and Therapy*. 2005; 43:1391–1424. [PubMed: 15885654]
- Lykken DT, Venables PH. Direct measurement of skin conductance: a proposal for standardization. *Psychophysiology*. 1971; 8:656–672. [PubMed: 5116830]
- McNally RJ. Mechanisms of exposure therapy: how neuroscience can improve psychological treatments for anxiety disorders. *Clinical Psychology Review*. 2007; 27:750–759. [PubMed: 17292521]
- McSweeney FK, Swindell S. Common processes may contribute to extinction and habituation. *Journal of General Psychology*. 2002; 129:364–400. [PubMed: 12494990]
- Milad MR, Rauch SL, Pitman RK, Quirk GJ. Fear extinction in rats: implications for human brain imaging and anxiety disorders. *Biological Psychology*. 2006; 73:61–71. [PubMed: 16476517]
- Milad MR, Wright CI, Orr SP, Pitman RK, Quirk GJ, Rauch SL. Recall of fear extinction in humans activates the ventromedial prefrontal cortex and hippocampus in concert. *Biological Psychiatry*. 2007; 62:446–454. [PubMed: 17217927]
- Mineka S, Oehlberg K. The relevance of recent developments in classical conditioning to understanding the etiology and maintenance of anxiety disorders. *Acta Psychologica*. 2008; 127:567–580. [PubMed: 18226795]
- Muris P, Merckelbach H. A comparison of two spider fear questionnaires. *J Behav Ther Exp Psychiatry*. 1996; 27:241–244. [PubMed: 8959425]

- Nofzinger EA, Buysse DJ, Germain A, Carter C, Luna B, Price JC, Meltzer CC, Miewald JM, Reynolds CF 3rd, Kupfer DJ. Increased activation of anterior paralimbic and executive cortex from waking to rapid eye movement sleep in depression. *Archives of General Psychiatry*. 2004; 61:695–702. [PubMed: 15237081]
- Orr SP, Metzger LJ, Lasko NB, Macklin ML, Peri T, Pitman RK. De novo conditioning in trauma-exposed individuals with and without posttraumatic stress disorder. *Journal of Abnormal Psychology*. 2000; 109:290–298. [PubMed: 10895567]
- Pace-Schott EF, Milad MR, Orr SP, Rauch SL, Stickgold R, Pitman RK. Sleep promotes generalization of extinction of conditioned fear. *Sleep*. 2009; 32:19–26. [PubMed: 19189775]
- Pace-Schott EF, Shepherd E, Spencer RM, Marcello M, Tucker M, Propper RE, Stickgold R. Napping promotes inter-session habituation to emotional stimuli. *Neurobiology of Learning and Memory*. 2011; 95:24–36. [PubMed: 20969968]
- Phelps EA, Delgado MR, Nearing KI, LeDoux JE. Extinction learning in humans: role of the amygdala and vmPFC. *Neuron*. 2004; 43:897–905. [PubMed: 15363399]
- Quirk GJ, Mueller D. Neural mechanisms of extinction learning and retrieval. *Neuropsychopharmacology*. 2008; 33:56–72. [PubMed: 17882236]
- Rachman S. Fears born and bred: non-associative fear acquisition? *Behaviour Research and Therapy*. 2002; 40:121–126. [PubMed: 11814177]
- Rodriguez BI, Craske MG, Mineka S, Hladek D. Context-specificity of relapse: effects of therapist and environmental context on return of fear. *Behav Res Ther*. 1999; 37:845–862. [PubMed: 10458048]
- Rowe MK, Craske MG. Effects of varied-stimulus exposure training on fear reduction and return of fear. *Behaviour Research and Therapy*. 1998; 36:719–734. [PubMed: 9682527]
- Santini E, Muller RU, Quirk GJ. Consolidation of extinction learning involves transfer from NMDA-independent to NMDA-dependent memory. *Journal of Neuroscience*. 2001; 21:9009–9017. [PubMed: 11698611]
- Schweckendiek J, Klucken T, Merz CJ, Tabbert K, Walter B, Ambach W, Vaitl D, Stark R. Weaving the (neuronal) web: fear learning in spider phobia. *Neuroimage*. 2011; 54:681–688. [PubMed: 20673801]
- Spielberger, CD.; Gorsuch, RL.; Lushene, RE. *Manual for the State-Trait Anxiety Inventory (Self-Evaluation Questionnaire)*. Palo Alto: Consulting Psychologists Press; 1990.
- Storsve AB, McNally GP, Richardson R. US habituation, like CS extinction, produces a decrement in conditioned fear responding that is NMDA dependent and subject to renewal and reinstatement. *Neurobiology of Learning and Memory*. 2010; 93:463–471. [PubMed: 20074653]
- Szymanski J, O'Donohue W. Fear of Spiders Questionnaire. *Journal of Behavior Therapy and Experimental Psychiatry*. 1995; 26:31–34. [PubMed: 7642758]
- van Overveld M, de Jong PJ, Peters ML, Cavanagh K, Davey GCL. Disgust propensity and disgust sensitivity: separate constructs that are differentially related to specific fears. *Personality and Individual Differences*. 2006; 41:1241–1252.
- Vansteenwegen D, Vervliet B, Iberico C, Baeyens F, Van den Bergh O, Hermans D. The repeated confrontation with videotapes of spiders in multiple contexts attenuates renewal of fear in spider-anxious students. *Behaviour Research and Therapy*. 2007; 45:1169–1179. [PubMed: 17045959]
- Walker MP. The role of sleep in cognition and emotion. *Annals of the New York Academy of Sciences*. 2009; 1156:168–197. [PubMed: 19338508]
- Wessa M, Flor H. Failure of extinction of fear responses in posttraumatic stress disorder: evidence from second-order conditioning. *American Journal of Psychiatry*. 2007; 164:1684–1692. [PubMed: 17974933]

Figure 1.

Experimental protocol showing experimental phases and putative emotional memory processes accounting for changes across phases. In S1, spider videos 13–14 were added in order to maximize extinction learning following (Vansteenwegen et al., 2007). However data for these videos were not analyzed so as to accommodate mixed ANOVA design with nested within-subject factors

Figure 2. Composite negative ratings (summed Disgust, Fear and Unpleasantness ratings). A. Experimental (Sleep and Wake) groups. B. Control (Morning and Evening) Groups. Arrows and boxes superimposed on histogram indicate significance levels for post-hoc means comparisons of phases in separate repeated measures ANOVAs for each group. Arrows indicate means comparisons that produced differing results in the two groups when analyzed separately. Boxes indicate Extinction Learning in S1 (see Supplementary Results) and Response Return in S2. Bars are standard error of the mean. † p < 0.10, * p < 0.05, ** p < 0.01, *** p < 0.001

Figure 3.

Video SCR computed by subtracting the mean SCL during the first 1 sec following video onset (the latency for SCR) from the highest SCL achieved during the following 29 sec of video (during which the noise stimulus never occurred). A. Experimental (Sleep and Wake) groups. B. Control (Morning and Evening) Groups. Arrows indicate means comparisons that produced differing results in the two groups analyzed separately. Boxes indicate Extinction Learning in S1 (see Supplementary Results) and Response Return in S2. Bars are standard error of the mean. † $p < 0.10$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, $\mu\text{S}^{1/2}$ square-root transformed SCR in micro-Siemens

Figure 4.

Noise SCR to 10 msec, 83 dB white noise played through headphones during 60% of videos and ISIs. A. Noise-SCR-in-Video in experimental (Sleep and Wake) groups. B. Noise SCR-in-ISI in experimental groups. C. Noise-SCR-in-Video in control (Morning and Evening) groups. D. SCR-in-ISI in control groups. Arrows indicate means comparisons that produced differing results in the two experimental groups analyzed separately. Boxes indicate Extinction Learning in S1 (see Supplementary Results) and Response Return in S2. Bars are standard error of the mean. † $p < 0.10$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.01$, $\mu S^{1/2}$ square-root transformed SCR in micro-Siemens.. (Because noise-containing portions of physiological recordings were intact for all Evening subjects, N=12 vs. 11 for other physiological measures in this group.)

Figure 5.

Heart-Rate acceleration (HRA) in response to spider video relative to preceding ISI. Heart Rate sampled from both ISI and video during initial period in which noise stimulus did not occur (first 2 and 30 sec respectively). A. Experimental (Sleep and Wake) groups. B. Control (Morning and Evening) Groups. Arrows indicate means comparisons that produced differing results in the two groups analyzed separately. Boxes indicate Extinction Learning in S1 (see Supplementary Results) and Response Return in S2. Bars are standard error of the mean. † $p < 0.10$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, BPM beats per minute

Figure 6.

Corrugator supercilii EMG in response to spider video relative to preceding ISI. EMG sampled from both ISI and video during initial period in which noise stimulus did not occur (first 2 and 30 sec respectively). A. Experimental (Sleep and Wake) groups. B. Control (Morning and Evening) Groups. Arrows indicate Sensitization. Box indicates Response Return. Bars are standard error of the mean. † $p < 0.10$, * $p < 0.05$, ** $p < 0.01$, mV millivolts

Table 1

Demographic, self-reported habitual sleep, sleepiness and substance use, subjective sleep duration and psychological traits in the Sleep and Wake groups.

Characteristic	Sleep (SD)	Wake (SD)	Morning	Evening	F(3,62) [†]
N	20	23	11	12	
Age	20.1 (1.5)	20.2 (2.1)	19.3 (1.4)	19.5 (1.1)	1.12
FSQ	101.4 (12.4)	107.1 (14.9)	110.8 (10.5)	110.3 (13.6)	1.72
SPQ	23.7 (3.3)	24.5 (4.3)	23.3 (2.4)	24.0 (3.2)	0.38
Habitual TST (hr)	7.6 (0.9)	7.4 (1.1)	7.7 (0.8)	7.7 (0.9)	0.38
Habitual SOL (min)	17.3 (10.9)	24.2 (23.7)	24.6 (15.3)	19.2 (13.5)	0.75
ESS	7.50 (3.29)	8.30 (4.42)	9.82 (4.36)	9.08 (3.80)	0.92
PSQI	4.80 (1.58)	5.09 (2.97)	5.67 (2.40)	4.92 (1.83)	0.30 ^a
MEQ	39.50 (7.56)	44.83 (10.50)	43.00 (10.34)	39.13 (10.08)	1.52 ^b
STAI-Trait	41.05 (8.41)	40.96 (10.11)	36.86 (11.49)	44.25 (10.11)	1.08
Disgust propensity	24.80 (5.14)	25.70 (5.87)	26.82 (4.75)	24.5 (3.09)	0.53
Disgust sensitivity	18.70 (4.51)	19.46 (7.88)	21.91 (5.84)	19.96 (3.48)	0.70
* NEO-PLR Neuroticism	103.63 (17.87)	101.52 (21.13)	95.64 (17.91)	101.67 (20.96)	0.75
NEO-PLR Extraversion	118.40 (21.68)	128.28 (19.92)	138.45 (15.15)	123.92 (15.10)	2.79 [*]
NEO-PLR Openness	120.40 (14.54)	117.87 (18.71)	116.46 (21.09)	114.67 (15.61)	0.30
NEO-PLR Agreeableness	120.75 (17.40)	116.30 (19.41)	110.09 (18.53)	116.25 (12.70)	0.87
NEO-PLR Conscientiousness	111.20 (16.68)	116.74 (21.54)	120.55 (20.13)	108.83 (25.98)	0.86
Habitual daily caffeine (serv.)	1.5 (1.2)	1.5 (1.2)	1.1 (1.0)	1.0 (0.7)	0.44 ^c
Habitual weekly EtOH (serv.)	3.3 (4.8)	2.4 (2.6)	3.6 (2.9)	3.8 (2.6)	0.50 ^d
Diary sleep Day -1 (min)	489 (91)	458 (35)	455 (20)	555 (78)	6.87 ^{***e}
Diary sleep Day -2 (min)	490 (68)	496 (98)	447 (109)	517 (116)	1.03 ^f
Diary inter-session sleep	431 (25)				
(N) Criteria deviations	(3) Head injury (1) Head injury & panic disorder & night terror (1) "Sleep pill"	(3) Head injury (2) Ritalin (1) Head injury & Insomnia		(1) Head injury & Ritalin & Depression (1) Fluoxetine	

* p < .05, Morning > Sleep (p < .01) and Evening (p < .05)

$p < .001$, Evening > Morning ($p < .0001$), Wake ($p < .0001$) and Sleep ($p = .008$)

^f Lower-case letters reflect smaller samples due to participant omission of data points:

^aF(3,60) Morning N=9;

^bF(3,61) Morning N=10;

^cF(3,61) Wake N=22;

^dF(3,57) Wake N=21, Morning N=10, Evening N=10;

^eF(3,60) Sleep N=19, Wake N=22;

^fF(3,59) Sleep N=19, Wake N=22, Morning N=10