Draft Genome Sequence of *Staphylococcus aureus* 118 (ST772), a Major Disease Clone from India

Sushma Prabhakara,^a Supriya Khedkar,^b Ramya Malarini Loganathan,^c S. Chandana,^c Malali Gowda,^c Gayathri Arakere,^a and Aswin Sai Narain Seshasayee^b

Society for Innovation and Development, Indian Institute of Science Campus, Bangalore, India^a; National Centre for Biological Sciences, TIFR, GKVK, Bangalore, India^b; and Centre for Cellular and Molecular Platforms, GKVK, Bangalore, India^c

We report the draft genome sequence of an ST772 *Staphylococcus aureus* disease isolate carrying staphylococcal cassette chromosome *mec* (SCC*mec*) type V from a pyomyositis patient. Our *de novo* short read assembly is ~2.8 Mb and encodes a unique Panton-Valentine leukocidin (PVL) phage with structural genes similar to those of ϕ 7247PVL and novel lysogenic genes at the N termini.

Community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA) has become a major disease threat in the community as well as in hospitals (5, 8). Until 2004, hospital-associated MRSA, carrying the staphylococcal cassette chromosome *mec* (SCC*mec*) types III and IIIA, was common in India (2). Now, these are being replaced by CA-MRSA strains carrying SCC*mec* types IV (EMRSA-15) and V (ST772) in Indian hospitals (6, 9). These newer variants also carry several toxins and virulence factors, including Panton-Valentine leukocidin (PVL). ST772, called the Bengal Bay clone by some, was previously reported only in Bangladesh, India, and Malaysia but is now being reported in England, Ireland, and several other European countries (1, 4, 10).

We obtained genomic DNA from an *S. aureus* isolate belonging to ST772 (designated strain 118) from a pyomyositis patient (7), fragmented it to 300 to 400 bp, and sequenced these fragments on an Illumina HiSeq 1000 sequencer, thus obtaining ~53 million 75-mer reads (~26.5 million pairs). For further analysis, we selected reads based on the following three criteria: (i) average read quality of ≥ 35 , (b) minimum base quality of ≥ 30 , and (c) no undetermined base. This gave us ~6.4 million read pairs, with an overall coverage of ~350. Since coverage of $>50\times$ to $100\times$ might compromise the quality of assemblies produced by the software Velvet (11), we randomly sampled ~15% of reads from these data and used them for the assembly process. The assembly was performed on three sets of independently sampled read sets to ensure consistency. In addition, we also performed a similar sequencing of *S. aureus* USA300, for which a reference genome sequence is already available.

The draft assembly of the genome of *S. aureus* 118 (ST772) covers ~2.81 million base pairs. These bases are covered by 67, 72, and 73 contigs in the three assemblies attempted. Paired-end mapping of all raw reads to the contigs predicts an insert size distribution which is similar to that obtained by mapping our USA300 reads to its fully sequenced reference sequence. Finally, ~94 to 95% of the bases in our contigs fall within alignments to fully sequenced *S. aureus* genomes (BLASTN; E value $< 10^{-100}$).

We identified 2,524 protein-coding genes predicted by Glimmer in two of the three assemblies constructed here. Of these, 2,459 (97%) have a counterpart with significant similarity (phmer, <http://hmmer.janelia.org>; E value $< 10^{-10}$; $\geq 40\%$ global sequence identity) in at least one fully sequenced *S. aureus* genome. We also detected an ~42-kb PVL phage sequence by using signa-

ture sequences published earlier (3). Comparison of this sequence with five other sequenced PVL phages (ϕ PVL, ϕ 108PVL, ϕ SLT, ϕ Sa2958, and ϕ 7247PVL) showed that it is closest to the ϕ 7247PVL phage from *S. aureus* ST59 (12). However, the best-aligning regions are largely restricted to the structural regions, with considerable variations in the lysogeny region.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at GenBank under the accession number [AJGE000000000](https://www.ncbi.nlm.nih.gov/nuccore/AJGE000000000). The version described in this paper is the first version. The raw Illumina sequencing reads have been deposited with the Short Read Archive and are available under the accession number [SRX118450](https://www.ncbi.nlm.nih.gov/sra/SRR118450).

ACKNOWLEDGMENTS

This work was supported by the following: a grant (BT/IN/New Indigo/16/GA/2010) from the Department of Biotechnology, the Ramanujan Fellowship (SR/S2/RJN-49/2010) from the Department of Science and Technology, Government of India, and a fellowship [09/860(0122)/2011-EMR-I] from the Council for Scientific & Industrial Research (CSIR), India.

Sequencing was done at the Centre for Cellular and Molecular Platforms (C-CAMP).

REFERENCES

1. Afroz S, et al. 2008. Genetic characterization of *Staphylococcus aureus* isolates carrying Panton-Valentine leukocidin genes in Bangladesh. *Jpn. J. Infect. Dis.* 61:393–396.
2. Arakere G, Nadig S, Macaden R, Amarnath SK, Raghunath D. 2005. Genotyping of methicillin-resistant *Staphylococcus aureus* strains from two hospitals in Bangalore, South India. *J. Clin. Microbiol.* 43:3198–3202.
3. Boakes E, et al. 2011. Distinct bacteriophages encoding Panton-Valentine leukocidin (PVL) among international methicillin-resistant *Staphylococcus aureus* clones harboring PVL. *J. Clin. Microbiol.* 49: 684–692.
4. Brennan GI, et al. 2012. Emergence of hospital- and community-

Received 28 March 2012 Accepted 30 April 2012

Address correspondence to Gayathri Arakere, gayathri.arakere@gmail.com, or Aswin Sai Narain Seshasayee, aswin@ncbs.res.in.

S.P. and S.K. contributed equally to this study.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00480-12

- associated Panton-Valentine leukocidin-positive methicillin-resistant *Staphylococcus aureus* genotype ST772-MRSA-V in Ireland and detailed investigation of a ST772-MRSA-V cluster in a neonatal intensive care unit. *J. Clin. Microbiol.* 50:841–847.
5. DeLeo FR, Otto M, Kreiswirth BN, Chambers HF. 2010. Community-associated methicillin-resistant *Staphylococcus aureus*. *Lancet* 375:1557–1568.
 6. D'Souza N, Rodrigues C, Mehta A. 2010. Molecular characterization of methicillin-resistant *Staphylococcus aureus* with emergence of epidemic clones of sequence type (ShT) 22 and ST 772 in Mumbai, India. *J. Clin. Microbiol.* 48:1806–1811.
 7. Kulkarni G, et al. 2009. Community-acquired methicillin-resistant *Staphylococcus aureus* pyomyositis with myelitis: a rare occurrence with diverse presentation. *Neurol. India* 57:653.
 8. Monecke S, et al. 2011. A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PLoS One* 6:e17936. doi:10.1371/journal.pone.0017936.
 9. Nadig S, et al. 2012. *Staphylococcus aureus* eye infections in two Indian hospitals: emergence of ST772 as a major clone. *Clin. Ophthalmol.* 6:165–173.
 10. Neela V, Ehsanollah GR, Zambari S, Van Belkum A, Mariana NS. 2009. Prevalence of Pantan-Valentine leukocidin genes among carriage and invasive *Staphylococcus aureus* isolates in Malaysia. *Int. J. Infect. Dis.* 13:e131–e132.
 11. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
 12. Zhang M, et al. 2011. Identification of the third type of PVL phage in ST59 methicillin-resistant *Staphylococcus aureus* (MRSA) strains. *FEMS Microbiol. Lett.* 323:20–28.