Published in final edited form as:

Biochemistry. 2009 July 28; 48(29): 6932–6939. doi:10.1021/bi900713j.

Hydrogen Bond Coupling in the Ketosteroid Isomerase Active Site^{†,‡}

Paul A. Sigala[§], Jose M. M. Caaveiro^{⊥,‡}, Dagmar Ringe[⊥], Gregory A. Petsko[⊥], and Daniel Herschlag^{§,*}

[§]Department of Biochemistry, Stanford University, Stanford, California 94305

[⊥]Departments of Biochemistry and Chemistry and Rosenstiel Basic Medical Sciences Research Center, Brandeis University, Waltham, Massachusetts 02454

Abstract

Hydrogen bond networks are key elements of biological structure and function. Nevertheless, their structural properties are challenging to assess within complex macromolecules. Hydrogen-bonded protons are not observed in the vast majority of protein X-ray structures, and static crystallographic models provide limited information regarding the dynamical coupling within hydrogen bond networks. We have brought together 1.1 – 1.3 Å resolution X-ray crystallography, ¹H NMR, site-directed mutagenesis, and deuterium isotope effects on the geometry and chemical shifts of hydrogen-bonded protons to probe the conformational coupling of hydrogen bonds donated by Y16 and D103 in the oxyanion hole of bacterial ketosteroid isomerase. Our results suggest a robust physical coupling of the equilibrium structures of these two hydrogen bonds such that a lengthening of one hydrogen bond by as little as 0.01 Å results in a shortening of the neighbor by a similar magnitude. Furthermore, the structural rearrangements detected by NMR in response to mutations within the active site hydrogen bond network can be explained on the basis of the observed coupling. The results herein elucidate fundamental structural properties of hydrogen bonds within the idiosyncratic environment of an enzyme active site and provide a foundation for future experimental and computational explorations of the role of coupled motions within hydrogen bond networks.

Hydrogen bonds are central to the structure and function of biological macromolecules. Despite the ubiquitous presence of hydrogen bonds in biology and their extensive study in small molecules (1, 2), incisive dissection of hydrogen bond properties within proteins and other large biomolecules remains challenging. Indeed, electron density for hydrogen-bonded protons is not typically observed in high-resolution X-ray diffraction studies of proteins. Thus, hydrogen bond formation must be indirectly assessed from the proximity of putative donor and acceptor heavy nuclei observed crystallographically. Furthermore, static structural

[†]This work was funded by an NSF grant to D.H., D.R., and G.A.P. (MCB-0641393). P.A.S. was supported in part by HHMI and Gerald Lieberman Predoctoral Fellowships. Portions of this research were conducted at the Stanford Synchrotron Radiation Laboratory, which is supported by the Department of Energy and the National Institutes of Health.

[‡]Coordinates for the pKSI^{D103N/D40N}•equilenin crystal structure have been deposited in the RCSB Protein Data Bank as entry 3FZW.

*To whom correspondence should be addressed: herschla@stanford.edu, Phone: (650) 723-9442, Fax: (650) 723-6783.

[#]Present address: Laboratory of Physical Biochemistry, Graduate School of Frontier Sciences, University of Tokyo, Japan.

SUPPORTING INFORMATION AVAILABLE

UV absorbance spectra for tKDI^{D103N/D40N}•3,4,5-F₃-phenol and tKDI^{Y57F/D40N}•3,4,5-F₃-phenol complexes (Figure S1), a structural model of the ordered water molecule observed in the oxyanion hole of pKSI^{D103N/D40N}•equilenin (Figure S2), and assignment of the downfield peaks observed in spectra of tKSI^{D103N/D40N} and tKSI^{Y57F/D40N} bound to 3,4-dinitrophenol. This material is available free of charge via the Internet at <http://pubs.acs.org>.

models provide limited insight regarding the structural plasticity and conformational dynamics of hydrogen bonds, whose distances can vary by up to ~ 1 Å based on the specific properties of the donor and acceptor groups and the local structural environment (1, 3–6).

Networks of adjoining hydrogen bonds commonly form in proteins, where they span secondary structures, link disparate domains, connect active site residues, position bound ligands, and stabilize charge rearrangement in reacting substrates (7). Recent studies suggests that the unique structural and dynamical properties of hydrogen bond networks play important roles in coupling the motions of distal groups for allostery and signal propagation (6, 8–10) and in fine-tuning the positions of protein and ligand groups, especially those involved in proton-coupled electron transfer (11–13). While our understanding of the dynamic motions with protein structural elements spanning a broad range of distance and time scales is rapidly increasing (14–18), direct assessment of the physical coupling between groups remains challenging (19, 20).

Bacterial ketosteroid isomerase (KSI)¹ from *Pseudomonas putida* (pKSI) and *Commamonas testosteroni* (tKSI) has provided a powerful experimental system to dissect the physical properties of key active site hydrogen bonds. KSI catalyzes the isomerization of steroid substrates via formation of a dienolate intermediate (Scheme 1A) (21). Localized negative charge on the oxygen of the reaction intermediate is stabilized by hydrogen bonds donated by Y16 (pKSI numbering) and protonated D103 (22, 23), which form a conserved hydrogen bond network within an active-site oxyanion hole via the additional hydrogen bond donated by Y57 to the hydroxyl of Y16 (Figures 1A & 1B) (24). The substantial rate decreases observed upon mutation of Y16 (25, 26) and D103 (23, 27, 28) and results from linear free energy relationship studies of KSI (4, 29) suggest important functional roles for these hydrogen bonds.

The steroid, equilenin, and single-ring phenolates bind in the KSI active site, accept 2.5 – 2.6 Å hydrogen bonds from Y16 and D103 (22, 29, 30), and serve as analogs of the dienolate reaction intermediate (Scheme 1B). The hydrogen-bonded protons donated by Y16 and D103 to the oxygen of KSI-bound phenolates appear as far-downfield peaks (14 – 18 ppm) in ¹H NMR spectra, providing a spectroscopic probe with which to dissect hydrogen bond properties. We previously showed that increasing oxyanion charge localization, analogous to the charge rearrangement during steroid isomerization, results in a shortening of these hydrogen bonds by ~ 0.02 Å per unit increase in oxyanion pK_a (29). The introduction of structural constraints within the oxyanion hole, however, can prevent such hydrogen bond shortening, highlighting the ability of binding interactions within enzyme active sites to alter hydrogen bond properties from those in solution (4).

Herein we use X-ray crystallography, site-directed mutagenesis, NMR, and hydrogen/deuterium geometric isotope effects to address two critical questions regarding the physical coupling of Y16 and D103. First, are the Y16 and D103 hydrogen bonds conformationally coupled, such that a structural perturbation to one is propagated to the other? And second, what is the distance scale of the observed coupling? Our results suggest a robust physical coupling of the equilibrium structures of the Y16 and D103 hydrogen bonds, with a lengthening of one by as little as 0.01 Å resulting in a shortening of the neighboring hydrogen bond by a similar magnitude.

¹Abbreviations: KSI, ketosteroid isomerase; pKSI, KSI from *Pseudomonas putida*; tKSI, KSI from *Commamonas testosteroni*; PDB, Protein Data Bank

MATERIALS AND METHODS

Materials

All reagents were of the highest purity commercially available. 3-F-5-CF₃-phenol, 3,4,5-F₃-phenol, and equilenin were purchased from Oakwood Products, Matrix Scientific, and Steraloids, respectively. DMSO-*d*₆ and D₂O were purchased from Cambridge Isotopes. All buffers were prepared with reagent-grade chemicals or better.

KSI Mutagenesis, Expression and Purification

QuikChange site-directed mutagenesis was used to introduce the D40N, D103N, and Y57F mutations, which were confirmed by sequencing miniprep DNA from DH5α cells on an ABI3100 capillary sequencer. KSI was expressed and purified as previously described (29). Final purity was >99%, as estimated from a Coomassiestained SDA-PAGE gel. Protein concentrations were determined using the calculated molar extinction coefficient in 6 M guanidine chloride (31).

Absorbance Spectra of KSI•phenolate Complexes

To confirm that phenols bind as ionized phenolates to KSI containing the D103N or Y57F mutations, absorbance spectra of 3,4,5-F₃-phenol were acquired in a microcuvette with a UVikon 9310 absorbance spectrophotometer. Spectra of 50 μM 3,4,5-F₃-phenol in 10 mM HCl (pH 2) and 10 mM NaOH (pH 12) were compared with the spectrum of the same concentration phenol in 40 mM potassium phosphate (KP_i) buffer, pH 7.2, in the presence of 300 μM tKSI^{D103N/D40N} or tKSI^{Y57F/D40N} (phenol >95% bound, data not shown), after subtraction of the free enzyme spectrum. Spectra for both mutants indicated binding as the ionized phenolate.

KSI X-ray Crystallography

pKSI^{D103N/D40N}•equilenin co-crystals in space group C2 were obtained using the hanging drop vapor diffusion method by mixing 2 μl of pKSI^{D103N/D40N} at 25 mg/ml pre-incubated with equilenin (KSI:equilenin molar ratio was 1:1.2) and 2 μl of reservoir solution (1.4 M ammonium sulfate, 6.5 % (v/v) 2-propanol, at pH 7.0). Rod-shaped crystals appeared after one week of incubation at room temperature. Cryo-protection was achieved by soaking the crystals in a solution of mother liquor containing 20% glycerol. Crystals were shock-cooled by immersion into liquid nitrogen. Diffraction data from single crystals maintained at 100 K were collected at beamline BL9-1 of the Stanford Synchrotron Radiation Laboratory (Stanford Linear Accelerator Center, Stanford, CA, USA). Data were integrated and scaled using the HKL2000 program package. An initial model of the protein was obtained by the molecular replacement method with the program PHASER (32), using the coordinates of a previously reported structure of pKSI^{D40N} (PDB code: 2PZV). Refinement and manual rebuilding were carried out using the programs REFMAC5 (33) and COOT (34), respectively. Model quality was assessed using the program PROCHECK (35) and the Ramachandran plot showed no residues outside the allowed regions. Coordinate uncertainty was estimated using the diffraction-component precision index (DPI) developed by Cruickshank (36) and carried out in REFMAC5. Structural coordinates have been deposited in the RCSB Protein Data Bank (<http://www.rcsb.org/pdb>) under accession code 3FZW.

KSI NMR Spectroscopy

¹H NMR spectra of KSI were acquired at the Stanford Magnetic Resonance Laboratory on an 800-MHz Varian ^{UNITY}INOVA spectrometer running VNMR v6.1C and equipped with a 5-mm, triple-resonance, gradient ¹H(¹³C/¹⁵N) probe. NMR samples consisted of 0.5 mM tKSI^{D40N} and 0.5 – 5.0 mM substituted phenol in 40 mM KP_i buffer (pH 7.2), 1 mM EDTA,

and 10% (v/v) DMSO- d_6 (which served as the lock solvent and prevented freezing at subzero temperatures) in 5 mM Shigemi symmetrical microtubes at -3.0 ± 0.5 °C. One-dimensional proton spectra were acquired using the 1331 binomial pulse sequence (37) to suppress the water signal, with a spectral width of 30 ppm (carrier frequency set on the water resonance) and an excitation maximum of 14 – 18 ppm. Data were collected for 512 – 5120 scans and processed using a 10-Hz line broadening and baseline correction applied over the peaks of interest. Chemical shifts were referenced internally to the water resonance (5.1 at -3.0 °C) and externally to a sample of sodium 3-trimethylsilylpropionate-2,2,3,3- d_4 (0 ppm) in the same buffer conditions (38). Hydrogen bond O $\cdots$ O and H $\cdots$ O distances were estimated from the observed chemical shifts for the detected hydrogen-bonded protons using the correlation functions given in references (39) and (40), respectively.

RESULTS AND DISCUSSION

To assess the presence of physical coupling between the Y16 and D103 hydrogen bonds in the KSI oxyanion hole, we first took a mutagenesis approach. We hypothesized that a structural perturbation introduced at either the D103 or Y16 hydrogen bonds via mutation might be accompanied by a structural rearrangement in the adjoining hydrogen bond neighbor if these interactions were indeed physically coupled via hydrogen bond formation to the common oxyanion acceptor. If no rearrangement in the adjoining, unmodified hydrogen bond neighbor were detected, then direct physical coupling of these hydrogen bonds might either be absent or overwhelmed by larger-scale individual or collective motions of groups within the KSI active site.

The mutations chosen to site-specifically perturb either the D103 or Y16 hydrogen bonds formed to the oxyanion of equilenin or phenolates were D103N and Y57F, which have minor effects on active site structure and result in small (3 – 30 fold) reductions in KSI activity (k_{cat}). These mutants (in the D40N background)² bind ionized phenolates and maintain two hydrogen bonds to the phenolate or equilenin oxyanion (Figure S1, supporting information). We expected the N $\cdots$ O hydrogen bond in the D103N mutant to be lengthened relative to unmodified D103 on the basis of the higher pK_a of an Asn side-chain amide (~ 17) (41) relative to the hydroxyl of D103 (~ 8) (27) and hence greater pK_a mismatch with the bound oxyanion (5, 29). We similarly expected the Y57F mutation, which ablates a hydrogen bond to Y16 and presumably weakens its ability to donate a hydrogen bond, to result in a lengthening of the Y16 $\cdots$ oxyanion hydrogen bond.

We first determined a high-resolution X-ray structure of pKSI^{D103N/D40N}•equilenin to directly assess the structural consequences of the D103N mutation and any propagated effects on the Y16•equilenin O $\cdots$ O distance relative to the previously published 1.1 Å resolution pKSI•equilenin structure (PDB 1OH0). We then exploited the exquisite sensitivity of NMR and the ability to spectroscopically observe the hydrogen bonded protons of Y16 and D103 to test for propagated effects from the D103N and Y57F mutations that would signal the presence of physical coupling within the oxyanion hole hydrogen bond network. Finally, we applied an isotopic substitution method to directly probe conformational coupling in the unmodified hydrogen bond network of tKSI^{D40N} in the absence of the potential indirect effects from oxyanion hole mutations.

Structure of pKSI^{D103N/D40N}•equilenin at 1.3 Å Resolution

A previously-published 1.1 Å resolution X-ray structure of pKSI•equilenin (22) identified short average O $\cdots$ O distances of 2.51 Å and 2.56 Å for the hydrogen bonds donated by

²The D40N mutation mimics the protonated general base, D40, present in the KSI-intermediate complex.

D103 and Y16, respectively, to the oxyanion of bound equilenin (Table 1). To assess the effect of the D103N mutation on these hydrogen bond distances, we determined the X-ray structure of pKSI^{D103N/D40N}•equilenin. Data collection and refinement statistics are given in Table 2. The overall KSI structure obtained at 1.3 Å resolution is nearly identical to that observed previously for pKSI•equilenin, with a root-mean-square deviation between the two structures of 0.3 Å for backbone atoms. The electron density map of the pKSI^{D103N/D40N}•equilenin structure, contoured in Figure 1A at 1.5 σ , shows well-defined density for the modeled atomic positions within the active site, with no indication of alternative ligand or side-chain conformations. We observed electron density for an ordered water molecule in the oxyanion hole within hydrogen bonding distance (3.27 Å) of the equilenin oxyanion (Figure S2). This water was not reported in the pKSI•equilenin structure (PDB 1OH0).

The refined orientation of bound equilenin is very similar to that observed previously for pKSI•equilenin, with equilenin positioned in both active sites to accept hydrogen bonds from Y16 and D103 or N103 (Figure 1B). Subtle displacements in the positions of the equilenin oxyanion and the N103 side-chain in pKSI^{D103N/D40N}•equilenin, however, result in a lengthened average N•••O distance of 2.89 Å relative to the 2.51 Å O•••O distance observed for D103 in pKSI•equilenin (Table 1). This length difference of 0.38 Å is clearly distinguishable at the stated resolutions and 0.03 – 0.05 Å estimated coordinate error for the two structures (see Table 1 and Materials and Methods).

Is lengthening of the hydrogen bond donated by residue 103 to the equilenin oxyanion accompanied by a change in the length of the Y16•equilenin hydrogen bond, as expected for physically coupled interactions? The average observed O•••O distance for the Y16•equilenin hydrogen bond in the pKSI^{D103N/D40N}•equilenin structure is 2.51 Å, which is 0.05 Å shorter than the corresponding distance observed for KSI•equilenin (Table 1). This apparent shortening is consistent with a modest coupling of these hydrogen bonds. However, the observed 0.05 Å change is similar to the 0.03 – 0.05 Å estimated coordinate uncertainty of the two structures, rendering a definitive conclusion about the scale of coupling problematic. Furthermore, the small change in O•••O distance may mask a larger change in the position of the hydrogen-bonded proton of Y16. As stated above, hydrogen-bonded protons are not typically observed in high-resolution X-ray structures, even structures obtained at nearly 1.0 Å resolution. To overcome this limitation, we next turned to ¹H NMR to directly detect the hydrogen-bonded protons in the oxyanion hole and assess changes in their positions in response to the D103N and Y57F mutations.

NMR Assessment of Oxyanion Hole Hydrogen Bond Coupling via the D103N and Y57F Mutations

Extensive studies of O-H•••O hydrogen bonds in small molecule complexes by neutron diffraction and solid state NMR have established a strong correlation between the observed chemical shift of a hydrogen-bonded proton and the O•••O and H•••O hydrogen bond distances (39, 40). Thus, changes in the observed chemical shifts of hydrogen-bonded protons can be used to estimate changes in the O•••O and H•••O hydrogen bond distances (42), as previously reported for KSI•phenolate complexes (4, 29).

To assess whether structural perturbations to the D103 or Y16 hydrogen bonds due to the D103N and Y57F mutations, respectively, result in propagated rearrangements within the adjoining, unmodified hydrogen bond, we acquired NMR spectra of 3,4-dinitrophenolate bound to tKSI^{D40N}, tKSI^{D103N/D40N}, and tKSI^{Y57F/D40N}. As shown in Figure 2, two far-downfield peaks at 14.0 and 14.2 ppm are observed for tKSI^{D40N}•3,4-dinitrophenolate. These peaks have previously been assigned to the hydrogen-bonded protons donated by D103 and Y16 to the phenolate oxygen, based on H-H NOESY spectra and oxyanion hole

mutations (29). These experiments, however, cannot distinguish which of the two peaks arises from Y16 and which is from D103; the conclusions herein do not require further assignment of these peaks.

Lengthening of the D103 hydrogen bond via the D103N mutation resulted in a ~1 ppm downfield shift in the NMR peak position corresponding to the remaining, unmutated hydrogen-bonded proton of Y16 (Figure 2, see further description of the peak assignments in the mutant spectra in Supporting Information). This chemical shift change is consistent with a ~0.05 contraction in the O...O and H...O hydrogen bond distances, based on the correlation functions given in references (40) and (39), and very similar to the 0.05 Å shortening observed in the pKSI^{D103N/D40N}•equilenin X-ray structure described above. The Y57F mutation maintains two O-H...O hydrogen bonds to the bound oxyanion but is expected to lengthen the Y16 hydrogen bond (see above). This mutation resulted in one peak moving ~2.4 ppm upfield and the other moving ~0.6 ppm downfield relative to tKSI^{D40N}•3,4-dinitrophenolate (Figure 2). These chemical shift changes are expected for a 0.13 Å lengthening of the Y16 hydrogen bond and a concomitant 0.03 Å shortening of the D103 hydrogen bond. For hydrogen bonds donated to the same acceptor, the shortening of one in response to lengthening of the other has previously been termed “anti-cooperative”, based on studies of small molecules (43, 44).

These changes suggest a modest scale of conformational coupling between the hydrogen bonds within the KSI oxyanion hole, such that a structural perturbation to one is propagated to its connected hydrogen bond neighbors. Nevertheless, structural rearrangements within nearby groups that are in van der Waals contact with Y16, D103, and the bound phenolate, in response to these mutations, could be the source of the observed spectral shifts rather than direct propagation of conformational changes through the hydrogen bond network. The presence of an ordered water molecule in mutant oxyanion holes (observed in the pKSI^{D103N/D40N}•equilenin structure described above) that is not observed in the pKSI•equilenin structure containing the unmodified oxyanion hole may further complicate the comparison of hydrogen bond properties between the mutant and wild-type oxyanion holes. We therefore applied an isotopic substitution method to directly probe hydrogen bond coupling within the intact hydrogen bond network in the KSI oxyanion hole with minimal secondary perturbation.

Direct Detection of Physical Coupling Between the Y16 and D103 Hydrogen Bonds

Extensive structural studies of hydrogen bonds in small molecules have revealed that short hydrogen bonds with O...O distances less than ~2.7 Å are subtly lengthened upon substitution of the bridging hydrogen by deuterium or tritium (1, 2, 45, 46). These changes have been ascribed to anharmonicity in the shape of the hydrogen bond potential energy well that arises from a shallow barrier for proton transfer. In such cases the lower zero-point energy for deuterium results in a shorter O-D bond than the original O-H bond by 0.01 – 0.02 Å and an increase in the O...O distance by a similar magnitude (2, 47). These changes are shown schematically in Figure 3A. Subtle changes in hydrogen bond length upon isotopic substitution are detectable by NMR and have been reported for small molecules in aprotic organic solvents (2, 47, 48), in ubiquitin (49), and in the active site of chymotrypsin (50).

For two short small molecule hydrogen bonds donated to a common acceptor, deuterium substitution of one hydrogen bond has been observed to result in an increased chemical shift of the proton bridging the neighboring, unsubstituted hydrogen bond. This scenario is shown schematically in Figure 3B (43, 51, 52). These observations are consistent with a lengthening of the first hydrogen bond upon deuterium substitution (due to a shorter O-D sigma bond) and a shortening of the adjoining, unmodified hydrogen bond (due to an

increase in the O-H bond length which leads to increased proton chemical shift). The propagation of such geometric perturbations through hydrogen bond networks as a consequence of deuterium substitution provides direct evidence for the robust physical coupling of the equilibrium conformations of adjoining hydrogen bonds. This coupling may derive from an increase in negative charge localization on the acceptor oxygen that occurs upon lengthening of one hydrogen bond from deuterium substitution such that the proton bridging the neighboring hydrogen bond moves towards the acceptor oxygen and the H•••O-acceptor distance decreases (Figure 3B). Transmission of geometric isotope effects within hydrogen bond networks in proteins has previously been shown for cytosine deaminase (53) and photoactive yellow protein (6).

We applied a similar deuterium substitution strategy to directly probe physical coupling between the Y16 and D103 hydrogen bonds in the unmodified KSI oxyanion hole, as follows. As the D₂O content of a KSI solution increases, the hydrogen-bonded protons in the oxyanion hole exchange with deuterium to populate four possible isotopologues, shown in Figure 4A. Although exchange of a hydrogen-bonded proton for a deuteron renders that hydrogen bond undetectable by ¹H NMR, its adjoining hydrogen-bonded neighbor remains detectable if its bridging proton has not exchanged with solvent deuterons. This is the case for the central HD and DH isotopologues shown in Figure 4A. Given the short 2.5 – 2.6 Å hydrogen bond O•••O distances previously reported for KSI^{D40N}-bound phenolates (29), we expected deuterium substitution of either hydrogen bond to lengthen it by 0.01 – 0.02 Å based on the geometric perturbations reported for similar-length hydrogen bonds in small molecules (45). If the structures of the Y16 and D103 hydrogen bonds were indeed physically coupled, as suggested by the mutagenesis results presented earlier, then deuteration of one hydrogen bond would be expected to result in a shortening of the adjoining hydrogen bond neighbor and more downfield chemical shift for its bridging proton. Thus, in the case of coupling, we expected to observe a second, more downfield resonance for each of the Y16 and D103 hydrogen-bonded proton peaks (four peaks total) in solutions of increasing D₂O, with the new peaks corresponding to the HD isotopologues in Figure 4A. Alternatively, if the Y16 and D103 hydrogen bonds were not strongly coupled, then we would expect to observe only two peaks that decreased in intensity with increasing D₂O content.

In the absence of solvent D₂O, two peaks are observed for the tKSI^{D40N}•3-F-5-CF₃-phenolate complex (Figure 4B), as expected for the exclusive presence of the HH isotopologue. With increasing D₂O content and population of the HD hydrogen bond isotopologues, a second, more-downfield peak is indeed observed for each peak. The intensity of the new HD peaks increases relative to the HH peaks with increasing D₂O, as the isotopologue equilibrium shifts in favor of the HD species. The observation of four peaks indicates that the chemical shift and hence the length of the D103 or Y16 hydrogen bond is altered upon structural perturbation of the neighboring hydrogen bond via deuterium substitution. This result provides direct evidence that the equilibrium structures of the oxyanion hole hydrogen bonds are physically coupled.

For both downfield peaks, the new resonance that appears with increasing D₂O is shifted ~0.2 ppm downfield relative to the HH isotopologue peak (Figure 4B). This 0.2 ppm shift is expected for shortening of the D103 and Y16 hydrogen bond H•••O and O•••O distances by ~0.01 Å. This estimated 0.01 Å shortening detected in the hydrogen bond whose proton has not exchanged is on the same scale as the ~0.01 Å lengthening expected in the hydrogen bond neighbor that has been deuterated (Figure 4A). The similar scale of the primary and propagated distance perturbations provides evidence for a robust degree of coupling of the Y16 and D103 hydrogen bond structures. This coupling is very similar to that observed in the structurally similar oxyanion hole of photoactive yellow protein (6).

While the observed geometric isotope effects provide evidence for robust coupling of hydrogen bond conformations on the ~ 0.01 Å scale (i.e. the primary and propagated distance changes are both ~ 0.01 Å), the smaller propagated effects observed upon mutation suggest that coupling on a larger scale may be attenuated by structural constraints and rearrangements within the active site. Indeed, the 0.4 Å lengthening due to the D103N mutation results in only a ~ 0.05 Å change in the Y16 hydrogen bond length and the 0.13 Å lengthening of the Y16 hydrogen bond from the Y57F mutation leads to only a ~ 0.03 Å estimated shortening of the D103 hydrogen bond.

Conclusions and Implications

The ability to detect coupled changes in oxyanion hole hydrogen bond distances on the ~ 0.01 Å scale indicates that such subtle rearrangements are not swamped out by larger-scale atomic displacements within the active site from individual or collective motions of protein groups. This observation is consistent with previous studies that have suggested a tightly packed environment within the KSI active site and limited dynamic mobility of groups relative to other regions of the enzyme (4, 54). Such limited mobility may enable the KSI active site to discriminate subtle geometrical changes within reacting substrates and contribute to selective transition state stabilization, as previously proposed (4).

While the results herein provide direct evidence for robust physical coupling of the Y16 and D103 hydrogen bond conformations on the 0.01 Å scale, the full scale of physical coupling and the attendant energetic consequences remain challenging to unravel. The D103N and Y57F mutations that perturb hydrogen bond distances within the oxyanion hole result in small (3 – 30 fold) reductions in k_{cat} , depending on the bacterial source of the enzyme (25, 27, 28). Furthermore, the energetic effects of the D103A and Y16F (55) or D103L and Y16F mutations (56) appear to be additive rather than cooperative, as might have been expected from substantial coupling of the two hydrogen bonds. These mutations, however, ablate one of the two oxyanion hole hydrogen bonds and result in neutral binding of phenolic transition state analogs (25, 57). Thus, the connection between the physical and energetic properties of the mutant and wild-type oxyanion holes is complex.

Disentangling the physical and energetic properties of hydrogen bonds within enzyme active sites remains an ongoing challenge. The results herein provide a foundation for future experimental and computational tests and comparisons of the role of coupled motions within hydrogen bond networks.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Corey Liu for assistance with NMR experiments and J. Brauman, H. Limbach, A. Mildvan, and P. Tolstoy for helpful discussions.

REFERENCES

1. Jeffrey, GA. An Introduction to Hydrogen Bonding. New York: Oxford University Press; 1997.
2. Hibbert F, Emsley J. Hydrogen bonding and chemical reactivity. Adv. Phys. Org. Chem. 1990; 26:255–379.
3. Steiner T, Saenger W. Lengthening of the covalent O-H bond in O-H---O hydrogen bonds re-examined from low-temperature neutron diffraction data of organic compounds. Acta Crystallogr. B. 1994; 50:348–357.

4. Sigala PA, Kraut DA, Caaveiro JM, Pybus B, Ruben EA, Ringe D, Petsko GA, Herschlag D. Testing geometrical discrimination within an enzyme active site: constrained hydrogen bonding in the ketosteroid isomerase oxyanion hole. *J. Am. Chem. Soc.* 2008; 130:13696–13708. [PubMed: 18808119]
5. Gilli P, Bertolasi V, Ferretti V, Gilli G. Covalent nature of the strong homonuclear hydrogen bond. Study of the O-H--O System by crystal structure correlation methods. *J. Am. Chem. Soc.* 1994; 116:909–915.
6. Sigala PA, Tsuchida MA, Herschlag D. Hydrogen bond dynamics in the active site of photoactive yellow protein. *Proc. Natl. Acad. Sci. U. S. A.* 2009 in press.
7. Fersht, AR. *Structure and Mechanism in Protein Science*. 2 ed.. New York: W.H. Freeman and Company; 1999.
8. Rodriguez JC, Zeng YH, Wilks A, Rivera M. The hydrogen-bonding network in heme oxygenase also functions as a modulator of enzyme dynamics: Chaotic motions upon disrupting the H-bond network in heme oxygenase from *Pseudomonas aeruginosa*. *J. Am. Chem. Soc.* 2007; 129:11730–11742. [PubMed: 17764179]
9. Bouvignies G, Bernado P, Meier S, Cho K, Grzesiek S, Bruschweiler R, Blackledge M. Identification of slow correlated motions in proteins using residual dipolar and hydrogen-bond scalar couplings. *Proc. Natl. Acad. Sci. U. S. A.* 2005; 102:13885–13890. [PubMed: 16172390]
10. Peralvarez-Marín A, Lorenz-Fonfria VA, Bourdelande JL, Querol E, Kandori H, Padros E. Inter-helical hydrogen bonds are essential elements for intra-protein signal transduction: The role of Asp115 in bacteriorhodopsin transport function. *J. Mol. Biol.* 2007; 368:666–676. [PubMed: 17367807]
11. Derege PJF, Williams SA, Therien MJ. Direct evaluation of electronic coupling mediated by hydrogen-bonds - Implications for biological electron-transfer. *Science*. 1995; 269:1409–1413. [PubMed: 7660123]
12. Reece SY, Hodgkiss JM, Stubbe J, Nocera DG. Proton-coupled electron transfer: the mechanistic underpinning for radical transport and catalysis in biology. *Philos. Trans. Royal Soc. B.* 2006; 361:1351–1364.
13. Hatcher E, Soudackov AV, Hammes-Schiffer S. Proton-coupled electron transfer in soybean lipoxygenase. *J. Am. Chem. Soc.* 2004; 126:5763–5775. [PubMed: 15125669]
14. Boehr DD, Dyson HJ, Wright PE. An NMR perspective on enzyme dynamics. *Chem. Rev.* 2006; 106:3055–3079. [PubMed: 16895318]
15. Henzler-Wildman KA, Lei M, Thai V, Kerns SJ, Karplus M, Kern D. A hierarchy of timescales in protein dynamics is linked to enzyme catalysis. *Nature*. 2007; 450:913–916. [PubMed: 18026087]
16. Karplus M, Kuriyan J. Molecular dynamics and protein function. *Proc. Natl. Acad. Sci. U. S. A.* 2005; 102:6679–6685. [PubMed: 15870208]
17. Michalet X, Weiss S, Jager M. Single-molecule fluorescence studies of protein folding and conformational dynamics. *Chem. Rev.* 2006; 106:1785–1813. [PubMed: 16683755]
18. Wales TE, Engen JR. Hydrogen exchange mass spectrometry for the analysis of protein dynamics. *Mass Spectrom. Rev.* 2006; 25:158–170. [PubMed: 16208684]
19. Agarwal PK, Billeter SR, Rajagopalan PT, Benkovic SJ, Hammes-Schiffer S. Network of coupled promoting motions in enzyme catalysis. *Proc. Natl. Acad. Sci. U. S. A.* 2002; 99:2794–2799. [PubMed: 11867722]
20. Daniel RM, Dunn RV, Finney JL, Smith JC. The role of dynamics in enzyme activity. *Annu. Rev. Biophys. Biomol. Struct.* 2003; 32:69–92. [PubMed: 12471064]
21. Pollack RM. Enzymatic mechanisms for catalysis of enolization: ketosteroid isomerase. *Bioorg. Chem.* 2004; 32:341–353. [PubMed: 15381400]
22. Kim SW, Cha SS, Cho HS, Kim JS, Ha NC, Cho MJ, Joo S, Kim KK, Choi KY, Oh BH. High-resolution crystal structures of delta5-3-ketosteroid isomerase with and without a reaction intermediate analogue. *Biochemistry*. 1997; 36:14030. [PubMed: 9369474]
23. Wu ZR, Ebrahimian S, Zawrotny ME, Thornburg LD, Perez-Alvarado GC, Brothers P, Pollack RM, Summers MF. Solution structure of 3-oxo-delta5-steroid isomerase. *Science*. 1997; 276:415–418. [PubMed: 9103200]

24. Kim DH, Jang DS, Nam GH, Choi G, Kim JS, Ha NC, Kim MS, Oh BH, Choi KY. Contribution of the hydrogen-bond network involving a tyrosine triad in the active site to the structure and function of a highly proficient ketosteroid isomerase from *Pseudomonas putida* biotype B. *Biochemistry*. 2000; 39:4581–4589. [PubMed: 10769113]
25. Kuliopulos A, Mildvan AS, Shortle D, Talalay P. Kinetic and ultraviolet spectroscopic studies of active-site mutants of delta-5-3-ketosteroid isomerase. *Biochemistry*. 1989; 28:149–159. [PubMed: 2706241]
26. Nam GH, Jang DS, Cha SS, Lee TH, Kim DH, Hong BH, Yun YS, Oh BH, Choi KY. Maintenance of alpha-helical structures by phenyl rings in the active-site tyrosine triad contributes to catalysis and stability of ketosteroid isomerase from *Pseudomonas putida* biotype B. *Biochemistry*. 2001; 40:13529–13537. [PubMed: 11695900]
27. Thornburg LD, Henot F, Bash DP, Hawkinson DC, Bartel SD, Pollack RM. Electrophilic assistance by Asp-99 of 3-oxo-Delta 5-steroid isomerase. *Biochemistry*. 1998; 37:10499–10506. [PubMed: 9671521]
28. Kim SW, Joo S, Choi G, Cho HS, Oh BH, Choi KY. Mutational analysis of the three cysteines and active-site aspartic acid 103 of ketosteroid isomerase from *Pseudomonas putida* biotype B. *J. Bacteriol.* 1997; 179:7742. [PubMed: 9401033]
29. Kraut DA, Sigala PA, Pybus B, Liu CW, Ringe D, Petsko GA, Herschlag D. Testing electrostatic complementarity in enzyme catalysis: Hydrogen bonding in the ketosteroid isomerase oxyanion hole. *PLoS Biol.* 2006; 4:e99. [PubMed: 16602823]
30. Petrounia IP, Pollack RM. Substituent effects on the binding of phenols to the D38N mutant of 3-oxo-delta5-steroid isomerase. A probe for the nature of hydrogen bonding to the intermediate. *Biochemistry*. 1998; 37:700–705. [PubMed: 9425094]
31. Gill SC, von Hippel PH. Calculation of protein extinction coefficients from amino acid sequence data. *Anal. Biochem.* 1989; 182:319–326. [PubMed: 2610349]
32. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser crystallographic software. *J. Appl. Crystallogr.* 2007; 40:658–674. [PubMed: 19461840]
33. Murshudov GN, Vagin AA, Dodson EJ. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr. D.* 1997; 53:240–255. [PubMed: 15299926]
34. Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. *Acta Crystallogr. D.* 2004; 60:2126–2132. [PubMed: 15572765]
35. Laskowski RA, Macarthur MW, Moss DS, Thornton JM. Procheck - a program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.* 1993; 26:283–291.
36. Cruickshank DWJ. Remarks about protein structure precision. *Acta Crystallogr. D.* 1999; 55:583–601. [PubMed: 10089455]
37. Turner DL. Binomial solvent suppression. *J. Magn. Reson.* 1983; 54:146–148.
38. Wishart DS, Bigam CG, Yao J, Abildgaard F, Dyson HJ, Oldfield E, Markley JL, Sykes BD. ¹H, ¹³C and ¹⁵N chemical shift referencing in biomolecular NMR. *J. Biomol. NMR.* 1995; 6:135–140. [PubMed: 8589602]
39. Harris TK, Mildvan AS. High-precision measurement of hydrogen bond lengths in proteins by nuclear magnetic resonance methods. *Proteins.* 1999; 35:275–282. [PubMed: 10328262]
40. Jeffrey GA, Yeon Y. The correlation between hydrogen-bond lengths and proton chemical shifts in crystals. *Acta Crystallogr. B.* 1986; 42:410–413.
41. Jencks, WP.; Regenstein, J. Ionization Constants of Acids and Bases. In: Fasman, GD., editor. *Handbooks of Biochemistry and Molecular Biology*. Cleveland: CRC Press; 1976. p. 305-351.
42. McDermott, A.; Rydenour, CF. Proton Chemical Shift Measurements in Biological Solids. In: Grant, DM.; Harris, RK., editors. *Encyclopaedia of Nuclear Magnetic Resonance*. Hoboken: Wiley; 1996. p. 3820-3824.
43. Tolstoy PM, Schah-Mohammed P, Smirnov SN, Golubev NS, Denisov GS, Limbach HH. Characterization of fluxional hydrogen-bonded complexes of acetic acid and acetate by NMR: geometries and isotope and solvent effects. *J. Am. Chem. Soc.* 2004; 126:5621–5634. [PubMed: 15113234]
44. Steiner T. The hydrogen bond in the solid state. *Angew. Chem. Int. Ed.* 2002; 41:48–76.

45. Ichikawa M. Hydrogen-bond geometry and its isotope effect in crystals with OHO bonds - revisited. *J. Molec. Struct.* 2000; 552:63–70.
46. Ubbelohde AR, Gallagher KJ. Acid-base effects in hydrogen bonds in crystals. *Acta Crystallogr.* 1955; 8:71–83.
47. Altman LJ, Laungani D, Gunnarsson G, Wennerstrom H, Forsen S. Proton, deuterium, and tritium nuclear magnetic resonance of intramolecular hydrogen bonds. Isotope effects and the shape of the potential energy function. *J. Am. Chem. Soc.* 1978; 100:8264–8266.
48. Smirnov SN, Golubev NS, Denisov GS, Benedict H, Schah-Mohammedi P, Limbach HH. Hydrogen deuterium isotope effects on the NMR chemical shifts and geometries of intermolecular low-barrier hydrogen-bonded complexes. *J. Am. Chem. Soc.* 1996; 118:4094–4101.
49. Jaravine VA, Cordier F, Grzesiek S. Quantification of H/D isotope effects on protein hydrogen-bonds by ^1H - ^1H and ^1H - ^{13}C couplings and peptide group ^{15}N and ^{13}C chemical shifts. *J. Biomol. NMR.* 2004; 29:309–318. [PubMed: 15213429]
50. Westler WM, Frey PA, Lin J, Wemmer DE, Morimoto H, Williams PG, Markley JL. Evidence for a strong hydrogen bond in the catalytic dyad of transition-state analogue inhibitor complexes of chymotrypsin from proton-triton NMR isotope shifts. *J. Am. Chem. Soc.* 2002; 124:4196–4197. [PubMed: 11960433]
51. Bolvig S, Hansen PE. Isotope effects on chemical shifts as an analytical tool in structural studies of intramolecular hydrogen bonded compounds. *Curr. Org. Chem.* 2000; 4:19–54.
52. Detering C, Tolstoy PM, Golubev NS, Denisov GS, Limbach HH. Vicinal H/D isotope effects in NMR spectra of complexes with coupled hydrogen bonds: Phosphoric acids. *Dokl. Phys. Chem.* 2001; 379:191–193.
53. Liu A, Lu Z, Wang J, Yao L, Li Y, Yan H. NMR detection of bifurcated hydrogen bonds in large proteins. *J. Am. Chem. Soc.* 2008; 130:2428–2429. [PubMed: 18247612]
54. Zhao Q, Li YK, Mildvan AS, Talalay P. Ultraviolet spectroscopic evidence for decreased motion of the active site tyrosine residue of delta 5-3-ketosteroid isomerase by steroid binding. *Biochemistry.* 1995; 34:6562. [PubMed: 7756287]
55. Pollack RM, Thornburg LD, Wu ZR, Summers MF. Mechanistic insights from the three-dimensional structure of 3-oxo-Delta(5)-steroid isomerase. *Arch. Biochem. Biophys.* 1999; 370:9–15. [PubMed: 10496971]
56. Jang DS, Cha HJ, Cha SS, Hong BH, Ha NC, Lee JY, Oh BH, Lee HS, Choi KY. Structural double-mutant cycle analysis of a hydrogen bond network in ketosteroid isomerase from *Pseudomonas putida* biotype B. *Biochem. J.* 2004; 382:967–973. [PubMed: 15228388]
57. Petrounia IP, Blotny G, Pollack RM. Binding of 2-naphthols to D38E mutants of 3-oxo-Delta 5-steroid isomerase: variation of ligand ionization state with the nature of the electrophilic component. *Biochemistry.* 2000; 39:110. [PubMed: 10625485]

Figure 1.

Crystal structure of equilinenin bound at the active site of pKSI^{D103N/D40N}. (A) Sigma-A weighted $2F_o - F_c$ electron density map (contoured at 1.5σ) showing equilinenin bound in the KSI oxanion hole, positioned to receive hydrogen bonds from Tyr16 and Asn103. (B) Superposition of the 1.3 Å resolution pKSI^{D103N/D40N}•equilinenin (carbon atoms in yellow) and 1.1 Å resolution pKSI•equilinenin (PDB code 1OH0, carbon atoms in green).

Figure 2. Downfield region of ¹H NMR spectra of KSI hydrogen bond mutants bound to 3,4-dinitrophenolate. Downfield NMR peaks corresponding to oxyanion hole hydrogen bonded protons are shown in red. All proteins carried the D40N mutation, as explained in the text. The spectrum of tKSI^{D40N}•3,4-dinitrophenolate was previously published (29).

Figure 3.

Schematic depictions of hydrogen bond changes upon deuterium substitution (A) For short hydrogen bonds with a shallow barrier to proton transfer, anharmonicity in the shape of the hydrogen bond potential energy well and the lower zero-point energy for deuterium results in a shorter O-D bond than the original O-H bond by 0.01 – 0.02 Å and a lengthening of the O...O distance by a similar magnitude (2, 45, 47). This figure is for illustrative purposes only and is not meant to convey the shape of the potential energy well for KSI oxyanion hole hydrogen bonds. (B) Conformational changes within a hydrogen bond network upon isotopic substitution. Lengthening of a hydrogen bond upon deuterium substitution leads to shortening of a neighboring hydrogen bond to the same acceptor.

Figure 4.

NMR detection of conformational coupling between the hydrogen bonds donated by Tyr16 and Asp103 in the KSI oxyanion hole. (A) Hydrogen/deuterium isotopologues of hydrogen bonds in the KSI oxyanion hole. (B) ^1H NMR spectra of $\text{tKSI}^{D40N} \cdot 3\text{-F-5-CF}_3\text{-phenolate}$ complexes in solutions containing increasing amounts of D_2O . The spectrum in 0% D_2O was previously published (29).

Scheme 1.

(A) Mechanism of KSI-Catalyzed Steroid Isomerization (B) Schematic Depiction of the Reaction Intermediate Analog, equilenin, Bound at the KSI Active Site.

TABLE 1

Refined Hydrogen Bond O•••O Distances to Equilenin in the KSI Oxyanion Hole

Structure (PDB) ^a	Resolution (Å) ^b	Tyr16-equilenin (Å)	Asp/Asn103-equilenin (Å)
pKSI•equilenin (1OH0)	1.1	2.55	2.54
		2.56	2.48
pKSI ^{D103N/D40N} •equilenin (3FZW)	1.3	2.50	2.90
		2.52	2.88

^aThe asymmetric unit of both structures contained two independently-refined monomers.^bCoordinate errors are estimated to be 0.03 Å (1OH0) and 0.05 Å (3FZW) based on (36).

TABLE 2

Crystallographic Data and Refinement Statistics for pKSI^{D103N/D40N}•equilenin

Data Set	
Resolution Range (Å)	55.05–1.32
Space Group	C2
a, Å	86.9
b, Å	71.2
c, Å	50.5
α, °	90.0
β, °	89.9
γ, °	90.0
Number Unique Reflections	68,149
Completeness, %	94.0
Multiplicity	2.9
R_{merge} % ^a	8.8
$I/\sigma_{\text{overall}}$ ($I/\sigma_{\text{high res}}$)	10.3 (2.4)
Refinement Statistics	
No. Residues	250
No. Waters	239
R_{work} % ^b	14.9
R_{free} % ^c	18.3
rmsd bond, Å	0.023
rmsd angle, °	2.1

$$^a R_{\text{merge}} = \frac{\sum_{\text{hkl}} \sum_i |I(\text{hkl})_i - [I(\text{hkl})]|}{\sum_{\text{hkl}} \sum_i I(\text{hkl})_i}$$

$$^b R_{\text{work}} = \frac{\sum_{\text{hkl}} |F(\text{hkl})_{\text{O}} - [F(\text{hkl})_{\text{C}}]|}{\sum_{\text{hkl}} F(\text{hkl})_{\text{O}}}$$

^c R_{free} was calculated exactly as R_{work} where $F(\text{hkl})_{\text{O}}$ were taken from 5% of the data not included in refinement.