Published in final edited form as:

Women Health. 2012 July ; 52(5): 472–484. doi:10.1080/03630242.2012.687442.

HIV Status and Gender: A Brief Report from Heterosexual Couples in Thailand

Li Li, Ph.D.^{a,*}, Li-Jung Liang, Ph.D.^b, Sung-Jae Lee, Ph.D.^a, and Shu C. Farmer, Ph.D.^a

^aUniversity of California, Los Angeles, Semel Institute Center for Community Health

^bUniversity of California, Los Angeles, Department of Medicine Statistics Core

Abstract

Although the impact of HIV falls on both partners of a married couple, the burden of stress may not be necessarily shared evenly. This study examined the relations among HIV status, gender and depressive symptoms among 152 married or cohabitating couples living with HIV in the Northern and Northeastern regions of Thailand. Depressive symptoms were assessed using a 15-item depressive symptom screening test that was developed and used previously in Thailand. Among the 152 couples, 59% were couples in which both members were people living with HIV (PLH) (seroconcordant; both PLH couples), 28% had only female members with HIV (serodiscordant; female PLH couples), and 13% had only male members with HIV (serodiscordant; male PLH couples). The prevalence of depressive symptoms between seroconcordant and serodiscordant groups was similar. However, female PLH reported significantly higher levels of depressive symptoms, regardless of their partners' HIV status. Future prevention programs focusing on serodiscordant couples should be planned to target HIV risk, as well as emphasis on mental health, with a particular focus on women's increased susceptibility to negative mental health outcomes.

Keywords

HIV; serodiscordant couples; depressive symptoms; Thailand

INTRODUCTION

As HIV has become a manageable illness with the recent advancement of antiretroviral treatment (ART), the focus of survival has gradually shifted to improving everyday lives for people living with HIV (PLH) and their family members. More and more married and cohabitating couples living with HIV, either those in which both partners (seroconcordant) or only one partner is affected (serodiscordant), face challenges associated with HIV daily, which could affect their mental health and quality of life (Van Der Straten et al., 1998; Persson, Barton, & Richards, 2006; de Walque, 2007; Rispel et al., 2011). Previous studies have documented that serodiscordant status adds pressure to a relationship in terms of fears of transmission, guilt and anxiety which could be emotionally exhausting, taking on caretaking abilities, barriers to pregnancy, status disclosure, and HIV stigma that affects the relationship (Van der Straten, 1998; Beckerman, 2002; Kalichman et al., 2002; Klein et al., 2003; Allen et al., 2003; Bunnell et al., 2005; Persson, Barton, & Richards, 2006; Beyeza-Kashesya et al., 2009; Talley & Bettencourt, 2010; Eyawo et al., 2010; Pereira et al., 2011).

*Corresponding Author: Li Li, Ph.D., UCLA Semel Institute - Center for Community Health, 10920 Wilshire Blvd., Suite 350, Los Angeles, CA 90024, U.S.A. (310) 794-2446, lililili@ucla.edu.

Conflict of Interests

The authors declare that they have no conflict of interest.

These unique hurdles create added stress that could affect mental health (Siegal & Lekas, 2002; Pomeroy, Green, & Van Laningham, 2002; Gordon-Garofolo & Rubin, 2004; Jarman et al., 2005; Bradley, Remien, & Dolezal, 2008; Plattner & Meiring, 2006).

Although the impact of HIV falls on both partners of a married or cohabitating couple, the burden of stress may not be necessarily shared evenly. Past studies have been done on the assumption that men are thought to be the index case in most relationships, and most social marketing and awareness campaigns have been focused on men (Eyawo et al., 2010). Although some studies on this topic have focused on the HIV status of the couples (Allen et al., 2003; Persson, Barton, & Richards, 2006; Persson & Richards, 2008; Persson et al., 2009; Bouhnik et al., 2007; Eaton et al., 2009; Kelly et al., 2011) or compiled them as a whole (Hernandez et al., 2009), very few studies have included gender in the equation. A study by Kennedy and colleagues reported that women were the most stressed when in a discordant relationship, regardless of their HIV status (Kennedy, 1995); and according to Stevens, women were most fearful when they were HIV-infected and navigating a serodiscordant relationship (Stevens & Galvao, 2007). This study examined the relationships between HIV status, gender and depressive symptoms among 152 married or cohabitating couples living with HIV in Thailand. We hypothesized that women would report higher depressive symptoms, compared to men. A better understanding of couples' mental health status in relation to gender and HIV status may help service providers develop more targeted, effective coping strategies for the population.

METHODS

Study Setting and Sample

The participants in this study were derived from a randomized controlled family intervention trial conducted in the Northern and Northeastern regions of Thailand (Li et al., 2010). The data were collected in 2007 from four district hospitals in the two regions (two district hospitals per region). Eligibility criteria included: 1) having at least one PLH in the household; 2) having at least one family member who was aware of the HIV status of the PLH; 3) having at least one school-aged child living in the household; and 4) the ability to provide written informed consent from the PLH and family members. Project recruiters screened 484 HIV-affected families at the participating hospitals. During monthly support group meetings, flyers were handed out to potential participants. Those who were interested in participating in the study were directed to contact a recruiter who screened and determined the eligibility of the participant. Of the 484 families screened, 431 families (89%) met the eligibility criteria. Of the 431 eligible families, 410 consented to participate (refusal rate = 4.9%). The PLH and one adult family member, preferably the other member of the couple, from each family were recruited. Because many family members were not the PLH's husband or wife, or cohabiting partners, 152 couples were identified and included in the study.

Following written and signed informed consent, a trained interviewer administered the assessment to the participants using Computer Assisted Personal Interview (CAPI). All participants received 300 Baht (equivalent to approximately \$10 US dollars at the time of the study) for their assessment participation. Approval of this study was obtained from the Institutional Review Board (IRB) at the University of California, Los Angeles, and the Thailand Ministry of Public Health Ethical Review Committee for Research in Human Subjects.

Instruments

Depressive symptoms were assessed using a 15-item depressive symptom screening test that was developed and used previously in Thailand (Thai Department of Mental Health, 2004). These questions asked about problems that had bothered participants in the past week (e.g., feeling depressed most of the time; feelings of hopelessness or worthlessness; loss of self-confidence), with response categories from 0 (not at all) to 3 (usually [5–7 days a week]). A summative composite scale was developed, with a range of 0 to 45 and an excellent internal consistency (Cronbach's $\alpha = 0.91$). We also collected demographic information, including age and gender of the participant, and household income.

Data Analysis

Demographic characteristics were summarized using descriptive statistics and frequencies. Groups were compared using ANOVA or chi-square test for continuous and categorical variables, respectively. A linear mixed-effects regression model with family-level random effects was used to compare the depressive symptom scores in the seroconcordant group (both PLH couples) with the two serodiscordant groups (female PLH couples and male PLH couples). Pre-selected covariates of interest included age, gender, annual household income, group (male PLH couples, female PLH couples, and both PLH), and group-by-gender interaction. These covariates were included in the analyses based on our *a priori* knowledge that they might be potential confounders in models using depressive symptoms as an outcome. The family-level random intercept was included to account for dependence within families. The comparisons of interest were made through model contrasts, and Akaike's Information Criterion (AIC) fit statistic was used to compare models. All statistical analyses were conducted with the SAS System for Windows (Version 9.2), and all the graphs were generated using the publicly available statistical software R (R Development Core Team, 2008).

RESULTS

Of the 152 couples in this study, about 80% were HIV-positive; 59% were couples in which both members were PLH (seroconcordant; both PLH couple), 28% of the couples had female members with HIV (serodiscordant; female PLH couple), and 13% of the couples had male members with HIV (serodiscordant; male PLH couple) (Table 1). Sixty-five percent of male participants vs. 83.5% of female participants were aged 40 years or younger. About half of the study participants reported their annual income to be 35,000 Baht or less (~\$ 1100 US dollars). The average age of the participants in the seroconcordant group was 36.7 years (SD = 5.0), which was younger than that of the participants in the other two groups (mean = 37.5 vs. 38.9 years for male PLH couple vs. female PLH couple groups, respectively; $P = 0.12$). The mean difference in age within a family for the female PLH couple group was higher. The average annual household income for the seroconcordant group was about 55,000 Baht (~\$1770 US dollars), and that for the male PLH couple group was about 600 Baht (~\$19 US dollars) higher than that for the female PLH couple group. No significant differences were observed for age and annual household income among the three groups.

Results from the mixed-effects regression model indicated that age and household income were not significantly associated with depressive symptoms ($P = 0.83$, and 0.10 , respectively) (Table 2). The distribution of depressive symptom scores between serodiscordant and seroconcordant groups were similar (Figure 1a; estimated difference = 0.809 , SE = 1.071 ; $P = 0.45$), and the distributions of depressive symptoms did not differ significantly between the two serodiscordant groups (Figure 1b; estimated difference = 3.021 , SE = 1.698 ; $P = 0.08$). In the final model where the group-by-gender interaction was

added, we found that women reported significantly more depressive symptoms than men in the seroconcordant group (both PLH couples: estimated difference=2.657, SE=1.117; $P=0.019$). Similarly, in the serodiscordant group (Female PLH couples), women reported significantly more depressive symptoms than their male partners (Estimated difference = 5.174, SE = 1.683; $P=0.003$). In the serodiscordant group (Male PLH couples), men reported slightly more depressive symptoms than their female partners (Estimated difference = -1.638, SE = 2.425; $P=0.501$), but the difference was not statistically significant (Table 2).

DISCUSSION

To our knowledge, this is one of the first studies to examine depressive symptoms in PLH by gender, as well as serostatus of the couples. Our findings have important implications for future prevention strategies focusing on the mental health of couples living with HIV. Female PLH in Thailand reported significantly more depressive symptoms, regardless of their partners' HIV status. This finding highlights that the burden of the HIV epidemic may differentially affect the sexes, perhaps reflecting women's and men's different roles in household and family activities. HIV poses a significant burden and increased demand of care giving for many families globally. In many parts of the world, the unequal division of labor within the household may put an additional burden of care on women, who predominantly fulfill the role as primary caregivers (MacNeil, 1996; Olenja, 1999; Bunting, 2001; Joseph & Bhatti, 2005; Kipp et al., 2007). For example, when a husband becomes ill, the wife provides care and takes on additional duties to support the family (Bharat & Aggleton, 1999; Joseph & Bhatti, 2005; Revenson et al., 2005). When women fall sick, other women in the family care for them and their children. The evidence from caregiving studies suggests that female caregivers are more likely to experience burden than are male caregivers (Miller & Cafasso, 1992; Chappell & Reid, 2002; Joseph & Bhatti, 2005; Revenson et al., 2005; Pinquart & Sorensen, 2006; Vithayachockitikhun, 2006; Kipp et al., 2007; Golobof et al., 2011). Women often take on the responsibility for household work and for caring for children, elderly and sick family members (Moser, 1989; du Guerny & Sjoberg, 1993; Revenson et al., 2005; Kipp et al., 2007; Dawuni, 2008). The situation could become more stressful when the caregiving woman herself is infected, as she struggles to balance her own health concerns with the demands of her family and relatives. Previous studies reported that, in spite of their HIV infection, family members expected women to remain in the role of primary family and child caregiver (Hackl et al., 1997; Bunting, 2001; Joseph & Bhatti, 2005; Kipp et al., 2007). The veritable role of gender prescribing a woman's place in society magnifies their vulnerability to the extent that they may be more negatively affected by the HIV epidemic.

Our study underscores the important notion that while HIV infection plays a role in adding stress in a family or couple, it is not spread evenly when gender is taken into account. In our study, the highest level of depressive symptoms was reported by the serodiscordant couples in which the male partner was infected by HIV. This finding is not surprising, because men in many cultures, including Thailand, are likely to engage in income-generating work that is often valued and recognized. They normally earn more money, are better educated and better represented in their communities than women (Whyte, 2001; Vithayachockitikhun, 2006). Their illness can directly affect family productivity and income; thus the quality of life of all members suffers (Russell, 2004). Men in this study reported the lowest level of depressive symptoms when their wife or partner was infected but they were not (serodiscordant female PLH). One interpretation could be that when women are infected, they may demand less care from their husbands or partners. They might sacrifice their caring needs for other family members, whereas when men are infected, the burden falls more heavily on their female partners.

Women face many unique additional HIV-related challenges, compared to their male counterparts. These unique challenges are often intensified by ways in dealing with stress and illness. Generally speaking, gender is a factor in the exposure, experience, and coping of stress (Davis, Burleson, & Kruszewski, 2009; Zwicker & DeLongis, 2010). For example, women are more vulnerable than men to be depressed (Boyd & Weissman, 1981; Baum & Grunberg, 1991; Wisniewski et al., 2005; Remien et al., 2006; Valverde et al., 2007; Gordillo et al., 2009; Pereira & Canavarro, 2011). Women are at a disadvantage in terms of being more prone to life stressors impacting their quality of life. Studies are replete with findings of women possessing lower QOL scores than men (e.g., poorer psychosocial adaptation, low levels of psychological adjustment/well-being, and higher visibility of psychopathological symptoms), suggesting gender inequity in HIV infection (Cederfjall et al., 2001; Wisniewski et al., 2005; Chandra et al., 2008; Solomon et al., 2008; Pereira & Canavarro, 2011). On the other hand, having depressive symptoms may impair women's abilities to cope with their own infection and provide care for the illness of a partner.

We identified several limitations in this study. The first limitation pertained to the eligibility criteria. To be eligible for the study, PLH had to have at least one family member who knew about their HIV status. These selection criteria limited the generalizability of the study findings to those whose HIV status was not disclosed. Also, although we included potential confounders (e.g., age, gender, income), a range of other possible factors that might have been related to depressive symptoms were not included in the study, potentially resulting in uncontrolled confounding. Second, the study focused on depressive symptoms rather than depression, so the findings may not be relevant to a diagnosis of major depression. Third, because we had fewer couples with male PLH, compared to female PLH, we may have had inadequate statistical power to detect meaningful differences between these two groups as statistically significant. Finally, given that this was a cross-sectional study, temporal relations could not be assessed (eg, depressive symptoms may have preceded HIV infection) so that we cannot make casual inferences about depressive symptoms and serodiscordance for HIV.

Despite these limitations, our findings have significant implications for future prevention programs focusing on serodiscordant couples and depressive symptoms. Couples in serodiscordant relationships face many ongoing challenges, which also heighten the threat to their mental health. Our study showed that among this vulnerable group, women are even more susceptible to depressive symptoms, compared to men. This key finding is consistent with other studies; demonstrating a common parallel on gender differences and HIV. HIV-positive women experience high levels of distress (Sanders, 2008; Jarman et al., 2005), initiating a desire for psychological protection in their relationships (Jarman et al., 2005). A study on HIV-negative women suggests gender inequality arises in serodiscordant relationships, escalating women's vulnerability to HIV infection (Persson & Richards, 2008). And finally the risk of HIV progression has been correlated to HIV-positive women with extensive depressive symptoms (Antelman et al., 2007). Studying gender differences within individual responses to HIV has implications for care, support and treatment programs, especially for addressing the needs of women. To simply emphasize the traditional caring role of women in AIDS patient care might lead to wider gender inequality, leaving women even more vulnerable to the HIV epidemic. Future prevention programs focusing on serodiscordant couples should be planned to target HIV risk, as well as place a heavy emphasis on mental health, with a particular focus on women's increased susceptibility to negative mental health outcomes.

Acknowledgments

This article was completed with the support of the National Institute of Nursing Research (grant NINR R01-NR009922) and the National Institute of Mental Health (grant NIMH 5K01MH085503). We thank our research coordinators, hospital directors, and health officers in Chiang Rai province (Mae Chan and Chiang Saen district hospitals) and Nakhon Ratchasima province (Pak Chong and Khonburi district hospitals). We thank our collaborators at the Thai Ministry of Public Health, Bureau of Epidemiology for their contributions to the study.

References

- Allen S, Meinen-Derr J, Kautzman M, Zulu I, Trask S, Fideli U, Musonda R, Kasolo F, Gao F, Haworth A. Sexual behavior of HIV discordant couples after HIV counseling and testing. *AIDS*. 2003; 17:733–740. [PubMed: 12646797]
- Antelman G, Kaaya S, Wei R, Mbwanbo J, Msamanga GI, Fawzi WW, Fawzi MC. Depressive symptoms increase risk of HIV disease progression and mortality among women in Tanzania. *J Acquir Immune Defic Syndr*. 2007; 44:470–77. [PubMed: 17179766]
- Baum A, Grunberg NE. Gender, stress, and health. *Health Psychol*. 1991; 10:80–85. [PubMed: 2055213]
- Beckerman NL. Couples coping with discordant HIV status. *AIDS Patient Care STDS*. 2002; 16:55–59. [PubMed: 11874636]
- Beyeza-Kashesya J, Kaharuza F, Mirembe F, Neema S, Ekstrom AM, Kulane A. The dilemma of safe-sex and having children: Challenges facing HIV sero-discordant couples in Uganda. *Afr Health Sci*. 2009; 9:2–12. [PubMed: 20842236]
- Bharat S, Aggleton P. Facing the challenge: Household responses to HIV/AIDS in Mumbai, India. *AIDS Care*. 1999; 11:31–34. [PubMed: 10434981]
- Bouhnik AD, Preau M, Lert F, Peretti-Watel P, Schiltz MA, Obadia Y, Spire B. VESPA Study Group. Unsafe sex in regular partnerships among heterosexual persons living with HIV: Evidence from a large representative sample of individuals attending outpatients services in France (ANRS-EN12-VESPA Study). *AIDS*. 2007; 21(Suppl 1):S57–62. [PubMed: 17159589]
- Boyd JH, Weissman M. Epidemiology of affective disorders. *Arch Gen Psychiatry*. 1981; 28:1039–46. [PubMed: 7025783]
- Bradley MV, Remien RH, Dolezal C. Depression Symptoms and Sexual HIV Risk Behavior Among Serodiscordant Couples. *Psychosom Med*. 2008; 70:186–91. [PubMed: 18256344]
- Bunnell RE, Nassozi J, Marum E, Mubangizi J, Malamba S, Dillon B, Kalule J, Bahizi J, Musoke N, Mermin JH. Living with discordance: knowledge, challenges, and prevention strategies of HIV-discordant couples in Uganda. *AIDS Care*. 2005; 17:999–1012. [PubMed: 16176896]
- Bunting SM. Sustaining the relationship: women's caregiving in the context of HIV disease. *Health Care Women Int*. 2001; 22:131–48. [PubMed: 11813792]
- Cederfjall C, Langius-Eklöf A, Lidman K, Wredling R. Gender differences in perceived health-related quality of life among patients with HIV infection. *AIDS Patient Care STDS*. 2001; 15:31–9. [PubMed: 11177586]
- Chandra PS V, Satyanarayana A, Satishchandra P, Satish KS, Kumar M. Do men and women with HIV differ in their quality of life? A study from South India. *AIDS Behav*. 2009; 13:110–7. [PubMed: 18654846]
- Chappell NL, Reid RC. Burden and well-being among caregivers: examining the distinction. *Gerontologist*. 2002; 42:772–80. [PubMed: 12451158]
- Davis, MC.; Burleson, MH.; Kruszewski, DM. Gender: Its relationship to stressor exposure, cognitive appraisal/coping processes, stress responses, and health outcomes. In: Contrade, RJ.; Baum, A., editors. *The Handbook of Stress Science: Biology, Psychology, and Health*. New York, NY: Springer Publishing Company; 2009. p. 247–61.
- Dawuni JJ. The gendered face of HIV/AIDS: The move towards policy implementation in Ghana. *Studies in Gender and Development in Africa*. 2008; 1:78–94.
- de Walque D. Sero-Discordant Couples in Five African Countries: Implications for Prevention Strategies. *Popul Dev Rev*. 2007; 33:501–23.

- du Guerny J, Sjoberg E. Inter-relationship between gender relations and the HIV/AIDS epidemic: some possible considerations for policies and programmes. *AIDS*. 1993; 7:1027–34. [PubMed: 8397938]
- Eaton LA, West TV, Kenny DA, Kalichman SC. HIV transmission risk among HIV seroconcordant and serodiscordant couples: Dyadic processes of partner selection. *AIDS Behav*. 2009; 13:185–95. [PubMed: 18953645]
- Eyawo O, de Walque D, Ford N, Gakii G, Lester RT, Mills EJ. HIV status in discordant couples in sub-Saharan Africa: a systematic review and meta-analysis. *Lancet Infect Dis*. 2010; 10:770–7. [PubMed: 20926347]
- Golobof A, Weine S, Bahromov M, Luo J. The roles of labor migrants' wives in HIV/AIDS risk and prevention in Tajikistan. *AIDS Care*. 2011; 23:91–7. [PubMed: 21218281]
- Gordillo V, Fekete EM, Platteau T, Antoni MH, Schneiderman N, Nöstlinger C. Eurosupport Study Group. Emotional support and gender in people living with HIV: Effects on psychological well-being. *J Behav Med*. 2009; 32:523–31. [PubMed: 19543823]
- Gordon-Garofalo VL, Rubin A. Evaluation of a psychoeducational group for seronegative partners and spouses of persons with HIV/AIDS. *Res Soc Work Pract*. 2004; 14:14–26.
- Hackl KL, Somlai AM, Kelly JA, Kalichman SC. Women living with HIV/AIDS: the dual challenges of being a patient and caregiver. *Health Soc Work*. 1997; 22:53–62. [PubMed: 9021418]
- Hernandez V, del Romero J, Garcia S, Rodríguez C, del Amo J, Castilla J. Reducing sexual risk behavior among steady heterosexual serodiscordant couples in testing and counseling program. *Sex Transm Dis*. 2009; 36:621–8. [PubMed: 19955873]
- Jarman M, Walsh S, de Lacey G. Keeping safe, keeping connected: A qualitative study of HIV-positive women's experiences of partner relationships. *Psychol Health*. 2005; 20:533–51.
- Joseph EB, Bhatti RS. Psychosocial problems and coping patterns of HIV seropositive wives of men with HIV/AIDS. *Soc Work Health Care*. 2005; 39:29–47. [PubMed: 15774383]
- Kalichman SC, Rompa D, Luke W, Austin J. HIV transmission risk behaviors among HIV positive persons in serodiscordant relationships. *Int J STD AIDS*. 2002; 13:677–82. [PubMed: 12396537]
- Kelly A, Kupul M, Frankland A, Worth H, Nosi S, Mek A, Kepa B, Walizopa L, Emori R, Pirpir L, Akuani F, Cangah B, Siba P. Living serodiscordantly in Papua New Guinea: Practices of HIV-positive people on ART by serostatus of regular heterosexual partner. *AIDS Care*. 2011; 23:734–40. [PubMed: 21390883]
- Kennedy CA. Gender differences in HIV-related psychological distress in heterosexual couples. *AIDS Care*. 1995; 7(Suppl 1):33–8.
- Kipp W, Tindyebwa D, Karamagi E, Rubaale T. How much should we expect? Family caregiving of AIDS patients in rural Uganda. *J Transcult Nurs*. 2007; 18:358–65. [PubMed: 17911576]
- Klein J, Pena JE, Thornton MH, Sauer MV. Understanding the Motivations, Concerns, and Desires of Human Immunodeficiency Virus 1 Serodiscordant Couples Wishing to Have Children Through Assisted Reproduction. *Obstet Gynecol*. 2003; 101:987–94. [PubMed: 12738162]
- Li L, Lee SJ, Jiraphongsa C, Khumtong S, Iamsirithaworn S, Thammawijaya P, Rotheram-Borus MJ. Improving Health and Mental Health for People Living with HIV/AIDS: 12-month Assessment of a Behavioral Intervention in Thailand. *Am J Public Health*. 2010; 100:2418–25. [PubMed: 20966372]
- MacNeil JM. Use of culture care theory with Baganda women as AIDS caregivers. *J Transcult Nurs*. 1996; 7:14–20. [PubMed: 8974412]
- Miller B, Cafasso L. Gender differences in caregiving: fact or artifact? *Gerontologist*. 1992; 32:447–53.
- Moser C. Gender planning in the Third World: meeting practical and strategic gender needs. *World Dev*. 1989; 11:1801.
- Olenja JM. Assessing community attitude towards home-based care for people with AIDS (PWAs) in Kenya. *J Community Health*. 1999; 24:187–199. [PubMed: 10399651]
- Pereira M, Dattilio FM, Canavarro MC, Narciso I. Couple-focused interventions for HIV-serodiscordant couples during transition to motherhood. *Contemp Fam Ther*. 2011; 33:143–60.

- Pereira M, Canavarro MC. Gender and age differences in quality of life and the impact of psychopathological symptoms among HIV-infected patients. *AIDS Behav.* 2011; 15:1857–69. [PubMed: 21431413]
- Persson, A.; Richards, W.; Barton, D.; Reakes, K. Monograph 1/2009. Vol. 2. Sydney: National Centre in HIV Social Research, University of New South Wales; 2009. Men and women living heterosexually with HIV. The straightpoz study.
- Persson A, Richards W. Vulnerability, gender and “proxy negativity”: Women in relationships with HIV-positive men in Australia. *Soc Sci Med.* 2008; 67:799–807. [PubMed: 18562064]
- Persson, A.; Barton, D.; Richards, W. Monograph 2/2006. Vol. 1. Sydney: National Centre in HIV Social Research, University of New South Wales; 2006. Men and women living heterosexually with HIV. The straightpoz study.
- Pinquart M, Sorensen S. Gender differences in caregiver stressors, social resources, and health: An updated meta-analysis. *J Gerontol B Psychol Sci Soc Sci.* 2006; 61B:33–45.
- Pomeroy EC, Green DL, Van Laningham L. Couples Who Care: The Effectiveness of a Psychoeducational Group Intervention for HIV Serodiscordant Couples. *Res Soc Work Pract.* 2002; 12:238–52.
- Plattner IE, Meiring N. Living with HIV: The psychological relevance of meaning making. *AIDS Care.* 2006; 18:241–5. [PubMed: 16546785]
- R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing; Vienna, Austria: 2008. URL <http://www.R-project.org>
- Remien RH, Exner T, Kertzner RM, Ehrhardt AA, Rotheram-Borus MJ, Johnson MO, Weinhardt LS, Kittel LE, Goldstein RB, Pinto RM, Morin SF, Chesney MA, Lightfoot M, Gore-Felton C, Dodge B, Kelly JA. NIMH Healthy Living Project Trial Group. Depressive symptomatology among HIV-positive women in the era of HAART: A stress and coping model. *Am J Community Psychol.* 2006; 38:275–85. [PubMed: 16967343]
- Revenson, TA.; Abraido-Lanza, AF.; Majerovitz, SD.; Jordan, C. Couples coping with chronic illness: What's gender got to do with it?. In: Revenson, TA.; Kayser, K.; Bodenmann, G., editors. *Couples coping with stress: Emerging perspectives on dyadic coping.* Washington, DC: American Psychological Association; 2000. p. 137–56.
- Rispel LC, Cloete A, Metcalf CA, Moody K, Caswell G. ‘It [HIV] is part of the relationship’: exploring communication among HIV-serodiscordant couples in South Africa and Tanzania. *Cult Health Sex.* 2012; 14:257–68. [PubMed: 22077668]
- Russell S. The economic burden of illness for households in developing countries: A review of studies focusing on malaria, tuberculosis, and human immunodeficiency virus/acquired immunodeficiency syndrome. *Am J Trop Med Hyg.* 2004; 71(Suppl 2):47–155.
- Sanders LB. Women’s Voices: The lived experience of pregnancy and motherhood after diagnosis with HIV. *J Assoc Nurses AIDS Care.* 2008; 19:47–57. [PubMed: 18191768]
- Siegal K, Lekas H. AIDS as a chronic illness: psychosocial implications. *AIDS.* 2002; 16(Suppl 4):69–76.
- Solomon S, Venkatesh KK, Brown L, Verma P, Cecelia AJ, Daly C, Mahendra VS, Cheng N, Kumarasamy N, Mayer KH. Gender-related differences in quality of life domains of persons living with HIV/AIDS in South India in the era prior to greater access to antiretroviral therapy. *AIDS Patient Care STDS.* 2008; 22:999–1005. [PubMed: 19072106]
- Stevens PE, Galvao L. “He won’t use condoms”: HIV-infected women’s struggles in primary relationships with serodiscordant partners. *Am J Public Health.* 2007; 97:1015–22. [PubMed: 17463377]
- Talley AE, Bettencourt BA. A relationship-oriented model of HIV-related stigma derived from a review of the HIV-affected couples literature. *AIDS Behav.* 2010; 14:72–86. [PubMed: 19037718]
- Thai Department of Mental Health. [Accessed June 10, 2011] Depression Screening Test, Ministry of Public Health, Thailand. 2004. <http://www.dmh.go.th/test/depress/asheet.asp?qid=1>
- Tukey, JW. *Exploratory Data Analysis.* Addison-Wesley; Reading, Massachusetts: 1977.
- Valverde EE, Purcell DW, Waldrop-Valverde D, Malow R, Knowlton AR, Gómez CA, Farrell N, Latka MH. for the INSPIRE Study Team. Correlates of depression among HIV-positive women

- and men who inject drugs. *J Acquir Immune Defic Syndr*. 2007; 46(Suppl 2):S96–100. [PubMed: 18089990]
- Van der Straten A, Vernon KA, Knight KR, Gómez CA, Padian NS. Managing HIV among serodiscordant heterosexual couples: serostatus, stigma and sex. *AIDS Care*. 1998; 10:533–48. [PubMed: 9828951]
- Van der Straten A. Couple Communication, sexual coercion and HIV risk reduction in Kigali, Rwanda. *AIDS*. 1998; 9:935–44. [PubMed: 7576330]
- Vithayachockitikhun N. Family caregiving of persons living with HIV/AIDS in Thailand: caregiving burden, and outcome measure. *Int J Nurs Pract*. 2006; 12:123–8. [PubMed: 16674778]
- Whyte, SR. Community participation in essential national health research (part 2). In: Neufeld, V.; Arbuthnot, N., editors. *Forging Links for Health Research: Perspectives from the Council on Health Research for Development*. Ottawa, ON, Canada: International Development Research Centre; 2001. p. 81-108.
- Wisniewski AB, Apel S, Selnes OA, Nath A, McArthur JC, Dobs AS. Depressive symptoms, quality of life, and neuropsychological performance in HIV/AIDS: The impact of gender and injection drug use. *J Neurovirol*. 2005; 11:138–43. [PubMed: 16036792]
- Zwicker A, DeLongis A. Gender, stress, and coping. *Handbook of Gender Research in Psychology*. 2010; 6:495–515.

Figure 1. Boxplots of depressive symptoms (a) serodiscordant vs. seroconcordant couples and (b) male PLH vs., female PLH vs. both. A boxplot (Tukey, 1977) is a convenient way of graphically depicting groups of numerical data through their five-number summaries (minimum, lower quartile, median, upper quartile, and maximum).

Table 1**Baseline Characteristics of Study Sample**

	N (%)
Number of Couples	152
Couple Type	
Male PLH Couple	20 (13.2)
Female PLH Couple	42 (27.6)
Both PLH	90 (59.2)
Age (Years)	
Male	
30	14 (9.2)
31–40	85 (56.0)
41–50	43 (28.3)
> 50	10 (6.6)
Female	
30	27 (17.9)
31–40	99 (65.6)
41–50	23 (15.2)
> 50	2 (1.3)
Annual Income	
<=15000 Baht (~\$480)	34 (22.4)
15001–35000 Baht (~\$480–\$1100)	40 (26.3)
35001–55000 Baht (~\$1000–\$1770)	51 (33.6)
> 55000 Baht (~\$1770)	27 (17.8)

Table 2

Estimates and Standard Errors (SE) from Mixed-Effects Regression Model for Depressive Symptoms

Outcome=Depressive Symptoms	Estimate (SE)	P
Age	0.015 (0.070)	0.831
Household Income (per 1,000)	−0.015 (0.009)	0.099
Main Effects		
Gender (Female-Male)	2.064 (1.063)	0.054
Group		0.202
Male PLH Couples	2.319 (1.544)	
Female PLH Couples	−0.702 (1.163)	
Both PLH Couples	Ref	
Interaction: Gender-by-Group		0.071
Comparisons of Interest		
Serodiscordant vs. Seroconcordant Couples	0.809 (1.071)	0.452
Male PLH Couples: Female vs. Male	−1.638 (2.425)	0.501
Female PLH Couples: Female vs. Male	5.174 (1.683)	0.003
Both PLH Couples: Female vs. Male	2.657 (1.117)	0.019