Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2012 July ; 21(7): 1143–1148. doi:10.1158/1055-9965.EPI-11-1030.

Correlation of LINE-1 methylation levels in patient-matched buffy coat, serum, buccal cell and bladder tumor tissue DNA samples

Dana van Bommel¹, Petra Lenz², Linda M Liao¹, Dalsu Baris¹, Lawrence R. Sternberg³, Andrew Warner³, Alison Johnson⁴, Michael Jones⁵, Masatoshi Kida⁶, Molly Schwenn⁷, Alan R. Schned⁸, Debra T. Silverman¹, Nathaniel Rothman¹, and Lee E. Moore^{1,9}

¹Occupational and Environmental Epidemiology Branch, Epidemiology and Biostatistics Program, National Cancer Institute, NIH, DHHS, Bethesda, Maryland, USA

²Division of Cancer Epidemiology and Genetics, Science Applications International Corporation-Frederick, Inc., National Cancer Institute-Frederick, Frederick, MD, 21702

³Pathology/Histotechnology Laboratory, Laboratory Animal Sciences Program, Science Applications International Corporation-Frederick, Inc., National Cancer Institute-Frederick, Frederick, MD

⁴Vermont Cancer Registry, Burlington, Vermont, USA, Department of Pathology

⁵Department of Pathology and Laboratory Medicine, Maine Medical Center Portland, Maine, USA

⁶Department of Pathology, University of Vermont College of Medicine, Burlington, Vermont, USA

⁷Maine Cancer Registry, Augusta, Maine, USA

⁸Dartmouth Medical School, Hanover, New Hampshire USA

Abstract

Background—Evidence suggests that global methylation levels in blood cell DNA may be a biomarker for cancer risk. To date, most studies have used genomic DNA isolated from blood or urine as a surrogate marker of global DNA methylation levels in bladder tumor tissue.

Methods—A subset of 50 bladder cancer cases was selected from the New England Bladder Case-Control Study. Genomic DNA was isolated from buffy coat, buccal cells, serum and formalin fixed paraffin embedded tissue for each participant. DNA methylation at four CpG sites within the *LINE-1* repetitive element was quantified using pyrosequencing and expressed as a mean methylation level across sites.

Results—Overall, the mean percent (%) *LINE-1* 5-methylcytosine (%5MeC) level was highest in serum (80.47% ± 1.44) and lowest in bladder tumor DNA (61.36% ± 12.74); and levels varied significantly across tissue types (p=0.001). An inverse association between *LINE-1* mean %5MeC and tumor stage (p=0.001) and grade (p=0.002) was observed. A moderate correlation between patient-matched serum and buffy coat DNA *LINE-1* %5-MeC levels was found (r=0.32, p=0.03) but levels were uncorrelated among other matched genomic DNA samples.

Conclusions—The mean promoter *LINE-1* %5MeC measurements were correlated between buffy coat and serum DNA samples. No correlation was observed between genomic DNA sources and tumor tissues; however a significant inverse association between tumor percent *LINE-1* methylation and tumor stage/grade was found.

⁹To whom correspondence may be addressed: Lee E. Moore, Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, 6120 Executive Blvd., EPS Rm. 8102, Bethesda, MD 20852, Phone: 301-496-6427; FAX: 301-402-1819; moorele@mail.nih.gov.

Impact—*LINE-1* methylation in measured case blood DNA did not reflect that observed in bladder tumor tissue but may represent other factors associated with carcinogenesis.

Introduction

Global hypomethylation is thought to contribute to carcinogenesis by increasing genome instability while gene-specific hypermethylation is associated with gene silencing (1,2). Bladder cancer studies have shown that chromosomal instability is widespread, occurs early, and is associated with chromosome 9 aberrations, the first and most frequently altered chromosome in bladder cancer. One of the earliest epigenetic changes associated with genomic instability is believed to be global hypomethylation. Methylation at CpG sites within Long Interspersed Nucleotide Repeats (*LINE-1*) in serum, buffy coat, buccal cell, and urine DNA have been analyzed as biomarkers of total global methylation status in cancer studies (3-8), however, the usefulness of genomic DNA *LINE-1* methylation as a biomarker remains uncertain, and is the subject of continued investigation (9,10).

High performance capillary electrophoresis of global methylation and PCR-based analysis of *LINE-1* repeat sequences in blood DNA have been used as biomarkers of bladder cancer risk. In three case-control studies, inverse associations were observed using post-diagnostic blood samples (4,5,11). It is unclear whether differences in global methylation among cases and controls in genomic DNA represent an epigenetic biomarker of recent exposures, a general marker of cancer susceptibility (such as chromosomal alterations), or differences related to carcinogenesis. It is also unknown how methylation levels in tumor DNA correlate with those in surrogate tissues, and within the same individuals.

To understand associations between *LINE-1* methylation levels across tissue types and in bladder tumor tissue, we compared levels from matched DNA specimens to determine if % 5-methyl cytosine (%5MeC) levels in the *LINE-1* promoter region correlated in DNA isolated from buccal cells, buffy coat, serum, and formalin-fixed paraffin embedded (FFPE) bladder tumor tissue from the same individuals. A secondary aim was to determine which peripheral DNA source best reflected *LINE-1* methylation levels in tumor tissue.

Methods

Study Population

The New England Bladder Cancer Study (NEB) is a population-based case-control study conducted in Maine, Vermont, and New Hampshire (12). Cases, aged 30-79 years with histologically confirmed urothelial carcinoma, were included (12). Rapid case ascertainment was conducted through pathology departments and hospital/state registries. We interviewed 1,213 bladder cancer cases. Participants were interviewed at home by a trained interviewer using a detailed computer-assisted personal interview (CAPI). Information collected included general demographics, occupational and residential histories, tobacco and hair coloring use, family history of cancer, and diet. Biological samples were collected, including blood and buccal cells. FFPE bladder tumor tissue was obtained from hospital pathologists. Fifty cases were selected from Maine and Vermont from whom buffy coat, serum, buccal cell, and FFPE tumor tissue was available.

Laboratory Methods

Genomic DNA was extracted from samples using standard phenol-chloroform methods. FFPE tissue blocks, considered representative of the patient's primary tumor, and hematoxylin and eosin (H&E)-stained diagnostic slides, were obtained from hospitals where cases were treated. Serial, five-micron tissue sections were placed onto charged slides. One slide was stained with H&E and scanned on an Aperio ScanScope® XT at 20x (Aperio

Technologies). Digital images were transferred into an Aperio Spectrum™ database and harvest regions identified by a study pathologist via Aperio ImageScope® annotation software. Tumor tissue was macrodissected and placed into 1.5ml centrifuge tubes. 175 μ L of microwave retrieval solution (0.1% guanidium thiocyanate and 0.1M NaOH), (13) was added, heated to 100°C for 10 min followed by 10 minutes at 100°C using the microwave module of a Milestone RH-1 tissue processor (Hacker Instruments and Industries, Inc.). Samples were cooled for 5 min, placed into a Thermomixer R (Eppendorf, North America) for 10 min at 65 °C at 1,400 RPM. One μ L of RNase (100 mg/mL) was added, samples were incubated for 30 min at 55 °C and 5 min at 65 °C without shaking. Samples were centrifuged (5 min/14,000g /4°C). DNA was isolated from supernatant using the phenol-based AutoGenprep 245T DNA extraction kit according to instructions. DNA yield and purity were determined by NanoDrop 1000 spectrophotometer (NanoDrop technologies). Bisulfite modification of DNA (0.25-0.5 μ g/ μ L) was conducted using the EZ DNA Methylation Kit (Zymo Research, Orange CA), according to the manufacturer's instructions, and stored at - 20 °C. An unmethylated cytosine site was included in the PCR amplification product as a positive, internal bisulfite conversion control.

LINE-1 methylation levels were quantified using pyrosequencing (EpigenDx Worcester, MA) (14-16), to examine the methylation status at four CpG sites in the LINE-1 promoter (-492 to -419 bp from ATG). Bisulfite-treated DNA was eluted to 20 μ L. Each 50 μ L PCR contained the bisulfite treated DNA, 10X PCR buffer, 3.0 mM MgCl₂, 200 μ M dNTPs, 0.2 μ M primers, 1.25 U DNA polymerase (HotStar, Qiagen Inc., Alameda, CA) 1.25 U. A biotinylated primer was used to capture one single-stranded DNA template for pyrosequencing, (17,18). PCR products (10 μ L) were sequenced using the Pyrosequencing PSQ96 HS System (Biotage, Kungsgatan, Sweden). Loci were analyzed as a T/C SNP using QCpG software (Biotage, Kungsgatan, Sweden). Each plate contained unmethylated, partially methylated, and heavily methylated DNAs (SssI treated DNA) for quality control.

Statistical Analysis

STATA (version 10.0) was used for data analysis. The study was designed to have 90% power to detect a 1.27% difference in mean methylation levels with a standard deviation of ± 3.0 and 80% power to observe a correlation of $r=0.4$ between mean methylation (N=50). The quality and quantity of extracted DNA from FFPE tumor tissue limited our bladder tumor sample size to N=34. Within that sample size (N=34), we had 80% power to detect a 1.27% difference in mean methylation levels with a SD ± 3.0 and 80% power to observe a correlation of $r=0.5$ between mean methylation levels.

Triplicate %5MeC measurements were averaged across four CpG dinucleotides in the LINE-1 promoter and expressed as a methylation index (percent 5-Methyl Cytosine, %5MeC). Initially, samples with a coefficient of variation (CV) > 10% (n=3) or a percent unconverted internal cytosine control > 15% (n=19) were excluded. Sensitivity analyses were conducted excluding samples with a CV > 5% (n=5) and those with unconverted cytosine >15% (internal cytosine control; n=19). The difference in calculated mean values was <1% for both criterion. Therefore, we included all samples regardless of the percent unconverted cytosine and excluded only samples with CVs >10% from the analysis. Data were distributed non-normally. Both non parametric (Kruskal-Wallis) and parametric tests (Student's t tests/ordered logistic regression) were conducted on log transformed *LINE-1* methylation levels. Correlation coefficients were evaluated using Pearson's test on transformed continuous. Correlations using other cut-points (medians, tertiles, and quartiles) were also examined.

Results

A sufficient quality and quantity of genomic DNA was extracted from all buccal, serum, and buffy coat samples; and 34 FFPE tumor tissue blocks (68%). Pyrosequencing data were obtained on 184 samples. Three samples were excluded (buffy coat: $n=1$; serum: $n=2$) with CVs $>10\%$. Table 1 summarizes case characteristics and the mean percent *LINE-1* methylation by genomic DNA source. Case demographics were similar to NEB cases overall (12). More subjects were white (94%), male (80%), 55 years old (70%), and smokers (70%).

Initially, we analyzed the mean *LINE-1* methylation CVs by DNA source. The lowest variance was observed for buccal cell (CV= 4.24 ± 0.89 ; range: 1.39-5.82), and buffy coat DNA (CV= 4.54 ± 0.65 ; range: 3.20-6.14) (data not shown). Variation was highest for tumor tissue (CV= 4.26 ± 1.59 ; range: 2.17-8.71) and serum DNA (CV= 5.06 ± 4.51 ; range: 2.65-12.06). The mean CVs of *LINE-1* methylation levels across DNA sources per individual were similar, with (CV= 4.41 ± 0.63 ; range: 3.52-7.04) and without the inclusion of tumor DNAs (CV= 4.39 ± 0.53 ; range: 3.35-5.63).

Mean *LINE-1* methylation levels were highest in serum ($80.47\% \pm 1.44$) followed by buffy coat ($79.36\% \pm 1.34$) and buccal cell DNA ($77.09\% \pm 2.09$) (Table 1). Bladder tumor DNA had the lowest *LINE-1* %5MeC and the most variation across samples ($61.36\% \pm 12.74$). Mean *LINE-1* methylation levels differed significantly across DNA source both with ($p=0.001$) and without ($p=0.0001$) inclusion of tumor levels. When stratified by case characteristics, buffy coat *LINE-1* methylation levels differed by gender, with males having higher methylation than females ($p=0.03$). We did not observe significant differences by the other DNA sources analyzed; however tumor DNA *LINE-1* methylation levels were inversely associated with stage ($p=0.001$) and grade ($p=0.002$).

In Table 2, correlations between *LINE-1* methylation levels are presented between DNA pairs extracted from different sources, and by tumor stage/grade. By DNA source, a moderate correlation was observed between patient-matched serum and buffy coat DNAs ($r=0.32$, $p=0.03$). *LINE-1* methylation levels were uncorrelated among other matched genomic DNA samples, or with tumor tissue. Grouping methylation levels by medians, tertiles, or quartiles, did not improve correlations between DNA sources (data not shown). Tumor tissue %5MeC levels showed a strong inverse association with tumor stage ($r=-0.56$, $p=0.0005$) and grade ($r=-0.52$, $p=0.002$).

Discussion

CpG methylation in repetitive regions of the genome is becoming widely used to estimate global methylation (reviewed in 10). Yet, few studies have compared %5MeC within repetitive genomic regions in DNA from matched sources. The NEB study provided an opportunity to evaluate these comparisons. In this study of 50 patients, associations were not observed between *LINE-1* mean %5MeC measured across DNA types. No correlation was observed between genomic DNA sources and tumor tissues. *LINE-1* %5MeC in buffy coat DNA was higher among males compared to females; and a significant inverse association between tumor %5MeC levels and tumor stage and grade was found. This difference between males and females is consistent with several recent studies of *LINE-1* in blood DNA (4-6,19-21), but not all (22). Similar to published studies, associations between smoking or age and tissue methylation were not observed (4-6,11, 19, 23,24).

Our analysis also supports reports that levels of *LINE-1* methylation in tumor tissue decreases with stage and grade. One study by Choi et al., (25) examined different interspersed repeats (*LINE-1* and *ALU*) and tandem repeats (Sat- α , NBL-2, and D4Z4) in

blood among 10 individually-matched tissue DNAs in 4 cancer-free individuals undergoing autopsy. In contrast to work presented here, the investigators reported *LINE-1* levels were consistent between tissues (n=10) and individuals (n=4). An inverse association between *LINE-1* %5MeC and matched adjacent normal urothelium and bladder tumor samples from cases was also observed; however correlations were not provided. The study by Choi et al., was one of the first studies demonstrating that tumor tissue methylation levels were lower than normal urothelial tissue and blood cell DNA. This study was one of the first to report substantial variability and lack of comparability in methylation outcome using different DNA sequences to estimate global methylation.

In the current study, the strongest correlation in methylation levels was observed between buffy coat and serum, both originating from blood. A recent study compared different blood cell types among 48 matched DNAs and noted similar moderate correlations within *LINE-1* methylation of blood cell subtypes (26). It has been suggested that a proportion of serum DNA is actually blood cell DNA although the exact proportion remains uncertain in bladder cancer patients. Other literature suggests that serum DNA originates from lysed, malignant, haemopoietic apoptotic, and/or necrotic circulating cells in cancer patients (27).

The potential for *LINE-1* methylation as a predictive biomarker of bladder cancer risk is an active area of research. Three case-control studies report associations between blood cell methylation and bladder cancer risk using post-diagnostic samples; but it is unclear whether a similar relationship is observed before diagnosis (8, 28). Such pre-diagnostic studies will elucidate whether hypomethylation in *LINE-1* is an independent risk factor for bladder cancer, or a question of reverse causality. Environmental exposures such as smoking may impact *LINE-1* methylation and its utility as a biomarker. Smoking modified the association between *LINE-1* levels and bladder cancer risk in two studies (4, 11), but not in a third study (5). These data suggest the need for researchers to continue modeling environmental exposures (when possible) during development of disease specific biomarkers.

It has been hypothesized that *LINE-1* hypomethylation in case blood DNA may reflect levels in bladder tumor tissue. Our study suggests that they are not directly related. Rather, it appears that hypomethylation of *LINE-1* in genomic DNA reflects other factors associated with bladder carcinogenesis such as repositioning and enhancement of gene expression. Hypomethylation of a specific *LINE-1* promoter has been shown to activate an alternate transcript of the *MET* oncogene in bladder tumors and normal bladder urothelium (29,30). It is unclear whether the *LINE-1* methylation levels in genomic DNA actually change expression of specific functional *LINE-1* elements and other cancer genes; however these findings provide a mechanism through which *LINE-1* methylation could modify cancer susceptibility.

Limitations of this study include lack of inclusion of healthy controls, limiting the analysis of bladder cancer risk factors such as smoking exclusively to cases. Healthy tissue nor exfoliated cells were collected in this cohort, therefore we were unable to compare *LINE-1* methylation levels across the four types of DNA isolated from healthy controls. This study was sufficiently powered to evaluate the study aims of examining modest correlations between paired DNA samples, however a larger sample of cases would have allowed for additional subgroup analyses. For example, the low proportion of non-smoking cases limited power to estimate differences by smoking status. Strengths of this study include the study design; 50 patients with matched genomic and tumor tissue DNAs for comparison. Isolation of DNA from FFPE tissue did limit our analysis to 34 tissue case samples. Yet even with the smaller sample size, we had sufficient power to compare %5MeC from measured in tissue to %5MeC measured in buffy coat, buccal, and serum. Another strength of the study was the use of Pyrosequencing; an accurate and reproducible method used to measure %5MeC. The

role of incomplete bisulfite conversion and high CVs across replicates was evaluated through sensitivity analyses, strengthening the overall interpretation of our results.

To our knowledge, this is the first study to systematically compare *LINE-1* methylation across these specific types of matched DNAs. The mean promoter *LINE-1* %5MeC measurements were correlated between buffy coat and serum DNA samples but not with tumor or buccal cell DNA. Although genomic *LINE-1* levels were not directly related to those observed in bladder tumor tissue, current case-control studies appear to support an association between global hypomethylation in blood and cancer risk; however the possibility of reverse causality cannot be eliminated. Future studies should include case-control and prospective studies of pre-diagnostic blood samples, to elucidate temporal associations between *LINE-1* methylation and bladder cancer risk.

Acknowledgments

Funding: This project has been funded in whole or in part with federal funds from the National Cancer Institute, National Institutes of Health, under Contract No. HHSN261200800001E. Mention of trade names, commercial products, or organizations does not imply endorsement by the U.S. Government.

References

1. Issa JP. Aging, DNA methylation and cancer. *Crit Rev Oncol Hematol*. 1999; 32:31–43. [PubMed: 10586353]
2. Esteller M, Herman JG. Cancer as an epigenetic disease: DNA methylation and chromatin alterations in human tumors. *J Pathol*. 2002; 196:1–7. [PubMed: 11748635]
3. Pufulete M, Al-Ghnam R, Leather AJ, Appleby P, Gout S, Terry C, et al. Folate status, genomic DNA hypomethylation, and risk of colorectal adenoma and cancer: a case control study. *Gastroenterology*. 2003; 124(5):1240–8. [PubMed: 12730865]
4. Wilhelm CS, Kelsey KT, Butler R, Plaza S, Gagne L, Sense MS, et al. Implications of *LINE1* methylation for bladder cancer risk in women. *Clin Cancer Res*. 2010; 16(5):1682–9. [PubMed: 20179218]
5. Cash HL, Tao L, Yuan JM, Marsit CJ, Houseman EA, Xiang YB, et al. *Line-1* hypomethylation is associated with bladder cancer risk among nonsmoking Chinese. *Int J Cancer*. 2011 Mar 28.
6. Hsuing DT, Marsit C, Houseman E, Eddy K, Furniss C, McClean M, et al. Global methylation level in whole blood as a biomarker in head and neck squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev*. 2007; 16:108–14. [PubMed: 17220338]
7. Hou L, Wang H, Sartori S, Gawron A, Lissowska J, Bollati V, et al. Blood leukocyte DNA hypomethylation and gastric cancer risk in a high-risk Polish population. *Int J Cancer*. 2010; 127:1866–74. [PubMed: 20099281]
8. Choi JY, James SR, Link PA, McCann SE, Hong CC, Davis W, et al. Association between global DNA hypomethylation in leukocytes and risk of breast cancer. *Carcinogenesis*. 2009; 30:1889–97. [PubMed: 19584139]
9. Terry MB, Delgado-Cruzata L, Vin-Raviv N, Wu HC, Santella RM. DNA methylation in white blood cells Association with risk factors in epidemiologic studies. *Epigenetics*. 2011; 6(7):828–837. [PubMed: 21636973]
10. Nelson HN, Marsit CJ, Kelsey KT. Global methylation in exposure biology and translational medical science. *Environ Health Perspectives*. Jun.2011 epub.
11. Moore LE, Pfeiffer RM, Poscablo C, Real FX, Kogevinas M, Silverman D, et al. Genomic DNA as a biomarker for bladder cancer susceptibility in the Spanish Bladder Cancer Study: a case-control study. *Lancet Oncol*. 2008; 9:359–366. [PubMed: 18339581]
12. Baris D, Karagas M, Verrill, Johnson A, Andrew AS, Marsit CJ. A case-control study of smoking and bladder cancer risk: emergent patterns over time. *J Natl Cancer Inst*. 2010; 101(22):1553–61. [PubMed: 19917915]

13. Shi SR, Datar R, Liu C, Wu L, Zhang Z, Cote RJ, et al. DNA extraction from archival formalin-fixed, paraffin-embedded tissues: heat-induced retrieval in alkaline solution. *Histochem Cell Biol.* 2004; 122:211–18. [PubMed: 15322858]
14. Yang AS, Estecio MR, Doshi K, Kondo Y, Tajara EH, Issa JP. A simple method for estimating global DNA methylation using bisulfite PCR of repetitive DNA elements. *Nucleic Acids Res.* 2004; 32:e38. [PubMed: 14973332]
15. Liu T, Zhang X, So CK, et al. Regulation of Cdx2 expression by promoter methylation, and effects of Cdx2 transfection on morphology and gene expression of human esophageal epithelial cells. *Carcinogenesis.* 2007; 28:488–96. [PubMed: 16990345]
16. Brakensiek K, Wingen LU, Langer F, Kreipe H, Lehmann U. Quantitative high-resolution CpG island mapping with Pyrosequencing reveals disease-specific methylation patterns of the CDKN2B gene in myelodysplastic syndrome and myeloid leukemia. *Clin Chem.* 2007; 53:17–23. [PubMed: 17095538]
17. England R, Pettersson M. Pyro Q-CpG™ : quantitative analysis of methylation in multiple CpG sites by Pyrosequencing. *Nature Methods.* 2005; 2:1–2.
18. Royo JL, Hidalgo M, Ruiz A. Pyrosequencing protocol using a universal biotinylated primer for mutation detection and SNP genotyping. *Nat Protoc.* 2007; 2:1734–9. [PubMed: 17641638]
19. Zhu ZZ, Hou L, Bollati V, Tarantini L, Marinelli B, Cantone L, et al. Predictors of global methylation levels in blood DNA of healthy subjects: a combined analysis. *Int J Epidemiol.* 2010 Sep 15.
20. El-Maari O, Walier M, Behne F, van Uum J, Singer H, Diaz-Lacava A, et al. Methylation at global LINE-1 repeats in human blood are affected by gender but not by age or natural hormone cycles. *PLoS One.* 2001; 6(1):e16252.
21. Zhang FF, Cardarelli R, Carroll J, Fulda KG, Kaur M, Gonzalez K, et al. Significant differences in global genomic DNA methylation by gender and race/ethnicity in peripheral blood. *Epigenetics.* 2011; 6(5):623–9. [PubMed: 21739720]
22. Chalitchagorn K, Shuangshoti S, Hourpai N, et al. Distinctive pattern of LINE-1 methylation level in normal tissues and the association with carcinogenesis. *Oncogene.* 2004; 23:8841–46. [PubMed: 15480421]
23. Rusiecki JA, Baccarilli A, Bollati V, Tarantini L, Moore LE, Bonefeld-Jorgensen EC. Global DNA hypomethylation is associated with high serum-persistent organic pollutants in Greenlandic Inuit. *Environ Health Perspect.* 2008; 116(1):1547–1552. [PubMed: 19057709]
24. Figueirido JC, Grau MV, Wallace K, Levine AJ, Shen L, Hamdan R, et al. Global DNA hypomethylation (LINE-1) in the normal colon and lifestyle characteristics and dietary and genetic factors. *Cancer Epidemiol Biomarkers Prev.* 2009; 18(4):1041–9. [PubMed: 19336559]
25. Choi SH, Worswick S, Byun HM, Shear T, Soussa JC, Wolff EM, et al. Changes in DNA methylation of tandem DNA repeats are different from interspersed repeats in cancer. *Int J Cancer.* 2009; 125:723–729. [PubMed: 19437537]
26. Wu HC, Delgado-Cruzata L, Flom JD, Kappil M, Reffis JS, Liao Y, et al. Global methylation profiles in DNA from different blood cell types. *Epigenetics.* 2011; 6(1):76–85.
27. Aung KL, Board RE, Ellison G, Donald E, Ward T, Clack G, et al. Current status and future potential of somatic mutation testing from circulating free DNA in patients with solid tumours. *HUGO J.* 2010; 4:11–21. [PubMed: 22132062]
28. Schultz WA. Do DNA methylation changes also occur in blood? *Lancet Oncol.* 2008; 9(4):312–313. [PubMed: 18374285]
29. Wolff EM, Hyang-Min B, Han HF, Sharma S, Nickols PW, Siegmund KD, et al. Hypomethylation of a LINE-1 promoter activates an alternate transcript of the MET oncogene in bladders with cancer. *PLoS Genet.* 2010; 6(4):e1000917. [PubMed: 20421991]
30. Wolff EM, Chihara Y, Pan F, Weisenberger DJ, Siegmund KD, Sugano K, et al. Unique DNA methylation patterns distinguish non-invasive and invasive urothelial cancers and establish an epigenetic field defect in premalignant tissue. *Cancer Res.* 2010; 70(20):8169–7178. [PubMed: 20841482]

Table 1
Mean LINE1 Methylation Levels (%) in Matched Tissue DNAs by Patient/Tumor Characteristics

Characteristics of Cases	Buccal		Bufty Coat		Serum		Bladder Tumor Tissue		p-value *** (without tissue)
	Cases	Mean±SD	Cases	Mean±SD	Cases	Mean±SD	Cases	Mean±SD	p-value ***
Sex	50	77.09±2.09	49	79.36±1.34	48	80.47±1.44	34	61.36±12.7	0.0001
Female	10	76.15±2.21	9	78.51±1.24	9	79.72±1.30	6	58.86±15.33	0.002
Male	40	77.3±2.07	40	79.56±1.30	39	80.64±1.43	28	61.9±12.37	0.0001
p-value **		0.10		0.03		0.08		0.56	
Age	19	77.14±1.83	18	79.23±1.47	18	80.05±1.43	15	63.51±13.78	0.0001
<55	16	77.41±2.21	16	79.72±0.78	16	80.82±1.57	8	64.5±7.15	0.0001
55-64	10	76.88±2.10	10	79.02±1.85	10	80.76±1.49	7	52.65±16.5	0.006
65-74	5	76.27±3.01	5	79.44±1.28	4	80.21±0.18	4	62.27±2.91	0.06
p-trend **		0.44		0.95		0.38		0.38	
Race	47	77.07±2.13	46	79.45±1.33	45	80.55±1.43	31	60.58±13.01	0.001
White	3	77.34±1.71	3	78.08±0.87	3	79.19±1.23	3	69.42±5.68	0.18
Non-white		0.82		0.09		0.11		0.29	
p-value **									
Smoking	14	76.95±1.69	13	79.06±1.17	13	80.68±1.38	10	59.22±9.40	0.0001
Never	16	77.57±1.71	16	79.52±1.51	15	80.30±1.37	9	59.85±17.66	0.0002
Former *	20	76.79±2.60	20	79.44±1.34	20	80.45±1.59	15	63.70±11.70	0.0001
Current		0.72		0.47		0.70		0.48	
p-trend **									
Ever *	36	77.13±2.25	36	79.48±1.40	35	80.39±1.48	24	63.26±13.98	0.0001
p-value **		0.80		0.35		0.53		0.74	
Tumor									

Characteristics of Cases	Buccal		Buffy Coat		Serum		Bladder Tumor Tissue		p-value *** (without tissue)
	Cases	Mean±SD	Cases	Mean±SD	Cases	Mean±SD	Cases	Mean±SD	p-value ***
Grade									
1	29	76.77±2.42	29	79.35±1.33	28	80.61±1.31	20	66.31±9.03	0.0001
2	8	77.47±2.24	8	78.63±1.67	8	80.89±2.10	5	63.71±2.91	0.005
3	11	77.97±1.47	11	79.82±0.94	11	79.74±1.25	8	50.66±15.65	0.0003
Unknown	1	74.97	1	80.75	1	81.02	1	36.35	N/A
Missing	1		-		-		-		N/A
p-trend**		0.09		0.52		0.14		0.002	
Tumor Stage									
Ta	39	76.98±2.19	39	79.25±1.38	38	80.68±1.45	26	64.98±8.62	0.0001
T1	6	78.43±1.14	6	79.48±10.70	6	79.29±0.92	4	53.33±18.57	0.02
T2a-T3a	4	76.43±1.94	4	80.27±1.31	4	80.21±1.41	4	45.86±17.32	0.15
missing	1	75.68	-		-		-		0.14
p-trend**		0.78		0.17		0.14		0.001	

* includes one occasional smoker

** p-value for Student's t test for dichotomous variables, p-value for ordered logistic regression analysis of categorical variables.

*** p-value for Kruskal Wallis Test

Pearson's correlation coefficients (p-values) of mean LINE1 methylation levels in case-matched DNAs and tumor stage and grade

Table 2

DNA Source	Buccal Cell	Buffy Coat	Serum	Bladder Tumor Tissue	Stage
Buffy Coat	0.04 (0.76)				
Serum	0.04 (0.78)	0.32 (0.03)			
Bladder Tumor Tissue	-0.08 (0.67)	-0.13 (0.46)	0.08 (0.64)		
Tumor Characteristic					
Stage	0.04 (0.78)	0.20 (0.17)	-0.22 (0.14)	-0.56 (0.0005)	
Grade	0.24 (0.09)	0.09 (0.52)	-0.22 (0.14)	-0.52 (0.002)	0.69 (<0.00001) *

* Tumor stage and grade classifications are predicted to be highly correlated