Published in final edited form as:

Immunogenetics. 2011 July ; 63(7): 397–408. doi:10.1007/s00251-011-0533-9.

Molecular bases of genetic diversity and evolution of the immunoglobulin heavy chain variable region (*IGHV*) gene locus in leporids

Ana Pinheiro^{1,2}, Dennis Lanning³, Paulo C. Alves^{1,2}, Rose G. Mage⁴, Katherine L. Knight³, Wessel van der Loo¹, and Pedro J. Esteves^{1,5}

¹Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal

²Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, 4169-007 Porto, Portugal

³Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, IL 60153, USA

⁴Laboratory of Immunology, NIAID, National Institutes of Health, Bethesda, MD 20892, USA

⁵Centro de Investigação em Tecnologias da Saúde, IPSN, CESPU, 4585-116 Gandra, Portugal

Abstract

The rabbit has long been a model for studies of the immune system. Work using rabbits contributed both to the battle against infectious diseases such as rabies and syphilis, and to our knowledge of antibodies' structure, function, and regulated expression. With the description of rabbit Ig allotypes, the discovery of different gene segments encoding immunoglobulins became possible. This challenged the “one gene-one protein” dogma. The observation that rabbit allotypic specificities of the variable regions were present on IgM and IgG molecules also led to the hypothesis of Ig class switching. Rabbit allotypes contributed to the documentation of phenomena such as allelic exclusion and imbalance in production of allelic gene products. During the last 30 years, the rabbit Ig allotypes revealed a number of unique features, setting them apart from mice, humans and other mammals. Here, we review the most relevant findings concerning the rabbit *IGHV*. Among these are the preferential usage of one *VH* gene in *VDJ* rearrangements, the existence of trans-species polymorphism in the *IGHV* locus revealed by serology and confirmed by sequencing *IGHV* genes in *Lepus*, the unusually large genetic distances between allelic lineages and the fact that the antibody repertoire is diversified in this species only after birth. The Whole Genome Sequence of rabbit, plus re-sequencing of additional strains and related genera, will allow further evolutionary investigations of antibody variation. Continued research will help define the roles that genetic, allelic and population diversity at antibody loci may play in host-parasite interactions.

Keywords

leporids; rabbit; *IGHV*; evolution; immunoglobulin repertoire

Historic perspective

Studies of the rabbit immune system have greatly contributed to our knowledge of the structure, function and regulation of antibodies. During the late 19th and early 20th centuries, the use of rabbit in immunological research was crucial to the development of the rabies vaccine by Louis Pasteur and to the battle against syphilis (reviewed in Dubiski, 1987). In the first half of the 20th century, the foundations of molecular immunology were laid with almost exclusive use of the rabbit. The antigenic polymorphism of serum components was described in this species as early as 1902, although the antigens involved were characterized only 50 years later (see Kelus and Gell, 1967). In 1956, Oudin demonstrated and defined allotypy of immunoglobulins in the rabbit. Allotypes were proven to be hereditary traits in rabbit by Dubiski and coworkers (1959) prior to the establishment of the two major linkage groups “*a*” and “*b*”, corresponding to the H and light chain loci, respectively (Oudin, 1960), and even before the heterodimeric structure of the antibody molecule was clearly established.

The existence of genetic markers of the different gene fragments encoding the antibody H chain was unique to this species and the rabbit Ig allotypes challenged the “one gene-one protein” dogma. Indeed, Todd (1963) and others found that rabbit allotypic specificities of the variable regions were present on both IgM and IgG molecules, suggesting that the same *VH* gene segment can be translocated to different constant region genes. This observation opened the road to confirmation of the concept of multiple “germline” *VH* gene segments that can be joined to a limited number of genes encoding the different heavy chain constant regions (Dreyer and Bennett, 1965), and led to the hypothesis of Ig class switching (Kearny et al., 1976; van der Loo et al., 1979; reviewed in Severinson et al., 1982). Hamers and co-workers (1966) reported cis-expression of the *VH* and *CH* genes by using allelic markers on both protein domains. The markers in the CH2 domain were re-discovered by Dubiski (1969) and renamed the e14 and e15 allotypes. Mage and co-workers (1971) reported the first of a number of crossing-over events observed by laboratories during breeding. Findings of genetic recombination between the genes controlling the rabbit *VH* and *CH* (estimated recombination frequency of 0.1%; (Mage, 1979; Kelus and Steinberg, 1991)) confirmed the model of Dreyer and Bennett (1965).

Documentation of the current *V-D-J-C* model therefore originated through studies of the rabbit allotypes. In addition to genetic linkage of *VH* and *CH* genes, and the expression of apparently identical *VH* regions on different classes of Ig, it was through studies of rabbit immunoglobulin markers that phenomena such as allelic exclusion and imbalance in production of allelic gene products were described (e.g. Davie et al., 1971; Loo and Kelus, 1978; Schmale et al., 1969; Wolf et al, 1971). During the last 30 years, the rabbit Ig allotypes revealed a number of unique features, setting them apart from mice, humans and other mammals. Among these are the preferential usage of one *VH* gene in *VDJ* rearrangements, the evidence for the existence of trans-species polymorphism in the *IGHV*, *IGHG* and *IGKC* loci, the unusually large genetic distances between the allelic lineages and the fact that the antibody repertoire is diversified in this species only after birth.

Lagomorph taxonomy

The order LAGOMORPHA comprises two families: *OCHOTONIDAE* and *LEPORIDAE*. The family *OCHOTONIDAE* includes 30 species restricted to the genus *Ochotona* (Pikas). The family *LEPORIDAE* can be divided into two groups: hares and rabbits. According to Chapman and Flux (1990), the hare group encompasses a single genus, *Lepus*, whereas the rabbit group includes 10 genera (*Brachylagus*, *Bunolagus*, *Caprolagus*, *Nesolagus*, *Oryctolagus*, *Pentalagus*, *Poelagus*, *Pronolagus*, *Romerolagus* and *Sylvilagus*) and 25 species (Alves and Hacklander,

2008). There is not a consensus estimate of the divergence time between rabbits and hares. Analyses of several nuclear and mitochondrial markers suggest that the genera *Oryctolagus* and *Lepus* diverged between 6 and 20 Mya (Biju-Duval et al., 1991; Halanich and Robinson, 1999; Mathee et al., 2004) (Fig. 1), although fossil data suggests an earlier divergence (between 2.5 and 3.5 million years ago (Mya) (Lopez-Martinez, 2008).

The genus *Oryctolagus* is monospecific (*O. cuniculus*). In the Iberian Peninsula, where the fossil record suggests the European rabbit originated (Pages, 1980; Lopez-Martinez, 1989, 2008; Corbet, 1994), two morphologically differentiated subspecies of European rabbit have been distinguished: *O. cuniculus algirus* and *O. cuniculus cuniculus* (Cabrera, 1914). These two subspecies diverged ~1.8 Mya (reviewed in Carneiro et al., 2009). *O. c. algirus* inhabits the southwestern Iberian Peninsula, while *O. c. cuniculus* is present in the northeastern Iberian Peninsula. *O. c. cuniculus* later expanded its range North towards France, likely after the last glacial peak (Queney et al., 2001), where it still remains present. The Pleistocene Ice ages forced many temperate species to retreat into southern refugia, leading to high levels of diversity and endemism in these areas (Hewitt 1996). The Iberian Peninsula was one of these refugia in Europe (Taberlet et al., 1998). This is reflected in the significantly lower genetic diversity of the wild French *O. c. cuniculus* populations compared to the Iberian populations (van der Loo et al., 1991, 1999; Queney et al., 2001; Esteves et al., 2004; Ferrand and Branco, 2007; Surridge et al., 2008; Carneiro et al., 2011). The European rabbit gene pool has been manipulated through a recent single domestication event of French origin, and therefore all domestic rabbits belong to subspecies *O. c. cuniculus* (reviewed in Ferrand and Branco 2007, Carneiro et al., 2011). Today, through mediated anthropic dispersal, the European rabbit is present in Continental Europe, England, Australia, New Zealand, North and South America, and North Africa.

For other leporids, the study of immunoglobulin genes is mainly restricted to the genera *Lepus* and *Sylvilagus* (Cazenave et al., 1977, Teherani and Mandy, 1976a, 1976b; Teherani et al., 1979; Teherani et al., 1982; Bouton and van der Loo, 1997; Esteves et al., 2002a, 2005, 2006). *Lepus* is a polytypic, cosmopolitan genus, comprising 24 to 30 currently recognized species (Corbet and Hill, 1980; Flux and Angermann, 1990; Alves and Hacklander, 2008), that, like *Sylvilagus*, most probably originated in North America (Lopez-Martinez 2008, Mathee et al., 2004). *Lepus* species that have been used in immunogenetic studies include *L. americanus*, *L. timidus*, *L. granatensis* and *L. europaeus*. *L. americanus* is distributed throughout North America and occupies a basal position in the *Lepus* group (Mathee et al., 2004). *L. granatensis* is endemic to the Iberian Peninsula, covering the whole peninsula except the Northeast where it is replaced by *L. europaeus*, which spans a wide range throughout Europe, with several introduced populations in different regions, like South America, Australia, New Zealand and several islands including Barbados, Reunion and Falklands (reviewed in Alves and Hacklander, 2008). *L. timidus* is a Palearctic relict species with an extensive distribution area in Europe and Asia, from Ireland and Scandinavia in the west, across Siberia, Mongolia and China to the northern islands of Japan in the east. *Sylvilagus* is restricted to the American continent and comprises seventeen species (reviewed in Alves and Hacklander, 2008). Within this genus, the most studied species is *S. floridanus*, a species that occupies Southern and Western North America.

Rabbit preferentially expresses *IGHV1*

The rabbit *IGHV* allotypes are highly divergent and behave as Mendelian alleles (Oudin, 1956, 1960; Dubiski et al., 1959; Dray et al., 1963; Kim and Dray, 1972; Mage et al., 1984a). Serological surveys of domestic rabbits defined three allotypic lineages, the so-called *VH*a allotypes *a1*, *a2* and *a3*. *VH* allotypic markers have not been found in other species, and the Mendelian behavior of rabbit *IGHV* allotypes was puzzling for many years,

as rabbit serological data revealed a complex situation where homozygous *VHa* rabbits were found to express distinct *VH* genes. Some of them were devoid of *VHa* allotype-specific determinants (the so-called *VHa* negative or *VHn*, alias *VHx*, *VHy* and *VHz*; Horng et al. 1976). The *VHa* positive molecules (i.e. displaying motifs of *VHa*-characteristic determinants) also showed variation which revealed the existence of multiple *VHa*-related gene fragments (Brezin and Cazenave 1975, Mage et al. 1976, van der Loo et al. 1977). This was later confirmed by genomic mapping of the rabbit *VH* region which revealed the existence of over 200 *VH* elements in the germline (Currier et al., 1988; Gallarda et al., 1985; Ros et al., 2004). This may, in part, explain why for many years it was so difficult to understand the presence of allelic allotypic markers on most serum immunoglobulins of all classes (and probably the skepticism of researchers less acquainted with serological methods).

An important contribution to understanding the mechanisms underlying preferential *VDJ* rearrangement in the rabbit was the study of a rabbit strain, called “Alicia”, detected as a mutant during breeding at the Basel Institute, Switzerland (Kelus and Weiss, 1986). The mutant rabbits derived from a *VH1a2* parental rabbit in which a 10 Kb segment of genomic DNA containing the *VH1* gene was deleted (Knight and Becker, 1990, Allegrucci et al., 1990, Ros et al. 2004). In contrast to normal individuals of the *a2* lineage, the young homozygous *ali/ali* mutant Alicia rabbits produced only trace amounts of *a2* molecules and their serum contained mostly Ig resulting from rearranged genes not encoding *VHa* allotype-associated epitopes (*VHa* negative or *VHn* genes) (Kelus and Weiss, 1986; Di Pietro et al., 1990; Chen et al., 1993). As the genes encoding the *VHn* allotypes map at least 100 Kb upstream of *VH1* (Mage et al., 2006), it is intriguing that *VDJ* rearrangements of these appeared to be more prevalent than *VDJ* rearrangements of the *D*-proximal functional *VHa* positive *VH4* gene in young Alicia rabbits. Further analysis showed that the *VH4* gene is in fact the predominantly rearranged *VH* gene found in the bone marrow of young Alicia rabbits, but for some reason most *VH4*-utilizing B-cells are eliminated. Thus, the immunoglobulins of the young rabbits are produced by B-cells that utilize *VHn*. As the young “Alicia” rabbit ages, the proportions of B cells expressing serological *a2* specificities increase (Kelus and Weiss, 1986; Di Pietro et al., 1990; Allegrucci et al., 1990; Chen et al., 1993; Pospisil et al., 1995). Analysis of nucleotide sequences of the promoter region showed that more than 80% of the *VDJ* rearrangements in older Alicia rabbits utilize either the functional *VH4* or *VH7* genes localized upstream of *VH1*. The *VH4* and the *VH7* genes have 7 (out of 11) specific nucleotides associated with the allotype *a2*, while the other nucleotides that characterize *a2* are gained through somatic gene conversion using *VH9* or a *VH9*-like germline gene as donor (Sehgal et al., 1998; Zhu et al., 1999).

The study of this mutant rabbit strain showed that the Mendelian inheritance of the *VHa* allotypes in normal rabbit is explained by the preferential usage of only one *VH* gene in *VDJ* rearrangements, i.e. the *D*-proximal *VH* gene segment *VH1* (see Figure 2), which is deleted from the “ali” genome (Knight and Becker, 1990; Knight, 1992, Allegrucci et al., 1990, Ros et al. 2004). Despite having more than 200 *VH* genes (Currier et al., 1988; Gallarda et al., 1985; Ros et al., 2004), over 50% are “non-functional” and, apart from *VH1*, an even smaller fraction seems to encode α -positive proteins (Ros et al., 2004). About 80 to 90% of circulating Ig molecules are derived from the *VH1* gene and express the *VHa* allotypic markers (e.g. Kindt, 1975; Margolies et al., 1977). The *VH* regions of the remaining 10–20% of Ig molecules are encoded by the *VHn* genes (Kim and Dray, 1973; Roux, 1981), which are localized at least 100 Kb upstream of *VH1* (Mage et al., 2006). Thus far, the basis for the preferential usage of the *VH1* gene in *VDJ* rearrangements remains unanswered. Enhanced chromatin accessibility of this region of the DNA may be responsible (Mage et al., 2006). Furthermore, we can speculate that the retention and

occasional usage of the *VHn* genes in 10–20% of *VDJ* rearrangements may represent an evolutionary relic of Lagomorphs ancestral immunologic response to pathogens.

Genetic diversity at the heavy chain variable region *IGHV* locus (a locus)

In the domestic rabbit, the three serologically defined allotypic lineages, *VHa* allotypes *a1*, *a2* and *a3*, are highly divergent ($\pm$ 20% amino acid sequence differences). The allelic specificities of *a1* and *a2* are correlated with several amino acid differences in framework regions 1 (positions 5, 8, 11, 13, 14, 17 and 18) and 3 (positions 74, 76, 79, 80, 83, 84, 96 and 97) (Fig. 3) (Tonnelle et al., 1983; Mage et al., 1984a; Knight and Becker, 1990). Serologically distinct *VHa* allotypes were discovered in wild rabbits and named *VHa100*–*VHa109* by Cazenave and coworkers (reviewed in Cazenave et al., 1987). Partial protein sequences of purified antibodies of the *VHa100* type (Tonnelle et al., 1983), suggest these are related to the more recently described and fully sequenced *VHa* allele(s) corresponding to the *a4* lineage (Esteves et al., 2004). Studies by double immunodiffusion of sera from domestic breeds and wild rabbits of Continental Europe (North of Pyrenees Mountains), Great Britain and overseas showed the presence of allotypes *a1*, *a2* and *a3*, as well as a rare allotype with partial reaction against *a3*-specific antiserum, found in the French populations. In all studied populations, the gene frequencies of *a1* and *a3* were similar ($\pm$ 40%) and higher than *a2* ($\pm$ 20%) (van der Loo, 1993). Interestingly, these allelic frequencies reflect the “pecking order” of their relative expression in heterozygous animals (Mage, 1967; Lummus et al., 1967). Quite surprisingly, the study of wild rabbit populations from the Iberian Peninsula belonging to the subspecies *O. c. algirus* and *O. c. cuniculus* revealed a different picture with a much higher prevalence of the “a-blank” allele. Also, a gradient in the gene frequencies of this allotype was noted, decreasing from southwestern to northeastern populations (Fig. 4) (Esteves et al., 2004). Sequencing of rearranged *VH* genes expressed in *O. c. algirus* rabbits that were typed as “a-blank” showed that these rabbits preferentially use *VH* genes that, although clearly related to the known *VHa* genes, defined a new major allotypic lineage, designated *a4*. The *a4* sequences display the rabbit *VHa* hallmark residues together with a number of unprecedented amino acid changes in FR2 and FR3, which may explain the lack of reactivity with known alloantisera (Fig. 3). The net protein distances between the *VH-a4* and the *VH-a1*, *a2*, and *a3* lineages, was 20%, 29%, and 21% respectively, similar to the net protein distances between *VH-a1*, *a2*, and *a3* lineages. As such, a new *VHa* allele was found in Iberian *O. c. algirus* populations, constituting the fourth of the distantly related lineages of the rabbit *VHa*-locus, one of which, the *a4* lineage, seems to be endemic in the Iberian range (Esteves et al., 2004). The data for the *VHa* lineages also suggested that some lineages, namely *VHa a1*, *a2* and *a4*, showed higher evolutionary rates than the *VHa a3* and *VHn* lineages. This increased evolutionary rate may account for the accumulation of a considerable fraction of the amino acid differences characterizing the *a4* lineage after the separation of the two *O. cuniculus* subspecies, explaining its association with subspecies *O. c. algirus* (Esteves et al., 2004).

Studies on the *IGHV* locus diversity for other leporids are limited to a few data obtained for some *Lepus* species, namely *L. americanus*, *L. europaeus* and *L. granatensis*. For *Lepus americanus*, only serological analyses have been conducted; these showed cross-reaction with rabbit anti-*a1*, -*a2* and -*a3* antisera and the absence of individuals with no cross-reaction to rabbit antisera (De Poorter, 1984). As for *L. europaeus* and *L. granatensis*, a serological analysis of several populations using allo-antisera against rabbit *a1*, *a2* and *a3* allotypes revealed only two phenotypes: partial reaction to anti-*a2* antisera and no reaction to any rabbit antiserum (Table 1) (Esteves, 2003). Sequences of expressed *VH* genes were obtained for only five *Lepus* specimens of both species, three of which expressed *a2*-cross-reacting serum proteins, and these could be grouped into two different lineages (see below „The trans-specific polymorphism of leporids *IGHV*). Extending the study of *IGHV*

diversity to more *Lepus* individuals and species could unravel more *VH* lineages, as shown by the rabbit *VH1-a4* lineage found only in *O. c. algirus*.

The trans-specific polymorphism of leporid *IGHV*

Such large interallelic distances as those observed at the rabbit *VHa* locus can be the outcome of unusually long allele persistence times and/or increased mutant recruitment rates. The first indications that the different *VH1* polymorphisms could be trans-specific came from studies showing serological cross-reactivity of *Lepus* sera with rabbit *VHa*-allotype-specific allo-antisera (van der Loo et al., 1977, Horng et al., 1980, van der Loo, 1987). Indeed, for *Lepus americanus* large scale serological analysis, comprising several hundreds of specimens, showed cross-reaction with rabbit anti-a1, -a2 and -a3 antisera, revealing at least three alleles with hierarchical frequency distributions similar to those observed for feral rabbits (De Poorter, 1984; van der Loo, 1987). Su and Nei (1999) compared the extent of sequence divergence between the rabbit *a1*, *a2* and *a3* allotypes with that between human and mouse *VH* gene sequences and concluded that, assuming a “normal” mutation rate, the rabbit *VH1* polymorphism has persisted for about 50 My. Since the *Oryctolagus* and *Lepus* genera diverged 12 Mya (Matthee et al., 2004), the allelic lineages present in one leporid species should be more related to some of the alleles expressed in the other species than to their conspecific allelic counterparts. Serological cross-reactivity can depend upon a small number of amino acid replacements, and allo-antisera may reveal plesiomorphic character states (i.e., which became antigenic in the recipient strain when they were altered), and so the hypothesis that the *VH1* polymorphism preceded the rabbit-hare split remained controversial until sequence data was obtained.

Focusing on this question, Esteves and co-workers (2005) sequenced rearranged *VH* gene products from five *Lepus* specimens of both *L. granatensis* and *L. europaeus* species, three of which expressed *a2*-cross-reacting serum proteins, and compared them to known rabbit *VH* sequences of different allotypes. Within the *Lepus VH* genes, two lineages were observed, one of which (*a2L*) showed some of the most outstanding characteristics of the rabbit *VHa* genes, furthermore possessing five out of eleven amino acid residues that characterize the allotype *VHa2* (Mage et al., 1984a). This lineage was only obtained from specimens showing rabbit *a2* antisera cross-reaction. A second lineage displayed more similarity to the rabbit *VHn* gene fragments (sL). Phylogenetic inference methods clearly placed the hare *a2L* sequences into the same monophyletic group with the rabbit *a2* sequences (Fig. 5), thus showing that these sequences were more closely related to the rabbit *VH1-a2* allele than *VH1-a2* was to its allelic counterparts *VH1-a1* and *VH1-a3*. Also, the genetic distance measured between different rabbit genomic *VHa* sequences was approximately 1.5 times larger than between the *Lepus a2L* consensus and genes of the rabbit *a2* allotype. Together, these findings strongly suggest that the allotype split predates the *Oryctolagus-Lepus* species split, supporting previous evidence of the trans-species nature of the *VH1* polymorphism. The alternative interpretation, that the evolutionary rate of the rabbit *a2* lineage has been significantly slower than at the other lineages, was not supported by the data, as the *a2* sequences were rather more derived compared to the other allotypes, in particular, the *a3* lineage.

The large inter-allelic differences at the *VH1* locus

Regarding the two different hypotheses invoked to explain the very large inter-allelic distances observed at the *IGHV1* locus, the results obtained seem to be contradictory. On the one hand, the association of specific alleles with subspecific markers suggests that the large differences between *a4* and domestic alleles (*a1*, *a2* and *a3*) may have accumulated after the separation of the subspecies, with different evolutionary rates among lineages. On the other

hand, the confirmation of the trans-specific nature of the *VH1* polymorphism between *Oryctolagus* and *Lepus* supports the hypothesis that very long lineage persistence times have contributed to allelic divergence. The evidence obtained points to two major conclusions, 1) the allelic lineages can be maintained in the genome for a long time and 2) mutation rates can differ between allelic lineages. The origin of the selection forces that regulate the *VH1* polymorphism is unknown. However, the requirement for exogenous factors, such as the intestinal microbiota, for diversification of the primary antibody repertoire in rabbits suggests that this mechanism might impose constraints on the evolution of the *VH1* alleles.

Diversification of the primary Ab repertoire in the rabbit

Among vertebrate species, two different strategies to generate the primary Ab repertoire have been adopted. Human and mouse preferentially use combinatorial rearrangements of a large number of *V*, *D*, and *J* gene segments. In contrast, several species possess or use only a limited number of germline *V* segments. Birds, together with several mammalian species (rabbit, sheep, pig and bovine), use strategies of primary repertoire development that overcome this limitation. In these species, the primary antibody repertoire is diversified post-rearrangement by one or both of two mechanisms: somatic gene conversion and somatic hypermutation.

In chicken, somatic gene conversion is the major mechanism of diversification of immunoglobulin heavy and light chain variable regions. A set of pseudogenes is used as donors and the unique rearranged *V(D)J* gene acts as acceptor (Reynaud et al., 1985; 1989). In sheep, B cell diversification occurs in early development in the ileal Peyer's patches and results from somatic hypermutation rather than somatic gene conversion (Reynaud et al., 1991; 1995). Both processes are used to diversify the antibody repertoire in cattle (Parron et al., 1996; Lucier et al., 1998).

Becker and Knight (1990) showed that the rearrangement process in the rabbit involves a single preferentially rearranged *VH* gene that is diversified mainly through somatic gene conversion, using other upstream *VH* genes as donors (Becker and Knight, 1990; reviewed in Knight and Crane, 1994). Somatic hypermutation also takes place, distributing point mutations throughout the entire *VDJ* gene (Short et al., 1991; Weinstein et al., 1994, Sehgal et al., 2000). In rabbits, B cells generated in the fetal liver and bone marrow subsequently migrate to the appendix and other gut-associated lymphoid tissues (GALT), where they proliferate and are intensively diversified. The *VH* genes upstream of *VH1* contribute to the expressed *VH1* diversification by somatic gene conversion. They evolved to some extent in concert, which resulted in the presence of the allotypic motifs on other *VH* gene segments of the Ig heavy chain locus (haplotype polymorphism).

In contrast to species such as chickens (e.g. Reynaud et al., 1989), sheep (Reynaud et al., 1995) and cattle (e.g. Lucier et al., 1998), the diversification of the primary antibody repertoire in rabbits is not developmentally programmed. It has been shown that the peripheral Ab repertoire is diversified between 4 weeks and 2 months after birth (Cooper et al., 1968; Weinstein et al., 1994). Surgical removal of the appendix and sacculus rotundus on the day of birth and the Peyer's patches at three weeks of age (Cooper et al., 1968) more recently showed that GALT is essential for the generation of the primary antibody repertoire (Vadaj et al., 1998). It has also been shown that exogenous factors, such as the intestinal microbiota, are required for Ab repertoire diversification in the rabbit (Lanning et al., 2000a, 2000b; Sehgal et al., 2002). The rabbit is a species with altricial young (naked and blind at birth). Other lagomorph species, such as hares (genus *Lepus*) have precocial young. The unique situation of post-natal diversification in rabbit could be related to the shortening of gestation time (four weeks in rabbit and six weeks in hares). Considering the differences in

reproductive biology between the genera *Oryctolagus* and *Lepus*, it would be interesting to investigate whether in *Lepus* the Ab repertoire is diversified neonatally or during foetal life.

Evolution of *VH* gene families

The study of *VH* genes from several vertebrate species showed that *VH* genes from the same species could belong to different groups. In tetrapods, the *IGHV* sequences have been classified into three clans, clans I, II and III (Schroeder et al. 1990; Kirkham et al. 1992; Das et al., 2008). The human and mouse possess *VH* genes that are very diversified and may cluster with either clans I, II or III (polyphyletic), whereas *VH* genes of chicken, rabbit, horse, and artiodactyls (cattle, sheep, and swine) are monophyletic. Chicken, rabbit and swine *VH* genes each form distinct clusters within group III, whereas those of horse, cattle and sheep each form distinct clusters within group II. Since *VH* genes of different artiodactyl species cluster within different groups (II and III), their common ancestor must have possessed *VH* genes from both groups (Sitnikova and Su, 1998). Thus, the presence of *IGHV* genes of only one clan in several tetrapod species reflects an evolutionary loss of *IGHV* genes.

Monofunctional multigene families are generally believed to undergo processes of genetic exchange, like gene conversion and unequal crossover, which homogenize the DNA sequences (Smith et al., 1971; Smith, 1974; Zimmer et al., 1980). This has been called “concerted” evolution. The concerted evolution model has been invoked to explain the evolution of *VH* genes (Hood et al., 1975; Ohta, 1983). However, the studies of *VH* gene families in human and mouse (Gojobori and Nei, 1984; Tutter and Riblet, 1989), and studies on much longer evolutionary time scales, showed that the pattern of *VH* gene evolution could be better explained by the so-called “birth-and-death” model of evolution (Ota and Nei, 1994; Nei et al., 1997; Sitnikova and Su, 1998; Das et al., 2008), in which the number of genes in a family (or “library”) are allowed to expand and contract. This model is similar to the “accordion model” of MHC evolution proposed by Klein et al., (1993) and postulates that, depending upon the need to protect the host from ever-changing groups of parasites, some *VH* gene libraries are duplicated and can diverge functionally, while others become pseudogenes and/or are deleted from the genome. The end result of this process is a mixture of divergent and highly homologous groups of genes, and the maintenance of a substantial number of pseudogenes (Ota and Nei, 1994; Nei et al., 1997). According to this model, the tetrapod ancestor possessed three *VH* gene groups (I, II, and III) and library contraction events occurred in several lineages independently. In species that have inherited only one group of *VH* genes, showing low levels of *IGHV* gene sequence variation, antibody diversification is mainly due to somatic hypermutation and/or somatic gene conversion. It appears that *VH* gene library contraction is associated with the development of a specific organ for extensive somatic diversification of the Ab repertoire (bursa of Fabricius in chicken; ileal Peyer's patches in sheep, cattle and probably swine and horse; appendix in rabbit).

Genome Sequencing and future prospects

Whole genome sequencing (WGS) at ~7× coverage and assembly of a high quality draft rabbit genome sequence was recently completed at the Broad Institute, Boston (OryCun2.0). The ENCODE Project, with ~1% of rabbit genomic sequence from a different, normal NZW animal, includes some genes of immunological interest but none of the regions with immunoglobulin heavy or light chain loci. The Thorbecke rabbit strain chosen by Broad for WGS was partially inbred. Although the rabbits accepted skin grafts, they were still segregating for the *VH*a allotypes a1 and a2. The rabbit chosen for sequencing was heterozygous at the immunoglobulin heavy chain locus. The *IGH* locus is not assigned to a

chromosome in the Broad assembly. The *IGHV* allele encoding a1 (VH1a1) is located in unplaced scaffold chrUn0742 and the *IGHV* allele encoding a2 (VH1a2) is located in unplaced scaffold chrUn0439. Unfortunately, there are missing traces or traces of low quality between the telomere and the *IGH*. Until new sequences are obtained, the current available traces from Broad OryCun2 may not allow further assembly. There are ~75 additional unplaced scaffolds that appear to contain *VH* genes.

Little is known about the evolution of the *IGHV* genes in the Lagomorph group. Thus far, only the rabbit *O. cuniculus* has been extensively studied and some sequences of *IGHV* genes have been obtained for two *Lepus* species, *L. europaeus* and *L. granatensis*. The study of these species has yielded some interesting results and led to new insights on the evolution of *IGHV* genes in the Lagomorph group. Also, the few data obtained so far for the Lagomorph *IGHG* hinge region and CH2 domain has shown some particular patterns of genetic diversity possibly linked to resistance against pathogens (Esteves et al., 2002a; Esteves et al., 2002b; Esteves et al., 2006). Extending our knowledge about both the *IGHV* genes and *IGH* constant regions to other Lagomorph species would certainly contribute further to our understanding of the evolution of *IGHV* genes and the roles that genetic diversity of immunoglobulins may play in host resistance to pathogens.

Acknowledgments

This work was supported by grants of Foundation for Science and Technology- Portugal to A.P. and P.J.E. (SFRH/BD/71252/2010 and SFRH/BPD/27021/2006, respectively, and the intramural program of the NIAID, NIH (R.G.M.).

References

- Allegrucci M, Newman BA, Young-Cooper GO, Alexander CB, Meier D, Kelus AS, Mage RG. Altered phenotypic expression of immunoglobulin heavy-chain variable-region (VH) genes in Alicia rabbits probably reflects a small deletion in the VH genes closest to the joining region. *Proc Natl Acad Sci U S A*. 1990; 87(14):5444–5448. [PubMed: 2115171]
- Alves, PC.; Hackländer, K. Lagomorph species: Geographical distribution and conservation status. In: Alves, PC.; Ferrand, N.; Hackländer, K., editors. *Lagomorph Biology*. Springer; Berlin Heidelberg: 2008. p. 395-405.
- Becker RS, Knight KL. Somatic diversification of immunoglobulin heavy chain VDJ genes: evidence for somatic gene conversion in rabbits. *Cell*. 1990; 63(5):987–997. [PubMed: 2124176]
- Biju-Duval C, Ennafaa H, Dennebouy N, Monnerot M, Mignotte F, Soriguer RC, Gaaïed AE, Hili AE, Mounolou J-C. Mitochondrial DNA evolution in lagomorphs: Origin of systematic heteroplasmy and organization of diversity in European rabbits. *Journal of Molecular Evolution*. 1991; 33(1):92–102.
- Bouton C, van der Loo W. The trans-species nature of rabbit b locus polymorphism is supported by studies on the snow-shoe hare. *Immunogenetics*. 1997; 45(6):444–446. [PubMed: 9089105]
- Brezin C, Cazenave PA. Cross reaction between allotype Aa1 of rabbit immunoglobulins and antibodies derived against allotype Aa3: participation of variants of Aa1 specificity in this cross reaction. *Immunochemistry*. 1975; 12(3):241–247. [PubMed: 811544]
- Cabrera. *Fauna Ibérica*. Museo Nacional de Ciencias Naturales de Madrid; Madrid: 1914.
- Carneiro M, Afonso S, Gerales A, Garreau H, Bolet G, Boucher S, Tircazes A, Queney G, Nachman MW, Ferrand N. The Genetic Structure of Domestic Rabbits. *Molecular Biology and Evolution*. 2011 (Epub ahead of print) doi:10.1093/molbev/msr003.
- Carneiro M, Ferrand N, Nachman MW. Recombination and Speciation: Loci Near Centromeres Are More Differentiated Than Loci Near Telomeres Between Subspecies of the European Rabbit (*Oryctolagus cuniculus*). *Genetics*. 2009; 181(2):593–606. [PubMed: 19015539]

- Cazenave, PA.; Bennamar, A.; Sogn, JA.; Kindt, TJ. Immunoglobulin genes in feral population. In: Dubiski, S., editor. The rabbit in contemporary immunological research. Longman Scientific & Technical; 1987. p. 148
- Cazenave PA, Brézin C, Puget A, Mandy W. Phylogeny of the e allotypic system of rabbit immunoglobulins: study of determinants recognized on ochotona IgG by anti-e15 antisera. *Ann Immunol.* 1977; 128(1–2):323–327.
- Chapman, J.; Flux, J. Introduction and overview of the Lagomorphs. In: Chapman, J.; Flux, J., editors. Rabbits, Hares and Pikas, Status, survey and conservation action plan UICN/SSC Lagomorph Specialist Group. Switzerland, Gland: 1990. p. 1–6.
- Chen HT, Alexander CB, Young-Cooper GO, Mage RG. VH gene expression and regulation in the mutant Alicia rabbit. Rescue of VH α 2 allotype expression. *J Immunol.* 1993; 150(7):2783–2793. [PubMed: 8454855]
- Cooper MD, Perey DY, Gabrielsen AE, Sutherland DE, McKneally MF, Good RA. Production of an antibody deficiency syndrome in rabbits by neonatal removal of organized intestinal lymphoid tissues. *Int Arch Allergy Appl Immunol.* 1968; 33(1):65–88. [PubMed: 4171257]
- Corbet, GB. Taxonomy and origins. In: Tompson, HV.; King, CM., editors. The european rabbit. Oxford Science Publications; Oxford: 1994. p. 1–6.
- Corbet, GB.; Hill, JE. A World list of mammalian species. Oxford University Press; Oxford: 1980.
- Currier SJ, Gallarda JL, Knight KL. Partial molecular genetic map of the rabbit VH chromosomal region. *J Immunol.* 1988; 140(5):1651–1659. [PubMed: 3126232]
- Das S, Nozawa M, Klein J, Nei M. Evolutionary dynamics of the immunoglobulin heavy chain variable region genes in vertebrates. *Immunogenetics.* 2008; 60(1):47–55. [PubMed: 18196235]
- Davie JM, Paul WE, Mage RG, Goldman MB. Membrane-associated immunoglobulin of rabbit peripheral blood lymphocytes: allelic exclusion at the b locus. *Proc Natl Acad Sci U S A.* 1971; 68(2):430–434. [PubMed: 4100380]
- De Poorter, M. An experimental test of predictions from different hypothesis of self regulation in the snowshoe hare (*Lepus americanus* Erxleben, 1777). Vrije University; Brussels: 1984.
- DiPietro LA, Short JA, Zhai SK, Kelus AS, Meier D, Knight KL. Limited number of immunoglobulin VH regions expressed in the mutant rabbit “Alicia”. *Eur J Immunol.* 1990; 20(6):1401–1404. [PubMed: 1973389]
- Dray S, Young GO, Nisonoff A. Distribution of Allotypic Specificities among Rabbit Gamma-Globulin Molecules Genetically Defined at Two Loci. *Nature.* 1963; 199:52–55. [PubMed: 14047942]
- Dreyer WJ, Bennett JC. The molecular basis of antibody formation: a paradox. *Proc Natl Acad Sci U S A.* 1965; 54(3):864–869. [PubMed: 4160200]
- Dubiski S. Immunochemistry and genetics of a “new” allotypic specificity Ae14 of rabbit gamma-G immunoglobulins: recombination in somatic cells. *J Immunol.* 1969; 103(1):120–128. [PubMed: 4389618]
- Dubiski, S. Foreword. In: Dubiski, S., editor. The rabbit in contemporary immunological research. Longman Scientific & Technical, John Wiley; 1987. p. ix–xii.
- Dubiski S, Dudziak Z, Skalba D. Serum groups in rabbits. *Immunology.* 1959; 2(1):84–92. [PubMed: 13640683]
- Esteves, PJ. Molecular and population genetic analysis of polymorphism at the antibody loci IgGCH2 and IgVH in lagomorphs. Faculdade de Ciências Universidade do Porto; Porto: 2003.
- Esteves PJ, Alves PC, Ferrand N, van der Loo W. Hotspot variation at the CH2–CH3 interface of leporid IgG antibodies (*Oryctolagus*, *Sylvilagus* and *Lepus*). *Eur J Immunogenet.* 2002a; 29(6): 529–535. [PubMed: 12437613]
- Esteves PJ, Alves PC, Ferrand N, van der Loo W. Restriction fragment alleles of the rabbit IGHG genes with reference to the rabbit IGHGCH2 or e locus polymorphism. *Anim Genet.* 2002b; 33(4): 309–311. [PubMed: 12139512]
- Esteves PJ, Carmo C, Godinho R, van der Loo W. Genetic diversity at the hinge region of the unique immunoglobulin heavy gamma (IGHG) gene in leporids (*Oryctolagus*, *Sylvilagus* and *Lepus*). *Int J Immunogenet.* 2006; 33(3):171–177. [PubMed: 16712647]

- Esteves PJ, Lanning D, Ferrand N, Knight KL, Zhai SK, van der Loo W. Allelic variation at the VHa locus in natural populations of rabbit (*Oryctolagus cuniculus*, L.). *J Immunol.* 2004; 172(2):1044–1053. [PubMed: 14707078]
- Esteves PJ, Lanning D, Ferrand N, Knight KL, Zhai SK, van der Loo W. The evolution of the immunoglobulin heavy chain variable region (IgVH) in leporids: an unusual case of transspecies polymorphism. *Immunogenetics.* 2005; 57(11):874–882. [PubMed: 16247606]
- Ferrand, N.; Branco, M. The evolutionary history of the european rabbit (*Oryctolagus cuniculus*): major patterns of population differentiation and geographic expansion inferred from protein polymorphism. In: Weiss, S.; Ferrand, N., editors. *Phylogeography of Southern Europe Refugia*. Springer; 2007. p. 207-235.
- Flux, JEC.; Angermann, R. The hares and jackrabbits. In: Chapman, JA.; Flux, JEC., editors. *Rabbits, Hares and Pikas: Status Conservation Action Plan*. International Union for Conservation of Nature and Natural Resources; Gland, Switzerland: 1990. p. 61-94.
- Gallarda JL, Gleason KS, Knight KL. Organization of rabbit immunoglobulin genes. I. Structure and multiplicity of germ-line VH genes. *J Immunol.* 1985; 135(6):4222–4228. [PubMed: 2999240]
- Gojobori T, Nei M. Concerted evolution of the immunoglobulin VH gene family. *Mol Biol Evol.* 1984; 1(2):195–212. [PubMed: 6443795]
- Halanych KM, Robinson TJ. Multiple substitutions affect the phylogenetic utility of cytochrome b and 12S rDNA data: examining a rapid radiation in leporid (Lagomorpha) evolution. *J Mol Evol.* 1999; 48(3):369–379. [PubMed: 10093227]
- Hewitt GM. Some genetic consequences of ice ages, and their role, in divergence and speciation. *Biological Journal of the Linnean Society.* 1996; 58(3):247–276.
- Hood L, Campbell JH, Elgin SC. The organization, expression, and evolution of antibody genes and other multigene families. *Annu Rev Genet.* 1975; 9:305–353. [PubMed: 813561]
- Horng WJ, Knight KL, Dray S. Heavy chain variable region allotypic sub-specificities of rabbit immunoglobulins. I. Identification of three subpopulations of a1 IgG molecules. *J Immunol.* 1976; 116(1):117–125. [PubMed: 54383]
- Horng WJ, Papagiannes E, Dray S, Rodkey LS. Expression of cross-reacting determinants of the immunoglobulin heavy chain variable region a3 allotype in *Oryctolagus* and *Lepus*. *Mol Immunol.* 1980; 17(1):111–117. [PubMed: 6153750]
- Kearney JF, Cooper MD, Lawton AR. B cell differentiation induced by lipopolysaccharide. IV. Development of immunoglobulin class restriction in precursors of IgG-synthesizing cells. *J Immunol.* 1976; 117(5 Pt 1):1567–72. [PubMed: 794414]
- Kelus AS, Gell PG. Immunoglobulin allotypes of experimental animals. *Prog Allergy.* 1967; 11:141–184. [PubMed: 4860380]
- Kelus AS, Steinberg CM. Is there a high rate of mitotic recombination between the loci encoding immunoglobulin VH and CH regions in gonial cells? *Immunogenetics.* 1991; 33(4):255–259. [PubMed: 1902823]
- Kelus AS, Weiss S. Mutation affecting the expression of immunoglobulin variable regions in the rabbit. *Proc Natl Acad Sci U S A.* 1986; 83(13):4883–4886. [PubMed: 3014517]
- Kim BS, Dray S. Identification and genetic control of allotypic specificities on two variable region subgroups of rabbit immunoglobulin heavy chains. *Eur J Immunol.* 1972; 2(6):509–514. [PubMed: 4120276]
- Kim BS, Dray S. Expression of the a, x, and y variable region genes of heavy chains among IgG, IgM, and IgA molecules of normal and a locus allotype-suppressed rabbits. *J Immunol.* 1973; 111(3):750–760. [PubMed: 4200366]
- Kindt TJ. Rabbit immunoglobulin allotypes: structure, immunology, and genetics. *Adv Immunol.* 1975; 21:35–86. [PubMed: 50001]
- Kirkham PM, Mortari F, Newton JA, Schroeder HW Jr. Immunoglobulin VH clan and family identity predicts variable domain structure and may influence antigen binding. *EMBO J.* 1992; 11(2):603–609. [PubMed: 1537339]
- Klein J, Ono H, Klein D, O'hUigin C. The accordion model of MHC evolution. *Progress in immunology.* 1993; 8:137–143.

- Knight KL. Restricted VH gene usage and generation of antibody diversity in rabbit. *Annu Rev Immunol.* 1992; 10:593–616. [PubMed: 1590997]
- Knight KL, Becker RS. Molecular basis of the allelic inheritance of rabbit immunoglobulin VH allotypes: implications for the generation of antibody diversity. *Cell.* 1990; 60(6):963–970. [PubMed: 2317867]
- Knight KL, Crane MA. Generating the antibody repertoire in rabbit. *Adv Immunol.* 1994; 56:179–218. [PubMed: 8073947]
- Lanning D, Sethupathi P, Rhee KJ, Zhai SK, Knight KL. Intestinal microflora and diversification of the rabbit antibody repertoire. *J Immunol.* 2000a; 165(4):2012–2019. [PubMed: 10925284]
- Lanning D, Zhu X, Zhai SK, Knight KL. Development of the antibody repertoire in rabbit: gut-associated lymphoid tissue, microbes, and selection. *Immunol Rev.* 2000b; 175:214–228. [PubMed: 10933605]
- Loor F, Kelus AS. Allelic exclusion in the B lineage cells of the rabbit. *Eur J Immunol.* 1978; 8(5): 315–324. [PubMed: 99318]
- Lopez-Martinez, N. Revisión sistemática y biostratigráfica de los Lagomorpha (Mammalia) del Terciario y Cuaternario de España vol 3. Memorias del Museo Paleontológico de la Universidad de Zaragoza. Diputación General de Aragón; 1989.
- Lopez-Martinez, N. The Lagomorph Fossil Record and the Origin of the European Rabbit. In: Alves, PC.; Ferrand, N.; Hackländer, K., editors. *Lagomorph Biology*. Springer; Berlin Heidelberg: 2008. p. 27–46.
- Lucier MR, Thompson RE, Waire J, Lin AW, Osborne BA, Goldsby RA. Multiple sites of V lambda diversification in cattle. *J Immunol.* 1998; 161(10):5438–5444. [PubMed: 9820519]
- Lumms Z, Cebra JJ, Mage R. Correspondence of the relative cellular distribution and serum concentration of allelic allotypic markers in normal and “allotype-suppressed” heterozygous rabbits. *J Immunol.* 1967; 99(4):737–743. [PubMed: 4963793]
- Mage R, Young GO, Dray S. An effect upon the regulation of gene expression: allotype suppression at the a locus in heterozygous offspring of immunized rabbits. *J Immunol.* 1967; 98(3):502–509. [PubMed: 4164057]
- Mage RG. A new look at the biological and genetic significance of rabbit heavy chain allotypes. *Ann Immunol (Paris).* 1979; 130(2):105–114. [PubMed: 111599]
- Mage RG, Bernstein KE, McCartney-Francis N, Alexander CB, Young-Cooper GO, Padlan EA, Cohen GH. The structural and genetic basis for expression of normal and latent VH_a allotypes of the rabbit. *Mol Immunol.* 1984; 21(11):1067–1081. [PubMed: 6083445]
- Mage RG, Lanning D, Knight KL. B cell and antibody repertoire development in rabbits: the requirement of gut-associated lymphoid tissues. *Dev Comp Immunol.* 2006; 30(1–2):137–153. [PubMed: 16098588]
- Mage RG, Young-Cooper GO, Alexander C. Genetic control of variable and constant regions of immunoglobulin heavy chains. *Nat New Biol.* 1971; 230(10):63–64. [PubMed: 5279812]
- Mage RG, Young-Cooper GO, Rejnek J, Ansari AA, Alexander CB, Appella E, Carta-Sorcini M, Landucci-Tosi S, Tosi RM. Rabbit Immunoglobulin Allotypes: complexities of their genetics, expression, structural basis and evolution. *Cold Spring Harbor Symp. Quant. Biol.* 1976; 41:677–686.
- Margolies MN, Cannon LE, Kindt TJ, Fraser B. The structural basis of rabbit VH allotypes: serologic studies on all H chains with defined amino acid sequence. *J Immunol.* 1977; 119(1):287–294. [PubMed: 406325]
- Matthee CA, van Vuuren BJ, Bell D, Robinson TJ. A molecular supermatrix of the rabbits and hares (Leporidae) allows for the identification of five intercontinental exchanges during the Miocene. *Syst Biol.* 2004; 53(3):433–447. [PubMed: 15503672]
- Nei M, Gu X, Sitnikova T. Evolution by the birth-and-death process in multigene families of the vertebrate immune system. *Proc Natl Acad Sci U S A.* 1997; 94(15):7799–7806. [PubMed: 9223266]
- Ohta T. On the evolution of multigene families. *Theor Popul Biol.* 1983; 23(2):216–240. [PubMed: 6612633]

- Ota T, Nei M. Divergent evolution and evolution by the birth-and-death process in the immunoglobulin VH gene family. *Mol Biol Evol.* 1994; 11(3):469–482. [PubMed: 8015440]
- Oudin J. The allotype of certain blood protein antigens. *C R Hebd Seances Acad Sci.* 1956; 242(21): 2606–2608.
- Oudin J. Allotopy of rabbit serum proteins. I. Immuno-chemical analysis leading to the individualization of seven main allotypes. *J Exp Med.* 1960; 112:107–124. [PubMed: 13731716]
- Pages MV. Essai de reconstitution de l'histoire du lapin de garenne en Europe. *Bull Mens Off Natl Chasse Sp. Scien. Techn.* 1980:13–21.
- Parnig CL, Hansal S, Goldsby RA, Osborne BA. Gene conversion contributes to Ig light chain diversity in cattle. *J Immunol.* 1996; 157(12):5478–5486. [PubMed: 8955197]
- Pospisil R, Young-Cooper GO, Mage RG. Preferential expansion and survival of B lymphocytes based on VH framework 1 and framework 3 expression: “positive” selection in appendix of normal and VH-mutant rabbits. *Proc Natl Acad Sci U S A.* 1995; 92(15):6961–6965. [PubMed: 7624353]
- Queney G, Ferrand N, Weiss S, Mougél F, Monnerot M. Stationary Distributions of Microsatellite Loci Between Divergent Population Groups of the European Rabbit (*Oryctolagus cuniculus*). *Molecular Biology and Evolution.* 2001; 18(12):2169–2178. [PubMed: 11719566]
- Reynaud CA, Anquez V, Dahan A, Weill JC. A single rearrangement event generates most of the chicken immunoglobulin light chain diversity. *Cell.* 1985; 40(2):283–291. [PubMed: 3917859]
- Reynaud CA, Dahan A, Anquez V, Weill JC. Somatic hyperconversion diversifies the single Vh gene of the chicken with a high incidence in the D region. *Cell.* 1989; 59(1):171–183. [PubMed: 2507167]
- Reynaud CA, Garcia C, Hein WR, Weill JC. Hypermutation generating the sheep immunoglobulin repertoire is an antigen-independent process. *Cell.* 1995; 80(1):115–125. [PubMed: 7813007]
- Reynaud CA, Mackay CR, Muller RG, Weill JC. Somatic generation of diversity in a mammalian primary lymphoid organ: the sheep ileal Peyer's patches. *Cell.* 1991; 64(5):995–1005. [PubMed: 1900459]
- Ros F, Puels J, Reichenberger N, van Schooten W, Buelow R, Platzer J. Sequence analysis of 0.5 Mb of the rabbit germline immunoglobulin heavy chain locus. *Gene.* 2004; 330:49–59. [PubMed: 15087123]
- Roux KH. A fourth heavy chain variable region subgroup, w, with 2 variants defined by an induced auto-antiserum in the rabbit. *J Immunol.* 1981; 127(2):626–632. [PubMed: 6166680]
- Schmale J, Costea N, Dray S, Heller P, Yakulis V. Allelic exclusion of light chain allotypes in rabbit IgM cold agglutinins. *Proc Soc Exp Biol Med.* 1969; 130(1):48–50. [PubMed: 4179405]
- Schroeder HW Jr. Hillson JL, Perlmutter RM. Structure and evolution of mammalian VH families. *Int Immunol.* 1990; 2(1):41–50. [PubMed: 2128464]
- Sehgal D, Mage RG, Schiaffella E. VH mutant rabbits lacking the VH1a2 gene develop a2+ B cells in the appendix by gene conversion-like alteration of a rearranged VH4 gene. *J Immunol.* 1998; 160(3):1246–1255. [PubMed: 9570541]
- Sehgal D, Obiakor H, Mage RG. Distinct clonal Ig diversification patterns in young appendix compared to antigen-specific splenic clones. *J Immunol.* 2002; 168(11):5424–5433. [PubMed: 12023335]
- Sehgal D, Schiaffella E, Anderson AO, Mage RG. Generation of heterogeneous rabbit anti-DNP antibodies by gene conversion and hypermutation of rearranged VL and VH genes during clonal expansion of B cells in splenic germinal centers. *Eur J Immunol.* 2000; 30(12):3634–3644. [PubMed: 11169406]
- Severinson E, Bergstedt-Lindqvist S, van der Loo W, Fernandez C. Characterization of the IgG response induced by polyclonal B cell activators. *Immunol Rev.* 1982; 67:73–85. [PubMed: 6751997]
- Short JA, Sethupathi P, Zhai SK, Knight KL. VDJ genes in VHα2 allotype-suppressed rabbits. Limited germline VH gene usage and accumulation of somatic mutations in D regions. *J Immunol.* 1991; 147(11):4014–4018. [PubMed: 1940383]
- Sitnikova T, Su C. Coevolution of immunoglobulin heavy- and light-chain variable-region gene families. *Mol Biol Evol.* 1998; 15(6):617–625. [PubMed: 9615443]

- Smith GP. Unequal crossover and the evolution of multigene families. *Cold Spring Harb Symp Quant Biol.* 1974; 38:507–513. [PubMed: 4524771]
- Smith GP, Hood L, Fitch WM. Antibody diversity. *Annu Rev Biochem.* 1971; 40:969–1012.
- Su C, Nei M. Fifty-million-year-old polymorphism at an immunoglobulin variable region gene locus in the rabbit evolutionary lineage. *Proc Natl Acad Sci U S A.* 1999; 96(17):9710–9715. [PubMed: 10449759]
- Surridge AK, van der Loo W, Abrantes J, Carneiro M, Hewitt GM, Esteves PJ. Diversity and evolutionary history of the MHC DQA gene in leporids. *Immunogenetics.* 2008; 60(9):515–525. [PubMed: 18584169]
- Taberlet P, Fumagalli L, Wust-Saucy A-G, Cosson J-F. Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology.* 1998; 7(4):453–464. [PubMed: 9628000]
- Teherani J, Capra JD, Aggarwal S, Mandy WJ. Amino acid sequence analysis of group e allotype-related peptides derived from lagomorph IgG. *Eur J Immunol.* 1979; 9(9):690–695. [PubMed: 92412]
- Teherani J, Capra JD, Mandy WJ. Amino acid sequence of the CH2 domain from various lagomorph IgGs. *Mol Immunol.* 1982; 19(7):841–846. [PubMed: 7121466]
- Teherani J, Mandy WJ. Constant region IgG allotypes in cottontail rabbits: group E allelic polymorphism. *Immunochemistry.* 1976a; 13(3):221–227. [PubMed: 819355]
- Teherani J, Mandy WJ. Constant region IgG allotypes in hares: group e allelic polymorphism. *Immunochemistry.* 1976b; 13(5):401–406. [PubMed: 939582]
- Todd CW. Allotypy in rabbit 19S protein. *Biochem Biophys Res Commun.* 1963; 11:170–175. [PubMed: 13985246]
- Tonnelle C, Cazenave PA, Brezin C, Moinier D, Fougereau M. Structural correlates to the rabbit immunoglobulin heavy chain a100 allotype. *Mol Immunol.* 1983; 20(7):753–761. [PubMed: 6412072]
- Tutter A, Riblet R. Evolution of the immunoglobulin heavy chain variable region (Igh-V) locus in the genus *Mus*. *Immunogenetics.* 1989; 30(5):315–329. [PubMed: 2509340]
- Vajdy M, Sethupathi P, Knight KL. Dependence of antibody somatic diversification on gut-associated lymphoid tissue in rabbits. *J Immunol.* 1998; 160(6):2725–2729. [PubMed: 9510172]
- van der Loo W. Population genetical studies on the adaptive significance of the immunoglobulin allotypes in wild rabbit. In: Dubiski, S., editor. *The Rabbit in Contemporary Immunological Research*. Longman Scientific & Technical, John Wiley; 1987. p. 101-126.
- van der Loo W. Variance analysis of immunoglobulin alleles in natural populations of rabbit (*Oryctolagus cuniculus*): the extensive interallelic divergence at the b locus could be the outcome of overdominance-type selection. *Genetics.* 1993; 135(1):171–187. [PubMed: 8224818]
- van der Loo W, Arthur CP, Richardson BJ, Wallage-Drees M, Hamers R. Nonrandom allele associations between unlinked protein loci: are the polymorphisms of the immunoglobulin constant regions adaptive? *Proc Natl Acad Sci U S A.* 1987; 84(9):3075–3079. [PubMed: 3106983]
- van der Loo W, De Baetselier P, Hamers-Casterman C, Hamers R. Evidence for quasi-silent germline genes coding for phylogenetically ancient determinants of the rabbit a locus allotypes. *Eur J Immunol.* 1977; 7(1):15–22. [PubMed: 66147]
- van der Loo W, Ferrand N, Soriguer RC. Estimation of gene diversity at the b locus of the constant region of the immunoglobulin light chain in natural populations of European rabbit (*Oryctolagus cuniculus*) in Portugal, Andalusia and on the Azorean Islands. *Genetics.* 1991; 127(4):789–799. [PubMed: 1903121]
- van der Loo W, Gronowicz ES, Strober S, Herzenberg LA. Cell differentiation in the presence of cytochalasin B: studies on the “switch” to IgG secretion after polyclonal B cell activation. *J Immunol.* 1979; 122(4):1203–8. [PubMed: 376713]
- van der Loo W, Mougél F, Sanchez MS, Bouton C, Castien E, Fonseca A, Ferrand N, Soriguer R, Monnerot M. Cytonuclear disequilibria in wild populations of rabbit (*Oryctolagus cuniculus* L.) suggest unequal allele turnover rates at the b locus (IGKC1). *Immunogenetics.* 1999; 49(7–8): 629–643. [PubMed: 10369921]

- Weinstein PD, Anderson AO, Mage RG. Rabbit IgH sequences in appendix germinal centers: VH diversification by gene conversion-like and hypermutation mechanisms. *Immunity*. 1994; 1(8): 647–659. [PubMed: 7600292]
- Wolf B, Janeway CA Jr. Coombs RR, Catty D, Gell PG, Kelus AS. Immunoglobulin determinants on the lymphocytes of normal rabbits. 3. As4 and As6 determinants on individual lymphocytes and the concept of allelic exclusion. *Immunology*. 1971; 20(6):931–944. [PubMed: 4104284]
- Zhu X, Boonthum A, Zhai SK, Knight KL. B lymphocyte selection and age-related changes in VH gene usage in mutant Alicia rabbits. *J Immunol*. 1999; 163(6):3313–3320. [PubMed: 10477601]
- Zimmer EA, Martin SL, Beverley SM, Kan YW, Wilson AC. Rapid duplication and loss of genes coding for the alpha chains of hemoglobin. *Proc Natl Acad Sci U S A*. 1980; 77(4):2158–2162. [PubMed: 6929543]

Fig. 1. Evolutionary topology reflecting the relationships within the Lagomorpha group based on a molecular super matrix (adapted from Matthee et al., 2004).

Fig. 2.

Rabbit *IGHV* (regions from *VH* haplotype (a2 F–I)) for which nucleotide sequence is available from sequenced, assembled BAC clones. Depicted are *V* genes, *D* and *J* regions and also *Cμ*. *VH* functional genes are represented in white rectangles, *VH* possible pseudogenes in grey and *VH* pseudogenes in black, identified according to criteria described in Ros et al. (2004). The *D*-proximal *VH* gene, *VH1*, used in the majority of *VDJ* rearrangements is indicated. Identified above are the non-overlapping BAC clones 38A2 and 225P18 (GenBank Accession Numbers AY386694 and AY386697), and 219D23 (GenBank Accession Number AY386695). The 3' end of BAC 225P18 overlaps the 5' end of BAC 219D23 (adapted from Ros et al., 2004).

Fig. 3.

Amino acid sequences of European rabbit (*O. cuniculus*) proteins (encoded by *VH1a1*–*a3*, *VHa4.1*–*4.2*, *VHx*, *VHy*, *VHz*) and hare (*VHa2L*, *VHsLe*, *VHsLg*) VH genes. Germline elements are *italicized* (Accession numbers: *VH1a1*-M93171; *VH1a2*-M93172; *VH1a3*-M93173; *VHx*-L03846; *VHy*-L03890; *VHz*-AF264469.). *VHa4.1*–*4.2*, *VHa2L*, *VHsLe* and *VHsLg* represent consensus protein sequences obtained by Esteves et al., 2004, 2005. Framework regions (FR1, FR2, FR3) and each CDR (CDR1 and CDR2) are defined according to the IMGT numbering system (also shown). CDRs are shaded. The amino acids that are characteristic of *a1*, *a2* and *a4* lineages are marked with black, grey and white arrows, respectively. Hallmark rabbit *VHa* residues are highlighted in bold and underlined. Note that the *Lepus* *VHa2L* shares these residues, as well as five of the *VH1a2* lineage characteristic amino acids. Dots (.) indicate identity with *VH1a1*, dashes (–) represent indels and question marks (?) represent ambiguous positions in consensus sequences.

Fig. 4.

Geographic distribution of *VHa* locus allotype frequencies, serologically determined for populations of European wild rabbit (*O. cuniculus*) from the Iberian Peninsula and France. The subspecies *O. c. algirus* occupies the Southwestern part of Iberia, which is indicated by *A*. Rabbits of the Northwestern areas of Iberia and of the rest of Europe (indicated by *B*) belong to the subspecies *O. c. cuniculus*. The contact zone between the two subspecies is indicated in gray. The coloring of the disks reflects the relative allele frequencies per locality, analyzed as a two-allele locus with *a-blank* (white) and *a-positive* (black) alleles. Locality abbreviations are as follows: prt - Portimão; san - Santarém; alc - Alcochete; ida - Idanha; cab - Cabreira; LI - Las Lomas; hue - Huelva; don - Doñana; cre - Ciudad Real; inf - Infantado; vv - Vila Viçosa; am - Amoladeras; tol - Toledo; bra - Bragança; tar-Tarragona; alt- Alicante; nav - Navarra; lle - Lerida; per - Perpigan; cam - Camargue; ver - Versailles; arj - Arjuzanx; gra - Grax; hel - Helene.

Fig. 5.

Neighbour-joining tree for *Oryctolagus cuniculus*, *Lepus europaeus* (Le), and *Lepus granatensis* (Lg) VH genes, showing the transspecies polymorphism of VHa2. VH genes from human class 3 (VH3) determine the root of the tree. Numbers represent the bootstrap values (BP, in bold), and the confidence probability values (CP, in italics). Expressed VH genes (cDNA) are labeled by black (Le), gray (Lg), and white (O. cuniculus) circles. Germline VHa gene sequences (VH1–VH4) of rabbit allotypes a1, a2, and a3 are marked by black triangles. White squares indicate VHa germline genes. Black squares indicate human VH3 germline genes. “With kind permission from Springer Science+Business Media: Immunogenetics, The evolution of the immunoglobulin heavy chain variable region (IgVH) in leporids: an unusual case of transspecies polymorphism., 57(11), 2005, 874, Esteves PJ, Lanning D, Ferrand N, Knight KL, Zhai SK, van der Loo W, Figure 1.”

Gene frequencies obtained in populations of *Lepus granatensis* and *L. europaeus* at IGHV. *Lepus* individuals were tested for cross-reaction with rabbit anti -a1, -a2 and -a3 antisera. Serotype codes are shown in parentheses: *I* indicates complete identity with rabbit allo-antisera, *p* partial identity and *O* no reaction (Esteves, 2003).

Table 1

	n	a1 (100)	a2 (010)	a3 (001)	a1v (p00)	a2v (0p0)	a3v (00p)	blank (000)
<i>Lepus granatensis</i>								
Portugal								
Bragança	13	-	-	-	-	0.08	-	0.92
Santarém	20	-	-	-	-	0.41	-	0.59
Idanha	20	-	-	-	-	0.29	-	0.71
Pancas	20	-	-	-	-	0.45	-	0.55
Aljustrel	20	-	-	-	-	0.55	-	0.45
Alcochete	7	-	-	-	-	0.08	-	0.92
Spain								
Granada	20	-	-	-	-	0.19	-	0.81
<hr/>								
average unweighted	120	-	-	-	-	0.29	-	0.71
<hr/>								
<i>Lepus europaeus</i>								
Lass (France)	20	-	-	-	-	0.5	-	0.5