Two TaqI RFLPs in the human von Willebrand factor gene

F. Bernardi, G. Marchetti, V. Bertagnolo, L. Faggioli and L. del Senno

Centro Studi Biochimici sul Morbo di Cooley, Università di Ferrara, Via L. Borsari, 46-44100 Ferrara, Italy

SOURCE/DESCRIPTION: A 2 Kb SacI fragment of the von Willebrand factor cDNA inserted in pGEM2

POLYMORPHISMS: Two TaqI polymorphisms with alleles of 3.3 (A1) and 2.6 (A2) and of 4.5 (B1) and 2.3 Kb (B2) respectively (see figure). Several constant bands are present.

FREQUENCIES: Estimated in 18 unrelated Italian subjects, 0.51 (A1) and 0.49 (A2); 0.95 (B1) and 0.05 (B2).

POLYMORPHIC FOR: BglII (Verweij et al. 1985), BamHI (Nishino and Lynch 1986).

CHROMOSOMAL LOCALISATION: 12p12-12pter (Ginsburg et al. 1985).

MENDELIAN INHERITANCE: Codominant segregation shown in four families.

PROBE AVAILABILITY: Available without restriction. Write to Professor D.C. Lynch.

OTHER COMMENTS: Detectable at wash stringency 2XSSC 60°C.

REFERENCE: Lynch et al. (1985) Cell 41 49. Ginsburg et al. (1985) Science 228 1405. Verweij et al. (1985) N.A.R. 13 8289. Nishino and Lynch (1986) N.A.R. 14 4697.

ACKNOWLEDGEMENTS: Supported by P.F. Ing. Gen. e Basi Mol. Mal. Ered. CNR Cont.n. 8400877. We thank Prof. D.C. Lynch for providing the vW cDNA.