Published in final edited form as:

Angew Chem Int Ed Engl. 2011 May 2; 50(19): 4419–4422. doi:10.1002/anie.201007685.

Automatic Molecular Weaving Prototyped Using Single-Stranded DNA**

Tanashaya Ciengshin, Ruojie Sha, Dr., and Nadrian C. Seeman, Prof.*

Department of Chemistry New York University New York, NY 10003 (USA)

Keywords

DNA Topology; Nanotechnology; Automatic Weaving; Self-assembly; L-Nucleotides

One of the goals of molecular-based nanoscience is the organization of matter into strong molecular structures that display the advantageous properties of analogous macroscopic structures.^[1] A woven fabric is an example of such a macroscopic structure, but deliberately braided woven molecules have not been reported, because nodes of designated^[2] and alternating signs must be placed specifically. Owing to its double helical structure, DNA is an ideal programmable molecule to build synthetic topological targets.^[3] A half-turn of DNA, about six nucleotide pairs, corresponds to a node or a crossing point in a knot or a catenane.^[4] Deliberate trefoil knots,^[5,6] a figure-8 knot,^[7] polyhedral catenanes,^[8,9] specifically linked electrophoretic mobility standards,^[10] and Borromean rings^[11] are examples of previous DNA topological constructs. Nevertheless, a woven arrangement requires an even greater level of control over the placement of nodes. Here, we have prototyped a planar woven arrangement, using the B-DNA conformation for all nodes by strategically combining D-nucleotides and L-nucleotides. This work represents the first step on the way to automatic molecular-scale weaving. A previous use of L-nucleotides in DNA nanotechnology has been reported,^[12] indicating that opposite-handed deviations from ideal structures occur in uniformly L-nucleotide DNA. However, there is no prior report of using a combination of D-nucleotides and L-nucleotides in the same strands for topological or nanotechnological purposes.

An image of a woven arrangement is illustrated in Figure 1a. This Roman mosaic in Conimbraga, Portugal, shows a simple braid. The arrangement contains alternating positive and negative nodes, as emphasized in Figure 1b. The node signs are generated relative to a group of vertical helix axes; were the picture to be rotated 90° within its plane, the signs would switch. In Figure 1c, each node has been drawn flanked by six horizontal lines, representing the base pairs of a half-turn of DNA.

The nodes produced by conventional right-handed double helical B-DNA are negative nodes.^[2] In the cases where positive nodes have been needed in the past,^[6,7,11] left-handed Z-DNA has been used to produce them. The disadvantages of using Z-DNA include the lack

** We thank Drs. Ortho Flint and Yoel Ohayon for instruction in using Knotilus. This research was supported by grant GM-29554 from NIGMS, grants CTS-0608889 and CCF-0726378 from the NSF, grant W911FF-08-C-0057 from ARO, via Pegasus Corporation, MURI W911NF-07-1-0439 from ARO, grants N000140910181 and N000140911118 from ONR and a grant from the W.M. Keck Foundation to N.C.S.

*Fax: (+1) 212-995-4475 ned.seeman@nyu.edu.

Supporting information for this article is available. The experimental details of DNA strand preparation, radioactive phosphorylation, ligation, restriction analysis, denaturing polyacrylamide gel electrophoresis and Ferguson analysis are described. Linking analysis of the product is shown.

of sequence diversity that typically characterizes Z-DNA-forming sequences^[13] and the difficulty of forming Z-DNA within very short segments. In addition, Z-DNA is not exactly a mirror image of right-handed B-DNA, exhibiting both a different helical twist and a different pitch from B-DNA, besides the fact that it contains D-nucleotides.^[14] L-nucleotides do not have these drawbacks, and we use them here in this new topological application of DNA. Thus, the negative (right-handed) nodes built here are made using conventional D-deoxyribose nucleotide pairs, and the positive (left-handed) nodes are made using L-deoxyribose nucleotide pairs.

Figures 1d and 1e illustrate a schematic of the deliberately braided structure, compared with one made of exclusively D-nucleotide DNA. Figure 1d shows the interwoven self-assembled structure made from a mixture of D- and L-nucleotides and Figure 1e shows the structure that would result using only D-nucleotides. Each molecule consists of two cyclic strands, one red and one aqua. Each of the crossings corresponds to a half-turn of DNA, similar to the representations in Figure 1c. The molecules consist of three vertical double-helix-containing domains. The central domain in the molecule shown in Figure 1d contains L-nucleotides, and it is flanked by two domains containing D-nucleotides. All three domains in the molecule shown in Figure 1e contain D-nucleotides. The differences between the two molecules are striking: Figure 1d contains an interwoven catenane, whereas Figure 1e contains two juxtaposed cyclic strands that are not linked at all. A simplified image that depicts the backbone topology of the braided molecule in Figure 1d is shown in Figure 1f and a similar image showing the topology of Figure 1e is shown in Figure 1g. The unlinked molecules in Figures 1e and 1g constitute an ideal control for the braided molecules in Figures 1d and 1f.

An alternative to the planar braided arrangement shown in Figure 1a is the tube-like braided structure shown in Figures 1h and 1i, where the two edges meet one another at the rear of the image to form a cylindrical pattern. Figure 1h is a schematic showing how alternating positive (green) and negative (red) nodes can match up behind the plane, to form a cylindrical or tube-like structure; an example of such a cylindrical structure is shown in Figure 1i, a lamp base that consists of a tube-like weave. Remarkably, the planar arrangement targeted in this work requires more synthetic effort than the tube-like arrangement would require. This difficulty is a consequence of the polarity of the backbone strands and the need to enforce anti-parallel strands on each DNA half-turn domain. Both of the systems shown in Figure 1d and 1e contain strand polarity indicators: The bowtie-like structures on the left sides of the molecules represent 3', 3' linkages and the filled double circles on the right sides of the structures represent 5', 5' linkages. These unusual linkages are hardly unprecedented in synthetic DNA molecules,^[15,16] but they do make the synthesis of the strands containing them more involved than the synthesis of strands made exclusively of 5', 3' linkages.

Figure 2 shows the sequences of the molecules synthesized for this study. The base sequences are identical, but the topology in the braided molecule in Figure 2a is different from the molecule in Figure 2b. Cyclic Molecule 1 is longer than Cyclic Molecule 2, and both cyclic molecules contain unique restriction sites. When restricted, the two different components can be identified readily by the lengths of the product strands. In each of the two systems, Cyclic Molecule 1 results from ligating Strand 1 and Strand 2; Cyclic Molecule 2 results from ligating Strand 3 and Strand 4.

Figure 3 shows the results of ligation and analysis of the molecules depicted in Figure 2. The samples were both stained and radioactively labeled. Panels (a) and (c) show stained gels of the molecules in Figure 2a and 2b, respectively; panels (b) and (d) are the corresponding autoradiograms. There is a band corresponding to the woven DNA catenane at the top of

each lane in Figure 3a and 3b, except for lanes 7 and 8 of 3b, where nothing has been labeled. A variety of other products is visible. Lane 1 contains the ligation products and lane 2 contains the ligation products after treatment with exonucleases I and III. The top two bands, ascribed to the circular strands remain intact in both lanes. Lanes 3 and 4 contain the same materials, but only the long circular strand has been labeled. It is clear that the top circular band remains in the autoradiogram, but not the bottom. Exonuclease treatment results in the disappearance of the band ascribed to the long linear strand (3b). Lanes 5 and 6 are again treated the same way, except that the short circular strand is labeled; only the short circular and linear strands are visible in lane 5, and the short linear strand is digested in lane 6 (3b). Figures 3c and 3d contain the exact same treatments and labeled materials as Figures 3a and 3b, respectively, except that the complex of Figure 2b is used. The results are identical, except for the key difference of the absence of the bands corresponding to the woven DNA catenane.

Figure 4a is an autoradiogram containing the restriction analysis of the woven catenane. Lane 1 contains catenane isolated from the gels of Figure 3a/3b. The large circle is restricted in lane 2, leaving the small circle and the long linear molecule. The small circle is restricted in lane 3, leaving the large circle and the short linear molecule. Lanes 4 and 5 contain large and small circular markers. It is clear that the top bands consist of a catenane of the large and small circles. The absence of the top band in Figures 3c and 3d confirms that the L-nucleotides are necessary for the catenane to form, and that no accidental linking of the two circles occurs in the course of the ligation process. The 10% gels shown here do not resolve singly-linked from doubly-linked catenanes. A 6% gel shown in the supplementary information shows that a small amount of singly-linked catenane contaminates the primary doubly-linked product. L-nucleotides produce positive nodes using only a single half-turn of DNA; in previous work with positive nodes,^{6,7,11} two or three nodes were used, because a half-turn of Z-DNA within a larger context was of marginal stability.

We have demonstrated that it is possible to form a molecular weave with DNA. It is clear from this prototype that more complex braided or woven patterns are now feasible. The key difficulty of working with such structures on the molecular scale is the demonstration of their formation. The presence or absence of linked species is adequate for demonstrating the topology in the present case, but it would be inadequate to demonstrate the formation of a more complexly woven pattern, which would require direct structural evidence of the topology of the species. Given the small amount of contamination from singly-linked catenanes in this study, the presence of by-products seems likely when topological targets of greater complexity are attempted. Nevertheless, the value of building macroscopic materials woven on the molecular scale, even if they contain a few errors, should not be underestimated. These constructs should demonstrate at least some of the features of macroscopically woven materials, and are expected to be quite robust. In addition to DNA constructs themselves, it is important to realize that DNA appears capable of directing molecular topology when it is one rail of a ladder polymer (i.e., a polymer in which two polymers are joined unit by unit through a linking group). Recent work has demonstrated that it is possible to connect other polymers, such as nylon,^[17] to DNA, and new, strong materials could result either from the ladder polymer, or from the ladder with the DNA rail removed.

Methods Summary

DNA molecules used in this work were designed using the program *Sequin*^[18] and were synthesized by phosphoramidite techniques using L-phosphoramidites (ChemGenes, Wilmington, MA) when needed.^[19] 5', 5' and 3', 3' linkages were produced as described

previously.^[15,16] Radioactive phosphorylation, ligation, restriction and Ferguson analysis were performed as described in the Supplementary Information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- [1]. Holmes AE, Barcena H, Canary JW. *Adv. Supramol. Chem.* 2002; 8:42–78.
- [2]. White JH, Millett KC, Cozzarelli NR. *J. Mol. Biol.* 1987; 197:585–603. [PubMed: 3441012]
- [3]. Seeman NC. *Am. Math. Soc. Proc. Symp. Applied. Math.* 2009; 66:121–153.
- [4]. Seeman NC. *Molecular Engineering.* 1992; 2:297–307.
- [5]. Mueller JE, Du SM, Seeman NC. *J. Am. Chem. Soc.* 1991; 113:6306–6308.
- [6]. Du SM, Stollar BD, Seeman NC. *J. Am. Chem. Soc.* 1995; 117:1194–1200.
- [7]. Du SM, Seeman NC. *J. Am. Chem. Soc.* 1992; 114:9652–9655.
- [8]. Chen J, Seeman NC. *Nature.* 1991; 350:631–633. [PubMed: 2017259]
- [9]. Zhang Y, Seeman NC. *J. Am. Chem. Soc.* 1994; 116:1661–1669.
- [10]. Fu T-J, Tse-Dinh Y-C, Seeman NC. *J. Mol. Biol.* 1994; 236:91–105. [PubMed: 8107128]
- [11]. Mao C, Sun W, Seeman NC. *Nature.* 1997; 386:137–138. [PubMed: 9062186]
- [12]. Lin C, Ke Y, Li Z, Wang JH, Liu Y, Yan H. *NanoLett.* 2009; 9:433–436.
- [13]. Ho PS, Ellison MJ, Quigley GJ, Rich A. *EMBO J.* 1986; 5:2737–2744. [PubMed: 3780676]
- [14]. Rich A, Nordheim A, Wang AHJ. *Ann. Rev. Biochem.* 1984; 53:791–846. [PubMed: 6383204]
- [15]. Sha R, Liu F, Bruist MF, Seeman NC. *Biochem.* 1999; 38:2832–2841. [PubMed: 10052955]
- [16]. Sha R, Liu F, Millar DP, Seeman NC. *Chem. & Biol.* 2000; 7:743–751. [PubMed: 10980454]
- [17]. Zhu L, Lukeman PS, Canary J, Seeman NC. *J. Am. Chem. Soc.* 2003; 125:10178–10179. [PubMed: 12926933]
- [18]. Seeman NC. *J. Biomol. Struct. & Dyns.* 1990; 8:573–581.
- [19]. Caruthers MH. *Science.* 1985; 230:281–285. [PubMed: 3863253]
- [20]. Rankin S, Flint O. J. *Knot Theor. & Ramifications.* 2004; 13:151–173.
- [21]. Pentecost CD, Chichak KS, Peters AM, Cave GWV, Cantrill SJ, Stoddart JF. *Angew. Chemie Int. Ed.* 2007; 46:218–222.

Figure 1.

Representations of Braided Topologies. (a–c) *A Braided Structure in Conimbraga, Portugal.*

(a) This picture shows a section of a Roman mosaic illustrating a braided structure. (b) Relative to a vertical helix axis, nodes have been assigned a sign, + (green) or – (yellow), depending on the nature of the crossing. (c) The equivalence between a half-turn of DNA (6 nucleotide pairs) and a node in a knot is emphasized, by replacing node signs by a 6 horizontal lines that correspond to base pairs; the same color coding is used as in (b). (d–g) *The Molecules Made in this Work.* (d) and (e) The molecule in (d) is a prototype of a woven or braided catenane. The need to maintain antiparallelism in the DNA strands forming the double helical segments results in needing to make 3', 3' linkages, drawn as bowtie features on the left, and 5', 5' linkages, drawn as fused circles on the right. The molecule in (e) is the same as in (d), except it does not contain L-nucleotides. (f) and (g) are simplifications of these molecules, drawn by Knotilus^[20] where the nature of (f) as a toroidal Solomon link^[21] is evident. (h and i) *A Woven Tube.* (h) Positive nodes are in green and negative nodes are in red, relative to a vertical helix axis. 5' ends of strands are indicated by filled circles of the same color as the strands. 3' ends are indicated by arrowheads, but only on the right, for clarity. (i) A lamp base decorated in a woven tube.

Figure 2.

The Sequences of the Molecules Used. Backbone segments drawn in black contain D-nucleotides and backbone segments drawn in red contain L-nucleotides. The molecule in (a) is the braided molecule, and the molecule in (b) is the unbraided control. The 3', 3' linkages are indicated by arrowheads pointing at each other, and the 5', 5' linkages are indicated by pairs of filled circles. Topology-determining nodes are indicated at the center of each molecule, where the helical nature of the strands is indicated by crossings whose signs are shown. Restriction sites are color coded and node signs are indicated. Ligation sites are denoted by small arrows pointing at the backbones. The two cyclic molecules are of different lengths.

Figure 3.

Gels Showing the Ligation Products. Panels (a) and (c) are 10% denaturing gels stained with stains-all dye; the woven DNA catenane is loaded in panel (a) and the D-nucleotide control is in panel (c). Panels (b) and (d) are autoradiograms of the same gels shown in (a) and (c), respectively. In each panel the following contents are loaded: Lane M is a 10 base pair marker; in lane 1, both large and small circular DNA strands are radioactively labeled; in lane 2, the sample from lane 1 has been treated with exonucleases I and III; in lane 3, only the large circular DNA strand is radioactively labeled; in lane 4, the sample from lane 3 has been treated with exonucleases I and III; in lane 5, only the small circular DNA strand is radioactively labeled; in lane 6, the sample from lane 5 has been treated with exonucleases I and III; in lane 7, none of the DNA strands are radioactively labeled; in lane 8 the sample from Lane 7 has been treated with exonucleases I and III. LC DNA refers to the long circular DNA strand, SC refers to the short circular DNA strand, LL DNA refers to the long linear DNA strand, and SL DNA refers to the short linear DNA strand. The 100 nt position on the marker is emphasized.

Figure 4.

Restriction Analysis of the Woven DNA Catenane. (a) This is a 10% denaturing gel where lane M contains a 10 base pair marker ladder, lane 1 contains purified braided catenane DNA, lane 2 contains the braided catenane DNA after restriction of the large circular DNA strand with Hind III, lane 3 contains the braided catenane DNA after restriction of the small circular DNA strand with Pvu II, lane 4 contains a marker of the large circular strand, and lane 5 contains a marker of the small circular strand. The same abbreviations apply as in Figure 3. (b) The Ferguson plot of the key breakdown species on this gel.