Published in final edited form as:

Psychol Med. 2012 September ; 42(9): 1925–1935. doi:10.1017/S0033291711002790.

Predicting persistent alcohol problems: A prospective analysis from the Great Smoky Mountain Study

William Copeland, PhD^a, Adrian Angold, MRCPsych^a, Lilly Shanahan, PhD^b, Jessica Dreyfuss, BS, Innocent Dlamini, MA^c, and E. Jane Costello, PhD^a

^aDuke University Medical Center

^bUniversity of North Carolina, Greensboro

^cRegeneron Pharmaceuticals

Abstract

Objectives—Rates of alcohol disorders peak in late adolescence and decrease substantially into the mid-20s. Our aim was to identify risk factors that predict alcohol problems that persist into the mid 20s.

Methods—Data are from the prospective, population-based Great Smoky Mountains Study ($N=1,420$) which followed children through late adolescence and into young adulthood. Alcohol persisters were defined as subjects with an alcohol disorder (abuse or dependence) in late adolescence (ages 19 and 21) that continued to meet criteria for an alcohol disorder at the mid-20s assessment.

Results—The 3-month prevalence of having an alcohol disorder (abuse or dependence) decreased markedly from late adolescence into the mid-20s. A third of late adolescents with an alcohol disorder continued to meet criteria for an alcohol disorder in young adulthood (37 of 144 who met criteria in late adolescence). Risk factors for persister status included multiple alcohol abuse criteria during late adolescence, but no alcohol dependence criteria. Risk factors for persister status also included *associated features* of alcohol dependence such as craving alcohol and drinking to unconsciousness. Persister status was also associated with depression, cannabis dependence and illicit substance use, but not with other psychiatric disorders. Over 90% of late adolescents with 3 or more of the risk factors identified met criteria for a young adult alcohol disorder.

Conclusions—Symptoms of alcohol abuse, not dependence, best predict long-term persistence of alcohol problems. The set of risk factors identified may be a useful screen for selective and indicated prevention efforts.

Keywords

alcohol abuse; alcohol dependence; risk factors; adolescence; young adulthood

Introduction

Rates of both alcohol use and alcohol use disorder (henceforth “alcohol disorders”) increase markedly across adolescence and into the early twenties (Chen and Kandel, 1995, Hasin et

Address for correspondence: William E. Copeland, Ph.D., Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Box 3454, Durham NC 27710, William.Copeland@duke.edu, Phone: (919) 687-4686.

No other authors have any biomedical financial interests.

al., 2007, Harford et al., 2005). Onset of these problems is predicted by psychiatric problems (particularly conduct problems), parental substance history, negative life events, and low SES (e.g., (Guo et al., 2000, Kaplow et al., 2001, Kosterman et al., 2000, Pagan et al., 2006, Tarter, 2002, Webb et al., 1991)). During the mid-to-late twenties, prevalence rates appear to diverge, however: Alcohol *use* decreases modestly (Chen and Kandel, 1995) whereas alcohol *disorders* decrease precipitously (Harford et al., 2005, Grant et al., 2004). For example, in the 2001 National Household Survey on Drug Abuse, twelve-month prevalence rates for any alcohol disorders dropped 28% for males and 47% for females from ages 18–23 to ages 24–29 (Harford et al., 2005). Thus, desistance from alcohol disorders by the mid-to-late twenties is common. In turn, persistence is developmentally deviant and likely associated with the poorest long-term prognosis. To date, risk factors for persistence into the mid-20s are rarely studied, but could provide important information for selective and indicated prevention efforts.

Studies aimed at understanding how persistence of and recovery/desistance from alcohol disorders unfolds for most people require community, epidemiologic samples, because treatment studies only tap into select, biased samples. Indeed, only 24.1% of individuals with lifetime alcohol dependence and 7.0% individuals with lifetime alcohol abuse receive treatment (Hasin et al., 2007). A number of studies have focused on *natural recovery* from alcohol disorders without treatment (for review see (Smart, 2007)), but these studies typically excluded individuals with any treatment history and also suffered from sample selection biases inherent in using media solicitation (Rumpf et al., 2000). Another group of community, longitudinal studies has looked at general predictors of recovery and relapse across the lifespan (Dawson et al., 2007, Pirkola et al., 2006, Hasin et al., 1997, Schuckit et al., 2000, De Bruijn et al., 2006, Grant et al., 2001, Hasin et al., 1990). Only a few studies have focused on the period from late adolescence to young adulthood when desistance is normative. These studies have described desistance as *maturing out*, and focused on family and educational role transitions, including marriage and parenthood, and personality factors in order to understand this normative decrease in alcohol disorders (Staff et al., 2010, Bachman et al., 2002, Littlefield et al., 2009, Littlefield et al., 2010, Yamaguchi and Kandel, 1985).

In a recent review of results from longitudinal studies, Robert Zucker recommended “longitudinal analyses that incorporate models of offset/desistance as well as continuity and growth” p. 107 (Zucker, 2008) as next steps in understanding drinking behavior. To date, there is no broad risk profile that helps identify late adolescents whose alcohol problems will persist. The few studies of persistence implicate the phenomenology of use (including severity (Delucchi et al., 2008), levels of dependence symptoms (Delucchi et al., 2008), and age of first substance use (Chen and Kandel, 1998)), polysubstance use (Kandel and Raveis, 1989), legal problems (Delucchi et al., 2008), depressive symptoms (Chen and Kandel, 1998) and being male (Delucchi et al., 2008). Other risk areas such as individual abuse and dependence criteria and psychiatric diagnoses have not been studied.

Here, we use an epidemiologic, community sample followed from adolescence into the mid-20s in order to develop a risk profile for the persistence of alcohol disorders into young adulthood. We define persistence and desistance according to DSM diagnostic status; thus the focus is on *severity*. This approach is consistent with previous work suggesting that developmental subtypes tend to “differ in degree rather than kind” p. 229 (Jackson et al., 2006). Because age of onset, severity of problems, psychiatric comorbidities, and specific symptom profiles have been implicated in developmental subtypes of alcohol problems (Cloninger, 1988, Zucker et al., 1995, Maggs and Schulenberg, 2004/2005, Babor et al., 1992), we will also evaluate the relative roles of these factors in the prediction of persistence versus desistance.

Methods

Participants

The Great Smoky Mountains Study is a longitudinal study of the development of psychiatric disorder and need for mental health services in rural and urban youth (Costello et al., 1996, Costello et al., 2003). A representative sample of three cohorts of children, age 9, 11, and 13 at intake, was recruited from 11 counties in western North Carolina in 1993. Potential participants were selected from the population of some 12,000 children using a household equal probability, accelerated cohort design. The accelerated cohorts design means that over several years of data collection each cohort reaches a given age in a different year, thus controlling for cohort effects. A screening questionnaire (the externalizing problems scale of the Child Behavior Checklist) was administered to a parent (usually the mother) of the first stage sample (N=3,896). All children scoring above a predetermined cutpoint (the top 25% of the total scores), plus a 1-in-10 random sample of the rest (i.e., the remaining 75% of the total scores), were recruited for detailed interviews. All subjects were subsequently assigned a weight inversely proportional to their probability of selection, so that the results are representative of the population from which the sample was drawn and not biased from the oversampling procedure. Ninety-five percent of families contacted completed the telephone screen. About 8% of the area residents and the sample are African American, fewer than 1% are Hispanic, and 3% are American Indian. The ethnic/racial makeup of the sample is consistent with the area sampled. Of all subjects recruited, 80% (N=1420) agreed to participate.

This paper focuses on assessments completed when the subjects were ages 19, 21 (1999 to 2005) and in their mid-20s (2006 to 210), but also includes some variables assessed in childhood or adolescence that have previously been useful in predicting initial alcohol use and misuse (e.g., (Hawkins et al., 1992)). Participants were interviewed within 3 months of their birthday each year. Across all waves, 81.9% of all possible interviews were completed (74% to 94% at any particular wave).

Procedures

The parent (biological mother for 83% of interviews) and subject were interviewed by trained interviewers separately until the subject was 16, and subjects only thereafter. Before the interviews began, parent and child signed informed consent forms approved by the Duke University Medical Center Institutional Review Board. Each parent and child received an honorarium for their participation.

Assessment

All variables except criminal offenses were assessed using the *Child and Adolescent Psychiatric Assessment* (CAPA) until age 16, and its upward extension, the *Young Adult Psychiatric Assessment* (YAPA) thereafter (Angold and Costello, 2000, Angold et al., 1999). These structured interviews were coded by a trained interviewer and each interview was then checked by a supervisor. A detailed glossary provides the operational rules for each item assessed. The time frame for determining the presence of diagnostic criteria was the *three months* immediately prior to the interview to minimize forgetting and recall biases, although onset dates were also collected for all symptoms. Most variables were assessed during late adolescence (ages 19 or 21).

Scoring programs written in SAS by the senior authors combined information about the date of onset, duration, and intensity of each symptom to create diagnoses according to the fourth edition of the Diagnostic and Statistical Manual of the American Psychiatric Association. A symptom was counted as present if reported by either parent or child or both, as is standard

clinical practice. Two-week test-retest reliability of diagnoses is comparable to that of other highly-structured child psychiatric interviews (Angold and Costello, 1995). Cohen's kappa value for substance abuse/dependence was 1.0 and the intraclass correlation for substance abuse/dependence symptoms was 0.98. Construct validity as judged by 10 different criteria including comparison to well-established scales for detecting psychopathology, ability to predict mental health service use, consistency of diagnoses over time, similarity of rates and age and gender patterns of disorder to those found using other interview. Validity across all criteria was good to excellent (Angold and Costello, 2000).

Alcohol Persistence—Persistence and desistance were defined based upon alcohol abuse and dependence diagnostic status in late adolescence and young adulthood. All subjects who met DSM-IV criteria for alcohol abuse or dependence at ages 19 or 21 assessments were included in this analysis. Alcohol abuse was diagnosed without application of the dependence exclusion criterion, allowing participants to be positive for both alcohol abuse and/or dependence. Participants with a late adolescent disorder who continued to meet criteria for an alcohol disorder (abuse or dependence) at the age 24/25/26 assessment were labeled *persisters* and participants who did not continue to meet criteria at the later assessment were labeled *desisters*.

Educational and early family role transitions—To confirm patterns observed in past studies, role transitions were tested as predictors of alcohol desistance (Staff et al., 2010, Bachman et al., 2002, Littlefield et al., 2009, Littlefield et al., 2010, Yamaguchi and Kandel, 1985). Parenthood, marriage, and post-secondary education were assessed through self-report during subject interviews in late adolescence (ages 19 and 21).

Predictors of Persistent Problems—Most predictors were assessed at the two late adolescent assessments (ages 19 and 21), but a number of risk factors associated with initiation of alcohol use and misuse were included from childhood/adolescence. The phenomenology of alcohol use was assessed via age of onset, frequency of use, alcohol disorder before late adolescence, each symptom of late adolescent alcohol abuse and dependence, and associated features of problematic use (i.e., craving, use of alcohol to improve mood, narrowed substance repertoire, use in the mornings, blackouts, and unconsciousness). Service use for alcohol-related problems was identified using the Child and Adolescent Services Assessment (Ascher et al., 1996), a portion of the CAPA/YAPA that assesses for the use of 39 types of service use.

Four categories of psychiatric disorders that have been associated with alcohol disorders in past research (Hasin et al., 2007) were tested: late adolescent anxiety disorders (separation anxiety, generalized anxiety, social phobia, specific phobia, agoraphobia, panic disorder, obsessive-compulsive disorder, and posttraumatic stress disorder), depressive disorders (major depression, dysthymia, depressive disorder, NOS), antisocial personality disorder, and childhood/adolescent conduct disorder. We also tested other substance problems as predictors of persistence, including any childhood/adolescent substance-related impairment, cannabis use, cannabis abuse, cannabis dependence, and other illicit drug use (e.g., opioids, cocaine, mushrooms, sedatives).

The following risk factors have also been associated with initial risk for alcohol use and transition to impairing use (Guo et al., 2000, Kaplow et al., 2001, Kosterman et al., 2000, Pagan et al., 2006, Tarter, 2002, Webb et al., 1991): First, socioeconomic status of the participant, including falling below the federal poverty line, high school education only, and low occupational prestige as defined by the NORC General Social survey (Nakao and Treas, 1992). Second, criminal contact was assessed from official charges collected from the North Carolina Administrative Offices of the Court Database and self-reported police contact.

Third, two types of negative events were assessed: events involving violence (e.g., physical abuse, violent death of a loved one) and events involving loss (e.g., parental divorce, breaking up with a boy/girlfriend). Finally, parent functioning variables were assessed during parent interviews during the subject's childhood and adolescence, including low family SES, parental mental health treatments, parental treatment for alcohol/drug use, and parental criminality. Additional details about any predictor variables are available from the first author upon request.

Analyses

All associations were tested using weighted logistic regression models in a generalized estimating equations framework implemented by SAS PROC GENMOD. Robust variance (sandwich type) estimates were used to adjust the standard errors of the parameter estimates for the stratified design effects. Therefore, the parameters reported here are representative of the population from which the sample was drawn.

Results

Consistent with past research, the 3-month likelihood of having of alcohol disorders (abuse or dependence) increased markedly from early adolescence to late adolescence (defined as ages 19 and 21) and then decreased into the mid-20s (figure 1a). No alcohol disorders were diagnosed before age 13. Across all ages, males displayed higher rates of alcohol disorders than females (odds ratio=3.3, 95% CI=2.0, 5.5, $p<0.001$). There were no differences in the rate of having had an alcohol disorders by race/ethnicity ($p=0.91$). Weekly alcohol use rates increased in a manner similar to alcohol disorder rates across adolescence, however, weekly use rates did not decrease at the same rate as alcohol disorder rates in young adulthood (figure 1b).

Sixteen-point-two percent (unweighted $N = 175$) of the overall sample met criteria for an alcohol disorder at one of the two assessments in late adolescence. One-hundred forty-four of those subjects (or 82.3%) were interviewed again at the mid-20s assessment. The follow-up rates were similar for participants without an alcohol disorder in late adolescence indicating no differential attrition (82.3% vs. 79.7%, $p=0.63$). Of those with a late adolescent alcohol disorder, 33.9% (unweighted $N = 37$) continued to meet criteria for an alcohol disorder in young adulthood (again, either abuse or dependence), whereas 66.1% (unweighted $N = 107$) desisted. Because persisters and desisters only constituted a subset of the total sample, only moderate to large effects were detectable (odds ratios > 3.0) in between-group comparisons. Despite this, persisters had elevated rates of concurrent marijuana use (74.2% vs. 0.0%, $p<0.0001$), illicit drug use (34.2% vs. 2.4%, $p<0.0001$), and depressive (9.2% vs. 0.0%, $p<0.0001$), and anxiety disorders (15.7% vs. 2.6%, $p<0.0001$) as compared to desisters. Persistent alcohol problems placed subjects at risk for a range of impairing concurrent comorbidities.

Educational and Early Family Role Transition

Prior studies suggest family and to a lesser extent education-related transitions may prove affect risk for persistent alcohol problems. Three late adolescent (ages 19 and 21) transitions were tested: parenthood, marriage, and post-secondary education. None of these potential protective factors were significantly associated with persistence/desistance (parenthood: 11.6% vs. 18.2% (desister rate first), $p=0.45$; marriage: 6.4% vs. 16.3%, $p=0.14$; post-secondary education: 50.9% vs. 24.7%, $p=0.07$), but all three interacted significantly with sex (figure 2). Females were more likely to desist if they were a parent (only at trend level) or married, whereas males were more likely to desist if they completed at least 2 years of

college. In contrast, marriage during late adolescence was a risk factor for persistence in males.

Predictors of Persistent Alcohol Problems

Compared to subjects with no history of alcohol problems in either late adolescence or young adulthood, both persisters and desisters were elevated on almost every risk index in late adolescence. To identify a risk profile of persistent alcohol problems, rates of risk factors were compared between persisters and desisters only. Neither sex nor race/ethnicity was significantly associated with persistent alcohol problems, but both variables were included in all analyses as covariates to minimize potential confounding. The onset of alcohol use was not significantly earlier in persisters than in desisters (8.9 (SD=6.0) vs. 11.1 (SD=4.1), $p=0.13$) and persisters were not more likely to have had alcohol problems in childhood (16.6% for persisters vs. 6.2% for desisters, $p=0.25$). (Some predictors, like childhood alcohol problems, were significant predictors in univariate models, but not when accounting for sex and race.) Both groups displayed similar rates of alcohol dependence at late adolescence assessments (43.6% for persisters vs. 35.4% for desisters, $p=0.59$), but persisters had a higher rate of alcohol abuse than desisters (98.2% vs. 88.5%, $p=0.04$). Use of specialty services tailored to problematic alcohol use was vanishingly rare in both groups (less than 1%).

Table 1 compares rates of items used to code for late adolescent alcohol abuse and dependence criteria and associated features between persisters and desisters. This analysis tests the importance of specific aspects of alcohol abuse and dependence in predicting persistence/desistence. Of abuse symptoms, three were significantly more common in persisters than desisters: alcohol-related poor judgment, social problems with friends, and dangerous activities. No symptoms of alcohol dependence were significantly elevated in persisters. Three associated features of alcohol dependence, although not formal criteria, predicted long-term persistence: craving alcohol, a narrowed substance repertoire and drinking to unconsciousness. Two additional aspects of drinking behavior were tested: any recent binge drinking and frequency of binge drinking. Almost all subjects - desisters or persisters alike - had reported binge drinking behavior, but the mean number of recent binge drinking episodes was higher in the persisters than desisters (10.0 vs. 18.4, $p=.05$). When all significant alcohol-related behaviors were tested as predictors simultaneously, all alcohol abuse symptoms and a narrowed substance repertoire continued to predict persistent alcohol problems.

Table 2 compares persisters and desisters in terms of additional domains of late adolescent function, many of which are known to be associated with onset of alcohol use and misuse: psychiatric functioning, other substance problems, socioeconomic status, crime, negative events, and parent history. Unless otherwise indicated, the domains were measured in late adolescence. Of all non-substance psychiatric outcomes tested, only late adolescent depressive disorders predicted persistent alcohol problems. Both cannabis dependence and illicit drug use increased risk for persistent problems. Across other domains, self-reported police contact and exposure to violence increased risk for persistence. None of these indicators were required to be related to alcohol use. When these predictors were tested together, cannabis dependence, self-reported police contact, and exposure to violence continued to predict persistence.

Moderation by Sex

It is possible that predictors might have different effects for males or females as was observed for the life transition analysis. To test this moderation hypothesis, models were rerun including an interaction term between sex and each predictor. The number of

significant interaction terms was only slightly more than would be expected by chance (3 of 43 tests) suggesting limited evidence of sex-specific predictors.

Cumulative Risk

A final model was tested which included all significant alcohol-related and other predictors. All variables except illicit drug use continued to predict persistence status independently. These items were then combined into a summary risk scale. Figure 3 displays the relationship between this cumulative late adolescent risk scale and the likelihood of displaying persistent alcohol abuse or dependence problems in young adulthood. Over 90% of subjects with 3 or more risk factors met criteria for a young adult alcohol disorder. This cutoff balanced high specificity (92.9%) with good sensitivity (66.2%).

Discussion

The increase in rates of alcohol disorders seen in late adolescence is both dramatic and time-limited. Despite high rates of continued alcohol use and vanishingly low rates of service utilization, over half of late adolescents with an alcohol disorder had desisted by their mid-20s. Thus, desistance from alcohol disorders was the norm. The question for this study was why a substantial minority of people in their mid 20s have persistent alcohol disorders. This question can only be addressed in a large community sample with prospective assessment of alcohol misuse across adolescence and young adulthood.

This study supported the importance of family and educational transitions in understanding risk for long-term alcohol problems. Risk factors predicting persistent alcohol disorders in this study generally differed from risk factors for initial problems that have been identified in previous research as most risk factors for persistence were secondary to alcohol use. Several criteria for alcohol abuse in late adolescence predicted persistence of alcohol disorders into the mid 20s as did some associated features of dependence. No dependence criteria predicted persistence. A few risk factors that are typically associated with early alcohol misuse also predicted persistent problems including depression, police contact and exposure to violence. Aspects of parent functioning, SES and psychiatric problems that are potent predictors of early alcohol use and abuse did not predict persistent problems.

Only by combining results from prior studies have we previously been able to infer the discrepancy between trajectories of alcohol use and disorder rates into young adulthood (e.g., (Harford et al., 2005, Chen and Kandel, 1995, Grant et al., 2004)). When reported together in GSMS, alcohol use and disorder levels both peak in late adolescence, but the substantial desistance of alcohol disorders is not mirrored in similar decreases in weekly alcohol use rates, suggesting that desistance of alcohol disorders is not an artifact of diminished use. Despite licit exposure to alcohol throughout adulthood, other studies suggest that rates of alcohol abuse and dependence decrease to the point of being relatively rare (<3% in males and <1% in females) in later adulthood (Harford et al., 2005). Late adolescence appears to be a risky period for alcohol exposure. It is notable that highest rates of criminal behavior (Sampson and Laub, 2003, Farrington, 1983) and traffic infractions (Harrington and McBride, 1970) also occur during late adolescence and late adolescence immediately *precedes* a period of intensive cortical myelination that diminishes impulsivity and improves decision-making (Thompson et al., 2000, Casey et al., 2000). Risk for alcohol problems may be greatest immediately following initial exposure and attenuate with repeated exposure. To test this *habituation* hypothesis, one would have to compare conditional risk for alcohol problems following alcohol initiation in different developmental periods.

Many of the strongest risk factors for persistent alcohol problems were alcohol abuse criteria rather than the associated psychosocial risk factors often associated with initial alcohol use. It is not surprising that behaviors directly related to alcohol use and other substances replaced other psychosocial risk factors in terms of understanding long-term problematic alcohol use. Other typologies used to characterize alcohol trajectories have done so on the basis of age of onset, use severity, or associated problems such as conduct problem (Cloninger, 1988, Zucker et al., 1995, Maggs and Schulenberg, 2004/2005, Babor et al., 1992). With the exception of binge drinking frequency, this study found little support for these factors in predicting DSM-based trajectories.

Criteria for alcohol abuse but no criteria for alcohol dependence discriminated persistent alcohol problems from time-limited impairing use. This is consistent with item response analyses of abuse and dependence criteria which found that these criteria form a continuum of disorder severity with abuse and dependence criteria intermixed (Saha et al., 2006, Martin and Srihari, 2006). These findings suggest two recommendations for current efforts at diagnostic refinement. First, these findings support the DSM-V proposal to combine abuse and dependence criteria into a single alcohol-use disorder. Second, associated features of alcohol dependence such as craving, narrowed use repertoire, and drinking to unconsciousness may have better predictive validity than some of the dependence criteria themselves do. These features should be subject to further testing to determine whether they merit inclusion as criteria in the revised alcohol-use disorder.

The risk scale derived from this study for predicting persistent alcohol problems was proficient both at identifying true positive and true negatives - a rare balance for medical tests (Kraemer, 1992). By comparison, the first three consumption items of the AUDIT are a commonly used screener for alcohol disorders. In a recent study based on the NESARC database, the optimal AUDIT-C cutpoint achieved similar specificity (92.5%) and higher sensitivity (73.6%) for predicting *current* alcohol abuse or dependence (Dawson et al., 2005). The risk scale in the current study focused on status *five or more years later*. Persistence was uncommon for those with 2 or fewer risk factors, but nearly universal for those with 3 or more of the identified risk factors. The predictive value of any scale, however, requires replication before it can be widely recommend, but these initial results are promising, and the contents of the scale involve information routinely collected in the clinical evaluation of alcohol problems. A tool with such predictive value could serve as a useful screen for late adolescents at risk for a prolonged course of alcohol problems and prioritizing subjects for intervention – goals that are relevant for both clinicians and those developing selective and indicated interventions.

Caveats

The Great Smoky Mountains study has several strengths besides its longitudinal, prospective design. The key variables, alcohol abuse and dependence, were assessed prospectively from middle childhood through adolescence and young adulthood into adulthood. A population-based design minimized selection biases. A wide range of functional domains in young adulthood were assessed to test as possible predictors. Despite these strengths, the sample is not representative of the U.S. population (Native Americans overrepresented and African Americans and Latinos underrepresented).

The five year follow-up identified a substantial decline in alcohol problems, but the subsequent course of problematic alcohol use is unknown, introducing the possibility of misclassification (e.g., identified desisters may still meet criteria for alcohol abuse and dependence again in the future). Furthermore, the 3-month assessment period used in this study minimized forgetting and recalls biases, but also missed diagnostic cases outside of the assessment window. Despite these potential misclassification issues, strong predictors of

persistent alcohol problems were identified suggesting that any misclassification bias was minimal. It is possible to question our definition of alcohol problems as involving a diagnosis of either abuse or dependence. Other subtypes and classes have been proposed to characterize alcohol use, some theory-based and others based upon modeling techniques. It is possible that alternative group definitions might affect the profile of significant late adolescent predictors. Our definition, however, is consistent with the DSM-V proposed changes to alcohol disorders and those in this combined category in the current study displayed widespread psychiatric and substance comorbidities, confirming the severity of their condition. Finally, the nature of the research question limited the number of subjects included in the analyses. As such, it was only possible to detect moderate to large effects and more modest elevations in other areas were not detectable. This concern may be particularly relevant to the sex by predictor interaction analyses where only interactions of large effect were detectable. It is important to note that GSMS is one of only a few large prospective, community studies that can address this issue.

Conclusion

It is uncommon for late adolescents with alcohol problems to receive any mental health services and rarer still to receive best-practice services, yet most did not display alcohol problems 5 years later. The late adolescent spike in alcohol problems is time-limited. This is not to say that treatment is unnecessary. Alcohol problems are associated with distress and impairments even if the disease-course is brief. This does suggest, however, that persistence is abnormal. Such persistence is associated with multiple comorbidities and such highly comorbid conditions are among the most costly to society (Costello et al., 2007). Fortunately, this study suggests that late adolescents at risk for persistent problems may be identified by aspects of their alcohol use, other substance use, and other associated behaviors and targeted for intensive intervention.

Acknowledgments

The work presented here was supported by the National Institute of Mental Health (MH63970, MH63671, MH48085), the National Institute on Drug Abuse (DA/MH11301), and the William T Grant Foundation. No authors have any conflicts of interest to report. Mr. Dlamini's work on this manuscript predated his employment at Regeneron and is entirely unrelated to his work there.

References

- Angold A, Costello E. The Child and Adolescent Psychiatric Assessment (CAPA). *Journal of the American Academy of Child and Adolescent Psychiatry*. 2000; 39:39–48. [PubMed: 10638066]
- Angold A, Costello EJ. A test-retest reliability study of child-reported psychiatric symptoms and diagnoses using the Child and Adolescent Psychiatric Assessment (CAPA-C). *Psychological Medicine*. 1995; 25:755–762. [PubMed: 7480452]
- Angold, A.; Cox, A.; Prendergast, M.; Rutter, M.; Simonoff, E.; Costello, EJ.; Ascher, BH. *The Young Adult Psychiatric Assessment (YAPA)*. Durham, NC: Duke University Medical Center; 1999.
- Ascher BH, Farmer EMZ, Burns BJ, Angold A. The Child and Adolescent Services Assessment (CASA): Description and psychometrics. *Journal of Emotional and Behavioral Disorders*. 1996; 4:12–20.
- Babor TF, Hofmann M, Delboca FK, Hesselbrock V, Meyer RE, Dolinsky ZS, Rounsaville B. Types of alcoholics, I: Evidence of an empirically derived typology based on indicators of vulnerability and severity. *Archives of General Psychiatry*. 1992; 49:599–608. [PubMed: 1637250]
- Bachman, JG.; O'Malley, PM.; Sscheulenberg, JE.; Johnston, LD.; Bryant, AL.; Merline, AC. *The decline of substance use in young adulthood: changes in social activities, roles and beliefs*. Mahwah, NJ: Erlbaum; 2002.
- Casey BJ, Giedd JN, Thomas KM. Structural and functional brain development and its relation to cognitive development. *Biological Psychology*. 2000; 54:241. [PubMed: 11035225]

- Chen K, Kandel D. Predictors of cessation of marijuana use: an event history analysis. *Drug and Alcohol Dependence*. 1998; 50:109–121. [PubMed: 9649962]
- Chen K, Kandel DB. The natural history of drug use from adolescence to the mid-thirties in a general population sample. *American Journal of Public Health*. 1995; 85:41–47. [PubMed: 7832260]
- Cloninger, CR. National Institute on Drug Abuse Research Monograph Series 89:Biological Vulnerability to Drug Abuse. NIDA, Office of Science; 1988. Etiologic factors in substance abuse: An adoption study perspective.
- Costello EJ, Angold A, Burns B, Stangl D, Tweed D, Erkanli A, Worthman C. The Great Smoky Mountains Study of Youth: Goals, designs, methods, and the prevalence of DSM-III-R disorders. *Archives of General Psychiatry*. 1996; 53:1129–1136. [PubMed: 8956679]
- Costello EJ, Copeland W, Cowell A, Keeler G. Service Costs of Caring for Adolescents With Mental Illness in a Rural Community, 1993–2000. *American Journal of Psychiatry*. 2007; 164:36. [PubMed: 17202542]
- Costello EJ, Mustillo S, Erkanli A, Keeler G, Angold A. Prevalence and development of psychiatric disorders in childhood and adolescence. *Archives of General Psychiatry*. 2003; 60:837–844. [PubMed: 12912767]
- Dawson DA, Goldstein RB, Grant BF. Rates and Correlates of Relapse Among Individuals in Remission From DSM-IV Alcohol Dependence: A 3-Year Follow-Up. *Alcoholism: Clinical and Experimental Research*. 2007; 31:2036–2045.
- Dawson DA, Grant BF, Stinson FS, Zhou Y. Effectiveness of the Derived Alcohol Use Disorders Identification Test (AUDIT-C) in Screening for Alcohol Use Disorders and Risk Drinking in the US General Population. *Alcoholism: Clinical and Experimental Research*. 2005; 29:844–854.
- De Bruijn C, Van Den Brink W, De Graaf R, Vollebergh WAM. The three year course of alcohol use disorders in the general population: DSM-IV, ICD-10 and the Craving Withdrawal Model. *Addiction*. 2006; 101:385–392. [PubMed: 16499511]
- Delucchi KL, Matzger H, Weisner C. Alcohol in emerging adulthood: 7-year study of problem and dependent drinkers. *Addictive Behaviors*. 2008; 33:134–142. [PubMed: 17537582]
- Farrington, DP. Offending from 10 to 25 years of age. In: Vandusen, KT.; Mednick, SA., editors. *Prospective Studies of Crime and Delinquency*. Boston, MA: Kluwer-Nijhoff; 1983.
- Grant BF, Dawson DA, Stinson FS, Chou SP, Dufouf MC, Pickering RP. The 12-month prevalence and trends in DSM-IV alcohol abuse and dependence: United States, 1991–1992 and 2001–2002. *Drug and Alcohol Dependence*. 2004; 74:223–234. [PubMed: 15194200]
- Grant BF, Stinson FS, Harford T. The 5-year course of alcohol abuse among young adults. *Journal of Substance Abuse*. 2001; 13:229–238. [PubMed: 11693449]
- Guo J, Collins LM, Hill KG, Hawkins JD. Developmental pathways to alcohol abuse and dependence in young adulthood. *Journal of Studies on Alcohol*. 2000; 61:799–808. [PubMed: 11188485]
- Harford TC, Grant BF, Yi HY, Chen CM. Patterns of DSM-IV alcohol abuse and dependence criteria among adolescents and adults: results from the 2001 National Household Survey on Drug Abuse. *Alcohol Clin Exp Res*. 2005; 29:810–818. [PubMed: 15897727]
- Harrington DM, McBride RS. Traffic violations by type, age, sex, and marital status. *Accident Analysis & Prevention*. 1970; 2:67–79.
- Hasin D, Grant B, Endicott J. The natural history of alcohol abuse: implications for definitions of alcohol use disorders. *Am J Psychiatry*. 1990; 147:1537–1541. [PubMed: 2221170]
- Hasin DS, Stinson FS, Ogburn E, Grant BF. Prevalence, Correlates, Disability, and Comorbidity of DSM-IV Alcohol Abuse and Dependence in the United States: Results From the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2007; 64:830–842. [PubMed: 17606817]
- Hasin DS, Van Rossem R, McCloud S, Endicott J. Differentiating DSM-IV alcohol dependence and abuse by course: Community heavy drinkers. *Journal of Substance Abuse*. 1997; 9:127–135. [PubMed: 9494944]
- Hawkins JD, Catalano RF, Miller JY. Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychological Bulletin*. 1992; 112:64–105. [PubMed: 1529040]

- Jackson KM, O'Neill SE, Sher KJ. Characterizing Alcohol Dependence: Transitions During Young and Middle Adulthood. *Experimental and Clinical Psychopharmacology*. 2006; 14:228–244. [PubMed: 16756427]
- Kandel DB, Raveis VH. Cessation of illicit drug use in young adulthood. *Archives of General Psychiatry*. 1989; 46:109–116. [PubMed: 2913970]
- Kaplow JB, Curran PJ, Angold A, Costello EJ. The prospective relation between dimensions of anxiety and the initiation of adolescent alcohol use. *Journal of Clinical Child Psychology*. 2001; 30:316–326. [PubMed: 11501249]
- Kosterman R, Hawkins J, Guo J, Catalano R, Abbott R. The dynamics of alcohol and marijuana initiation: Patterns and predictors of first use in adolescence. *American Journal of Public Health*. 2000; 90:360–366. [PubMed: 10705852]
- Kraemer, HC. *Evaluating Medical Tests: Objective and Quantitative Guidelines*. Sage Publications, Inc; 1992.
- Littlefield A, Sher K, Wood P. Is “maturing out” of problematic alcohol involvement related to personality change? *Journal of Abnormal Psychology*. 2009; 118:360–374. [PubMed: 19413410]
- Littlefield AK, Sher KJ, Wood PK. A personality-based description of maturing out of alcohol problems: Extension with a Five-Factor model and robustness to modeling challenges. *Addictive Behaviors*. 2010; 35:948–954. [PubMed: 20598445]
- Maggs JL, Schulenberg JE. Trajectories of Alcohol Use During the Transition to Adulthood. *Alcohol Research & Health*. 2004/2005; 28:195–201.
- Martin A, Srihari V. Geometrically evident: framing studies using the graphic appraisal tool for epidemiology (GATE). 2006; 45:1521–1526.
- Nakao, K.; Treas, J. *GSS Methodological Report No 74*. Chicago, Illinois: National Opinion Research Center; 1992. The 1989 Socioeconomic Index of Occupations: Construction from the 1989 Occupational Prestige Scores.
- Pagan JL, Rose RJ, Viken RJ, Pulkkinen L, Kaprio J, Dick DM. Genetic and Environmental Influences on Stages of Alcohol Use Across Adolescence and into Young Adulthood. *Behavior Genetics*. 2006; 36:483. [PubMed: 16586152]
- Pirkola SP, Poikolainen K, Lonnqvist JK. Currently active and remitted alcohol dependence in a nationwide adult general population--results from the Finnish Health 2000 study. *Alcohol and Alcoholism*. 2006; 41:315–320. [PubMed: 16492724]
- Rumpf HJ, Bischof G, Hapke U, Meyer C, John U. Studies on natural recovery from alcohol dependence: sample selection bias by media solicitation? *Addiction*. 2000; 95:765–775. [PubMed: 10885051]
- Saha TD, Chou SP, Grant BF. Toward an alcohol use disorder continuum using item response theory: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Psychological Medicine*. 2006; 36:931–941. [PubMed: 16563205]
- Sampson R, Laub J. Life-course desisters? Trajectories of crime among delinquent boys followed to age 70. *Criminology*. 2003; 41:555–592.
- Schuckit MA, Smith TL, Landi NA. The 5-Year Clinical Course of High-Functioning Men With DSM-IV Alcohol Abuse or Dependence. *Am J Psychiatry*. 2000; 157:2028–2035. [PubMed: 11097971]
- Smart, R. Natural Recovery or Recovery without Treatment from Alcohol and Drug Problems as Seen from Survey Data. In: Klingemann, H.; Sobell, LC., editors. *Promoting Self-Change From Addictive Behaviors*. Springer; US: 2007.
- Staff J, Schulenberg JE, Maslowsky J, Bachman JG, O'Malley PM, Maggs JL, Johnston LD. Substance use changes and social role transitions: Proximal developmental effects on ongoing trajectories from late adolescence through early adulthood. *Development and Psychopathology*. 2010; 22:917–932. [PubMed: 20883590]
- Tarter RE. Etiology of adolescent substance abuse: a developmental perspective. *American Journal on Addictions*. 2002; 11:171–91. [PubMed: 12202010]
- Thompson PM, Giedd JN, Woods RP, Macdonald D, Evans AC, Toga AW. Growth patterns in the developing brain detected by using continuum mechanical tensor maps. *Nature*. 2000; 404:190–3. [PubMed: 10724172]

- Webb JA, Baer PE, McLaughlin RJ, McKelvey RS, Caid CD. Risk factors and their relation to initiation of alcohol use among early adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1991; 30:563–568. [PubMed: 1890089]
- Yamaguchi K, Kandel DB. On the Resolution of Role Incompatibility: A Life Event History Analysis of Family Roles and Marijuana Use. *American Journal of Sociology*. 1985; 90:1284–1325.
- Zucker RA. Anticipating problem alcohol use developmentally from childhood into middle adulthood: what have we learned? *Addiction*. 2008; 103:100–108. [PubMed: 18426543]
- Zucker, RA.; Fitzgerald, HE.; Moses, HD. Emergence of alcohol problems and the several alcoholisms: A developmental perspective on etiologic theory and life course trajectory. In: CICCHETTI, D.; COHEN, DJ., editors. *Developmental Psychopathology: Volume 2 - Risk, Disorder, and Adaptation*. New York, NY: John Wiley & Sons, Inc; 1995.

A)

Figure 1.

Three-month prevalence rates for a) alcohol abuse or dependence and b) at least weekly alcohol use. Assessment schedule and participation rates are available from table 1.

Figure 2.

Comparison of males and females rates of persistence and desistance for protective factors. Graphs compare rates for (a) 2 or more years of post-secondary education, (b) marriage, and (c) parenthood. All interaction terms were statistically significant (post-secondary education: $p=0.03$; marriage: not able to be estimated due to no cases of persistence in married females; parenthood: $p=0.04$). Post-secondary education was protective for males (OR=0.2, 95%CI 0.0, 0.9; $p=0.05$) but not for females (OR=3.6, 95%CI 0.5, 27.1; $p=0.20$). Marriage increase risk for persistence in males (OR=12.3, 95%CI 2.1, 71.5; $p=0.005$) but was protective in female (odds ratio could not be estimated as *no female persists were married*). Despite a significant interaction term, parenthood was not associated with persister/desister status for females (OR=0.2, 95%CI 0.0, 1.4; $p=.09$) or males (OR=2.7, 95%CI 0.5, 15.5; $p=.25$).

* Indicates significant group difference.

Figure 3.

Association between cumulative scale of late adolescent risk factors and 3-month mid 20s alcohol disorder rates. Controlling for sex and race/ethnicity, the number of late adolescent risk factors significantly predicted young adult persistence status (OR=8.4, 95%CI 4.5, 15.7; $p<.0001$).

Table 1

Alcohol-related predictors of alcohol abuse/dependence persistence from young adulthood to one's mid-20s

	Desisters		Persisters		Odds Ratio	95% CI	p
	%	SE	%	SE			
Predictor variables							
Alcohol abuse variables							
Disinhibited aggression	39.0	8.0	17.4	9.3	0.3	0.1–1.5	0.144
Disinhibited sexuality	35.3	7.7	34.7	11.6	1.0	0.3–3.5	0.959
Poor judgment	24.1	6.8	53.5	12.2	5.1	1.4–18.3	0.011
Social problems at home	7.8	4.3	11.3	7.1	1.6	0.3–9.5	0.609
Social problems with friends	0.9	0.6	21.1	10.5	36.7	5.2–257.2	0.000
Affects school/work functioning	9.1	5.0	3.2	1.7	0.3	0.1–1.6	0.161
Dangerous activities	35.5	7.6	68.2	10.8	4.0	1.2–13.2	0.023
Criminality	12.4	5.4	10.8	6.5	0.8	0.2–4.3	0.822
Trouble with the law	5.2	3.3	1.9	1.1	0.3	0.1–2.0	0.216
Alcohol dependence variables							
Tolerance	34.3	7.5	42.1	12.1	1.6	0.5–5.3	0.450
Excessive time spent	54.9	8.0	69.7	11.6	1.9	0.5–6.8	0.317
Overconsumption	51.9	8.0	48.2	12.3	0.9	0.3–2.8	0.821
Uncontrollable use	26.1	7.2	47.2	12.3	2.5	0.7–8.7	0.137
Attempts to cut back	43.1	8.0	51.3	12.3	1.3	0.4–4.3	0.619
Reduced activities	34.3	7.5	42.1	12.1	1.6	0.5–5.3	0.450
Physical problems	3.3	2.9	0.3	0.3	0.1	0.0–1.9	0.127
Other alcohol-related behaviors							
Craving for alcohol	5.8	3.1	23.1	10.6	5.0	1.1–22.0	0.035
Use of alcohol to improve mood	25.0	6.7	33.5	11.4	1.7	0.5–6.3	0.410
Narrowed substance repertoire	0.7	0.5	14.6	9.3	39.1	4.0–379.9	0.002
Use in the morning	6.8	3.7	8.3	6.9	1.1	0.1–9.3	0.946
Blackouts	13.6	4.8	26.1	10.8	2.5	0.6–10.2	0.200
Unconsciousness	25.6	6.9	55.7	12.2	3.9	1.1–14.1	0.042
Daily use	22.3	5.8	43.9	12.1	2.6	0.8–8.4	0.103

Withdrawal criterion for alcohol dependence was excluded due to low prevalence. All variables assesses in young adulthood (ages 19 and 21).

All variables assesses in young adulthood unless otherwise indicated.

Table 2

Predictors of alcohol abuse/dependence persistence from young adulthood to the mid-20s

	Desisters		Persisters				
Predictor variables	%	SE	%	SE	Odds Ratio	95% CI	p
Psychiatric functioning							
Anxiety disorders	13.0	5.3	9.2	7.0	0.9	0.1–6.6	0.923
Depressive disorders	3.1	1.1	11.8	7.6	4.8	1.1–21.4	0.039
Antisocial personality disorder	8.8	4.8	15.6	8.9	1.9	0.3–12.4	0.499
Child/Adolescent conduct disorder	12.3	5.1	20.0	8.8	1.6	0.4–6.9	0.529
Other Substance use/problems							
Child/Adolescent substance problems	18.0	5.5	28.1	10.9	1.7	0.5–6.2	0.396
Nicotine use	57.6	8.0	61.8	12.2	1.1	0.3–3.8	0.938
Cannabis use	71.0	7.3	84.6	9.3	2.3	0.5–10.9	0.281
Cannabis abuse	35.4	7.8	43.6	12.1	1.4	0.4–4.6	0.586
Cannabis dependence	7.5	3.5	27.7	10.8	8.8	2.1–36.8	0.003
Other illicit drug use	17.2	4.5	49.8	12.3	5.0	1.5–16.6	0.008
Socio-economic status variables							
Poverty	50.3	8.0	65.7	11.7	2.1	0.6–7.4	0.257
Low education	79.7	5.9	65.7	11.6	0.5	0.1–1.9	0.324
Low occupational status	34.0	7.6	34.3	11.6	1.0	0.3–3.5	0.960
Crime							
Official criminal offenses	51.2	8.0	60.2	12.1	1.2	0.4–4.0	0.787
Self-reported police contact	49.8	8.0	77.0	10.7	3.5	1.1–11.3	0.036
Negative events							
Exposure to violence	4.8	3.7	29.6	11.6	8.3	1.1–61.2	0.038
Negative events involving loss	30.8	7.5	24.6	10.6	0.7	0.2–2.9	0.666
Parental functioning							
Parental low socioeconomic status	23.1	6.4	18.3	9.1	0.7	0.1–3.5	0.651
Parental mental health problems	63.7	7.9	55.3	12.3	0.7	0.2–2.2	0.517
Parental substance problems	17.7	5.6	21.1	9.2	1.2	0.3–4.7	0.744

	Desisters		Persisters		Odds Ratio	95% CI	p
	%	SE	%	SE			
Predictor variables							
Parental criminality	45.7	8.0	44.3	12.2	0.9	0.3-2.7	0.797

All variables assessed in young adulthood (ages 19 and 21) unless otherwise indicated.