Published in final edited form as:

J Cell Physiol. 2012 December ; 227(12): 3857–3875. doi:10.1002/jcp.24100.

$\alpha_M\beta_2$ -integrin-intercellular adhesion molecule-1 interactions drive the flow-dependent trafficking of Guillain-Barré syndrome patient derived mononuclear leukocytes at the blood-nerve barrier *in vitro*

Nejla Yosef, M.S.¹ and Eroboghene E. Ubogu, M.D.^{1,*}

¹Neuromuscular Immunopathology Research Laboratory, Department of Neurology, Baylor College of Medicine, Houston, Texas, United States of America

Abstract

The mechanisms of hematogenous leukocyte trafficking at the human blood-nerve barrier (BNB) are largely unknown. Intercellular adhesion molecule-1 (ICAM-1) has been implicated in the pathogenesis of Guillain-Barré syndrome (GBS). We developed a cytokine-activated human *in vitro* BNB model using primary endoneurial endothelial cells. Endothelial treatment with 10 U/mL tissue necrosis factor- α and 20 U/mL interferon- γ resulted in *de novo* expression of proinflammatory chemokines CCL2, CXCL9, CXCL11 and CCL20, with increased expression of CXCL2-3, CXCL8 and CXCL10 relative to basal levels. Cytokine treatment induced/ enhanced ICAM-1, E- and P-selectin, vascular cell adhesion molecule-1 and the alternatively spliced pro-adhesive fibronectin variant, fibronectin connecting segment-1 expression in a time-dependent manner, without alterations in junctional adhesion molecule-A expression. Lymphocytes and monocytes from untreated GBS patients express ICAM-1 counterligands, α_M - and α_L -integrin, with differential regulation of α_M -integrin expression compared to healthy controls. Under flow conditions that mimic capillary hemodynamics *in vivo*, there was a >3-fold increase in total GBS patient and healthy control mononuclear leukocyte adhesion/ migration at the BNB following cytokine treatment relative to the untreated state. Function neutralizing monoclonal antibodies against human α_M -integrin (CD11b) and ICAM-1 reduced untreated GBS patient mononuclear leukocyte trafficking at the BNB by 59% and 64.2% respectively. Monoclonal antibodies against α_L -integrin (CD11a) and human intravenous immunoglobulin reduced total leukocyte adhesion/ migration by 22.8% and 17.6% respectively. This study demonstrates differential regulation of α_M -integrin on circulating mononuclear cells in GBS, as well as an important role for α_M -integrin-ICAM-1 interactions in pathogenic GBS patient leukocyte trafficking at the human BNB *in vitro*.

Keywords

Adhesion molecules; Blood-nerve barrier; Cytokines; Guillain-Barré syndrome; Integrins; Leukocytes

Address all correspondence to: Eroboghene E. Ubogu, M.D., Department of Neurology, Baylor College of Medicine, One Baylor Plaza, Mailstop NB302, Houston, TX 77030-3411, United States of America. Telephone Number: +1-713-798-4654, Facsimile Number: +1-713-798-3676, ubogu@bcm.edu.

Supplementary material

Supplementary material is available on-line.

INTRODUCTION

Guillain-Barré syndrome (GBS) is an immune-mediated disorder restricted to peripheral nerves and nerve roots characterized by varying degrees of weakness, sensory loss and autonomic dysfunction. GBS is the most common cause of rapidly progressive paralysis worldwide, and often results in significant morbidity or death from respiratory failure, cardiac arrhythmias or complications arising from prolonged immobility such as pulmonary embolism. Several variants exist, classified based on clinical and electrophysiological features. The most common variant in industrialized countries, acute inflammatory demyelinating polyradiculoneuropathy (AIDP) is characterized by intense hematogenous mononuclear leukocyte infiltration into the endoneurium of peripheral nerves and nerve roots, temporally associated with demyelination and axonal degeneration (Hartung et al., 2002; Hughes and Cornblath, 2005; Kieseier et al., 2004). These features are also observed in a representative animal model, experimental autoimmune neuritis (EAN) (Meyer zu Hörste et al., 2007; Xia et al., 2010a).

Observational studies of AIDP-affected patient nerve biopsies show that monocytes/macrophages are the predominant leukocytes present in affected lesions, followed by T-lymphocytes and B-lymphocytes. Macrophage-mediated segmental demyelination is the pathological hallmark of AIDP (Kiefer et al., 2001). CD4⁺ T-helper 1 (Th1) cells support peripheral nerve inflammation by secreting pro-inflammatory cytokines and chemokines that activate monocytes and induce further hematogenous leukocyte recruitment. CD8⁺ cytotoxic T-cells may directly damage myelin and axons. Antibodies secreted by circulating or infiltrated activated B-lymphocytes or plasma cells in further contribute to the immune-mediated attack on the peripheral nerves and nerve roots (Gold et al., 1999; Kieseier et al., 2004; Meyer zu Hörste et al., 2007). B-lymphocytes are also potent antigen-presenting cells capable of *in situ* T-cell activation in peripheral nerves (Dalakas, 2008). Elevated numbers of myeloid and plasmacytoid dendritic cells have also been described in the cerebrospinal fluid of AIDP patients (Press et al., 2005).

The precise mechanisms of leukocyte infiltration into human peripheral nerves and nerve roots are undetermined. Leukocyte infiltration across microvascular endothelium is a complex and coordinated process driven by shear forces and the interaction between selectins (such as P-selectin and E-selectin), chemokines, and cell adhesion molecules (such as intercellular adhesion molecule-1 [ICAM-1], vascular cell adhesion molecule-1 [VCAM-1], junctional adhesion molecule-A [JAM-A]) expressed on the endothelium; and carbohydrate ligands (such as P-selectin glycoprotein-1 and sialyl Lewis x), chemokine receptors, integrins (such as $\alpha_L\beta_2$ [also known as CD11a/CD18 or leukocyte function-associated antigen {LFA}-1], $\alpha_M\beta_2$ [also known as CD11b/CD18 or Mac-1], and $\alpha_4\beta_1$ [also known as CD49d/CD29 or very late antigen (VLA)-4 integrin] and matrix metalloproteases (MMPs) expressed by infiltrating leukocytes. The end result is the transmigration of leukocytes from the bloodstream to sites of infection, or tissue inflammation or injury (Alon and Ley, 2008; Cinamon et al., 2001; Man et al., 2007; Simon and Green, 2005).

Observational studies in GBS patient nerves, cerebrospinal fluid and sera, as well as in EAN have shown a potential causal association between selective cytokines (e.g. tissue necrosis factor [TNF]- α , interferon [IFN]- γ , interleukin (IL)-1 β , IL-2 and IL-12), chemokines (such as CCL2 [also known as monocyte chemoattractant protein {MCP}-1], CXCL10 [also known as IFN- γ inducible protein-10 kDa {IP-10}] and their receptors CCR2 and CXCR3; cell adhesion molecules (e.g. ICAM-1) and matrix metalloproteases (e.g. MMP-9) in disease progression and maximal clinical severity (Hadden et al., 2001; Kieseier et al., 2004; Kieseier et al., 2000; Kieseier et al., 2002; Lu and Zhu, 2010; Meyer zu Hörste et al., 2007; Nyati et al., 2010; Orlikowski et al., 2003; Press et al., 2003; Putzu et al., 2000; Xia et al.,

2010b). These data imply that hematogenous leukocyte infiltration into peripheral nerve and nerve root endoneurium is pathologically relevant in AIDP.

Endoneurial microvessels and the perineurium are responsible for regulating the molecular composition of the endoneurium; a process required for normal axonal impulse transmission. These anatomical components consist of specialized cells that possess intercellular tight junctions and lack fenestrations. Endoneurial microvessels restrict the passive diffusion of solutes, macromolecules and leukocytes from circulating blood. These microvessels can be considered as the blood-nerve barrier (BNB). The perineurium restricts the passive diffusion of interstitial fluid components from the epineurium (which has fenestrated macrovessels) into the endoneurium (Olsson, 1990; Reina et al., 2000; Reina et al., 2003). Hematogenous leukocytes need to interact with and migrate across the BNB in order to carry out effector functions in AIDP.

Using a recently established *in vitro* BNB model developed with *primary human endoneurial endothelial cells* (pHEndECs) that form this restrictive barrier *in vivo* (Yosef et al., 2010), we sought to determine mechanisms relevant to pathogenic leukocyte trafficking into peripheral nerves. We demonstrated that pro-inflammatory cytokines up-regulate or induce the expression of selective chemokines and cell adhesion molecules at the BNB, enhancing the total adhesion/ migration of untreated GBS (AIDP-variant) mononuclear leukocytes compared to the untreated BNB *in vitro*. Interestingly, cytokine treatment of the BNB significantly enhanced healthy control mononuclear leukocyte trafficking at the BNB, implying a more important role for endoneurial endothelial activation in leukocyte trafficking at the BNB than leukocyte activation state. ICAM-1 was responsible for cytokine-mediated GBS patient mononuclear leukocyte adhesion/ migration, dependent on interactions with α_M -integrin (CD11b). Despite ubiquitous mononuclear leukocyte α_L -integrin (CD11a) expression, this molecule played a minor role in GBS leukocyte trafficking at the human BNB *in vitro*. Differential regulation of CD11b expression was observed in GBS mononuclear leukocyte subpopulations, suggesting selective trafficking of CD11b+ leukocytes into peripheral nerves and nerve roots *in vivo*. We provide insights into the effects of cytokine stimulation at the human BNB and the molecular determinants of pathogenic leukocyte trafficking that may be relevant to the pathogenesis of AIDP and other causes of peripheral nerve inflammation.

MATERIALS AND METHODS

Human BNB model

pHEndECs were isolated within 24 hours of death from the sciatic nerves of individuals with no known history of peripheral nervous system disease or systemic illness, by endoneurial stripping, enzymatic digestion and density centrifugation, as previously published (Yosef et al., 2010). The endothelial cells were cultured on rat tail collagen (RTC)-coated Cell BIND® tissue culture plates (Corning, Lowell, MA) expanded in growth media and cryopreserved at passages 3–5 using a controlled rate cryopreservation method. pHEndECs (passage 6–8) were seeded on RTC-coated Corning CellBIND® tissue culture plates, polyester membrane transwell inserts (Corning, Lowell, MA) or autoclaved borosilicate glass coverslips (VWR, Chester, PA) for subsequent experiments.

Patient selection and leukocyte isolation

Six consecutive patients with clinical, electrophysiological and cerebrospinal fluid evidence of GBS (AIDP variant) were evaluated by a board-certified electrodiagnostic and neuromuscular specialist (E.E.U.) at Ben Taub General Hospital and prospectively recruited into the study for whole blood donation. The clinical characteristics of these patients are

summarized in Table 1. There were 5 men and 1 woman with an average age of 43.3 years (range 26–62). Time from disease onset to blood sampling varied between 4–8 days (median 7 days). Mean cerebrospinal fluid white cell count was 7 (2–18, normal <10/ μ L) with a mean protein concentration of 96.4 mg/dL (range 52.2–146.0; normal 15–45). Five of the six patients were eventually treated with plasmapheresis and one with human intravenous immunoglobulin (IVIg). 4 age-matched individuals (3 men and 1 woman) with no history of chronic or recent systemic illness served as healthy controls.

Informed consent was obtained to perform a single blood draw via the antecubital vein prior to immune-modulatory treatment. The protocol was approved by the Baylor College of Medicine Institutional Review Board and Harris County Hospital District Research and Sponsored Programs Office. Peripheral blood mononuclear leukocytes (PBMLs) were isolated from 60–80 mL of whole heparinized blood per patient using Ficoll® density centrifugation, as previously published (Mahad et al., 2006; Ubogu et al., 2006a). Isolated PBMLs were enumerated using a hemacytometer, reconstituted at 10^7 cells/ mL and cryopreserved using a controlled rate process in Cell Freezing Medium-1X dimethylsulfoxide (Sigma-Aldrich, St. Louis, MO) and stored in liquid nitrogen for future use. Leukocyte viability was >99% following reconstitution in warmed media at 37°C using the 0.4% trypan blue exclusion test (Altman et al., 1993). Duration of storage had no effect on leukocyte viability based on this test.

Flow cytometry

To determine the effect of cytokine treatment on pHEndEC viability, we seeded pHEndECs at 15,000 cells/ cm^2 on 6-well RTC-coated Corning CellBIND® tissue culture plates and allowed them to grow to confluence over 4–5 days. Near-confluent cultures were treated with 0–100 U/mL human recombinant TNF- α and 0–200 U/mL IFN- γ (both from R&D Systems [Minneapolis, MN]: 1.1×10^5 U/ μ g TNF- α and 2.0×10^4 U/ μ g IFN- γ) in growth media for 24 hours and harvested using 0.05% trypsin-0.53 mM EDTA in 1X Hank's Balanced Salt Solution (Mediatech, Manassas, VA) for 30–60 seconds at 37°C in humidified incubator containing 95% air and 5% carbon dioxide. Harvested pHEndECs were washed and reconstituted at 500,000 cells/ mL per sample in 1% bovine serum albumin (BSA) in 1X phosphate buffer saline (PBS). 20 μ L of 7-Amino-actinomycin D (7-AAD) staining solution (BD Biosciences, San Jose, CA) was added to each vial and thoroughly mixed for 3 minutes to detect non-viable cells by flow cytometry. Data were collected using a FACS Canto flow cytometer (BD Biosciences, San Jose, CA), gated according to previously established endothelial cell side and forward scatter characteristics and analyzed relative to unstained controls using FlowJo® version 7.2.5 software (Tree Star, Ashland, OR). A minimum of 25,000 events were analyzed per sample.

To determine ICAM-1 expression (a known marker of endothelial cell activation) at the human BNB following cytokine treatment *in vitro*, we treated near confluent pHEndEC cultures with regular growth media or media containing the maximum concentration of TNF- α and IFN- γ that did not reduce cell viability for 24 hours. Endothelial cells were harvested and reconstituted in 100 μ L of 1:9 vol./vol. solution of 0.2 mg/mL normal goat immunoglobulin (Invitrogen, Carlsbad, CA) to block non-specific antibody binding, as previously described (Yosef et al., 2010). Monoclonal mouse anti-human ICAM-1 IgG2b antibodies (clone LB-2, Santa Cruz Biotechnology, Santa Cruz, CA) were added to each sample at 1 μ g per 10^6 cells in 1% BSA in 1X PBS for 15 minutes at room temperature (RT), followed by a wash and incubation with 2 μ g per 10^6 cells R-PE-conjugated goat anti-mouse IgG2b antibodies (SouthernBiotech, Birmingham, AL) in 1% BSA in 1X PBS for 15 minutes at RT in the dark. Following a final wash, data acquisition and analysis of unfixed pHEndECs were performed as above (in triplicate), with appropriate unstained and isotype controls.

To determine surface integrin expression on subpopulations of isolated PBMLs from the six untreated GBS patients and four healthy controls, cryopreserved vials were thawed to 37°C in a water bath, washed by centrifugation in assay buffer (Dulbecco's PBS +2% normal goat serum + 0.1% sodium azide) and reconstituted at 10^6 cells per test. The viability of cryopreserved GBS PBMLs was also determined by flow cytometry using 7-AAD, as previously described. Following non-specific antibody blockade with normal goat IgG for 15 minutes, PBMLs were incubated with 5 µg of mouse anti-human CD11a IgG2a (anti- α_L integrin, clone 38; AbD Serotec, Oxford, United Kingdom), 0.5 µg of mouse anti-human CD11b IgG1 (anti- α_M integrin, clone ICRF44; BioLegend, San Diego, CA) or 1.0 µg of mouse anti-human activated CD11b (aCD11b) IgG1-RPE (anti- α_M integrin, clone CBRM1/5; Santa Cruz Biotechnology, Santa Cruz, CA) in 100 µL of assay buffer containing 10^6 cells at RT for 15 minutes.

Following washes, PBMLs were incubated with secondary antibodies goat anti-mouse IgG2a-RPE (for α_L -integrin) or goat anti-mouse IgG1-RPE (for α_M -integrin) [both from SouthernBiotech, Birmingham, AL] at a dose of 0.5 µg per 10^6 cells (none with aCD11b) and a cocktail containing the following mouse anti-human monoclonal antibodies (all from BD Biosciences, San Jose, CA): CD3-FITC (clone UCHT1, 15 µL), CD4-PerCP (clone SK3, 15 µL) or CD8-PerCP (clone SK1, 15 µL), CD14-APC (clone MØP9, 5 µL) and CD19 APC-H7 (clone SJ25C1, 5 µL) in the dark at RT for 15 minutes. Following a final wash, acquisition and analysis of unfixed PBMLs by flow cytometry was performed as above, detecting 100,000 events per sample. Appropriate unstained and isotype controls were used as negative controls. Data were obtained at least in triplicate from three independent experiments per GBS patient/control and averaged to reduce potential sampling variations in leukocyte subpopulation CD marker and integrin expression. Data are displayed as a % of parent population or mean fluorescent intensity (MFI; a marker of surface receptor density).

Polymerase chain reaction (PCR)

PCR was performed to determine the effect of cytokine treatment on ICAM-1 gene transcript expression at the human BNB *in vitro*. Cartridge purified primers and all reagents and reaction kits were purchased from Invitrogen (Carlsbad, CA), unless otherwise stated. Total RNA was obtained from confluent cultured pHEndECs treated with 0–100 U/mL TNF- α and 0–200 U/mL IFN- γ for 24 hours using TRIzol® reagent according to the manufacturer's instructions. Total RNA was spectrophotometrically quantified using a nanodrop® ND-1000 UV/Vis spectrophotometer (NanoDrop Technologies Wilmington, DE) as previously published (Yosef et al., 2010). Primer sequences for ICAM-1 have been recently published (Yosef et al., 2010).

Reverse transcription and PCR amplification were carried out using a TC-412 Thermal Cycler (Techne, Burlington, NJ). Single stranded cDNA was synthesized using the Superscript II® Reverse Transcriptase kit according to the manufacturer's instructions, and PCR performed as previously published (Yosef et al., 2010). Omission of specific primers served as negative controls while glyceraldehyde-3-phosphate dehydrogenase (GAPDH) amplification served as an internal control. PCR products were resolved and visualized on 1% ultra-pure agarose gels stained with ethidium bromide, with DNA markers verifying size. Digital images were generated with the same exposure times and settings using an AlphaImager HP image documentation system (Cell Biosciences, Santa Clara, CA) attached to a Sony ICX267AL 1.39 Megapixel CCD camera. Complementary DNA expression was quantified by spot densitometry (based on integrated density values with background correction) relative to GAPDH expression using the AlphaView software program (version 3.2.2.0).

Transendothelial electrical resistance (TEER) measurements

Cytokine treatment may alter tight junction protein expression and reduce endothelial barrier resistance, facilitating leukocyte extravasation. In order to determine whether cytokine treatment altered human BNB resistance *in vitro*, TEER was directly measured on confluent pHEndECs seeded at 250,000/cm² in growth media on RTC-coated Corning polyester membrane transwell inserts (6.5 mm diameter, 3.0 µm pore size) in 24-well tissue culture plates on day 7 using an epithelial volttohmmeter with STX2 electrodes (World Precision Instruments, Sarasota, FL) as previously described (Yosef et al., 2010). Measurements were made following treatment with or without TNF-α and IFN-γ for 24 hours. Comparisons were made relative to untreated inserts to determine the effect of cytokine treatment on human *in vitro* BNB resistance.

Chemokine antibody array

To determine and quantify chemokine expression by the *in vitro* human BNB under basal conditions and following cytokine treatment, we extracted cytoplasmic proteins from confluent pHEndECs cultured on 6-well RTC-coated Corning CellBIND® tissue culture plates and performed a protein array using the RayBio® Human Chemokine Antibody Array 1 kit (Raybiotech, Norcross, GA) according to the manufacturer's instructions. Briefly, array membranes spotted with human chemokine antibodies were blocked and incubated with protein extracts from untreated and cytokine-treated pHEndECs diluted 1:10 in blocking buffer at RT for 2 hours. After washing, membranes were incubated with a cocktail of biotinylated human chemokine antibodies in blocking buffer at RT for 2 hours and then incubated with horseradish peroxidase (HRP)-conjugated streptavidin in blocking buffer overnight at 4°C following further washes.

Membranes were treated with a 1:1 mix of detection buffers for 2 minutes at RT, then exposed to the same x-ray film for 40 seconds and processed with an SRX-101 A film processor (Konica Minolta, Tokyo, Japan). Digital images of generated autoradiographs were taken using an AlphaImager HP image documentation system (Cell Biosciences, Santa Clara, CA) attached to a Sony ICX267AL 1.39 Megapixel CCD camera. Protein expression was determined by spot densitometry (based on integrated density values with background correction) relative to internal array controls using the AlphaView software program (version 3.2.2.0) to determine the effect of cytokine treatment on chemokine expression at the human BNB *in vitro*.

Western blot

Cytoplasmic proteins were extracted from pHEndEC cultures initially grown to confluence in 6-well RTC-coated Corning CellBIND® tissue culture plates, followed by treatment with a combination of TNF-α and IFN-γ for 0–48 hours, to determine cell adhesion molecule expression at the human BNB *in vitro*. Extraction was performed by gentle scraping in cold lysis buffer (10 mM KCl + 10 mM Hepes + 0.1 mM EDTA + 0.1 mM EGTA + 1mM DDT + 0.5 mM PMSF + 0.05% NP-40 in deionized distilled water) containing protease inhibitor cocktail (1:300 dilution; Sigma-Aldrich, St. Louis, MO) on ice, following by brief sonication and centrifugation at 18,000g for 20 minutes at 4°C. Protein extracts were quantified using the Bradford Coomassie assay and stored at –80°C. Equal amounts of protein (6 µg) were denatured, reduced and separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (10% slab gel) using the Bio-Rad MiniProtean III Apparatus (Hercules, CA), then electrotransferred to polyvinylidene fluoride membranes in transfer buffer (1:20 of 5X Towbin's buffer [144g Glycine, 30.25g Tris, 2g SDS, made up to 1L with distilled water] + 10% methanol in distilled water) using the Bio-Rad Trans-Blot Apparatus (Hercules, CA), according to manufacturer instructions.

Membranes were blocked with 5% non-fat milk (NFM) in Tris buffered saline + 0.05% Tween-20 (TBS-T) for 1–2 hours at RT, followed by incubation with primary antibodies diluted in TBS-T containing 1–5% NFM or 5% goat serum (1:100–1:200) overnight at 4°C. The following antibodies (all from Santa Cruz Biotechnology [Santa Cruz, CA] unless otherwise stated) were used: mouse anti-human E-selectin IgG1(clone A-10), rabbit anti-human P-selectin polyclonal IgG, mouse anti-human ICAM-1 IgG2b (clone LB-2), mouse anti-human VCAM-1 IgG1 (clone E-10), rabbit anti-human JAM-A polyclonal IgG (Invitrogen, Carlsbad, CA), mouse anti-human fibronectin connecting segment-1 (FN CS-1) IgM (clone 2Q607) and mouse anti-human GAPDH IgG1 (clone 0411). Following several washes, membranes were incubated with HRP-conjugated polyclonal goat anti-mouse IgG, goat anti-mouse IgM or polyclonal goat anti-rabbit IgG secondary antibodies (Jackson ImmunoResearch, West Grove, PA) diluted TBS-T/ 1% NFM (1:5,000–1:10,000 dilution) for 1–3 hours at RT. Negative controls were performed and binding specificity determined by omitting primary antibodies. Molecular weight markers verified protein size.

After washing, membranes were treated with Pierce® Enhanced Chemiluminescence western blotting substrate plus detection reagents (Thermo Scientific, Rockford, IL) and visualized by generating autoradiographs following film exposure from 1 minute to 2 hours using an SRX-101A film processor (Konica Minolta, Tokyo, Japan). Digital autoradiographs were taken and processed as described above. Cell adhesion molecule protein expression was quantified by spot densitometry (based on integrated density values with background correction) relative to GAPDH expression from concurrently developed films imaged at the same time point using the AlphaView software program. Data from three independent experiments (performed in duplicate) were averaged to determine the time-dependent effect of cytokines on cell adhesion molecule protein expression at the human BNB.

Indirect immunocytochemistry

pHEndECs were cultured to confluence on glutaraldehyde (GTA) cross-linked RTC-coated borosilicate glass coverslips placed in 24-well Corning costar tissue culture plates. Autoclaved glass coverslips were treated with a 1:1 mix of 0.2% GTA in 1X PBS and 50 µg/mL Type I RTC (BD Biosciences, San Jose, CA) in 0.02N acetic acid for 1 hour at RT. Following two washes with 1X PBS, coverslips were treated with 2% glycine for 5 minutes to neutralize residual GTA, washed again with PBS and gel-coated with RTC in ammonia vapor as previously described (Yosef et al., 2010).

Following further PBS washes, pHEndECs were seeded at >30,000 cells/ cm² for 4–5 days in growth media, then treated further with or without TNF-α and IFN-γ for 24 hours. Indirect immunocytochemistry was performed on 4% paraformaldehyde-fixed pHEndECs with the same antibodies used for western blotting, as previously published (Yosef et al., 2010). Isotype-matched FITC or Alexa Fluor 488-conjugated goat anti-mouse/ goat anti-rabbit antibodies (SouthernBiotech, Birmingham, AL or Invitrogen, Carlsbad, CA) were used as secondary antibodies. Mounted coverslips were visualized using an Axioskop Epifluorescent microscope equipped with an Axiocam MRc 5 digital camera (Carl Zeiss Microimaging, Thornwood, NY). Images were taken using the same exposure time settings, initially processed with the Zeiss Axiovision software and merged using Adobe Photoshop CS2 (Adobe Systems, San Jose, CA). Cell adhesion molecule surface expression following cytokine treatment was compared with untreated endothelial cells.

Flow-dependent leukocyte trafficking assays

We sought to determine mechanisms of pathogenic leukocyte trafficking at the BNB *in vitro*. We seeded pHEndECs at >25,000 cells/ cm² in the center of RTC-coated 35 mm Corning CellBIND® tissue culture plates initially demarcated with non-cytotoxic

transparent CultureWell™ silicone well dividers (Electron Microscopy Sciences, Hartfield, PA) for 24 hours. Four to six days after plating, confluent cells were treated with or without TNF- α and IFN- γ in growth media for 15–26 hours and placed at 37°C in a humidified incubator with 95% air, 5% carbon dioxide. Cryopreserved PBMLs from untreated GBS patients and controls were thawed to 37°C in a water bath, washed in assay buffer (1% BSA in RPMI-1640 media) by centrifugation at 300 g for 5 minutes at RT in an Allegra X-12R centrifuge (Beckman Coulter, Fullerton, CA) and resuspended at 10^7 cells/mL. 2×10^6 cells were aliquoted per Eppendorf tube and stored on ice prior to experiments (to limit temperature-dependent chemokine receptor internalization). Prior to use, aliquoted cells were further washed in assay buffer by centrifugation at 10,000 revolutions per minute for 30 seconds at RT in a Beckman Coulter benchtop centrifuge, warmed to 37°C for 5 minutes, resuspended at 500,000 cells/mL in assay media and placed in a 10 mL Luer-lock syringe (BD Biosciences, San Jose, CA).

PBMLs were infused into a parallel plate flow chamber (GlycoTech, Rockville, MD) attached with a silicone gasket (5.0 mm wide $\times$ 0.25 mm thick) that was firmly placed in each 35 mm tissue culture dish containing confluent pHEndECs (i.e. the human BNB model) using a Smith Medical (St. Paul, MN) Medfusion Protégé 3010 infusion pump at a rate of 4.5 mL/ hour and withdrawn at the same rate using a KD scientific (Holliston, MA) model 210 pump to achieve a constant linear flow velocity of 1 mm/s, an estimated flow rate in capillaries (Nishimura et al., 2006) for 30 minutes. The human leukocyte-BNB model flow system was mounted on a Zeiss Axiovert 40 Inverted Microscope. Leukocyte trafficking events on a frame of view of 650 μ m $\times$ 870 μ m were captured in real-time by taking brightfield digital images every second with a Zeiss AxioCam MRC 5 digital camera. Consecutive images were merged and videos generated at 10 frames per second using the National Institutes of Health Image J software program.

Adherent cells were defined as cells that firmly attached to the model BNB surface, and cells that attached, spread and moved on the model BNB surface without detaching, penetrating its surface or leaving the frame of view and remained phase bright throughout the duration of the experiment. Migrated cells were defined as cells that attached and underwent phase change from dark to bright (implying surface penetration of the model BNB) during the experiment (Gopalan et al., 2001; Man et al., 2009; Schreiber et al., 2007). The total number of adherent and migrated PBMLs was counted in three non-adjacent fields and averaged per experiment to account for observed regional variations in leukocyte trafficking. Comparisons were made between PBML adhesion/migration with (i.e. cytokine activated) or without cytokine treatment (i.e. basal) from each GBS patient and control, and expressed as a percentage relative to mean basal cell counts from the same patient.

In another series of experiments, 2 million GBS patient PBMLs/mL were pre-incubated for 5 minutes at 37°C with different concentrations of function-neutralizing mouse anti-human CD11a IgG2a (anti- α_L integrin, clone 38; AbD Serotec, Oxford, United Kingdom; 0.02–100 μ g/mL), mouse anti-human CD11b IgG1 (anti- α_M integrin, clone ICRF44; BioLegend, San Diego, CA; 0.2–20 μ g/mL), mouse anti-human ICAM-1 IgG2b (Santa Cruz, Santa Cruz, CA; 0.02–20 μ g/mL), human IVIg (Carimune® nanofiltered, CSL Behring, King of Prussia, PA; 20 mg/mL) and mouse IgG1 isotype antibody (clone 11711, R&D Systems, Minneapolis, MN, 0.2–20 μ g/mL) in assay media alone or in combination. Trafficking assays were performed as above using the cytokine-activated human BNB model. In some experiments, the BNB model endothelial cells were also treated with 1 μ g/mL mouse anti-human ICAM-1 or isotype antibody in assay media for 10 minutes at 37°C in an incubator. The total numbers of adherent/migrated PBMLs were quantified by video microscopy as described above and expressed as a percentage of mean uninhibited cell counts from the same untreated GBS patient studied.

Flow-dependent adhesion assay

In order to determine the GBS patient leukocyte subpopulations that firmly adheres to/ migrates across the BNB following cytokine treatment *in vitro*, we performed direct immunocytochemistry on tissue cultures plates with low autofluorescence after an *in vitro* adhesion assay. We cultured pHEndECs to confluence on RTC-coated ibiTreat, tissue culture treated, low walled 35-mm μ -dishes (ibidi USA, Verona, WI) over 5–6 days and treated with 10 U/mL TNF- α and 20 U/mL IFN- γ for 24 hours prior to experiments. Following two washes with warmed TEM buffer, 1 million PBMLs in 1 mL of TEM buffer were added to each dish and allowed to adhere at 37°C in a VWR incubating orbital shaker set at 60 revolutions/minute for 1 hour. Unbound leukocytes were aspirated and plates washed twice with Dulbecco's PBS, followed by fixation with 4% paraformaldehyde in 1X PBS for 10 minutes. After washing, dishes were permeabilized with ice cold acetone for 5 minutes at -20°C, washed, air-dried and blocked with 2% normal goat serum. Endothelial cells were lightly counterstained with 10 μ g/mL polyclonal rabbit anti-human zona occludens (ZO)-1 IgG and an Alexa-Fluor 488-conjugated goat anti-rabbit IgG (H+L) antibody (1:500 dilution, both antibodies from Invitrogen, Carlsbad, CA). A cocktail of containing FITC-conjugated mouse anti-human CD3, CD14 or CD19 antibodies with aCD11b-RPE (clone CBRM1/5) was added to independent dishes for 60 minutes in the dark at RT. After several washes, the dishes were treated with 0.45 μ M DAPI for 3 minutes at RT in the dark, washed further, followed by glass coverslips mounted in Vectashield placed in the middle of each dish. Stained dishes were processed as described previously in the immunocytochemistry section.

Statistical analysis

Statistical analyses (Student's t-test/ analysis of variance for parametric variables or Mann-Whitney's U-test for non-parametric variables, with data distribution curves verified by tests of skewness and Kurtosis) were performed using JMP® version 8 software (SAS Institute, Cary, NC), with statistical significance set at $p < 0.05$. Variations of the mean are expressed as standard errors.

RESULTS

10 U/mL TNF- α + 20 U/mL IFN- γ maximally stimulates the human *in vitro* BNB without compromising transendothelial electrical resistance

Flow cytometry studies demonstrated 95–97% pHEndEC viability in basal untreated cultures detected by lack of 7-AAD staining (Figure 1A). There was no significant increase in non-viable pHEndECs between basal, cells treated with 1U/mL TNF- α and 2 U/mL IFN- γ (Figure 1B) and cells treated with 10 U/mL TNF- α and 20 U/mL IFN- γ (Figure 1C). At higher doses, there was a progressive increase in non-viable cells detected (Figures 1D-F). These data suggest that significantly increased BNB endothelial cell apoptosis/ death may occur at 20 U/mL TNF- α and 40 U/mL IFN- γ *in vitro*.

Up-regulation in ICAM-1 expression is a well-described consequence of endothelial cell cytokine activation *in vitro* and *in vivo* (Ubogu et al., 2006a). By PCR, we demonstrated increased ICAM-1 mRNA expression by confluent pHEndECs relative to basal conditions following treatment with 10 U/ mL TNF- α and 20 U/mL IFN- γ for 24 hours, with reduced mRNA expression following treatment with 100 U/mL TNF- α and 200 U/mL IFN- γ relative to 10 U/ mL TNF- α and 20 U/mL IFN- γ -treated pHEndECs (Figure 2A). This data suggest a “U-shaped” dose-dependent pHEndEC ICAM-1 expression curve following pro-inflammatory cytokine treatment *in vitro*. To further confirm human BNB cytokine-mediated endothelial cell activation following treatment with 10 U/mL TNF- α and 20 U/mL IFN- γ , we demonstrated a mean 2.8-fold increase in ICAM-1-positive live unfixed

pHEndECs (mean 33.2% vs. 93.2%; $p=0.03$) associated with a 2.5-fold increase in MFI (mean 1625.3 vs. 4023.5) by flow cytometry relative to basal conditions (Figure 2B).

BNB compromise may occur during peripheral nerve inflammation (Kieseier et al., 2004). BNB breakdown could imply passive influx of activated leukocytes into the endoneurium in AIDP. Confluent pHEndECs on RTC-coated transwell inserts (with surface area of 0.33 cm²) showed a mean basal TEER of 116.7 Ω .cm², with a small increase following cytokine treatment ($p=0.002$, with 1U/mL TNF- α and 2 U/mL IFN- γ , and $p=0.001$, with 10 U/mL TNF- α and 20 U/mL IFN- γ) that returned to baseline following treatment with 100 U/mL TNF- α and 200 U/mL IFN- γ . There was no significant difference in TEER comparing 1U/mL TNF- α and 2 U/mL IFN- γ and 10 U/mL TNF- α and 20 U/mL IFN- γ treated confluent pHEndEC layers ($p=0.15$, Figure 2C). These data imply that pro-inflammatory cytokine treatment (TNF- α and IFN- γ) over a 100-fold range did not significantly compromise mean BNB TEER after 24 hours relative to untreated confluent endothelial layers *in vitro*.

These data imply that the *in vitro* human BNB model was significantly activated by 10 U/mL TNF- α and 20 U/mL IFN- γ without inducing cell death or compromising TEER relative to the untreated state. These cytokine concentrations within the range observed in the sera of GBS and septic patients *in vivo* (Kabir et al., 2003; Radhakrishnan et al., 2004). Cytokine treatment with 10 U/mL TNF- α and 20 U/mL IFN- γ was used for subsequent experiments designed to mimic pro-inflammatory events at the human BNB.

Cytokine treatment induces or up-regulates specific chemokines at the *in vitro* BNB

Cytokine treatment induced *de novo* cytoplasmic expression of CCL2, CCL20, CCL27, CXCL9 and CXCL11 at the BNB *in vitro*, as well as a 5.8-fold increase in CXCL2-3 ($p=0.0002$; there was no increase in CXCL1 alone), 4.4-fold increase in CXCL8 ($p<0.0001$) and 2.5-fold increase in CXCL10 ($p=0.0003$) expression. Induced expression of CCL2 is also a known marker of endothelial cell activation (Ubogu et al., 2006a). There were statistically significant, but less than 2-fold increases in CCL4 (1.7, $p=0.003$), CCL5 (1.4, $p=0.0004$), CCL23 (1.6, $p=0.0008$), CXCL5 (1.5, $p=0.0004$) and CXCL7 (1.7, $p=0.002$). CCL26 was reduced to 0.9-times its basal level ($p=0.02$). There was no statistically significant change in CCL22 and CCL24 levels ($p=0.06$ and 0.09 respectively), as shown in Figure 3. These observations suggest potential activation of the innate (i.e. neutrophil dependent: CXCL2-3, CXCL8 interacting with chemokine receptors CXCR1 and CXCR2; monocyte-dependent: CCL2 interacting with CCR2) and adaptive (i.e. T-cell-dependent: CXCL9, 10 and 11 interacting with CXCR3 on CD4+ T-helper 1 (Th1) cells, CCL20 interacting with CCR6 on CD4+ T-helper17 (Th17) cells and CCL27 interacting with CCR10 on activated T-cells) immune signaling pathways at the human *in vitro* BNB following physiological cytokine treatment.

Cytokine treatment induces or up-regulates specific cell adhesion molecules in a time-dependent manner at the *in vitro* BNB

In order to determine the time-dependent effect of cytokine treatment on endothelial cell adhesion molecule protein expression at the *in vitro* BNB, we performed western blot of pHEndEC protein lysates extracted from cells treated for 0–48 hours. There was increased E-selectin, P-selectin, ICAM-1, VCAM-1 and FN CS-1 expression by 12 hours following TNF- α and IFN- γ treatment that persisted at peak or near peak levels for up to 48 hours (Figure 4). There was no change in JAM-A expression (binds to leukocyte α_L -integrin), implying that this molecule was not regulated by physiological human BNB cytokine treatment *in vitro*. These data were further confirmed by indirect immunocytochemistry of confluent pHEndEC cultures (Figure 5).

CD11b integrin is differentially expressed on untreated GBS patient derived peripheral blood mononuclear leukocytes

Cryopreserved PBMLs from the 6 GBS patients and 4 controls had a mean viability of 97.0% (range 95.8 to 98.0%), consistent with viability data from freshly isolated PBMLs from normal healthy subjects following Ficoll separation from whole heparinized blood. Compared to healthy controls, there was an increase in the relative % of CD3+ CD4+ T-cells (54.8% vs. 40.8%) and reduction in % CD3+ CD8+ T-cells (42.1% vs. 52.4%) without significant change in the total % of T-cells (55.2% vs., 54.4%) in GBS-patient derived PBMLs (Figure 6A). This suggests a phenotypic shift consistent with the notion that GBS is a CD4+ T-cell dependent disorder. There were also significant increases in %CD14+ monocytes (7.3% vs. 4.3%) and CD19+ B-cells (24.5% vs. 14.4%) as shown in Figure 6A, implying effector roles of these leukocyte subpopulations in this untreated GBS patient cohort.

Interestingly, there were reductions in CD11b expression on untreated GBS patient-derived CD3+ CD4+ T-cells (mean 6.4% vs. 14.9%) and CD3+ CD8+ (mean 20.9% vs. 46.5%) compared with untreated controls (Figure 6B). There were no significant differences in % CD11b expression on CD14+ monocytes (mean 98.4% vs. 99.6%) and CD19+ B-cells (24.2% vs. 21.8%) between untreated GBS patients and healthy controls, as shown in Figure 6B. In addition, there were significant reductions in CD11b MFI in all untreated GBS-derived mononuclear leukocyte subpopulations compared to normal controls (Figure 7). These data raise the question whether there was selective migration of CD11b-high PBMLs into peripheral nerves and nerve roots in this cohort of GBS patients (thereby reducing relative proportions of these cells in the circulating leukocyte blood pool) or a negative regulatory role of CD11b in GBS.

Circulating PBMLs derived from GBS patients and healthy controls rarely express aCD11b. Less than 3% of CD14+ monocytes, 0.5% of CD19+ B-cells and 0.2% CD3+ CD4+ or CD8+ T-cells were aCD11b+ by flow cytometry. Representative dot plots demonstrating the differential expression of CD14+ monocytes (Figure 8), CD3+ CD4+ T-cells (Figure 9), CD3+ CD8+ T-cells (Figure 10) and CD19+ B-cells (Figure 11), including CD11b and aCD11b expression, in untreated GBS PBMLs and healthy controls are shown for comparative purposes.

Analysis of untreated GBS patient-derived PBMLs demonstrated >99% of CD14+ monocytes (range 99.40–99.93%) and >95% of CD 19+ B-lymphocytes (range 95.36–99.38%) expressed CD11a (α_L -integrin). Approximately 100% of CD3+ CD4+ (range 99.81–100%) and CD3+ CD8+ (range 99.83–100%) T-cells were also CD11a-positive. There was no significant difference in CD11a relative % and MFI compared with healthy controls (data not shown), suggesting that CD11a is not differentially regulated on GBS PBMLs. A representative dot plots depicting CD11a expression on PBMLs from an untreated GBS patient are shown in Figure 12. The ubiquitous expression of α_L -integrin on GBS PBML subpopulations suggests that this integrin could be pathogenically relevant to PBML trafficking across the human BNB.

Cytokine treatment enhances GBS patient PBML adhesion/migration at the *in vitro* BNB

The sequential process of leukocyte rolling, attachment, post-attachment locomotion, firm adhesion and transmigration was observed at the human *in vitro* BNB under flow. Initial adhesion occurred after 5 minutes from the start of the infusion and continued throughout the assay for at least 25 minutes. There were regional variations in adhesion/ migration, with foci of leukocytes aggregating in certain regions, seemingly attracting other PBMLs, as shown with GBS patient-derived PBMLs in Figure 13A–C. Compared to the untreated (i.e.

basal) BNB, cytokine treatment induced a mean 3.2-fold increase in GBS patient PBML adhesion/migration (range 1.4–5.9, $p=0.0001$) as shown in Figure 13D (and Supplementary videos 1 and 2). Similarly, cytokine treatment significantly enhanced healthy control PBML trafficking at the BNB under hydrodynamic forces relative to basal conditions (mean 3.3-fold; range 2.4–6.4, $p<0.0001$). Untreated GBS patient derived PBMLs adhered or migrated more than healthy controls under basal (61.9% increase, $p=0.02$) and cytokine treated (30.5% increase, $p=0.03$) BNB conditions. These data support the observation that GBS PBMLs are activated and “primed” towards trafficking at the BNB; however BNB activation is a stronger stimulus for trafficking *in vitro*.

GBS patient PBML adhesion/migration at the cytokine treated human *in vitro* BNB is dependent on α_M -integrin-ICAM-1 interactions

ICAM-1 expression had been previously demonstrated on endoneurial microvessels in GBS patient sural nerve biopsies (Putzu et al., 2000), suggesting an important role in disease pathogenesis. 1 $\mu\text{g/mL}$ anti-ICAM-1 antibody treatment to the BNB with concurrent PBML treatment at 10 ng per 10^6 cells for 5 minutes (with final concentration of 5 ng/mL during flow assay) resulted in a maximal mean 64.2% reduction in total PBML adhesion/migration relative to BNB cytokine activation without inhibitors (i.e. down to basal trafficking levels; $p<0.00001$). Surprisingly, antibodies against α_L -integrin (anti-CD11a) at 100 ng per 10^6 cells (final concentration of 50 ng/mL) resulted in a maximal mean 22.8% reduction in total PBML adhesion/migration ($p=0.007$). Antibodies against α_M -integrin (anti-CD11b) at 1 μg per 10^6 cells (final concentration of 500 ng/mL) resulted in a maximal mean 59% reduction in total GBS PBML adhesion/migration ($p=0.0001$), implying that this was the counter-ligand responsible for ICAM-1 mediated GBS leukocyte trafficking at the BNB *in vitro* (Figure 14, and Supplementary videos 3–5). Competitive inhibition occurred when maximal doses of anti-ICAM-1 and anti-CD11b antibodies were combined, resulting in less efficacious PBML adhesion/ migration blockade compared to either alone (mean 34.8% reduction, $p=0.01$ relative to anti-ICAM-1 and $p=0.02$ relative to anti-CD11b alone). This data suggest that a single function-neutralizing antibody is sufficient to block GBS PBML adhesion/ migration mediated by α_M -integrin-ICAM-1 interactions *in vitro*. Lower and higher anti-ICAM-1 and integrin antibody concentrations resulted in reduced blockade efficacy, implying a narrow therapeutic window (data not shown).

Human IVIg had a small, but statistically significant effect on GBS PBML adhesion/ migration at the BNB (mean 17.6% reduction; $p=0.02$ relative to cytokine-activated BNB without inhibitors) with a final concentration of 5 mg/mL during flow assay. The dose of IVIg was estimated based on currently recommended doses used to treat GBS patients (400 mg/kg/day, estimating that blood accounts for ~8% of body weight). Antibodies against ICAM-1 and α_M -integrin were more potent in modulating GBS PBML adhesion/migration at the cytokine activated *in vitro* BNB than IVIg ($p=0.0006$ relative to ICAM-1 and $p=0.001$ relative to CD11b), as shown in Figure 14. These data suggest that function-neutralizing antibodies against ICAM-1 and α_M -integrin competitively inhibit untreated pathogenic GBS patient PBML adhesion/migration more effectively than antibodies against α_L -integrin and human IVIg at the human *in vitro* BNB. We did not study the determinants of PBML trafficking at the BNB in normal healthy individuals, as hematogenous leukocytes are rarely observed in healthy persons based on studies in human and mouse peripheral nerves (Kieseier et al., 2002; Xia et al., 2010a).

GBS patient-derived CD14+ monocytes expressing activated CD11b preferentially adhere to the cytokine treated human *in vitro* BNB

Direct immunocytochemistry demonstrates adhesion/migration of untreated GBS-derived PBMLs to the *in vitro* BNB in clusters (Figure 15A) as demonstrated with video phase

contrast microscopy (Figure 13). The majority of adherent PBMLs expressed aCD11b (Figures 15B–D) in contrast to circulating PBMLs (as evidenced by flow cytometry data: Figures 8–11), implying chemoattractant-mediated CD11b-integrin activation during the trafficking process at the human BNB *in vitro*. Further studies demonstrated that CD14⁺ monocytes were the predominant adherent/migrated PBML subpopulation at the BNB (Figures 15E–H), consistent with observational data in human GBS and murine EAN peripheral nerves.

DISCUSSION

In this study, we provide insights into the response of the human BNB to local cytokine stimulus and show an important role for α_M -integrin-ICAM-1 interactions in GBS mononuclear leukocyte trafficking *in vitro*. Very few hematogenous leukocytes are present in the peripheral nerves of healthy individuals and mice (Kieseier et al., 2002; Xia et al., 2010a), so we focused on modeling pathogenic trafficking at the BNB *in vitro* using leukocytes derived from patients with organ-restricted peripheral nerve inflammation. There are phenotypic and functional differences between endothelial cells from different species and different tissues within the same species (Aird, 2007a; Aird, 2007b; Allt and Lawrenson, 2000; Bell and Weddell, 1984; Malmgren and Olsson, 1980; Poduslo et al., 1994; Yano et al., 2007; Yosef et al., 2010). Furthermore, as observed in peripheral nerves, significant differences exist between macrovascular and microvascular endothelium within the same tissue (Olsson, 1968; Yano et al., 2007). As a result, studies evaluating inflammatory signaling pathways relevant to the human BNB require human endoneurial endothelial cells.

Developing a physiological human *in vitro* BNB was essential to understanding the potential mechanisms of leukocyte-endothelial interactions during peripheral nerve inflammation *in vivo*. Elevated cytokine levels have been described in the sera and peripheral nerves of GBS patients and in EAN rodents. We demonstrated that 10 U/mL TNF- α and 20 U/mL IFN- γ sufficiently activated the human BNB model (based on ICAM-1 expression) without inducing endothelial cell death or compromising TEER, as similarly observed with an *in vitro* model of the human blood-brain barrier (Ubogu et al., 2006a). Pathogenic leukocyte infiltration across the BNB may not involve passive mechanisms secondary to cytokine-mediated endothelial injury.

Chemokines attract leukocytes to sites of inflammation *in vivo* and across concentration gradients *in vitro* (Ubogu et al., 2006b). Chemokines are also responsible for the arrest and activation of circulating leukocytes on vascular endothelium (inducing conformational changes between integrin subunits that convert them from a resting to an activated state capable of firm cell adhesion molecule binding) (Alon and Ley, 2008). We demonstrate induced CCL2 expression and a 2.5 fold increased CXCL10 expression by the BNB following cytokine treatment *in vitro*. CCL2 and CXCL10 have been implicated in the pathogenesis of GBS and EAN (Kieseier et al., 2000; Kieseier et al., 2002; Orlikowski et al., 2003; Press et al., 2003; Xia et al., 2010b). CCL2 signals via its receptor, CCR2, present on a majority of circulating monocytes/macrophages, while CXCL10 signals via its receptor CXCR3, preferentially expressed on CD4⁺ Th1 cells (Deshmane et al., 2009; Lacotte et al., 2009). Interestingly, other chemokine ligands that signal through CXCR3 (CXCL9 and CXCL11) were expressed *de novo* by the *in vitro* human BNB following cytokine treatment, suggesting a crucial role for effector T-cell trafficking at the cytokine activated BNB.

In addition, there was increased neutrophil-responsive CXCL2-3 and CXCL8 expression following cytokine treatment. Although neutrophils have been rarely described in GBS (Berciano et al., 1997; Rauschka et al., 2003), these chemokines could serve as arrest

chemokines necessary for integrin activation or participate in CD4⁺Th17 T-cell mediated signaling pathways, as suggested in central nervous system inflammation and possibly in chronic inflammatory demyelinating polyradiculoneuropathy (Carlson et al., 2008; Chi et al., 2010). To support the hypothesis that Th17-mediated signaling may be pathogenically relevant at the human BNB during inflammation, we demonstrated *de novo* expression of CCL20 which signals through CCR6, a chemokine receptor restricted to CD4⁺ Th17 cells (Kim, 2009). A small, but significant reduction in CCL26 (interacts with CCR3 present in CD4⁺ Th2 cells) was also observed. These data suggest that physiological doses of TNF- α and IFN- γ induce Th1 and Th17 chemokine expression with reduction in Th2 chemokine expression at the human BNB *in vitro*. The role of Th17-mediated inflammation at the human BNB requires further investigation. Chemokines expressed at high levels under basal conditions (e.g. CCL4 and CCL5) may participate in endoneurial homeostasis or immunosurveillance.

We demonstrated increased expression of E- and P-selectin, ICAM-1, VCAM-1 and FN CS-1 at the BNB following cytokine treatment. Selectins participate in leukocyte rolling on endothelial surfaces, while ICAM-1, VCAM-1 and FN CS-1 have been implicated in leukocyte adhesion and transmigration (Alon and Ley, 2008; Man et al., 2009; Man et al., 2007). JAM-A has been implicated in leukocyte adhesion and migration on cytokine-stimulated endothelium dependent on the nature of the inflammatory model (Nourshargh et al., 2006). Although pHEndECs constitutively expressed JAM-A, physiological cytokine treatment did not alter protein expression, implying this molecule may not participate in leukocyte trafficking at the BNB. Increased levels of soluble E-selectin, ICAM-1 and VCAM-1 have been described in the sera of GBS patients, with higher levels temporally associated with disease activity (Hadden et al., 2001). ICAM-1 (also known as CD54) is a cell surface glycoprotein and member of the immunoglobulin superfamily. Soluble ICAM-1 is a known marker of endothelial activation *in vivo*. Elevated levels of E-selectin may implicate shedding from endoneurial endothelial cells in GBS, and although the same may apply to ICAM-1 and VCAM-1, these adhesion molecules are also expressed on circulating leukocytes. ICAM-1 has been demonstrated on endoneurial microvessels in GBS and in EAN, while P-selectin has been demonstrated on endoneurial microvessels in EAN (Putzu et al., 2000; Tran et al., 2010). Blockade of ICAM-1 and VCAM-1 with monoclonal antibodies in active myelin-induced or adoptive transfer Lewis rat EAN altered disease course, implying potential efficacy in GBS (Archelos et al., 1993; Enders et al., 1998).

We demonstrated an important role for ICAM-1 in untreated GBS PBML adhesion/migration at the BNB under flow conditions. It interacts with either CD11a or α_L -integrin (expressed on all leukocyte subpopulations) or CD11b or α_M -integrin (expressed predominantly on monocytes/ macrophages, natural killer cells, neutrophils and a subset of CD8⁺ T-cells) non-covalently associated with CD18 (β_2 -integrin) to form heterodimers (Muller, 2011; Simon and Green, 2005). In addition to its role in cell adhesion, $\alpha_M\beta_2$ -integrin acts as a pattern recognition receptor that interacts with complement C3 fragment, iC3b, bound on the surface of microbes, resulting in complement-mediated phagocytosis. It can also bind foreign antigen directly resulting in non-opsonic phagocytosis (Le Cabec et al., 2002). Complement C3 activation products have been described on Schwann cell membranes in AIDP (Hafer-Macko et al., 1996; Nyland et al., 1981).

In contrast to CD11a, we observed differential expression of CD11b on GBS-derived PBMLs compared to healthy controls. We surprisingly observed that fewer CD3⁺ CD4⁺ and CD3⁺ CD8⁺ T-cells derived from this cohort of untreated GBS (AIDP) patients expressed CD11b. For all leukocyte subpopulations tested, CD11b MFI was also reduced in GBS patient leukocytes. These data could suggest that loss of CD11b expression on effector T-cells and reduced surface density expression on monocytes and B-cells is associated with

GBS pathogenesis. However, activation-dependent expression of $\alpha_M\beta_2$ -integrin on T-cells has also been described *in vitro* (including induced expression on CD4+ T-cells). The increased expression following re-stimulation of recently activated T-cells suggests an important role of this integrin in activated T-cell effector function (Wagner et al., 2001).

We demonstrated that α_M -integrin was the counterligand responsible for ICAM-1-dependent GBS PBML adhesion/ migration at the cytokine activated human BNB under capillary hydrodynamic forces *in vitro*, despite the ubiquitously high PBML α_L -integrin expression. Based on these observations, we propose that CD11b+ monocytes, T-cells and B-cells selectively adhere to and migrate across the BNB during the early stages of AIDP, with preferential trafficking of cells with the highest integrin surface density. This would reduce the relative % of CD11b+ cells or MFI or both in the circulating venous blood pool compared to healthy control patients.

We provide evidence that adherent/ migrated GBS PBMLs at the BNB express aCD11b, and CD14+ monocytes are the predominant leukocyte subpopulation *in vitro*, consistent with *in vivo* studies of infiltrated mononuclear cells in GBS and EAN. We demonstrate that PBMLs derived from circulating venous blood in GBS patients rarely express aCD11b, implying an importance of chemoattractant signals at the BNB as a pre-requisite for pathogenic trafficking. CCL2 and CXCL10 are chemokines that may facilitate GBS monocyte and T-cell trafficking at the human BNB respectively, based on *in vivo* observational studies. Not only do α_M -integrin-ICAM-1 interactions at the BNB provide insight to the pathogenic infiltration of monocytes/ macrophages and activated T-lymphocytes seen in GBS and EAN, $\alpha_M\beta_2$ -integrin provides a link between monocyte trafficking at the BNB and subsequent complement-mediated injury observed in GBS. Aberrant activation of the innate and adaptive immune systems could both result in α_M -integrin-ICAM-1 mediated leukocyte infiltration into the peripheral nerve endoneurium with consequential demyelination and axonal injury in GBS.

This study shows that α_L -integrin plays a relatively minor role in total untreated GBS PBML adhesion/migration at the *in vitro* human BNB. Lewis rat EAN studies demonstrated a role for $\alpha_L\beta_2$ -integrin in disease induction following active immunization with bovine spinal nerve root myelin, without effect in adoptive transfer EAN using P2-specific T-cells. That data suggested that $\alpha_L\beta_2$ -integrin acted primarily during EAN induction without significant impact on the effector phase (Archelos et al., 1994). Similarly, $\alpha_L\beta_2$ -integrin knockout and wild-type T-cells showed similar infiltration kinetics into the central nervous system and lymph nodes in murine experimental autoimmune encephalitis (EAE) (Dugger et al., 2009). Although $\alpha_L\beta_2$ -integrin is important for naïve T-cell adhesion to antigen presenting cells necessary for activation and effector cell proliferation, regulatory T-cell homeostasis and function have been shown to critically depend on this integrin as well (Wohler et al., 2009). Reduced percentages of circulating regulatory T-cells have been described in GBS (Harness and McCombe, 2008; Pritchard et al., 2007). Thus, blockade of $\alpha_L\beta_2$ -integrin may be ineffective, and potentially harmful in GBS (AIDP) patients.

Human IVIg inhibition of GBS patient PBML adhesion/migration at the *in vitro* BNB was less effective at a ~10,000-fold higher concentration relative to monoclonal α_M -integrin and ICAM-1 antibodies. IVIg had been shown to alter leukocyte rolling and adhesion on post-capillary venules in EAE-affected mice and multiple sclerosis patient derived leukocytes trafficking over non-cerebral endothelium *in vitro* (Lapointe et al., 2004). IVIg arrests disease progression in AIDP, and results in better outcomes relative to untreated children, with similar outcomes compared to plasma exchange in adults (Hughes et al., 2007). Although IVIg probably has several immune modulatory effects (Dalakas, 2004; Hartung, 2008), antibodies against specific cell adhesion molecules and their signaling counterligands

may be more effective in modulating pathogenic leukocyte trafficking at the BNB in AIDP, as demonstrated with α_M -integrin and ICAM-1 in this study.

There is evidence that circulating T-cells derived from GBS patients are activated, relative to healthy controls (Hadden et al., 2001; Harness and McCombe, 2008). In support of this, we determined that PBMLs derived from untreated GBS patients preferentially interacted with the basal and cytokine activated human BNB *in vitro*. However, cytokine activation with TNF- α and IFN- γ provided a stronger stimulus for leukocyte adhesion/ migration than endogenous PBML activation state, implying a crucial role of proinflammatory cytokine BNB activation during pathogenic leukocyte trafficking.

Despite the limitations of *in vitro* leukocyte-endothelial cell models, the flow-dependent human BNB assay provides an avenue to decipher the mediators of leukocyte trafficking and their signaling pathways potentially relevant to the pathogenesis of peripheral nerve inflammatory disorders such as GBS. Our data suggest an important role for α_M -integrin-ICAM-1 dependent signaling on GBS patient PBML trafficking at the BNB. Monocytes and lymphocytes were not separated in order to more accurately mimic inflammatory events at the BNB *in vivo*. A severe EAN model that closely recapitulates the clinical, electrophysiological and pathologic features of AIDP (Xia et al., 2010a) provides a murine model for *in vivo* testing of small molecular inhibitors or monoclonal antibodies against α_M -integrin or ICAM-1 during its effector phase, similar to untreated GBS patients in clinical practice (Xia et al., 2010a). Blockade of leukocyte trafficking at the BNB could lead to targeted, more efficacious therapies for peripheral neuroinflammation. Further studies are needed to determine the mediators of leukocyte subpopulation trafficking at the BNB in health and disease.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Grant information: Baylor College of Medicine New Investigator Start-up Program (2007–2011), Guillain-Barré syndrome/ Chronic Inflammatory Demyelinating Polyradiculoneuropathy Foundation International Research Grants (2007–2009 and 2010–2012), NIH Grants NS073702, NS075212 and NS078226 (all to EEU).

We will like to thank the GBS patients and healthy donors for donating blood for the leukocyte isolation and the Neurology Service, Ben Taub General Hospital for patient referral. Special thanks to Ms. April Durett, Center of Cell & Gene Therapy, Department of Pediatrics, Baylor College of Medicine for her expert technical assistance with flow cytometry, and Drs. C. Wayne Smith and Fong Lam, Children's Nutritional Research Center, Department of Pediatrics, Baylor College of Medicine, for initial assistance in setting up the leukocyte trafficking assay.

References

- Aird W. Phenotypic heterogeneity of the endothelium: I. Structure, function, and mechanisms. *Circ Res.* 2007a; 100(2):158–173. [PubMed: 17272818]
- Aird W. Phenotypic heterogeneity of the endothelium: II. Representative vascular beds. *Circ Res.* 2007b; 100(2):174–190. [PubMed: 17272819]
- Allt G, Lawrenson J. The blood-nerve barrier: enzymes, transporters and receptors--a comparison with the blood-brain barrier. *Brain Res Bull.* 2000; 52(1):1–12. [PubMed: 10779695]
- Alon R, Ley K. Cells on the run: shear-regulated integrin activation in leukocyte rolling and arrest on endothelial cells. *Curr Opin Cell Biol.* 2008; 20(5):525–532. [PubMed: 18499427]
- Altman SA, Randers L, Rao G. Comparison of trypan blue dye exclusion and fluorometric assays for mammalian cell viability determinations. *Biotechnol Prog.* 1993; 9(6):671–674. [PubMed: 7764357]

- Archelos JJ, Mäurer M, Jung S, Miyasaka M, Tamatani T, Toyka KV, Hartung HP. Inhibition of experimental autoimmune neuritis by an antibody to the lymphocyte function-associated antigen-1. *Lab Invest.* 1994; 70(5):667–675. [PubMed: 8196363]
- Archelos JJ, Mäurer M, Jung S, Toyka KV, Hartung HP. Suppression of experimental allergic neuritis by an antibody to the intracellular adhesion molecule ICAM-1. *Brain.* 1993; 116 (Pt 5):1043–1058. [PubMed: 7693297]
- Bell M, Weddell A. A descriptive study of the blood vessels of the sciatic nerve in the rat, man and other mammals. *Brain.* 1984; 107 (Pt 3):871–898. [PubMed: 6478181]
- Berciano J, Figols J, García A, Calle E, Illa I, Lafarga M, Berciano MT. Fulminant Guillain-Barré syndrome with universal inexcitability of peripheral nerves: a clinicopathological study. *Muscle Nerve.* 1997; 20(7):846–857. [PubMed: 9179157]
- Carlson T, Kroenke M, Rao P, Lane TE, Segal B. The Th17-ELR+ CXC chemokine pathway is essential for the development of central nervous system autoimmune disease. *J Exp Med.* 2008; 205(4):811–823. [PubMed: 18347102]
- Chi LJ, Xu WH, Zhang ZW, Huang HT, Zhang LM, Zhou J. Distribution of Th17 cells and Th1 cells in peripheral blood and cerebrospinal fluid in chronic inflammatory demyelinating polyradiculoneuropathy. *J Peripher Nerv Syst.* 2010; 15(4):345–356. [PubMed: 21199106]
- Cinamon G, Shinder V, Alon R. Shear forces promote lymphocyte migration across vascular endothelium bearing apical chemokines. *Nat Immunol.* 2001; 2(6):515–522. [PubMed: 11376338]
- Dalakas M. B cells as therapeutic targets in autoimmune neurological disorders. *Nat Clin Pract Neurol.* 2008; 4(10):557–567. [PubMed: 18813230]
- Dalakas MC. Intravenous immunoglobulin in autoimmune neuromuscular diseases. *JAMA.* 2004; 291(19):2367–2375. [PubMed: 15150209]
- Deshmane S, Kremlev S, Amini S, Sawaya B. Monocyte chemoattractant protein-1 (MCP-1): an overview. *J Interferon Cytokine Res.* 2009; 29(6):313–326. [PubMed: 19441883]
- Dugger KJ, Zinn KR, Weaver C, Bullard DC, Barnum SR. Effector and suppressor roles for LFA-1 during the development of experimental autoimmune encephalomyelitis. *J Neuroimmunol.* 2009; 206(1–2):22–27. [PubMed: 19010554]
- Enders U, Lobb R, Pepinsky R, Hartung H, Toyka K, Gold R. The role of the very late antigen-4 and its counterligand vascular cell adhesion molecule-1 in the pathogenesis of experimental autoimmune neuritis of the Lewis rat. *Brain.* 1998; 121 (Pt 7):1257–1266. [PubMed: 9679778]
- Gold R, Archelos J, Hartung H. Mechanisms of immune regulation in the peripheral nervous system. *Brain Pathol.* 1999; 9(2):343–360. [PubMed: 10219750]
- Gopalan P, Jones D, McIntire L, Smith C. Cell adhesion under hydrodynamic flow conditions. *Curr Protoc Immunol.* 2001; Chapter 7(Unit 7):29. [PubMed: 18432840]
- Hadden RD, Karch H, Hartung HP, Zielasek J, Weissbrich B, Schubert J, Weishaupt A, Cornblath DR, Swan AV, Hughes RA, Toyka KV. Group PESG-BST. Preceding infections, immune factors, and outcome in Guillain-Barré syndrome. *Neurology.* 2001; 56(6):758–765. [PubMed: 11274311]
- Hafer-Macko CE, Sheikh KA, Li CY, Ho TW, Cornblath DR, McKhann GM, Asbury AK, Griffin JW. Immune attack on the Schwann cell surface in acute inflammatory demyelinating polyneuropathy. *Ann Neurol.* 1996; 39(5):625–635. [PubMed: 8619548]
- Harness J, McCombe PA. Increased levels of activated T-cells and reduced levels of CD4/CD25+ cells in peripheral blood of Guillain-Barré syndrome patients compared to controls. *J Clin Neurosci.* 2008; 15(9):1031–1035. [PubMed: 18653346]
- Hartung H, Willison H, Kieseier B. Acute immunoinflammatory neuropathy: update on Guillain-Barré syndrome. *Curr Opin Neurol.* 2002; 15(5):571–577. [PubMed: 12352001]
- Hartung HP. Advances in the understanding of the mechanism of action of IVIg. *J Neurol.* 2008; 255(Suppl 3):3–6. [PubMed: 18685919]
- Hughes R, Cornblath D. Guillain-Barré syndrome. *Lancet.* 2005; 366(9497):1653–1666. [PubMed: 16271648]
- Hughes RA, Swan AV, Raphaël JC, Annane D, van Koningsveld R, van Doorn PA. Immunotherapy for Guillain-Barré syndrome: a systematic review. *Brain.* 2007; 130(Pt 9):2245–2257. [PubMed: 17337484]

- Kabir K, Keller H, Grass G, Minor T, Stueber F, Schroeder S, Putensen C, Paul C, Burger C, Rangger C, Neville L, Mathiak G. Cytokines and chemokines in serum and urine as early predictors to identify septic patients on intensive care unit. *Int J Mol Med*. 2003; 12(4):565–570. [PubMed: 12964035]
- Kiefer R, Kieseier BC, Stoll G, Hartung HP. The role of macrophages in immune-mediated damage to the peripheral nervous system. *Prog Neurobiol*. 2001; 64(2):109–127. [PubMed: 11240209]
- Kieseier B, Kiefer R, Gold R, Hemmer B, Willison H, Hartung H. Advances in understanding and treatment of immune-mediated disorders of the peripheral nervous system. *Muscle Nerve*. 2004; 30(2):131–156. [PubMed: 15266629]
- Kieseier B, Krivacic K, Jung S, Pischel H, Toyka K, Ransohoff R, Hartung H. Sequential expression of chemokines in experimental autoimmune neuritis. *J Neuroimmunol*. 2000; 110(1–2):121–129. [PubMed: 11024541]
- Kieseier B, Tani M, Mahad D, Oka N, Ho T, Woodroffe N, Griffin J, Toyka K, Ransohoff R, Hartung H. Chemokines and chemokine receptors in inflammatory demyelinating neuropathies: a central role for IP-10. *Brain*. 2002; 125(Pt 4):823–834. [PubMed: 11912115]
- Kim C. Migration and function of Th17 cells. *Inflamm Allergy Drug Targets*. 2009; 8(3):221–228. [PubMed: 19601882]
- Komarek V. Gross Anatomy. In: Hedrich, HJ.; Bullock, G., editors. *The Laboratory Mouse*. Amsterdam: Elsevier Academic Press; 2004. p. 117–132.
- Lacotte S, Brun S, Muller S, Dumortier H. CXCR3, inflammation, and autoimmune diseases. *Ann N Y Acad Sci*. 2009; 1173:310–317. [PubMed: 19758167]
- Lapointe BM, Herx LM, Gill V, Metz LM, Kubes P. IVIg therapy in brain inflammation: etiology-dependent differential effects on leucocyte recruitment. *Brain*. 2004; 127(Pt 12):2649–2656. [PubMed: 15355874]
- Le Cabec V, Carréno S, Moisan A, Bordier C, Maridonneau-Parini I. Complement receptor 3 (CD11b/CD18) mediates type I and type II phagocytosis during nonopsonic and opsonic phagocytosis, respectively. *J Immunol*. 2002; 169(4):2003–2009. [PubMed: 12165526]
- Lu MO, Zhu J. The role of cytokines in Guillain-Barré syndrome. *J Neurol*. 2011; 258(4):533–548. [PubMed: 21104265]
- Mahad D, Callahan M, Williams K, Ubogu E, Kivisäkk P, Tucky B, Kidd G, Kingsbury G, Chang A, Fox R, Mack M, Sniderman M, Ravid R, Staugaitis S, Stins M, Ransohoff R. Modulating CCR2 and CCL2 at the blood-brain barrier: relevance for multiple sclerosis pathogenesis. *Brain*. 2006; 129(Pt 1):212–223. [PubMed: 16230319]
- Malmgren L, Olsson Y. Differences between the peripheral and the central nervous system in permeability to sodium fluorescein. *J Comp Neurol*. 1980; 191(1):103–107. [PubMed: 7400388]
- Man S, Tucky B, Bagheri N, Li X, Kocher R, Ransohoff R. $\alpha 4$ Integrin/FN- CS1 mediated leukocyte adhesion to brain microvascular endothelial cells under flow conditions. *J Neuroimmunol*. 2009; 210(1–2):92–99. [PubMed: 19345424]
- Man S, Ubogu E, Ransohoff R. Inflammatory cell migration into the central nervous system: a few new twists on an old tale. *Brain Pathol*. 2007; 17(2):243–250. [PubMed: 17388955]
- Meyer zu Hörste G, Hartung H, Kieseier B. From bench to bedside--experimental rationale for immune-specific therapies in the inflamed peripheral nerve. *Nat Clin Pract Neurol*. 2007; 3(4):198–211. [PubMed: 17410107]
- Muller WA. Mechanisms of leukocyte transendothelial migration. *Annu Rev Pathol*. 2011; 6:323–344. [PubMed: 21073340]
- Nishimura N, Schaffer C, Friedman B, Tsai P, Lyden P, Kleinfeld D. Targeted insult to subsurface cortical blood vessels using ultrashort laser pulses: three models of stroke. *Nat Methods*. 2006; 3(2):99–108. [PubMed: 16432519]
- Nourshargh S, Krombach F, Dejana E. The role of JAM-A and PECAM-1 in modulating leukocyte infiltration in inflamed and ischemic tissues. *J Leukoc Biol*. 2006; 80(4):714–718. [PubMed: 16857733]
- Nyati KK, Prasad KN, Verma A, Paliwal VK. Correlation of matrix metalloproteinases-2 and -9 with proinflammatory cytokines in Guillain-Barré syndrome. *J Neurosci Res*. 2010; 88(16):3540–3546. [PubMed: 20936699]

- Nyland H, Matre R, Mørk S. Immunological characterization of sural nerve biopsies from patients with Guillain-Barré syndrome. *Ann Neurol*. 1981; 9(Suppl):80–86. [PubMed: 7224617]
- Olsson Y. Topographical differences in the vascular permeability of the peripheral nervous system. *Acta Neuropathol*. 1968; 10(1):26–33. [PubMed: 5640115]
- Olsson Y. Microenvironment of the peripheral nervous system under normal and pathological conditions. *Crit Rev Neurobiol*. 1990; 5(3):265–311. [PubMed: 2168810]
- Orlikowski D, Chazaud B, Plonquet A, Poron F, Sharshar T, Maison P, Raphaël J, Gherardi R, Créange A. Monocyte chemoattractant protein 1 and chemokine receptor CCR2 productions in Guillain-Barré syndrome and experimental autoimmune neuritis. *J Neuroimmunol*. 2003; 134(1–2):118–127. [PubMed: 12507779]
- Poduslo J, Curran G, Berg C. Macromolecular permeability across the blood-nerve and blood- brain barriers. *Proc Natl Acad Sci U S A*. 1994; 91(12):5705–5709. [PubMed: 8202551]
- Press R, Nennesmo I, Kouwenhoven M, Huang YM, Link H, Pashenkov M. Dendritic cells in the cerebrospinal fluid and peripheral nerves in Guillain-Barré syndrome and chronic inflammatory demyelinating polyradiculoneuropathy. *J Neuroimmunol*. 2005; 159(1–2):165–176. [PubMed: 15652416]
- Press R, Pashenkov M, Jin J, Link H. Aberrated levels of cerebrospinal fluid chemokines in Guillain-Barré syndrome and chronic inflammatory demyelinating polyradiculoneuropathy. *J Clin Immunol*. 2003; 23(4):259–267. [PubMed: 12959218]
- Pritchard J, Makowska A, Gregson NA, Hayday AC, Hughes RA. Reduced circulating CD4+CD25+ cell populations in Guillain-Barré syndrome. *J Neuroimmunol*. 2007; 183(1–2):232–238. [PubMed: 17239444]
- Putzu GA, Figarella-Branger D, Bouvier-Labit C, Liprandi A, Bianco N, Pellissier JF. Immunohistochemical localization of cytokines, C5b-9 and ICAM-1 in peripheral nerve of Guillain-Barré syndrome. *J Neurol Sci*. 2000; 174(1):16–21. [PubMed: 10704976]
- Radhakrishnan VV, Sumi MG, Reuben S, Mathai A, Nair MD. Serum tumour necrosis factor-alpha and soluble tumour necrosis factor receptors levels in patients with Guillain-Barre syndrome. *Acta Neurol Scand*. 2004; 109(1):71–74. [PubMed: 14653854]
- Rauschka H, Jellinger K, Lassmann H, Braier F, Schmidbauer M. Guillain-Barré syndrome with marked pleocytosis or a significant proportion of polymorphonuclear granulocytes in the cerebrospinal fluid: neuropathological investigation of five cases and review of differential diagnoses. *Eur J Neurol*. 2003; 10(5):479–486. [PubMed: 12940826]
- Reina M, López A, Villanueva M, de Andrés J, León G. Morphology of peripheral nerves, their sheaths, and their vascularization. *Rev Esp Anesthesiol Reanim*. 2000; 47(10):464–475. [PubMed: 11171467]
- Reina M, López A, Villanueva M, De Andrés J, Machés F. The blood-nerve barrier in peripheral nerves. *Rev Esp Anesthesiol Reanim*. 2003; 50(2):80–86. [PubMed: 12712870]
- Schreiber T, Shinder V, Cain D, Alon R, Sackstein R. Shear flow-dependent integration of apical and subendothelial chemokines in T-cell transmigration: implications for locomotion and the multistep paradigm. *Blood*. 2007; 109(4):1381–1386. [PubMed: 17038526]
- Simon SI, Green CE. Molecular mechanics and dynamics of leukocyte recruitment during inflammation. *Annu Rev Biomed Eng*. 2005; 7:151–185. [PubMed: 16004569]
- Tran GT, Hodgkinson SJ, Carter NM, Killingsworth M, Nomura M, Verma ND, Plain KM, Boyd R, Hall BM. Membrane attack complex of complement is not essential for immune mediated demyelination in experimental autoimmune neuritis. *J Neuroimmunol*. 2010; 229(1–2):98–106. [PubMed: 20850187]
- Ubogu E, Callahan M, Tucky B, Ransohoff R. Determinants of CCL5-driven mononuclear cell migration across the blood-brain barrier. Implications for therapeutically modulating neuroinflammation. *J Neuroimmunol*. 2006a; 179(1–2):132–144. [PubMed: 16857269]
- Ubogu E, Cossoy M, Ransohoff R. The expression and function of chemokines involved in CNS inflammation. *Trends Pharmacol Sci*. 2006b; 27(1):48–55. [PubMed: 16310865]
- Wagner C, Hänsch GM, Stegmaier S, Deneffle B, Hug F, Schoels M. The complement receptor 3, CR3 (CD11b/CD18), on T lymphocytes: activation-dependent up-regulation and regulatory function. *Eur J Immunol*. 2001; 31(4):1173–1180. [PubMed: 11298342]

- Wohler J, Bullard D, Schoeb T, Barnum S. LFA-1 is critical for regulatory T cell homeostasis and function. *Mol Immunol*. 2009; 46(11–12):2424–2428. [PubMed: 19428111]
- Xia R, Yosef N, Ubogu E. Clinical, electrophysiological and pathologic correlations in a severe murine experimental autoimmune neuritis model of Guillain-Barré syndrome. *J Neuroimmunol*. 2010a; 219(1–2):54–63. [PubMed: 20034679]
- Xia RH, Yosef N, Ubogu EE. Selective expression and cellular localization of pro-inflammatory chemokine ligand/receptor pairs in the sciatic nerves of a severe murine experimental autoimmune neuritis model of Guillain-Barré syndrome. *Neuropathol Appl Neurobiol*. 2010b; 36(5):388–398. [PubMed: 20500551]
- Yano K, Gale D, Massberg S, Cheruvu P, Monahan-Earley R, Morgan E, Haig D, von Andrian U, Dvorak A, Aird W. Phenotypic heterogeneity is an evolutionarily conserved feature of the endothelium. *Blood*. 2007; 109(2):613–615. [PubMed: 16990601]
- Yosef N, Xia R, Ubogu E. Development and characterization of a novel human in vitro blood-nerve barrier model using primary endoneurial endothelial cells. *J Neuropathol Exp Neurol*. 2010; 69(1): 82–97. [PubMed: 20010300]

Figure 1.

Effect of cytokine treatment on the human BNB endothelial cell viability (**A**) Representative flow cytometry dot plots show confluent pHEndEC % viability (7-AAD negative, number in left upper gate) under basal culture conditions. (**B, C**) There is no significant reduction in pHEndEC viability following treatment with 1–10 U/mL TNF- α and 2–20 U/mL IFN- γ for 24 hours, relative to basal conditions. (**D, E, F**) Progressive reduction in endothelial cell viability is observed at higher doses. Numbers indicate the percentage of total events in each gate. The upper right gate represents the % of non-viable cells.

Figure 2.

Effect of cytokine treatment on the human BNB (A) Representative ethidium bromide stained agarose gel with DNA size markers shows increased ICAM-1 c-DNA expression following treatment with 10 U/mL TNF- α and 20 U/mL IFN- γ , with reduction of expression to near basal levels following treatment with 100 U/mL TNF- α and 200 U/mL IFN- γ for 24 hours (1=basal, 2=10 U/mL TNF- α + 20 U/mL IFN- γ , 3=100 U/mL TNF- α + 200 U/mL IFN- γ) (B) Representative flow cytometry histogram demonstrates increased number of ICAM-positive live pHEndECs following treatment with 10 U/mL TNF- α and 20 U/mL IFN- γ for 24 hours, associated with increase in mean fluorescent intensity (MFI: Comp-PE-A: IgG2b) relative to basal conditions. (C) BNB TEER measurements show a small increase in endothelial resistance following low dose cytokine treatment for 24 hours, with reduction to basal values with a high dose TNF- α and IFN- γ . * indicates $p < 0.05$, N.S. indicates not significant relative to basal, untreated BNB.

Figure 3.

Chemokine expression at the BNB. **(A)** Chemokine antibody array autoradiograph shows the distribution of chemokines expressed by the BNB under basal conditions. The upper four and lower two spots represent internal array controls. **(B)** Chemokine antibody array autoradiograph shows the distribution of chemokines expressed by the BNB following cytokine treatment with 10 U/mL TNF- α and 20 U/mL IFN- γ . **(C)** Mean relative chemokine expression (referenced to mean integrated density values [IDV] of internal array controls) comparing the basal and cytokine activated BNB is shown, with *de novo* or increased expression of specific Th1 and Th17-related chemokines and reduced expression of a Th2-related chemokine (see results).

Figure 4.

Cell adhesion molecule protein expression at the BNB. Graphical representation of the time-dependent protein expression of (A) E-selectin, (B) P-selectin, (C) ICAM-1, (D) VCAM-1, (E) FN CS-1, (F) JAM-A, following cytokine treatment, relative to integrated density values (IDV) of GAPDH is shown. Data represents at least three independent experiments performed in duplicate. * indicates $p < 0.05$ relative to basal conditions, N.S. indicates not significant. Error bars depict standard errors of the means. (G) Representative autoradiographs show the time-dependent effect of cytokine treatment with 10 U/mL TNF- α and 20 U/mL IFN- γ at the human *in vitro* BNB, in the same time sequence as shown in the graphs.

Figure 5.

Surface expression of cell adhesion molecules at the BNB. Merged indirect fluorescent immunocytochemistry photomicrographs demonstrate increased expression (green [FITC or Alexa Fluor 488] immunoreactivity) of P-selectin, E-selectin, ICAM-1, VCAM-1 and FN CS-1 on the surface of fixed pHEndECs grown on GTA-crosslinked RTC-coated glass coverslips following cytokine activation (CA) with 10 U/mL TNF- α and 20 U/mL IFN- γ for 24 hours relative to basal (B) culture conditions. Nuclei are depicted in blue (DAPI). ICAM-1 was diffusely expressed on pHEndEC cytoplasm, while P-selectin, E-selectin, VCAM-1 and FN-CS-1 demonstrated a predominantly perinuclear expression pattern. There was no change in the cytoplasmic expressed JAM-A following cytokine activation. Scale bars represent 20 μ m for P-selectin, E-selectin, ICAM-1, VCAM-1 and FN CS-1 (B), 25 μ m for FN CS-1 (CA) and 50 μ m for JAM-A.

Figure 6.

Differential PBML CD11b expression in GBS. **(A)** Bar histograms demonstrate a relative increase in % CD3+ CD4+ T-cells and reduction in %CD3+ CD8+ T-cells in untreated GBS relative to controls in this cohort, without change in % of total CD3+ T-cells derived from isolated PBMLs. Increases in % CD14+ monocytes and % CD19+ B-cells are observed in GBS relative to controls. **(B)** Significant reduction in CD11b+ CD3+ CD4+ and CD3+ CD8+ T-cells is seen in untreated GBS compared with controls with no difference observed with CD14+ monocytes and CD19+ B-cells. * indicates $p < 0.05$ relative to healthy controls, ^ indicates not significant.

Figure 7. PBML CD11b surface density changes in GBS. **(A)** Bar histograms demonstrate reduction in CD14+ CD11b MFI (as a measure of surface receptor density) in untreated GBS-derived PBMLs relative to controls. **(B)** Significant reductions in CD11b MFI are also demonstrated on CD3+ CD4+ and CD3+ CD8+ T-cells and CD19+ B-cells. * indicates $p < 0.05$ relative to healthy controls.

Figure 8.

CD11b expression on CD14⁺ monocytes in GBS. Representative flow cytometry dot plots demonstrate gating properties for PBML in an untreated GBS patient (**A**) and age- and sex-matched healthy control (**F**) based on forward and side-scatter characteristics, with subsequent gating on CD14⁺ (**B, G**), CD14⁺ CD11b⁺ (**C, H**) and CD14⁺ aCD11b⁺ (**D, I**) monocytes. There is an increase in %CD14⁺ monocytes without alterations in CD11b expression in this cohort of GBS patients compared with controls. Activated CD11b-expressing monocytes are rarely observed in PBMLs derived from venous blood in GBS patients and controls. Gates are based on isotype controls in each sample. Numbers depict % parent population.

Figure 9.

CD11b expression on CD3+ CD4+ T-cells in GBS. Representative flow cytometry dot plots demonstrate gating properties for T-cells in an untreated GBS patient (**A**) and age- and sex-matched healthy control (**F**) based on CD3 expression, with subsequent gating on CD4+ (**B**, **G**), CD4+ CD11b+ (**C**, **H**) and CD4+ aCD11b+ (**D**, **I**) T-cells. There is an increase in %CD3+ CD4+ T-cells and reduction in CD11b+ cells without significant alterations in total % CD3 expression in this cohort of GBS patients compared with controls. Activated CD11b-expressing CD4+ T-cells are rarely observed in PBMLs derived from venous blood in GBS patients and controls. Gates are based on isotype controls in each sample. Numbers depict % parent population.

Figure 10.

CD11b expression on CD3+ CD8+ T-cells in GBS. Representative flow cytometry dot plots demonstrate gating properties for T-cells in an untreated GBS patient (**A**) and age- and sex-matched healthy control (**F**) based on CD3 expression, with subsequent gating on CD8+ (**B**, **G**), CD8+ CD11b+ (**C**, **H**) and CD8+ aCD11b+ (**D**, **I**) T-cells. There is a reduction in %CD3+ CD8+ T-cells and reduction in CD11b+ cells without significant alterations in total % CD3 expression in this cohort of GBS patients compared with controls. Activated CD11b-expressing CD8+ T-cells are rarely observed in PBMLs derived from venous blood in GBS patients and controls. Gates are based on isotype controls in each sample. Numbers depict % parent population.

Figure 11.

CD11b expression on CD19⁺ T-cells in GBS. Representative flow cytometry dot plots demonstrate gating properties for PBML in an untreated GBS patient (**A**) and age- and sex-matched healthy control (**F**) based on forward and side-scatter characteristics, with subsequent gating on CD19⁺ (**B, G**), CD19⁺ CD11b⁺ (**C, H**) and CD19⁺ aCD11b⁺ (**D, I**) monocytes. There is a significant increase in %CD19⁺ B-cells with a small, but statistically non-significant increase in mean CD11b expression in this cohort of GBS patients compared with controls. Activated CD11b-expressing B-cells are rarely observed in PBMLs derived from venous blood in GBS patients and controls. Gates are based on isotype controls in each sample. Numbers depict % parent population.

Figure 12.

CD11a expression on PBMLs in GBS. Representative flow cytometry dot plots demonstrate gating properties for PBML in an untreated GBS patient (A) based on forward and side-scatter characteristics, with subsequent gating on CD14+ monocytes (B), CD19+ B-cells (C) and CD3+ T-cells (D). Subsequent gating for each subpopulation demonstrates >99% CD11a+. Gates are based on isotype controls in each sample. Numbers depict % parent population.

Figure 13.

Flow-dependent GBS PBML trafficking at the BNB. (A) Representative digital phase contrast photomicrograph taken >20 minutes after the initiation of hydrodynamic flow show few attached PBMLs on the untreated human BNB model surface. (B) Representative digital phase contrast photomicrograph taken >20 minutes after the initiation of hydrodynamic flow show clusters of attached PBMLs on the cytokine activated human BNB model surface. PBMLs were from the same GBS patient shown in (A). White arrows indicate attached PBML clusters. Scale bar=100 μm . (C) A higher magnification photomicrograph demonstrates GBS PBML attached in clusters on the BNB surface under the influence of flow. Scale bar= 25 μm (D) Graphical representation of the mean numbers of adherent/ migrated GBS PBML per high power field (hpf; 870 $\times$ 650 μm) in 4 healthy controls and 6 GBS patients following BNB cytokine activation (CA), compared to the untreated, basal BNB. This graph demonstrates that GBS PBMLs significantly interact with the *in vitro* BNB at higher levels than healthy control PBMLs. The more significant inductive effect of 10 U/mL TNF- α and 20 U/mL IFN- γ on untreated GBS and healthy leukocyte trafficking at the human BNB *in vitro* is demonstrated. # indicates $p < 0.05$ comparing GBS PBML adhesion/ migration with controls. * indicates $p < 0.05$ comparing BNB cytokine activated PBML adhesion/ migration with basal trafficking.

Figure 14. Function-neutralizing inhibition studies at the BNB. Graphical representation of antibody blockade effect on untreated GBS PBML leukocyte adhesion/migration as a percentage of uninhibited numbers following BNB cytokine treatment is shown (see results). Monoclonal antibodies against ICAM-1 and CD11b (blocks α_M -integrin) were the most effective, implying an important role in GBS PBML trafficking at the BNB *in vitro*. * indicates $p<0.05$ relative to cytokine activated (CA), ** indicates $p<0.05$ comparing either ICAM-1 or CD11b blockade with combined ICAM-1/ CD11b inhibition, *** indicates $p<0.05$ comparing CD11a antibodies (blocks α_L -integrin) to either ICAM-1 or combined ICAM-1/ CD11a blockade. N.S. indicates not significant.

Figure 15.

Characterization of adherent/migrated GBS PBMLs at the BNB. (A) A representative digital photomicrograph demonstrates clusters of adherent PBML (white arrows) on the surface of the cytokine activated BNB stained with DAPI (blue) to detect nuclei. pHEndECs have larger nuclei than PBMLs. (B, C, D) A representative merged digital photomicrograph demonstrates clusters of PBMLs (DAPI, blue) on the pHEndEC surface (green ZO-1 counterstain, B). Majority of adherent cells express aCD11b (orange, C), verified by the merged photomicrograph (D). (E, F, G, H) CD14⁺ (green) aCD11b⁺ (orange) monocytes (DAPI, blue) are the predominant leukocyte subpopulation that adhere to the cytokine activated in vitro BNB, as demonstrated by individual and merged digital photomicrographs. Scale bar = 50 μ m for A, 20 μ m for B–D and 10 μ m for E–H.

Clinical characteristics of GBS (AIDP) patients

Table 1

Clinical characteristics of GBS (AIDP) patients. Basic demographics, clinical features and electrophysiological findings of the patient cohort are shown. All patients met current diagnostic criteria for AIDP and provided blood samples prior to treatment.

Age (years)	Sex	Duration of symptoms (days)	Clinical Features	Electrophysiological findings
48	Female	7	Mild quadriparesis, distal sensory loss, ataxic gait, areflexia	Mild-to-moderately severe demyelinating polyradiculoneuropathy with secondary axonal loss.
62	Male	5	Moderate quadriparesis, distal sensory loss, areflexia	Mild-to-moderately severe polyradiculoneuropathy. Distal conduction block.
26	Male	8	Moderate paraparesis, mild right> left upper extremity weakness, diffuse sensory loss in lower extremities and right upper extremity, areflexia	Moderate severe polyradiculopathy with ongoing denervation. Prolonged/absent F-waves. Proximal conduction block.
57	Male	7	Severe quadriparesis with bulbar dysfunction. Normal sensation. Dysautonomia. Hyporeflexia in upper extremities, absent reflexes in lower extremities. Fasciculations.	Moderately severe motor polyradiculopathy. Distal conduction block. Diffuse myokymia.
41	Male	4	Severe quadriparesis with diffuse sensory loss. Areflexia. Early respiratory failure	Markedly severe polyradiculopathy with very low motor amplitudes. Absent F-wave responses.
26	Male	8	Moderate distal weakness in hands and feet. Distal sensory loss. Hyporeflexia in upper extremities and knees. Ankle areflexia.	Mild-to-moderately severe polyradiculoneuropathy with secondary axonal loss. Distal and proximal conduction block. Absent F-waves