Published in final edited form as:

Org Lett. 2011 March 4; 13(5): 836–839. doi:10.1021/ol1027893.

2-Azapinanes: Aza Analogues of the Enantiomeric Pinyll Carbocation Intermediates in Pinene Biosynthesis

Juan A. Faraldos^{*,†,||}, Benson M. Kariuki[‡], and Robert M. Coates^{*,†}

Department of Chemistry, University of Illinois, Urbana-Champaign, IL 61801, USA, School of Chemistry, Cardiff University, Cardiff CF10 3AT, Wales, UK

Abstract

The enantiomeric 2-azapinanes, aza analogues of the pinyll carbocation intermediates in pinene biosynthesis, were synthesized from (–)- and (+)-*cis*-pinonic acids. The individual reactions in the 5-step sequence were Beckmann rearrangement of the pinonic acid oximes, cyclization to the N-acetyl lactams, hydrolysis to the NH-lactams, N-methylations, and LiAlH₄ reductions. The anti stereochemistry of the N-methyl groups in the salts with respect to the gem-dimethyl bridge was established by NOE measurements and by an X-ray diffraction analysis.

α - And β -pinene are widespread monoterpene olefins based on the bicyclo[3.1.1]heptane nucleus¹ and major constituents of conifer oleo resin and its turpentine distillate.² Like many monoterpenes, the pinenes occur naturally in both enantiomeric forms, often even in the same plant.^{1b,3} The availability of both enantiomers, together with the ring strain and steric bias imposed by the gem dimethyl bridge, have been exploited in the synthesis of enantio-pure organic compounds⁴ and in the development of pinane-based chiral auxiliaries for asymmetric synthesis.⁵

Considerable evidence supports a catalytic mechanism for pinene biosynthesis in which the substrate geranyl diphosphate (**1**) first undergoes a stereospecific allylic rearrangement to the key enzyme-bound intermediates (*R*)- or (*S*)-linalyl diphosphate (**2**), thereby allowing the required 2,3-bond rotation essential for further progress towards the enzymatic bicyclic

faraldosja@Cardiff.ac.uk; rmcoates@uiuc.edu.

[†]University of Illinois, Urbana-Champaign, IL, USA

^{||}Current address: Cardiff University, Cardiff, Wales, UK

[‡]Cardiff University, Cardiff, Wales, UK

Supporting Information Available. General methods, materials, instrumentation, and experimental details for all compounds, ¹H and ¹³C NMR spectra of all synthetic intermediates, 2-azapinane, and 2-azapinane salts; DEPT, HSQC, and NOE spectra of *ent*-5-TFA salt and X-ray crystallographic data for 5-HCl. This material is available free of charge via the Internet at <http://pubs.acs.org>.

products (Scheme 1).⁶ S_N' cyclizations of the endo, anti conformations (e.g. **2b**) followed by π -cyclizations of the enantiomeric α -terpinyl carbocations generate the bicyclic pinyl ions (C and *ent*-C) which undergo proton eliminations to form the pinene isomers (Scheme 1 and Figure 1).

Terpene synthase inhibitors which bind tightly in the active sites of these enzymes afford mechanistic insights, and in some instances, serve as useful aids for crystallization of the proteins yielding valuable three-dimensional structural data by X-ray diffraction analysis.⁷ Aza analogues of high-energy carbocation intermediates have found applications as mechanism-based inhibitors and active site probes for terpene synthases.^{7,8} For the purpose of characterizing monoterpene synthases, we have carried out the synthesis, characterization, and applications of 3-aza-2,3-dihydrogeranyl diphosphate, the enantiomeric 7-aza-7,8-dihydrolimonenes, 2-azabornanes, 2-azacamphanes, and azatricyclenes.⁹ In this paper we report the preparation and characterization of the enantiomeric 2-azapinanes (**5** and *ent*-**5**; Figure 1)¹⁰ potential inhibitors and active site probes for the pinene synthases.

The synthesis of [1*R*, 5*S*]-2-azapinane (**5**)¹¹ (Scheme 2) was accomplished in five steps (ca 46–50% overall yield) starting from known acetamide **8**,¹² which was available via the Beckmann rearrangement (H_2NOSO_3H , AcOH, reflux, 77–81%) of (–)-*cis*-pinonic acid (**6**)^{12,13} ($[\alpha]_D = -94.1^\circ$ (*c* 0.63, $CHCl_3$) [lit.^{13c} – 93.7 (*c* 4.60, $CHCl_3$), lit.¹⁴ – 94.2 (*c* 5–10, $CHCl_3$)] obtained by $KMnO_4$ oxidation of (–)- α -pinene.¹³ The optical purity of the starting (–)-*cis*-pinonic acid was estimated to be 99.0% assuming that the highest absolute value of specific rotation ($[\alpha]_D = +95^\circ$ ($CHCl_3$))^{13b, 15,16} reported for (+)- and (–)-*cis*-pinonic acids corresponds to 100% optically pure dextrorotatory enantiomer. An identical synthesis of the enantiomeric [1*S*, 5*R*]-2-azapinane (*ent*-**5**) was carried out in parallel 2 starting from (+)- α -pinene (**3**, $[\alpha]_D = +47.1^\circ$ (neat), 91% ee) by the same reactions and procedures illustrated in Scheme 2.

Considerable effort was required to develop a satisfactory procedure for the initial cyclization step **8** $\rightarrow$ **9**. Thus, refluxing solutions of amide acid **8** in Ac_2O ¹⁷ for 1 h, led only to the corresponding mixed pinonic acetic anhydride as revealed by ^{13}C NMR analysis, (δ_c 174.2 ($NHCO$), 169.7 (CH_2CO_2) and 168.2 ppm (CH_3CO_2). Although longer reaction time (17 h) effected partial cyclization to a 2:1 mixture of the mixed anhydride and *N*-acetyl lactam **9**, further progress towards the cyclized product was quite slow (2:3 mixture at 48 h). However addition of NaOAc and continuous removal of acetic acid during the reaction lead to increased conversion and satisfactory isolated yields of 77–80% after 72 h. Formation of the 6-membered ring in *N*-acetyl lactam **9** was accompanied by large downfield shifts of the bridgehead methine proton CH-N (4.03 $\rightarrow$ 4.80 ppm) and the *N*-acetyl CH_3 (1.95 $\rightarrow$ 2.57 ppm) in the 1H NMR spectra which seem attributable to an altered conformation of the acetyl carbonyl group.

Other known procedures for dehydrative cyclizations were unsuccessful in the case of amide acid **8**. Exposure to P_2O_5 in refluxing toluene¹⁸ led to the formation of pinonic anhydride as evidenced by the ^{13}C NMR carbonyl signals at 172.7 ($NHCO$), and 169.8 (O_2CO) ppm, without any evidence of cyclization. Similarly no cyclization was observed when the corresponding methyl ester of **8**¹⁹ was heated in refluxing xylenes. Experiments to effect cyclization of the related amino ester²⁰ through selective hydrolysis of the amide ester by means of amide *O*-alkylation²¹ to the corresponding imidate with triethyl oxonium fluoroborate according to a literature procedure [$Et_3O^+BF_4^-$ (1 eq.)/ CH_2Cl_2 followed by cold aq. $NaHCO_3$],²² lead only to extensive decomposition to uncharacterized material.

The remaining steps in the 2-azapinane synthesis proceeded readily and in high yield without extensive experimentation. Thus hydrolysis of the *N*-acetyl substituent was effected

with alumina suspended in hexane-EtOAc²³ in yields up to 98%. The detachment of the exocyclic carbonyl group was accompanied now by an upfield shift of the bridgehead N-CH to 3.26 ppm (ddd) in the proton NMR spectrum. *N*-Methylation was brought about by reactions with dimethyl sulfate²⁴ or methyl iodide (CsCO₃, acetone)²⁵ to give tertiary lactam **11** in 93–96% yield. Finally LiAlH₄ reduction of the lactam carbonyl afforded the highly volatile 2-azapinane (**5**) which was isolated and characterized as the trifluoroacetate salt **5**·TFA (80–84% yield over two steps) following a recently developed protocol for isolation of aziridine trifluoroacetate salts.^{9e} Alternatively 2-azapinane was isolated as its hydrochloride salt **5**·HCl in 78% overall yield after recrystallization from dry THF. The enantiomeric 2-azapinane (*ent*-**5**) and its salts *ent*-**5**·TFA (Figure 2) and *ent*-**5**·HCl were obtained from (+)- α -pinene by the same reaction sequence.

The structure of the 2-azapinane salts was confirmed and the anti configuration of the N-CH₃ group was established by 1-D, 2-D, and NOE NMR experiments with *ent*-**5** trifluoroacetate (Figure 2). The key proton resonances H3-anti (2.94 ppm), H3-syn (4.07), NH (9.21), H1 (3.53), H6-syn (1.82), 8-CH₃ (1.07) and 9-CH₃ (1.33) as well as the corresponding ¹³C NMR signals were assigned by means of DEPT, HSQC, NOE, and analysis of coupling patterns. The NOE data in Figure 2 were critical in assigning the more stable anti position of the NCH₃ generated in the salt formation. This conclusion was verified by an X-ray diffraction analysis of [1*R*, 5*S*]-2-azapinane hydrochloride salt (Figure 3) showing a bicyclic ring system with structural features very similar to those found in *N*-*tert*-butyloxycarbonyl-7-aza-4-oxatricyclo(4.3.1.0^{2,6})decan-3-one.²⁶ Although protonation of 2-azapinane with TFA and HCl most likely occurs initially opposite to the germinal dimethyl bridge in accord with the stereochemistry of reactions with the pinenes and their derivatives,^{1,5} equilibration through rapid proton transfers with the free amine evidently leads to the observed stable configuration of the isolated salts.

The 5-step synthetic schemes described provide efficient access to the [1*R*, 5*S*]- and [1*S*, 5*R*]-2-azapinanes (**5** and *ent*-**5**) in enantiomerically pure form. These novel aza analogues are now available for use as mechanism-based inhibitors of the pinene synthases, and as potential active site probes in the crystallography of these ubiquitous monoterpene cyclases. In addition, the enantiomerically pure bicyclic lactams [1*R*, 5*S*]- and [1*S*, 5*R*]-**9** as well as the 2-azapinanes are suitable starting materials for synthesis of novel heterocycles related to the pinanes, such as the 2-azapinane analogue **13** of pinane-thromboxane A₂. (**12**)²⁷ (Figure 4). A proposed synthesis could follow synthetic procedures developed earlier for the synthesis of pinane-thromboxane A₂.²⁸

Crystal Structure

The salt crystallizes in the chiral space group P2₁. Protonation occurs at the N2 position, with the chloride counter ion also included in the crystal structure (See Supporting Information). The cyclobutyl ring C1-C6-C5-C7 displays the longest C-C bonds (in the 1.542(3) – 1.560(3) range) and the most acute angles C-C-C angles (in the 85.24(16) – 88.50(17) range).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by a NIH grant to R.M.C. (GM 13956).

References

1. (a) Grayson DH. Nat Prod Rep. 2000; 17:385–419. [PubMed: 11014339] (b) Erman, WF. In Chemistry of the Monoterpenes Part A. Gassman, PG., editor. Vol. 11. Marcel Dekker; New York: 1985. p. 213-249. (c) Erman, WF. In Chemistry of the Monoterpenes Part B. Gassman, PG., editor. Vol. 11. Marcel Dekker; New York: 1985. p. 929-1033.
2. Derfer, JM.; Traynor, SG. The Chemistry of Turpentine. In: Zinkel, DF.; Russell, J., editors. Naval Stores: Production, Chemistry, Utilization. Vol. Chapter 8. Pulp Chemicals Association; New York: 1989.
3. Phillips MA, Savage TJ, Croteau R. Arch Biochem Biophys. 1999; 372:197–204. [PubMed: 10562434]
4. Il'ina IV, Volcho KP, Salakhutdinov NF. Russ J Org Chem. 2008; 44:1–23.
5. (a) Ramachandran PV. Aldrichim Acta. 2002; 35:23–35. (b) Cho BT. Aldrichim Acta. 2002; 35:3–16. (c) Matteson, DS. Stereo-directed Synthesis with Organoboranes. Springer-Verlag; Berlin: 1995.
6. Wise, ML.; Croteau, R. Comprehensive Natural Products Chemistry. In: Cane, DE., editor. Isoprenoids Including Carotenoids and Steroids. Vol. 2 Chapter 5. Elsevier; Amsterdam: 1999. p. 97-152.
7. Christianson DW. Chem Rev. 2006; 106:3412–3442. [PubMed: 16895335]
8. For reviews and leading references, see: (a) Narula AS, Rahier A, Benveniste P, Schuber F. J Am Chem Soc. 1981; 103:2408–2409. (b) Rahier A, Taton M, Benveniste P. Biochem Soc Trans. 1990:48–52. [PubMed: 2185086] (c) Poulter CD. Acc Chem Res. 1990; 23:70–77. (d) Abe I, Tomesch JC, Wattanasin S, Prestwich GD. Nat Prod Rep. 1994; 11:279–302. [PubMed: 15200015] (e) Rahier A, Taton M, Bouviernave P, Schmitt P, Beneviste P, Schuber F, Narula AS, Cattel L, Anding C, Place P. Lipids. 1986; 21:52–62.
9. (a) Szabo CM, Matsumura Y, Fukura S, Martin MB, Sanders JM, Sengupta S, Cieslak JA, Loftus TC, Lea CR, Lee H-J, Koohang A, Coates RM, Sagami H, Oldfield E. J Med Chem. 2002; 45:2185–2196. [PubMed: 12014956] (b) McGeady P, Pyun HJ, Coates RM, Croteau R. Arch Biochem Biophys. 1992; 299:63–72. [PubMed: 1444453] (c) Whittington DA, Wise ML, Urbansky M, Coates RM, Croteau RB, Christianson DW. Proc Natl Acad Sci USA. 2002; 99:15375–15380. [PubMed: 12432096] (d) Faraldos JA, Coates RM. unpublished results. (e) Hu H, Faraldos JA, Coates RM. J Am Chem Soc. 2009; 131:11998–12006. [PubMed: 19653649]
10. The systematic names and absolute stereochemistry of azapinanes 5 and *ent*-5 are [1*R*, 5*S*]- and [1*S*, 5*R*]-2,7,7-trimethyl-2-azabicyclo[3.1.1]heptane derived from (–)- and (+)- α -pinene, respectively.
11. For a summary of a synthesis of the isomeric 3-azapinane, see Gannon M, Postlewhite A, McElhinney RS. J Chem Res, Synop. 1979; 12:393.
12. Borges JR, Fernandez F, Garcia X, Huergueta AR, Lopez C. Org Prep Proc Int. 1997; 29:303–307.
13. (a) Delepine M. Bull Soc Chim Fr. 1936:1369–1382. (b) Wolk JL, Goldschmidt Z, Dunkelblum E. Synthesis. 1986:347–348. For purification of cis-pinonic acid see, (c) Muscio OJ Jr, Poulter CD. J Org Chem. 1974; 39:3288–3291.
14. Thoi LV. Ann Chim (Paris). 1955; 10:35–91.
15. Barbier P, Grignard V. Bull Soc Chim Fr. 1910:548–557.
16. Harispe M, Mea D. Bull Soc Chim Fr. 1962:1340–1343.
17. Pavia MR, Moos WH, Hershenson FM. J Org Chem. 1990; 55:560–564.
18. Lindsay Smith JR, Norman ROC, Rose ME, Curran ACW. J Chem Soc, Perkin Trans I. 1979:1185–1188.
19. Karpf M, Djerassi C. J Am Chem Soc. 1981; 103:302–306.
20. (a) Boeckman RK Jr, Perni RB. J Org Chem. 1986; 51:5486–5489. (b) Lyssenko KA, Lenev DA, Kostyanovsky RG. Tetrahedron. 2002; 58:8525–8537.
21. Meerwein H, Hinz G, Hofmann P, Kroning E, Pfiel E. J Prakt Chem. 1937; 147:257–285.
22. (a) Hanessian S. Tetrahedron Lett. 1967; 16:1549–1552. [PubMed: 6042551] (b) Keck GE, McLaws MD, Wager TT. Tetrahedron. 2000; 56:9875–9883.
23. Lin G, Midha KK, Hawes EM. J Heterocyclic Chem. 1991; 28:215–219.

24. Cushman M, Chen JK. *J Org Chem.* 1987; 52:1517–1521.
25. (a) Gaul C, Njardarson JT, Shan D, Dorn DC, Wu KD, Tong WP, Huang XY, Moore MAS, Danishefsky SJ. *J Am Chem Soc.* 2004; 126:11326–11337. [PubMed: 15355116] (b) Paquette LA, Wise LD. *J Am Chem Soc.* 1965; 87:1561–1566. (c) Fujita R, Yoshisuji T, Wakayanagi S, Wakamatsu H, Matsuzaki H. *Chem Pharm Bull.* 2006; 54:204–208. [PubMed: 16462064]
26. Basler B, Schuster O, Bach T. *J Org Chem.* 2005; 70:9798–9808. [PubMed: 16292808]
27. For a previous synthesis of an amino pinane-thromboxane A₂ analog see: (a) Mais DE, Knapp D, Halushka P, Ballard K, Hamanaka N. *Tetrahedron Lett.* 1984; 25:4207–4210. (b) Mais DE, DeHoll D, Sightler H, Halushka PV. *Eur J Pharm.* 1988; 148:309–315.
28. Nicolau KC, Magolda RL, Smith JB, Aharony D, Smith EF, Lefer AM. *Proc Natl Acad Sci USA.* 1979; 76:2566–2570. [PubMed: 288046]

Figure 1.

Enantiomeric 2-azapinane salts (**5 $\cdot$ H⁺** and **ent-5 $\cdot$ H⁺**), aza analogues of the pinyl carbocation intermediates (**C** and **ent-C**) in pinene biosynthesis.

Figure 2.

Key NOE correlations observed for [1*S*, 5*R*]-2-azapinane trifluoroacetate salt (*ent*-5•TFA): H1 → NH, H1 → H6-syn, H6-anti → NMe, NMe → H3-anti, NH → 8Me, H3-syn → NH, H3-syn → H4-syn.

Figure 3. ORTEP diagram of [1*R*, 5*S*]-2-azapinane hydrochloride (5·HCl) from the X-ray crystal structure determination showing 40% probability with the chloride counter ion omitted.

Figure 4.
Structures of pinane-thromboxane A₂ and its 2-azapinane analogue

Scheme 1.

Scheme 2.