Published in final edited form as:

Eur J Neurol. 2012 October ; 19(10): 1276–1277. doi:10.1111/j.1468-1331.2012.03746.x.

Estrogen in ischemic stroke: the Debate Continues

Bharti Manwani and Louise D. McCullough*

Departments of Neurology and Neuroscience, University of Connecticut Health Center, Farmington, CT

Sex differences abound in the epidemiological literature on ischemic stroke. Females enjoy a lower stroke incidence throughout most of the lifespan; however this benefit dissipates with aging [1]. Women over 80 have higher mortality and poorer functional outcomes compared to age-matched men after stroke. Over the next twenty years this disparity is expected to increase, and women will bear the brunt of stroke related death and disability. Much of the “female-protected” phenotype has been attributed to life-long exposure to estrogen, and the rise in stroke incidence occurs concomitantly with loss of ovarian function. Since a low estrogenic state (menopause) is associated with this transition into the higher stroke risk state, the link between estrogen and stroke has been an area of interest for many stroke investigators (reviewed recently in [2]). It is now well recognized that estradiol plays a vital role in the ischemic brain. Animal models of experimental stroke have shown that estrogen (17- β estradiol) is a robust neuroprotective agent in both males and females [3].

Despite the strong experimental evidence of estrogen's beneficial effects in pre-clinical stroke models, estrogen supplementation in clinical trials involving post-menopausal women have not been successful. In the Women's Health Initiative (WHI), administration of conjugated equine estrogen (0.625 mg/day) to post menopausal women without a history of stroke increased stroke incidence [4]. In the Women's Estrogen For Stroke Trial (WEST) (1mg/day) of 17 beta-estradiol surprisingly increased mortality and morbidity in post-menopausal women with a recent TIA or non-disabling stroke [5]. Although as noted by many, numerous issues make these trials somewhat difficult to interpret such as the initiation of therapy well after menopause, use in patients with established vascular disease, and administration of supraphysiological doses of estrogen, this has led to a reassessment of the efficacy of exogenous estrogen for the prevention of stroke in post-menopausal women. However, much less is known regarding estrogen effects in the acute injury period. In this issue of *The European Journal of Neurology*, Pappa and colleagues evaluated serum levels of endogenous estradiol in postmenopausal women with acute ischemic or hemorrhagic stroke [6].

In this study, serum levels of estradiol were evaluated in 302 postmenopausal women (age 73.6 $\pm$ 12.9 years; approximately 26.7 $\pm$ 10.6 years post menopause) admitted with acute ischemic stroke. Patients on steroid or hormone replacement therapy or women with transient ischemic attack or subarachnoid hemorrhage were excluded. This is a novel approach as these women were evaluated in the acute period, in which data is sorely lacking. Historically, most clinical studies have focused on the effects of chronic exogenous administration of estrogen on stroke incidence. The main question addressed was whether the levels of endogenous estradiol in elderly women after stroke were associated with changes in stroke severity or outcome. The authors utilized the National Institute of Health

*Corresponding Author: Department of Neurology, University of Connecticut Health Center, 263 Farmington Avenue, MC1840, Farmington, CT 06030 lmcullough@uconn.edu Phone number: (860) 679-3186 Fax number: (860) 679-1181.

Conflict of Interest: None

Stroke Scale (NIHSS) and modified Rankin Scale (mRS) at both admission and one month post stroke. Estradiol levels, measured acutely post stroke, were significantly higher in non-survivors as compared to survivors one month after stroke. They also found that endogenous estradiol levels were an independent predictor of stroke severity as quantified by NIHSS at the time of admission. In a multivariate analysis, estradiol levels continued to be a significant independent predictor of adverse functional outcomes and handicap (mRS 4) one-month post stroke even after removing confounders such as age and initial NIHSS. As hemorrhagic and ischemic stroke pathophysiology may differ, the authors also removed the 13% of patients with hemorrhage, and repeated the analysis (261 patients) and found similar results. Importantly, as testosterone can be peripherally converted to estrogen via the actions of aromatase in adipose tissue, the authors also measured BMIs and found no association between BMI and outcomes. These findings are consistent with recent reports showing increased mortality [7] and stroke risk (also associated with greater central adiposity) [8] in post menopausal women with higher estrogen levels.

The findings of this study suggest that high levels of endogenous estradiol are deleterious in acute stroke. The authors suggest that the concept of estrogen as a neuroprotective agent “should be challenged”. However, these results should be interpreted with caution, and certainly these results do not definitively answer the question of whether acute estrogen is detrimental or beneficial, as well noted by the authors. There is a clear association between estradiol levels and stroke outcomes, but this does not imply causation. The poorer outcomes seen in women with the highest levels estradiol levels may not be a direct biological effect but simply reflective of larger lesion sizes (infarct volumes) which were not assessed in this study. Indeed, raising endogenous estradiol levels may even reflect an endogenous protective mechanism, as has been seen in animal models in which up-regulation of aromatase occurs in the infarct area, and infarct damage worsens with loss or inhibition of aromatase in the acutely injured brain [9]. Another important limitation of this paper is that the authors do not mention the exact time points at which blood samples were collected from the patients, although the authors previous studies (using the same cohort) state that these were collected in the morning 2-3 days after onset [10]. However, biomarkers are known to vary widely, and can be especially sensitive to stroke acuity, post-stroke infection rates, medication use etc. many which were not controlled for in this initial study. Additionally, levels of other endogenous estrogens (i.e., estrone) were not assessed. Future studies examining serial samples and the utilization of an age-matched control group (for example, post menopausal women admitted to the hospital for reasons other than stroke) may provide some clarity on levels of estradiol in stress and sickness. Overall, these results although intriguing, and a large step forward, yet again demonstrate the many uncertainties related to the role of estrogen in postmenopausal women with ischemic stroke.

Acknowledgments

Funding source: NIH R01-NS055215 to L.D. McCullough and AHA predoctoral fellowship 11PRE7440068 to B. Manwani

References

1. Roger VL, et al. Heart Disease and Stroke Statistics--2011 Update: A Report From the American Heart Association. *Circulation*. 123(4):e18–e209. [PubMed: 21160056]
2. Lisabeth L, Bushnell C. Stroke risk in women: the role of menopause and hormone therapy. *Lancet Neurol*. 11(1):82–91. [PubMed: 22172623]
3. McCullough LD, Hurn PD. Estrogen and ischemic neuroprotection: an integrated view. *Trends Endocrinol Metab*. 2003; 14(5):228–35. [PubMed: 12826329]

4. Wassertheil-Smoller S, et al. Effect of estrogen plus progestin on stroke in postmenopausal women: the Women's Health Initiative: a randomized trial. *Jama*. 2003; 289(20):2673–84. [PubMed: 12771114]
5. Viscoli CM, et al. A clinical trial of estrogen-replacement therapy after ischemic stroke. *N Engl J Med*. 2001; 345(17):1243–9. [PubMed: 11680444]
6. Pappa T, et al. Estradiol levels predict adverse short term health outcomes in post-menopausal acute stroke women. *European Journal of Neurology*. XXX. In Press.
7. Maggio M, et al. Relationship between higher estradiol levels and 9-year mortality in older women: the Invecchiare in Chianti study. *J Am Geriatr Soc*. 2009; 57(10):1810–5. [PubMed: 19737330]
8. Lee JS, et al. Prospective study of endogenous circulating estradiol and risk of stroke in older women. *Arch Neurol*. 67(2):195–201. [PubMed: 20142527]
9. McCullough LD, et al. Aromatase cytochrome P450 and extragonadal estrogen play a role in ischemic neuroprotection. *J Neurosci*. 2003; 23(25):8701–5. [PubMed: 14507969]
10. Pappa T, et al. Severity and outcome of acute stroke in women: relation to adrenal sex steroid levels. *Metabolism*. 61(1):84–91. [PubMed: 21820139]