RESEARCH REPORTS

Clinical

A.C.R. Tanner^{1,3*}, A.L. Sonis⁴,
P. Lif Holgersson⁶, J.R. Starr^{2,5,7},
Y. Nunez⁸, C.A. Kressler^{1,3},
B.J. Paster^{1,3}, and I. Johansson⁶

¹Department of Molecular Genetics; ²Center for Clinical and Translational Research, The Forsyth Institute, Cambridge, MA, USA; ³Department of Oral Medicine, Infection and Immunity; ⁴Department of Developmental Biology; ⁵Department of Oral Health Policy and Epidemiology, Harvard School of Dental Medicine, Harvard University, Boston, MA, USA; ⁶Department of Odontology, Cariology Section, Faculty of Medicine, Umeå University, Umeå, Sweden; ⁷Department of Epidemiology, School of Public Health, University of Washington, Seattle, WA; and ⁸Seton Hall University, South Orange, NJ, USA; *corresponding author, annetanner@forsyth.org

J Dent Res 91(9):853-858, 2012

ABSTRACT

White-spot lesions (WSL) associated with orthodontic appliances are a cosmetic problem and increase risk for cavities. We characterized the microbiota of WSL, accounting for confounding due to gingivitis. Participants were 60 children with fixed appliances, aged between 10 and 19 yrs, half with WSL. Plaque samples were assayed by a 16S rRNA-based microarray (HOMIM) and by PCR. Mean gingival index was positively associated with WSL ($p = 0.018$). Taxa associated with WSL by microarray included *Granulicatella elegans* ($p = 0.01$), *Veillonellaceae* sp. HOT 155 ($p < 0.01$), and *Bifidobacterium* Cluster 1 ($p = 0.11$), and by qPCR, *Streptococcus mutans* ($p = 0.008$) and *Scardovia wiggsiae* ($p = 0.04$). Taxa associated with gingivitis by microarray included: *Gemella sanguinis* ($p = 0.002$), *Actinomyces* sp. HOT 448 ($p = 0.003$), *Prevotella* cluster IV ($p = 0.021$), and *Streptococcus* sp. HOT 071/070 ($p = 0.023$); and levels of *S. mutans* ($p = 0.02$) and *Bifidobacteriaceae* ($p = 0.012$) by qPCR. Species' associations with WSL were minimally changed with adjustment for gingivitis level. Partial least-squares discriminant analysis yielded good discrimination between children with and those without WSL. *Granulicatella*, *Veillonellaceae* and *Bifidobacteriaceae*, in addition to *S. mutans* and *S. wiggsiae*, were associated with the presence of WSL in adolescents undergoing orthodontic treatment. Many taxa showed a stronger association with gingivitis than with WSL.

KEY WORDS: orthodontic, adolescents, white-spot lesions, microbial ecology, *Scardovia wiggsiae*, HOMIM.

DOI: 10.1177/0022034512455031

Received April 10, 2012; Last revision June 20, 2012; Accepted June 21, 2012

A supplemental appendix to this article is published electronically only at <http://jdr.sagepub.com/supplemental>.

© International & American Associations for Dental Research

White-spot Lesions and Gingivitis Microbiotas in Orthodontic Patients

INTRODUCTION

Orthodontic treatment with fixed appliances increases the risk of development of decalcified white-spot lesions (WSL; Lovrov *et al.*, 2007; van der Veen *et al.*, 2010), which can occur within 6 mos of appliance placement (Tufekci *et al.*, 2011) (Fig. 1). While the majority of WSL re-mineralize when appliances are removed, WSL can progress to cavitation (van der Veen *et al.*, 2010) and are a cosmetic problem (Maxfield *et al.*, 2012). WSL result from increased plaque accumulation due to inadequate oral hygiene around orthodontic appliances (Chapman *et al.*, 2010), which also leads to gingivitis (Naranjo *et al.*, 2006; Rego *et al.*, 2010).

Evaluation of caries-associated bacteria in orthodontic patients has focused principally on *Streptococcus mutans* and lactobacilli (Boyar *et al.*, 1989; Ahn *et al.*, 2007). Plaque and WSL development under orthodontic bands observed by a multispecies microarray was associated with a diverse microbiota (Torlakovic *et al.*, 2012), including *Scardovia wiggsiae*, a newly recognized species (Downes *et al.*, 2011) associated with severe early childhood caries (ECC; Tanner *et al.*, 2011a,b). *Prevotella*, *Capnocytophaga*, *Selenomonas*, and *Fusobacterium* species have also been detected in childhood caries (Aas *et al.*, 2008; Gross *et al.*, 2010; Tanner *et al.*, 2011b). These latter taxa are more frequently associated with gingivitis, and their role in dental caries is unclear.

The goal of this study was to evaluate the microbiota of WSL in orthodontic patients as a model of initial caries development in children. We assessed the relationship between WSL and gingivitis and whether they share microbial risk factors.

MATERIALS & METHODS

Clinical Methods

Children aged between 10 and 19 yrs old, with stainless steel brackets bonded with non-fluoride-releasing adhesive (Dentsply GAC, Bohemia, NY, USA), were recruited from the dental department of Children's Hospital, Boston. Children were medically healthy and had not used antibiotics within the preceding 3 mos. The parent or guardian provided informed consent, and the children

Figure 1. White-spot lesions at gingival margins of lower left lateral incisors between fixed orthodontic bracket and gingival margin. Marked gingival inflammation, edema, and redness can be seen at the gingival margins of central and lateral lower incisors.

with orthodontia agreed to participate. Participants resided in water-fluoridated areas (1 ppm) and reported using fluoridated toothpastes. They were sequentially approached until 30 children with WSL and 30 without WSL had been recruited. The Institutional Review Boards of Children's Hospital, Harvard University, and The Forsyth Institute approved the study design, protocol, and informed consent. The clinical phase of the study was conducted between July 2009 and January 2010.

The numbers of decayed, missing, and filled teeth (DMFT) when appliances were placed were recorded. At a 6- to 12-month follow-up appointment, bracketed teeth were cleaned and examined for WSL adjacent to the bonded brackets by direct visualization with 2X magnification (dental loops), and from intra-oral photographs. Gingival and plaque indices were measured with a 1 to 4 score by quadrant. Each child completed a short survey. WSL were usually sampled from buccal anterior tooth surfaces, and from a matched site in children without WSL. One plaque sample was taken from each child by means of sterile toothpicks, and DNA from samples was purified with MasterPure kits (Epicentre Biotechnologies, Madison, WI, USA; Kanasi *et al.*, 2010b).

Microbiological Methods

Microarray Analysis

Samples were analyzed by microarray to approximately 300 bacterial taxa with the HOMIM assay (<http://mim.forsyth.org>) as described previously (Tanner *et al.*, 2011a; Torlakovic *et al.*, 2012). Probes to 100 taxa included in the current analyses were selected based on reactivity to at least one, but not to all samples.

Bacterial-specific PCR Analysis

Quantitative-PCR was performed to detect *S. mutans* (Psoter *et al.*, 2011) and *Bifidobacteriaceae* (Matsuki *et al.*, 2004). We developed qPCR for *S. wiggsiae* using primers designed using Unipro UGENE: forward, 5'-TGCCTGAAGCCCAGGACGTA-3', and reverse, 5'-TGTGTGGTGTGGTGAAGTGGACTTTAT-3'. The qPCR standard curve reaction mixture, total volume 5 μ L, consisted of Roche SYBR Green master mix (Atlanta, GA, USA) 2X (2.5 μ L), 20 μ M of each primer (0.4 μ L), PCR-grade water (0.1 μ L), and *S. wiggsiae* FO 424 genomic DNA

from 5 ng/ μ L to 5 fg/ μ L (2 μ L). The qPCR conditions were an initial denaturation of 95°C for 15 min, followed by 45 cycles at 95°C for 15 sec, 67°C for 30 sec, and 72°C for 40 sec, and generated a 766-bp amplicon. *S. wiggsiae* was 10-fold serially diluted, plated, and cultured anaerobically, and colonies were counted after 5 days. Aliquots of dilutions were assayed by qPCR for DNA quantitation.

We $\log_{10}$ -transformed the resulting DNA levels after adding 0.001 to *Bifidobacterium* and *S. wiggsiae* counts and 0.01 to the *S. mutans* counts. To derive qualitative qPCR results, we denoted the presence of taxa as *S. mutans* > 0.55, *S. wiggsiae* > 0.55, and *Bifidobacterium* > 7.8.

Personnel performing microbiology assays were not aware of the clinical status of samples.

Statistical Analyses

WSL and Bacteria

We used logistic regression to estimate associations between the presence of WSL and the presence of individual bacterial taxa (microarray or PCR) or levels of specific taxa (qPCR). If the statistical analyses involved any group with < 5 children, we used exact logistic regression. For all odds ratio (OR) estimates, we calculated 95% confidence intervals and p-values. Because the study was exploratory and potentially underpowered, we did not adjust these for multiple comparisons and do not interpret them as dichotomous significance tests (Thomas *et al.*, 1985; Rothman, 1990). For hypothesis generation, we report taxa that exhibited strong or moderately strong associations, with $p < 0.2$. To explore the possibility of confounding by bacterial associations with gingivitis, we refit all models including each child's mean gingivitis score as a covariate. We repeated the analysis after excluding individuals with *S. mutans* DNA < 5 ($n = 5$ and 8 for those with and without WSL, respectively).

Gingivitis and Bacteria

We fit logistic or exact logistic regression models to estimate the associations between gingivitis level (high, ≥ 2 ; or low, < 2) and the presence of each taxon. Secondly, we refit the models adjusting for the presence or absence of WSL.

The statistical analyses were performed with Stata software version 10.1 (StataCorp. 2007, *Stata Statistical Software: Release 10*. StataCorp LP, College Station, TX, USA).

Partial Least-squares Discriminant Analysis (OPLS-DA)

Multivariate partial least-squares discriminant analysis (OPLS-DA) was performed (SIMCA P+, version 12.0, Umetrics AB, Umeå, Sweden) as described previously (Kanasi *et al.*, 2010b; Lif Holgersson *et al.*, 2011). Partial least-squares (PLS), which defines the maximum separation between class members (here WSL), is suitable for data where the number of observations is smaller than the number of variables, and where the X variables co-vary. Dichotomous HOMIM and qPCR signals, plaque, gingivitis, and drinks between and with meals built the X-block, and the presence of WSL the Y-block (outcome). The criteria for including a HOMIM variable into the x-block were

Figure 2. Microbiota from HOMIM microarray analysis. **(A)** Microbiota associated with white-spot lesions (WSL) ordered according to detection in children with or without WSL. *Granulicatella elegans* and *Veillonellaceae* species were detected more frequently in samples from WSL than in those without WSL, whereas *Cardiobacterium hominis* was detected more frequently from children without WSL. The exact p values, odds ratios, and 95% confidence intervals are in Appendix Table 2. **(B)** Microbiota associated with gingivitis ordered according to detection in children with or without WSL, as in **(A)**. In these children, more taxa were associated with higher gingivitis than with WSL, including *Actinomyces* sp. HOT 448 and *Gemella sanguinis*. The exact p values, odds ratios, and 95% confidence intervals are in Appendix Table 3.

that the detection prevalence differed by $\geq 14\%$ or that the odds ratio to have WSL if having a species was > 2 or < 0.2 , or that it differed between groups at $p < 0.05$. All variables were auto-scaled to unit variance, and cross-validated prediction of Y was calculated.

RESULTS

The study population was comprised of 28% Hispanic, 34% White, 17% Asian, and 25% Black children, and most were born in the USA. Cases were, on average, over a year older than controls, with higher proportions of girls and Hispanic children (Appendix Table 1). Children with WSL had a higher mean gingival index, but did not differ in DMFT or plaque index, compared with children without WSL. Most children reported flossing. There were minimal differences in the diet (Appendix Table 1). Children reported averages of 1.6 mealtime and 1.8 between-meal beverages, and 1.3 nighttime snacks. In comparisons of mealtime with between-meal beverages, frequencies were, respectively, 20% and 7% milk, 30% and 37% juice, and

7% and 12% diet sodas, with no differences in water or sugar-containing sodas.

HOMIM Microarray Data

White-spot Lesions

Granulicatella elegans (OR 12.0, $p = 0.01$) and *Veillonellaceae* species HOT 155 (OR 4.6, $p = 0.01$) were detected more frequently in samples from WSL than in those without WSL (Fig. 2A, Appendix Table 2). Other taxa exhibiting moderately strong associations with the presence of WSL ($p < 0.2$) included *Bifidobacterium* Cluster I (OR = 5.8), *Selenomonas sputigena* (OR 2.3), *Prevotella* Cluster IV (OR 2.3), *Streptococcus* sp. HOT 071/070 (OR 2.5), *Prevotella melaninogenica* (OR 2.3), and *S. wiggsiae* (OR 5.7). *Cardiobacterium hominis* was associated with controls (OR = 0.3, $p = 0.02$). The magnitude of ORs and p values attenuated only slightly after adjustment for gingivitis levels or exclusion of children with *S. mutans* DNA > 5 (data not shown).

Figure 3. *S. mutans*, *S. wiggisiae*, and *Bifidobacterium* detected by qPCR. **(A)** Mean *S. mutans*, *S. wiggisiae*, and *Bifidobacterium* levels comparing children with and without white-spot lesions (WSL). Species levels are in pg of DNA/1 μ L sample, total sample size 100 μ L. Mean levels of *S. mutans* and *S. wiggisiae* were higher in children with WSL than in those without WSL. The odds ratios and 95% confidence intervals, and adjustment for gingivitis, are in Appendix Table 2. **(B)** Mean *S. mutans*, *S. wiggisiae*, and *Bifidobacterium* levels (by qPCR) comparing children with and without elevated gingivitis. Species levels are in pg of DNA/1 μ L sample, total sample size 100 μ L. *S. mutans* and *Bifidobacterium* were detected at higher levels in increased gingivitis. The odds ratios and 95% confidence intervals and adjustment for WSLs are in Appendix Table 3. **(C)** Detection frequencies of *S. mutans*, *S. wiggisiae*, and *Bifidobacterium* in children with and without white-spot lesions (WSL). *S. mutans* and *S. wiggisiae*, at a modest level, were detected more frequently in samples from children with WSL than in those without WSL at a comparable threshold of detection (0.55 pg DNA). *Bifidobacterium* were not associated with WSL at a detection threshold >7.8 ng DNA, reflecting the higher levels at which *Bifidobacterium* were detected in samples (Fig. 3A). The odds ratios and 95% confidence intervals and adjustment for gingivitis are in Appendix Table 2. **(D)** Detection frequencies of *S. mutans*, *S. wiggisiae*, and *Bifidobacterium* in children with and without elevated gingivitis. *S. mutans*, *S. wiggisiae*, and *Bifidobacterium* were detected more frequently in children with higher levels of gingivitis. The odds ratios and 95% confidence intervals and adjustment for WSL are in Appendix Table 3.

The *Lactobacillus* cluster I (*Lactobacillus casei*, *Lactobacillus paracasei*, *Lactobacillus rhamnosus*) was detected in a low proportion of children, whereas *Actinomyces* cluster I (*Actinomyces meyeri*, *Actinomyces odontolyticus*, *Actinomyces oricola*, *Actinomyces naeslundii* II) and *Actinomyces gerensceriae* were detected in over 65% children, but their detection and other *Actinomyces* did not differ between children with and those without WSL (Appendix Table 2). *Actinomyces* sp. HOT 448 was detected in 30% of children with WSL compared with 17% of those without WSL (OR 2.1). *S. mutans* and *Streptococcus sobrinus* were detected infrequently and were not WSL-associated in the microarray data. In children with low levels of *S. mutans*, as assessed by qPCR, *S. wiggisiae* by qPCR and *P. melaninogenica* and *Veillonellaceae* species (HOT 155) from HOMIM were associated with WSL with or without adjustment for gingivitis.

Gingivitis

Gemella sanguinis (OR 8.4, $p = 0.002$), *Actinomyces* sp. HOT 448 (OR 9.8, $p = 0.003$), *Prevotella* Cluster IV (OR 3.5, $p = 0.021$), *Streptococcus* sp. HOT 071/070 (OR 5.3, $p = 0.023$), *Streptococcus parasanguinis* (OR 3.2, $p = 0.035$), *Leptotrichia* sp. HOT 417/462 (OR 9.2, $p = 0.041$), and *Gemella haemolysans* (OR 3.0, $p = 0.042$) were associated with gingivitis (Fig. 2B, Appendix Table 3). Other taxa exhibiting moderate associations with more gingivitis ($p < 0.2$) included: *Dialister invisus* (OR 2.9), *Megasphaera micronuciformis* (OR 2.8), *Veillonellaceae* sp. HOT 155 (OR 2.7), *Selenomonas sputigena* (OR 2.6), *Selenomonas infelix* (OR 2.8), *Actinomyces* Cluster I (OR 2.6), *Prevotella pallens* (OR 4.5), *S. wiggisiae* (OR 6.1), *Leptotrichia hofstadii* (OR 6.1), and *Slackia exigua* (OR 2.3). Taxa in reduced gingivitis included *Fusobacterium periodonticum* (OR 0.04) and *Neisseria elongata* (OR 0.04). The magnitude of ORs and p values changed only slightly after adjustment for the presence of WSL. This adjustment strengthened the magnitude of ORs more often than it attenuated them.

PCR Data

qPCR detection of *S. wiggisiae* ranged from $> 10^2$ to 10^7 fg DNA (see Appendix Fig. for amplification and standard quantification plots). A 0.1-pg quantity of DNA was equivalent to 37.5 CFU. By qPCR, *S. mutans* and *S. wiggisiae* were associated with WSL in species levels, and with *S. mutans* in detection frequency (Fig. 3, Appendix Table 2). *S. mutans* and

Bifidobacterium were associated with higher gingivitis in levels and detection frequency (Fig. 3, Appendix Table 3). *S. wiggisiae* was associated with higher gingivitis only in detection frequency (Fig. 3, Appendix Tables 2 and 3).

PLS-DA Modeling

PLS-DA modeling identified a significant component that explained 40.7% and predicted 23.9% ($R^2 = 0.407$ and $Q^2 = 0.239$), although some overlap was observed between WSL and non-WSL children (Fig. 4A). *Cardiobacterium hominis* detection and drinking milk between meals were significantly associated with controls. Having WSL was significantly associated with the presence of *Granulicatella elegans*, *Veillonellaceae* sp. HOT 155, *S. wiggisiae* (qPCR), *S. mutans* (qPCR), *S. sputigena* and *Prevotella* Cluster IV, sugar drinks with meals, and gingival and plaque scores (Fig. 4B).

DISCUSSION

In this pilot study, children with WSL associated with fixed orthodontic appliances showed few differences in demographics and diet compared with control children without WSL. Microbiology findings confirmed the diversity of the microbiota of dental plaque and the relationship of *S. mutans* with fixed orthodontia previously observed in longitudinal studies of eleven (Boyar *et al.*, 1989) and eight (Torlakovic *et al.*, 2012) children. Our novel findings included the association of *Scardovia wiggsiae* with WSL in the presence and absence of *S. mutans*, and after adjustment for the presence of gingivitis, extending the association of caries with *S. wiggsiae* from severe ECC (Tanner *et al.*, 2011b).

The Human Oral Microbial Identification Microarray (HOMIM) facilitates detection of multiple taxa in plaque samples, but we observed few microbial differences between WSL- and non-WSL-associated plaques. *Granulicatella elegans* was associated with severe ECC by clonal (Kanasi *et al.*, 2010a), but not cultural or microarray, analysis of severe ECC (Tanner *et al.*, 2011a). *G. elegans*, *Veillonellaceae* sp. HOT 155, and *Bifidobacterium* Cluster I were not associated with developing WSL according to HOMIM (Torlakovic *et al.*, 2012), suggesting that association with WSL of these taxa requires further study. The increases in lactobacilli and actinomyces by culture (Boyar *et al.*, 1989; Kupietzky *et al.*, 2005), but not in the current study, may reflect differences between the microbiological assays. *Slackia exigua* was previously associated with severe ECC (Tanner *et al.*, 2011a), but in the current study, *S. exigua* was associated with gingivitis, consistent with its detection in periodontitis (Abiko *et al.*, 2010).

The microbiota of gingivitis with fixed orthodontic appliances has been described (Naranjo *et al.*, 2006; Rego *et al.*, 2010), but not with adjustment for WSL. We observed a strong association between gingivitis and detection of WSL, as we had observed with severe ECC (Tanner *et al.*, 2011b). *Gemella sanguinis*, *Actinomyces* sp. HOT 448, *Prevotella* cluster IV, *Streptococcus parasanguinis*, *Leptotrichia* sp. HOT 417/462, and *Gemella haemolysans* showed a stronger association with gingivitis than with WSL, suggesting that the primary disease association is with gingivitis. Other species we detected in association with gingivitis, including *Prevotella*, *Selenomonas*, *Actinomyces*, *Dialister*, *Slackia*, and *Parvimonas* species, have been reported in association with childhood caries (Aas *et al.*,

Figure 4. Partial least-squares (PLS) modeling plots. (A) PLS score plot of children with and without WSL. WSL children fell mainly toward the upper right, and non-WSL children mainly toward the lower left, with some overlap in the middle. The PLS model used WSL as the dependent variable, and the microarray, PCR, dietary, clinical, and demographic information as the independent matrix. Cross-validation was done by a systematic prediction of one 7th of the data by the remaining 6/7th of the data. R^2 and Q^2 -values give the capacity of the X-block to explain (R^2) and predict (Q^2) the outcome. (B) PLS column loading plot of children with and without WSL. Bars show mean PLS correlation coefficients with measurement error (and error bars representing 95% confidence interval) for children with and without WSL. Variables with highest correlation coefficients in each group are displayed. The variables with the strongest influence in the model were *C. hominis* and drinking milk between meals in non-WSL children, and *G. elegans*, *Veillonellaceae* [G] sp. HOT 155, drinking sugar-containing drinks with meals, *S. wiggsiae* (qPCR), *S. mutans* (qPCR), and higher levels of gingivitis in the WSL children.

2008; Gross *et al.*, 2010), associations that could reflect unmeasured confounding by gingivitis.

The multivariate PLS-DA modeling correctly identified most, but not all, children with WSL. The overlap may reflect similarities in the plaque and caries histories of children with and those without WSL. It may also reflect that WSL represents an early and reversible stage of caries, with only a subset of sites likely to progress if left untreated. This modeling approach is of particular value when there are more variables than study participants, and species that might be interdependent based on shared environmental requirements. Validity of the resulting model for classifying study participants will need testing in other samples.

Although the sample size was not unusually small, it could not take into account multiple hypothesis testing. Correction for

multiple comparisons would have reduced statistical power to a low level; none of the findings met the 'false discovery rate' threshold (Benjamini and Hochberg, 1995). Applying significance thresholds is one way of evaluating results but is not always useful in non-randomized studies (Rothman, 1990, 1998). The magnitude of the differences and their confidence intervals can still be interpreted in a preliminary fashion.

A strength of this study was the use of a microarray to detect multiple species and taxon-specific qPCR to improve sensitivity and quantitation of high-interest species. Including analyses for both WSL and gingivitis clarified clinical associations for some species. Multivariate modeling allowed for visualization of the differences between clinical groups. Limitations of this study include the cross-sectional design, so we could not determine whether the microbiota changed with or could predict development of WSL. Further, the use of a defined probe set may not have adequately detected key species in WSL ecology.

We conclude that there is a complex microbiota associated with WSL that can include *S. wiggsiae* and possibly *G. elegans*, *Veillonellaceae*, and *Bifidobacteriaceae*, in addition to *S. mutans*. Of clinical significance, procedures to prevent the development of WSL should include testing more bacterial taxa than mutans streptococci and *Lactobacillus* species, and consider the impact of treatment regimens on the diverse microbiota.

ACKNOWLEDGMENTS

We thank Winston Kuo and Alex Trachtenberg for assistance with qPCR assays, and Ralph Kent and Natalia Chalmers in project development. This work was conducted with support from USPHS grants DE-015847, DE-021796, and DE-007327 from the NIDCR (NIH), Throne-Holst's Foundation, the Bingham Trust, and Harvard Catalyst UL1 RR 025758, RC1 DE020549. The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article.

REFERENCES

- Aas JA, Griffen AL, Dardis SR, Lee AM, Olsen I, Dewhirst FE, *et al.* (2008). Bacteria of dental caries in primary and permanent teeth in children and young adults. *J Clin Microbiol* 46:1407-1417.
- Abiko Y, Sato T, Mayanagi G, Takahashi N (2010). Profiling of subgingival plaque biofilm microflora from periodontally healthy subjects and from subjects with periodontitis using quantitative real-time PCR. *J Periodontol* 45:389-395.
- Ahn SJ, Lim BS, Lee SJ (2007). Prevalence of cariogenic streptococci on incisor brackets detected by polymerase chain reaction. *Am J Orthod Dentofacial Orthop* 131:736-741.
- Benjamini Y, Hochberg Y (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* 57:289-300.
- Boyar RM, Thylstrup A, Holmen L, Bowden GH (1989). The microflora associated with the development of initial enamel decalcification below orthodontic bands *in vivo* in children living in a fluoridated-water area. *J Dent Res* 68:1734-1738.
- Chapman JA, Roberts WE, Eckert GJ, Kula KS, Gonzalez-Cabezas C (2010). Risk factors for incidence and severity of white spot lesions during treatment with fixed orthodontic appliances. *Am J Orthod Dentofacial Orthop* 138:188-194.
- Downes J, Mantzourani M, Beighton D, Hooper S, Wilson MJ, Nicholson A, *et al.* (2011). *Scardovia wiggsiae* sp. nov., isolated from the human oral cavity and clinical material, and emended descriptions of the genus *Scardovia* and *Scardovia inopinata*. *Int J Syst Evol Microbiol* 61(Pt 1):25-29.
- Gross EL, Leys EJ, Gasparovich SR, Firestone ND, Schwartzbaum JA, Janies DA, *et al.* (2010). Bacterial 16S sequence analysis of severe caries in young permanent teeth. *J Clin Microbiol* 48:4121-4128.
- Kanasi E, Dewhirst FE, Chalmers NI, Kent R Jr, Moore A, Hughes CV, *et al.* (2010a). Clonal analysis of the microbiota of severe early childhood caries. *Caries Res* 44:485-497.
- Kanasi E, Johansson I, Lu SC, Kressin NR, Nunn ME, Kent R Jr, *et al.* (2010b). Microbial risk markers for childhood caries in pediatricians' offices. *J Dent Res* 89:378-383.
- Kupietzky A, Majumdar AK, Shey Z, Binder R, Matheson PB (2005). Colony forming unit levels of salivary Lactobacilli and *Streptococcus mutans* in orthodontic patients. *J Clin Pediatr Dent* 30:51-53.
- Lif Holgersson P, Harnevik L, Hernell O, Tanner AC, Johansson I (2011). Mode of birth delivery affects oral microbiota in infants. *J Dent Res* 90:1183-1188.
- Lovrov S, Hertrich K, Hirschfelder U (2007). Enamel demineralization during fixed orthodontic treatment – incidence and correlation to various oral-hygiene parameters. *J Orofac Orthop* 68:353-363.
- Matsuki T, Watanabe K, Fujimoto J, Kado Y, Takada T, Matsumoto K, *et al.* (2004). Quantitative PCR with 16S rRNA-gene-targeted species-specific primers for analysis of human intestinal bifidobacteria. *Appl Environ Microbiol* 70:167-173.
- Maxfield BJ, Hamdan AM, Tufekci E, Shroff B, Best AM, Lindauer SJ (2012). Development of white spot lesions during orthodontic treatment: perceptions of patients, parents, orthodontists, and general dentists. *Am J Orthod Dentofacial Orthop* 141:337-344.
- Naranjo AA, Trivino ML, Jaramillo A, Betancourth M, Botero JE (2006). Changes in the subgingival microbiota and periodontal parameters before and 3 months after bracket placement. *Am J Orthod Dentofacial Orthop* 130:275.e17-e22.
- Psoter WJ, Ge Y, Russell SL, Chen Z, Katz RV, Jean-Charles G, *et al.* (2011). PCR detection of *Streptococcus mutans* and *Aggregatibacter actinomycetemcomitans* in dental plaque samples from Haitian adolescents. *Clin Oral Investig* 15:461-469.
- Rego RO, Oliveira CA, dos Santos-Pinto A, Jordan SF, Zambon JJ, Cirelli JA, *et al.* (2010). Clinical and microbiological studies of children and adolescents receiving orthodontic treatment. *Am J Dent* 23:317-323.
- Rothman KJ (1990). No adjustments are needed for multiple comparisons. *Epidemiology* 1:43-46.
- Rothman KJ (1998). Writing for epidemiology. *Epidemiology* 9:333-337.
- Tanner AC, Kent RL Jr, Lif Holgersson P, Hughes CV, Loo CY, Kanasi E, *et al.* (2011a). Microbiota of severe early childhood caries before and after therapy. *J Dent Res* 90:1298-1305.
- Tanner AC, Mathney JM, Kent RL, Chalmers NI, Hughes CV, Loo CY, *et al.* (2011b). Cultivable anaerobic microbiota of severe early childhood caries. *J Clin Microbiol* 49:1464-1474.
- Thomas DC, Siemiatycki J, Dewar R, Robins J, Goldberg M, Armstrong BG (1985). The problem of multiple inference in studies designed to generate hypotheses. *Am J Epidemiol* 122:1080-1095.
- Torlakovic L, Klepac-Ceraj V, Ogaard B, Cotton SL, Paster BJ, Olsen I (2012). Microbial community succession on developing lesions on human enamel. *J Oral Microbiol* 4. doi: 10.3402/jom.v4i0.16125. Epub 2012 Mar 14. (in press).
- Tufekci E, Dixon JS, Gunsolley JC, Lindauer SJ (2011). Prevalence of white spot lesions during orthodontic treatment with fixed appliances. *Angle Orthod* 81:206-210.
- van der Veen MH, Attin R, Schwestka-Polly R, Wiechmann D (2010). Caries outcomes after orthodontic treatment with fixed appliances: do lingual brackets make a difference? *Eur J Oral Sci* 118:298-303.