Molecular mechanisms underlying the fetal programming of adult disease

Thin Vo · Daniel B. Hardy

Received: 29 April 2012 / Accepted: 2 May 2012 / Published online: 24 May 2012
© The International CCN Society 2012

Abstract Adverse events in utero can be critical in determining quality of life and overall health. It is estimated that up to 50 % of metabolic syndrome diseases can be linked to an adverse fetal environment. However, the mechanisms linking impaired fetal development to these adult diseases remain elusive. This review uncovers some of the molecular mechanisms underlying how normal physiology may be impaired in fetal and postnatal life due to maternal insults in pregnancy. By understanding the mechanisms, which include epigenetic, transcriptional, endoplasmic reticulum (ER) stress, and reactive oxygen species (ROS), we also highlight how intervention in fetal and neonatal life may be able to prevent these diseases long-term.

Keywords Fetal Programming · Epigenetics · microRNA · Posttranslational Histone Modifications · DNA Methylation · ER Stress · Nuclear Receptors

Supported by: CIHR Operating Grant and Natural Sciences and Engineering Research Council of Canada

T. Vo · D. B. Hardy (✉)
The Department of Physiology & Pharmacology,
University of Western Ontario,
London, Ontario, Canada N6A 5C1
e-mail: daniel.hardy@schulich.uwo.ca

T. Vo · D. B. Hardy
The Department of Obstetrics & Gynecology,
University of Western Ontario,
London, Ontario, Canada N6A 5C1

T. Vo · D. B. Hardy
The Children's Health Research Institute,
London, Ontario, Canada N6C 2V5

T. Vo · D. B. Hardy
The Lawson Health Research Institute,
London, Ontario, Canada N6A 4V2

Introduction

The incidence of low birth weight babies (defined as ≤ 2500 g or 5.5lbs) worldwide is estimated to be 15.5 %, and that number is greatly underestimated (Lopez et al. 2006). As a general indicator of public health, it is imperative that we study the etiology and outcomes of the individuals that develop as low birth weight babies. Similarly, the incidences of non-communicable diseases such as heart disease, type II diabetes, hypertension, obesity and the metabolic syndrome are on the rise in North America (Ford et al. 2004; WRITING GROUP MEMBERS et al. 2010; McGuire 2011). More than one in three Americans are obese (Ogden et al. 2012), compared to one in four in Canada (McGuire 2011). This trend is beginning to appear all over the world, including in developing nations (Grundy 2008; Nestel et al. 2007). Moreover, diseases such as cardiovascular disease are a leading cause of death in the United States (WRITING GROUP MEMBERS et al. 2010). Although the prevalence of these chronic and non-communicable diseases puts tremendous strain on the health care system and society, intervention with diet or drugs can play a significant role to reduce their incidence. For example, a meta-analysis prospective study, using data from 58 clinical trials as well as nine cohort studies, indicates that in patients with vascular disease, a 1.8 mM reduction in LDL cholesterol by statins resulted in a 17 % reduction in stroke and a 60 % reduction in the risk of ischemic heart disease (Law et al. 2003). Current treatment for these diseases, in addition to adopting a healthy lifestyle via alterations in diet and promoting exercise, relies on the use of pharmaceuticals. Unfortunately, these treatments are not efficacious for all individuals; for example, in some patients statin treatment can lead to rhabdomyolysis and hepatitis-associated liver failure (Law et al. 2003). Therefore, additional strategies in disease prevention are warranted.

The ‘fetal origins’ or ‘Barker’ hypothesis suggests that impaired growth of the fetus during gestation strongly correlates to the development of chronic disease in later life (Hockaday and Yajnik 2003; Barker 1990). Epidemiological studies have demonstrated strong correlation between low birth weight infants and the development of type II diabetes, cardiovascular disease, and hypertension (Jaquet et al. 2000; Hales et al. 1991; Huxley et al. 2007; Nilsson et al. 1997; Curhan et al. 1996a; Curhan et al. 1996b). It is postulated that the fetus is physiologically ‘programmed’ in utero to adapt to its environment (Barker et al. 2002; Desai and Hales 1997; Bateson et al. 2004). However, this adaptation becomes maladaptive when the infant is exposed to a different postnatal environment.

Experiments of intrauterine growth restriction (IUGR) in animal models provide further evidence to support the hypothesis that impaired growth in utero via various maternal deficiencies leads to impairment of glucose, cholesterol, and triglyceride metabolism in adulthood (Langley et al. 1994; Dahri et al. 1991; Lucas et al. 1996; Sohi et al. 2011a). In utero deficiencies that can lead to impaired growth in humans and animals include hypoxia (Wang et al. 2009), deficiencies in essential vitamins and minerals (Lewis et al. 2001), diminished protein (Sohi et al. 2011a), caloric restriction (Woodall et al. 1996), and excess glucocorticoids (Benediktsson et al. 1993; Reynolds 2010). Although the correlation between impaired fetal growth and the risk for developing chronic disease in adulthood is undoubtedly strong, the mechanisms behind these programming effects are only beginning to be elucidated. Studies have only begun to scratch the surface in understanding the molecular events leading to permanent changes to short- and long-term physiology and pathophysiology. This review aims to look at the current literature to highlight the possible mechanisms

involved with ‘fetal programming.’ These mechanisms of interest include the role of epigenetics, nuclear receptors, reactive oxygen species (ROS) and markers of endoplasmic reticulum stress (ERS). Please see Fig. 1.

Epigenetics

The development of many complex and chronic diseases cannot be simply explained with genomic heritability alone (Manolio et al. 2009). Epigenetics has emerged as an important mechanism in adjusting the expression patterns of genes in a site and tissue specific manner as an adaptive response to insults during the developmental period. Epigenetic mechanisms essentially influence the long-term expression of a gene by altering the ability of the transcriptional machinery to interact with the chromatin environment. Moreover, they influence heritable changes in phenotype without altering the genetic sequence of an organism. Epigenetic changes can be both transient (Barth and Imhof 2010) and persist for long periods of time (Talens et al. 2010). Mechanisms of epigenetic action include DNA methylation, post-translational histone modifications, and more recently discovered microRNA-mediated repression and activation.

DNA methylation

One way the chromatin environment can be altered is due to direct DNA methylation, via the addition of a methyl group to CpG sites on the DNA by members of the DNA methyltransferase family. In addition, the presence of methionine, an essential amino acid, is also essential to DNA methylation as it is the ultimate methyl donor for many methylation reactions. Similarly, folate/folic acid is involved

Fig. 1 Proposed molecular mechanisms of how maternal protein restriction (MPR) in the rat diminishes the expression of hepatic LXR α and LXR-target genes in postnatal life. MPR-induced effects are highlighted in purple

in methionine metabolism and is required for methylation reactions and DNA synthesis. Consequently, altered dietary intake of such nutrients may significantly affect DNA methylation profiles, and ultimately gene expression (Waterland 2006; Kim et al. 1997; Wilson et al. 1984). Traditionally, an increase in methylation across CpG sites can impair initiation, elongation or termination of a gene. However, this may not always hold true as is the case with expression of insulin growth factor 2 (Igf-2), which is increased in IUGR offspring due to intragenic methylation of its promoter (Murrell et al. 2001). Alterations in DNA methylation may be caused by changes in an organism's environment in order to adapt, however in cases of IUGR these DNA methylation responses may be maladaptive due to differing environments in utero and postnatally.

Recently it was demonstrated that during maternal protein restriction (MPR) in mouse pregnancy, expression of the Liver X Receptor ($LXR\alpha$), a nuclear receptor involved in cholesterol homeostasis, is suppressed in the fetal liver as a result of methylation of the upstream region of the $LXR\alpha$ promoter (van Straten et al. 2010). However the effects in postnatal life were not investigated. Another recent study examining the effects of maternal protein restriction and maternal protein excess in porcine offspring found alterations in the expression patterns of a variety of genes involved in DNA methylation and methionine metabolism (Altmann et al. 2012). Beyond the realm of maternal protein restriction, Zhang et al. found that a high fat diet throughout pregnancy and lactation lead to alterations in methyl CpG binding protein-2, a protein involved in the silencing of genes via DNA methylation (Zhang et al. 2009). In addition, a study done by Nijland et al. (2010) that examined maternal nutrient restriction in baboon offspring found decreased methylation of CpG on the promoter of *PCK1* coupled with an increase in *PCK1* transcript in the neonate baboons (Nijland et al. 2010). Overexpression of PEPCCK, the product of *PCK1* translation, has been implicated in hyperglycemia and type II diabetes (Valera et al. 1994; Gomez-Valades et al. 2008). Another study in baboon offspring of maternal nutrient restricted mothers found altered patterns of global DNA methylation in the brain, liver, and kidney between nutrient restricted offspring and offspring whose mothers received adequate food intake (Unterberger et al. 2009).

In humans, a recent double blind randomized control trial examining the effects of micronutrient supplementation during the pre- and periconceptual period found different methylation patterns at differentially methylated regions between the supplement group and the placebo group (Cooper et al. 2012). Another study done in humans by Einstein et al. (2010) indicated hypermethylation of the *HNF4A* gene, a nuclear receptor implicated in type II diabetes (Yamagata et al. 1996), in IUGR infants.

The current evidence suggests that DNA methylation may be a key player in modulating phenotypes in response to the environment. This applies especially to the mother and its newborn in less than ideal environments (*i.e.* situations of famine, poor nutrition or high altitude).

Post-translational histone modifications

The second major epigenetic mechanism involves influencing the chromatin environment via a number of post-translational modifications, including methylation, acetylation, phosphorylation, ubiquitination and ADP-ribosylation of histones (Jenuwein and Allis 2001). The combinatorial nature of these covalent modifications reveal a "histone code", which may serve critical as an adaptive regulatory mechanism that can also influence gene expression in a tissue- and gene-specific manner at times of insult during development. Interestingly, these histone modifications occur and are maintained by a diverse range of histone modifying enzymes including families of histone acetylases and methyltransferases (Marmorstein and Trievel 2009), whose levels may also be altered as a result of a developmental insult. It is important to realize that the different prenatal insults that lead to IUGR offspring seem to have both common and distinct adaptive responses initiated via epigenetic mechanisms. Therefore IUGR offspring derived from different insults may differ or be similar due to global, tissue, or site-directed epigenetic modifications.

To date, there is very little known about the epigenetic alterations associated with expression of target genes in the fetal tissues, and how they are influence these genes throughout normal and abnormal fetal and postnatal development (Burdge et al. 2007a, b; Lillycrop et al. 2005; Rees et al. 2000). Using chromatin immunoprecipitation (ChIP), our laboratory recently demonstrated that MPR-induced IUGR rat male offspring had a decrease in postnatal Cyp7a1 expression, the critical enzyme involved in cholesterol catabolism, both short- and long-term (Sohi et al. 2011a). This was demonstrated to be associated with decreased recruitment of RNA polymerase II, enhanced trimethylation of Histone H3 (Lysine 9), and suppressed acetylation of histone H3 (Lysine 9, 14), all markers of chromatin silencing, within the LXRE region ($-128/-81$ bp) of the *Cyp7a1* promoter. In contrast, MPR female offspring had normal cholesterol, restored levels of Cyp7a1 expression, RNA polymerase II binding, acetylation and trimethylation of Histone H3 (Lysine 9, 14) within the same promoter region (Sohi et al. 2011a).

microRNAs

Aside from post-translational histone modifications, which may govern the long-term expression of genes, microRNAs

(miRNAs) may also play a key role in the fetal programming of cholesterol homeostasis. miRNAs are short, non-coding RNA molecules of 20–25 nucleotides in length that regulate gene expression via mRNA degradation and/or translational repression (Khorram et al. 2010; Xu et al. 2010). By regulating the expression of target genes, miRNAs alter a variety of physiological processes including cell cycle regulation, differentiation, metabolism, and aging (Xu et al. 2010). miRNAs silence gene expression by binding to the 3'-untranslated region (3'-UTR) with partial sequence homology to induce cleavage or repression of productive translation (Brennecke et al. 2005). Given their ability to bind 3'-UTR with partial sequence homology, a single miRNA may have multiple targets in the genome (Brennecke et al. 2005). Conversely, given the nature of miRNA targeting, a single mRNA transcript can theoretically be targeted by several miRNAs (Brennecke et al. 2005). While miRNAs likely play an important role in the etiology of adult diseases and cancer, their role in fetal development and programming remain elusive. Recent microarray studies employing primers for a variety of miRNAs have demonstrated that maternal nutrient restriction can permanently alter the expression of miRNAs in the aortas of newborn and aging rat offspring (Khorram et al. 2010). Moreover, circulating hypoxia-regulated miRNAs have been demonstrated to be increased in pregnant women with fetal growth restriction (Mouillet et al. 2010). The role and identification of miRNAs altering the expression of genes involved in the fetal programming of cholesterol, fatty acids, glucose, and insulin homeostasis remain to be identified, but are the subject of great interest.

Nuclear receptors

Nuclear receptors are part of a family of ligand-mediated transcription factors involved in regulating transcription of genes responsible for growth, development, and differentiation (Mangelsdorf et al. 1995; Repa and Mangelsdorf 2000). Potential ligands for nuclear receptors include hormones, oxysterols, and lipophilic vitamins. There are four main classes of nuclear receptors, classified by their ligand-binding properties, DNA-binding properties, and dimerization properties (Repa and Mangelsdorf 2000). The four classes of nuclear receptors all have a similar mechanism of action through the activation and repression on promoter elements of genes; with slight differences (*i.e.* translocation between nucleus and cytoplasm differs between Class I and Class II nuclear receptors). Humans are thought to have 48 nuclear receptors while rats and mice have 47 and 49, respectively (Zhang et al. 2004). Examples of nuclear receptors include the estrogen receptor (ER), glucocorticoid receptor (GR), progesterone receptor (PR), retinoic acid receptor (RAR), retinoic X receptor (RXR), thyroid

hormone receptor (TR), mineralocorticoid receptor (MR) and the liver X receptor (LXR). While some of these nuclear receptors have been implicated in IUGR and the development of chronic diseases, scientists have only begun to elucidate how nuclear receptors may be involved in the developmental origins of disease on a molecular level.

The liver X receptor

The LXRs (LXR α and LXR β), part of the 1H subfamily of nuclear receptors, have long been implicated in the homeostasis of cholesterol and triglycerides (Janowski et al. 1996; Lehmann et al. 1997). Although, both LXRs share similar homology, they are expressed in different tissue and are differentially regulated in terms of nuclear and cytosolic trafficking (Repa and Mangelsdorf 2000; Prufer and Boudreaux 2007). Furthermore, both LXRs must heterodimerize with the retinoid X receptor (RXR), prior to binding with DNA (Willy et al. 1995). LXR α is mainly expressed in the liver, adipose tissue, spleen, and lungs (Willy et al. 1995; Apfel et al. 1994), while LXR β is expressed ubiquitously (Song et al. 1994). Studies have also found that both isoforms may be involved in different pathways in the regulation of cholesterol and triglycerides (Lund et al. 2006). More recently, it has been found that LXR is also a glucose sensor and involved in the regulation of glucose homeostasis (Mitro et al. 2007b).

Given that LXR α plays a role in cholesterol (Repa and Mangelsdorf 2000), fatty acid (Repa and Mangelsdorf 2000; Lehmann et al. 1997) and glucose (Mitro et al. 2007b) homeostasis, it is an attractive candidate to elucidate the molecular mechanisms underlying the etiology of the metabolic syndrome. To date, few studies have demonstrated links between nutrition, hepatic LXRs and liver function. Both iron restriction and maternal protein restriction have been demonstrated to lead to decreased fetal LXR α (van Straten et al. 2010; Zhang et al. 2005), but less is known about the long-term effects in postnatal life. Our recent studies have demonstrated that in maternal protein restriction (MPR) in rats during pregnancy and lactation, the offspring are low birth weight offspring, with permanent elevation in circulating cholesterol and impaired glucose homeostasis (Sohi et al. 2011a; Vo et al. 2012). We have demonstrated that if MPR rat offspring were placed on a normal diet during lactation, hepatic LXR α was enhanced, preventing decreases in Cyp7a1 and hypercholesterolemia by three weeks of age (Sohi et al. 2011a). Further work on the role of LXR α , RXR, and other lipid-sensing nuclear receptors are warranted to understand the transcriptional mechanisms involved in cholesterol, glucose, and triglyceride homeostasis.

The glucocorticoid receptor

The glucocorticoid receptor is a transcription factor part of the 3C subfamily of nuclear receptors. GR has two isoforms in humans and is expressed in all tissue, with pleiotropic effects throughout the body (Oakley et al. 1996; Lu and Cidlowski 2005). In conjunction with its natural ligand cortisol, GR is vital for the anti-inflammatory response, stress response, metabolism, and development (Barnes 1998; Sapolsky et al. 2000; Rhen and Cidlowski 2005). After ligand binding, GR exerts its actions via two main mechanisms: direct binding to DNA (Mangelsdorf et al. 1995) or through protein-protein interactions with other transcription factors such as NF-kappa B (Ray and Prefontaine 1994).

It has become common practice in medicine to give preterm fetuses glucocorticoids as a therapeutic agent to accelerate lung maturation and prevent fetal respiratory distress syndrome (Liggins and Howie 1972; Young et al. 1980; Knight et al. 1994). Common corticosteroids administered antenatally to prevent fetal respiratory distress syndrome include dexamethasone and betamethasone. However, glucocorticoid exposure during pregnancy has been strongly implicated in fetal programming and the development of chronic disease (Benediktsson et al. 1993; Reynolds 2010; Seckl 2004). Besides direct administration of glucocorticoids to the fetus, maternal undernourishment may also exhibit a similar effect of increased glucocorticoids in the mother and fetus (Blondeau et al. 2001; Lesage et al. 2001; Habib et al. 2011). Previous studies have implicated altered expression of GR in the brain (Levitt et al. 1996) and liver (Nyirenda et al. 1998) in the programming of hypertension and impaired glucose homeostasis in rat offspring when administered dexamethasone antenatally. Similarly, nutrient restriction leads to increased expression of GR in various tissues of rat and sheep offspring (Bertram et al. 2001; Whorwood et al. 2001). In a recent study done by Valtat and colleagues, offspring of dams fed a calorie-restricted diet one week prior to delivery displayed increased corticosterone levels, decreased beta-cell mass, and impaired glucose tolerance (Valtat et al. 2011). However, in offspring where their pancreatic precursor cells lacked GR expression, the deleterious effects were attenuated (Valtat et al. 2011). Thus, GR signaling was observed to play a critical role in the programming of beta-cell dysfunction. Other studies have also demonstrated the important role of corticosteroids on beta-cell development and proliferation (Blondeau et al. 2001; Gesina et al. 2004; Dumortier et al. 2011).

Evidence has emerged supporting the fact that the effects of GR expression on development are likely to be influenced at least partially by epigenetic modifications (Thomassin et al. 2001; Stevens et al. 2010; Begum et al. 2012). In 2005, Lillycrop and colleagues found that maternal protein restriction lead to increased expression of GR, coupled with

decreased methylation of the GR promoter—decreased methylation of the promoter is presumed to lead to increased transcription of the gene (Lillycrop et al. 2005). Another study done by Lillycrop et al. (2007) found that methylation of the GR promoter in maternal protein restricted rat offspring may be due to a reduction in DNA methyltransferase-1 expression, leading to impaired methylation of DNA and histones (Lillycrop et al. 2007).

Interestingly, evidence even suggests these epigenetic modifications may even persist across generations. A study done by Burdge and colleagues (2007a, b) found that methylation of GR in adulthood persisted across the F1 generation into the F2 generation of offspring, likely through stable epigenetic modifications in the female gametes of F0 (Burdge et al. 2007a, b).

Peroxisome proliferator-activated receptors (PPARs)

The peroxisome proliferator-activated receptors (PPAR), part of the 1C subfamily of nuclear receptors, are another group of ligand-activated nuclear receptors that may play a vital role in fetal programming. Originally named for its role in peroxisome proliferation (Issemann and Green 1990), the PPARs are now known to be involved in wide variety physiological functions. Similar to the LXR, PPARs must heterodimerize with RXR prior to DNA binding and transcriptional regulation (Kliwer et al. 1992; Keller et al. 1993).

There are three isoforms of PPAR: PPAR α , PPAR β/δ and PPAR γ , which each have unique expression patterns (Michalik et al. 2006). The main functions of PPAR α are maintaining energy homeostasis (Lefebvre et al. 2006) and modulating inflammatory responses (Chinetti et al. 2000). PPAR β/δ is also involved in maintaining energy homeostasis and is required for the development and maintenance of various tissues including the placenta (Barak et al. 2002; Schaiff et al. 2007), muscle (Luquet et al. 2003; Angione et al. 2011), skin and brain (Peters et al. 2000). Finally, PPAR γ is involved in the differentiation and proper function of adipose tissue (Tontonoz et al. 1994; Imai et al. 2004; Farmer 2006). While there does not seem to be any clear endogenous ligands for the PPARs, it is thought that many different products of fatty acid metabolism are responsible for inducing the actions of these nuclear receptors (Michalik et al. 2006).

The role of PPARs in fetal programming involves a variety of tissues and organs, however altered muscle development and function appears to be a large contributing factor to insulin resistance and the metabolic syndrome based on previous studies. A study done by Bayol and colleagues in 2005 found that when pregnant rats were fed a “cafeteria diet” (high in fat and sugar), the offspring exhibited a 25 % reduction in muscle cross-sectional area concomitant with increased PPAR γ (Bayol et al. 2005). The authors concluded that increased expression of PPAR γ was a compensatory mechanism to preserve insulin sensitivity.

Moreover, another study done by Wang and colleagues in 2004 found that with targeted expression of PPAR β/δ in skeletal muscle, the mice were more resistant to developing obesity (Wang et al. 2004). Similarly, when PPAR β/δ was selectively ablated in skeletal myocytes, the mice were more prone to developing obesity and diabetes (Schuler et al. 2006). This is of great interest considering that hypoxia, associated with IUGR, increases PPAR δ expression in cultured C₂C₁₂ muscle myoblasts (Regnault et al. 2010). A more recent study has found that maternal obesity leads to altered insulin signaling in muscle of sheep offspring along with increased intramuscular triglycerides, however they did not examine the role of PPAR (Yan et al. 2011). Overall, these studies provide strong evidence for the role of PPAR and its interaction with muscle in the development of impaired glucose and fatty acid metabolism.

PPAR expression in adipose tissue may also play a key role in the programming of obesity. Muhlhausler and colleagues (2007) found that offspring of ewes fed a high nutrition diet (155 % versus 100 % control) lead to increased expression of PPAR γ along with other lipogenic and adipogenic factors in perirenal fat (Muhlhausler et al. 2007). Another group found increased PPAR γ expression in adipose tissue coupled with hypertension, and increased adiposity in offspring of mice dams fed an obesogenic diet (Samuelsson et al. 2008). Evidence has also emerged demonstrating that PPAR-mediated programming of visceral obesity in IUGR offspring may be sex-specific (Duffield et al. 2009).

The estrogen receptor (ER α and ER β)

Sex steroid hormones (e.g. estrogen acting through ER α) remain an important mediator in the fetal origins of adult disease. It is well established that the onset and severity of numerous adult onset diseases differ between exist in men and women. For example, men have higher 24-hour mean blood pressure, by approximately 6 to 10 mm Hg, compared to age-matched premenopausal women, but this trend reverses after women reach menopause (Reckelhoff 2001). In females, it is hypothesized that differences in sex hormones modulate regulatory systems leading to decreased hypertension and vascular dysfunction (Reckelhoff 2001). Given the widespread origins of the metabolic syndrome, it is difficult to assess whether gender and, more specifically, sex hormones influence physiological homeostatic mechanisms. Recent studies in animal models have suggested that perturbations to the maternal environment during pregnancy can lead to sex-specific, long-term consequences in postnatal life. For example, male offspring of rat dams exposed to 30 % global nutrient restriction during pregnancy develop hypertension earlier than their female counterparts, whereas severe protein restriction (5 % protein diet) during pregnancy results in programmed hypertension in both sexes (Woods

et al. 2004). Moreover, an MPR diet during gestation and lactation results in pancreatic β cell dysfunction and visceral obesity exclusively in the adult male offspring at postnatal day 130 (Guan et al. 2005; Petrik et al. 1999).

The sex specificity exists even at the epigenetic level in IUGR offspring. For example, uteroplacental insufficiency induced IUGR rats at postnatal day 21 have a global increase in the females and decrease in the males with respect to acetylation of Histone H3 (Lysine 9, 14) at in the hippocampus and white matter (Ke et al. 2006). Our studies have demonstrated that circulating cholesterol in MPR offspring was increased associated with impaired Cyp7a1 expression in both sexes at three weeks (pre-weaning), but persists only in the males at 4 months of age (Sohi et al. 2011a). This was associated with male-specific silencing of the promoter of *Cyp7a1*. While the mechanisms underlying these sex-specific programming effects remain unknown, it has been hypothesized that sex steroids (e.g. estrogen) protect the female against development of these disease processes, including elevated blood pressure (Ozaki et al. 2001). Evidence for this comes from the aromatase knockout (ArKO) mouse, which cannot synthesize endogenous estrogens due to disruption of the *Cyp19* gene (Hewitt et al. 2003). ArKO females challenged with a high cholesterol diet have higher circulating cholesterol and lower Cyp7a1 expression compared to wildtype females and males of either genotype (Hewitt et al. 2003), and estrogen replacement reversed the hepatic steatosis (Hewitt et al. 2004). However, the effects of estrogen may be only part of the reason for the sexual dimorphism observed in MPR offspring. Given that MPR males have reduced levels of circulating testosterone compared to control males (Chamson-Reig et al. 2009), it is possible that loss of this androgen may also underlie male-specific impairment of Cyp7a1 and cholesterol catabolism. In addition, MPR male offspring at 130 days of age have two-fold higher levels of circulating insulin (Chamson-Reig et al. 2009), which inhibits Cyp7a1 transcription in both rat hepatocytes and streptozotocin (STZ)-induced diabetic rats via decreases in the binding of transcription factors FoxO1 and Smad3 to the promoter of *Cyp7a1* (Li et al. 2008). Furthermore, studies have now demonstrated that sex steroid hormones can influence epigenetic mechanisms, including post-translational histone modifications (Leader et al. 2006). Therefore, gender-specific differences in the fetal programming of cholesterol homeostasis may be due in part to alterations in their “histone code”.

Mitochondrial function and the role of reactive oxygen species

Reactive oxygen species (ROS) are free radicals that are byproducts of normal physiological reduction-oxidation reactions carried out by eukaryotic cells. Furthermore, ROS are involved in many cellular and physiological functions

including: proliferation, differentiation, apoptosis, and morphogenesis (Covarrubias et al. 2008). Common ROS include: nitric oxide (NO), superoxide (O_2^-), and hydrogen peroxide (H_2O_2). Oxidative stress occurs when ROS levels exceed certain threshold levels and begin to impair physiological functions. Given that ROS signaling and the maintenance of appropriate ROS levels plays a vital role in a diverse range of cellular processes, especially development, it is likely that ROS may mediate some of the programming effects of IUGR. Some studies have already begun to shed some light on the correlation between IUGR and increased ROS in the fetus (Karowicz-Bilinska et al. 2002; Raab et al. 2009).

One of the main cellular organelles involved in the production and regulation of ROS levels is the mitochondria. Studies have demonstrated that IUGR can lead to increased oxidative stress in rat hepatic mitochondria and impaired hepatic mitochondrial function (Peterside et al. 2003). A similar result was found in the pancreases of IUGR rat offspring (Simmons 2005). Considering pancreatic islet cells have been shown to express much lower antioxidant enzymes compared to other tissues, islet cells are likely to be more prone to ROS-related damage (Lenzen et al. 1996; Tiedge et al. 1997). Consequently, ROS-mediated damage and oxidative stress is thought to be one of the leading contributions to β -cell damage (Robertson et al. 2003). Thus, if IUGR lead to increased ROS in the pancreas, this could lead to impaired development of β -cells and predispose the fetus to impaired glucose tolerance and diabetes in later life (Simmons 2005; Simmons et al. 2005).

Another area of the body in which oxidative stress appears to cause long-term impairment is the cardiovascular system. Previous studies have established links between impaired vascular endothelial function and the presence of ROS (Nakazono et al. 1991). A study done by Franco Mdo and colleagues (2002) found that offspring of nutrient-restricted dams developed hypertension concurrent with increased oxidative stress in mesenteric arterioles (Franco Mdo et al. 2002). In addition to vascular dysfunction, IUGR also seems to induce ROS-mediated damage in the heart (Franco Mdo et al. 2002). von Bergen and colleagues (2009) found that the offspring of ewes administered dexamethasone during pregnancy displayed increased H_2O_2 in cardiac mitochondria as well as increased catalase activity (von Bergen et al. 2009). A more recent study found that rat offspring suffering from prenatal hypoxia developed oxidative stress in the fetal heart by the end of pregnancy (Giussani et al. 2012). The offspring also developed impaired NO-dependent relaxation of peripheral arteries and altered contractility of the heart in adulthood (Giussani et al. 2012). Hence, oxidative stress might not only impair the programming of metabolic pathways in IUGR offspring, but vascular formation and function as well. Concurrent destruction of metabolic pathways and vascular function provide higher risk to the development of diseases such as the metabolic syndrome.

Endoplasmic reticulum stress

Maternal insults such as hypoxia and low nutrition can force developing cells and tissues to reduce protein synthesis both short and long-term. This makes sense considering that ~30 % of total placental oxygen consumption is used for the oxidative process of protein folding (Carter 2000). In placental insufficiency, the associated deprivation of amino acids and oxygen leads to an increase in ROS and ATP resulting in the accumulation of misfolded proteins (Yung et al. 2008). This in turn leads to activation of the endoplasmic reticulum (ER) stress pathway or the uncoupled protein response (UPR) (Yung et al. 2008). In high altitude pregnancies, whereby pO_2 is reduced in association with a fall in birthweight of 100 g/1000 m (Moore et al. 1998; Giussani et al. 2001; Keyes et al. 2003), ER stress and protein synthesis inhibition was activated (Yung et al. 2012). In fetal tissues, decreased maternal dietary protein has also been demonstrated to lead to protein synthesis (increased phosphorylated eIF2 $_{\alpha}$) in the rat liver (Parimi et al. 2004), while paradoxically in IUGR sheep messenger ribonucleic acid translation initiation occurred without increases eIF2 $_{\alpha}$ (Thorn et al. 2009). Bispham and colleagues (2005) found that maternal nutrient restriction led to offspring with more adipose tissue associated with increased expression of uncoupling protein (UCP)-2, a hallmark of ER stress and protein synthesis inhibition (Bispham et al. 2005). While ER stress is associated with short-term adaptations in fetal life to maternal insults, the long-term effects ER stress on overall offspring health is less understood. This is of great interest considering that IUGR is associated with long-term insulin resistance and obesity, and that various components of the ER stress pathway have been linked to insulin resistance. For example, activation of the uncoupled protein response (UPR) in obesity leads to increases in phospho-IRS1 (p-IRS1), a marker of insulin resistance (Ozcan et al. 2004). In addition, a loss of weight leading to increased insulin sensitivity was associated with a reduction in markers of UPR activation (Gregor et al. 2009). In the rat, we and others have previously demonstrated that maternal protein restriction (MPR) in pregnancy and lactation results in impaired fetal growth, decreased liver to bodyweight ratio, insulin resistance, and hypercholesterolemia in adulthood (postnatal day 130) (Sohi et al. 2011a; Chamson-Reig et al. 2009). This is associated with a significant decrease in the hepatic phosphorylation of AKT1 (Serine 473) and increased levels of p85 protein, both indicative of impaired insulin signaling (Sohi et al. 2012). This coincided with elevation of established ER stress markers in the liver, including an increase in X box binding protein 1 (XBP-1) mRNA splicing levels and elevated ER chaperones (Glucose regulated protein 94 and 78) (Sohi et al. 2012). It was also concomitant with attenuated protein synthesis inhibition (increased phosphorylated eIF2 $_{\alpha}$). Interestingly, fetal hepatic GRP94 and 78 protein levels were found

elevated in LP offspring at embryonic day 19, suggesting that ER stress may persist from fetal life into adulthood in low birth weight offspring. Future studies will be aimed at uncovering the underlying molecular mechanisms behind the long-term elevation of ER stress in this and other fetal programming models of insulin resistance and obesity.

Tissue plasticity: reversing the in utero origins of adult disease

The development of many organs occurs both pre- and postnatally. For example, in the liver, development consists of embryonic cell specification, budding, and then differentiation (Kung et al. 2010). Until birth, the liver has a major haematopoietic function (Gualdi et al. 1996), but by mid-gestation in rodents, the liver bud is formed containing bipotential progenitor cells that differentiate into either hepatocytes or ductal cells (Cascio and Zaret 1991). In the last 3 days of gestation in the rat, liver mass triples due to a high rate of fetal hepatocyte proliferation (Greengard et al. 1972), followed by a transition of fetal to adult rat hepatocytes in the first week of postnatal life (Gruppuso et al. 1997). Given that during this neonatal period there is a high rate of replication, neogenesis and apoptosis (Greengard et al. 1972) leading to extensive liver remodeling, this period represents a critical window for therapy designed to improve hepatic growth and function long-term. For example, it has been demonstrated in IUGR rats derived from uterine-ligated dams, that neonatal administration of Exendin-4TM (a GLP-1 analog) prevents the development of diabetes due to a restoration of the transcription factor Pdx-1, and ultimately β cell function (Stoffers et al. 2003). Moreover, Exendin-4TM treatment during this neonatal period also prevented the development of hepatic oxidative stress and insulin resistance (Raab et al. 2009). This indicates quite remarkably that neonatal intervention in rats can influence both pancreatic and liver development long-term. Therefore the goal of future studies is to understand how we can exploit this plasticity in organ development to correct the short- and long-term abnormalities resulting from an adverse in utero environment. While at birth the rat liver is less mature than the human liver (Kung et al. 2010), further insights into the reversibility of fetal programming effects on liver development offers promise in human IUGR pregnancies. Our recent studies indicate that restoration of maternal protein intake during lactation can rescue liver growth and prevent the development of hypercholesterolemia long-term (Sohi et al. 2011b). However the underlying epigenetic and transcriptional mechanisms are unknown. While LXR agonists have been demonstrated to activate acetylation of LXR-target promoters (Talukdar and Hillgartner 2006) and lower LDL cholesterol in atherosclerosis-prone adult mice

(Talukdar and Hillgartner 2006), their use in neonatal life is limited (Fluhr et al. 2005). Given Cyp7a1 expression is enhanced by histone hyperacetylation (Mitro et al. 2007a), it is conceivable that LXR agonists in vivo could boost the expression of LXR target genes, via increases in both LXR binding and histone acetylation surrounding the LXRE sites. Preliminary evidence from our laboratory suggest that three-week old MPR offspring treated with an LXR agonist (GW3695) from postnatal day 5 to 15 had decreased circulating cholesterol:HDL ratios compared to vehicle treated MPR offspring (Sohi et al. 2011b). Moreover, three-week old MPR offspring treated with GW3695 from postnatal day 5 to 15 had increased hepatic expression of LXR α and Cyp7a1, concomitant increased the recruitment of RNA polymerase II and acetylation of histone H3 (lysine 9,14) surrounding the *Cyp7a1* promoter by 3 weeks of age (Sohi et al. 2011b). While the effects of LXR and RXR agonists on cholesterol homeostasis still need to be assessed long-term, preliminary data suggests that LXR and other nuclear receptor agonists may play a promising role in reversing the long-term adverse effects of impaired fetal development on offspring health.

Another interesting paper by Langley-Evans (1997) found that offspring of maternal protein restricted rats were “rescued” from developing hypertension when dams were administered metyrapone, an inhibitor of corticosterone synthesis (Langley-Evans 1997). A sex-specific effect was also seen when a replacement dose of corticosterone was administered with metyrapone. Females responded to the replacement dose and developed hypertension, while males did not exhibit this effect. These results lend evidence to the role of corticosteroids in early programming and the possibility of reversal through pharmacological intervention.

Other studies have indicated that intervention with certain nutrients can reverse the insults to the fetus due to malnutrition. For instance, Jackson et al. (2002) found that when maternal protein restricted dams were supplemented with glycine, offspring hypertension was ameliorated, implicating that the availability of glycine is crucial for adequate development of the cardiovascular system (Jackson et al. 2002). Increased vitamin usage or ROS scavengers such as Vitamin C might also reverse ROS-mediated damage (Giussani et al. 2012).

Given that folic acid/folate is a crucial source of methyl donors for methylation; it may play a role in epigenetic-mediated effects long-term. Supplementation of folic acid following protein restriction appears to alter the growth curve of offspring depending on the fat and protein content of their diet (Burdge et al. 2008). Maternal folic acid/folate supplementation in maternal protein restricted dams also appears to reverse the programmed effects of hypertension in offspring (Torrens et al. 2006). Folic acid supplementation also attenuated altered mRNA expression of various genes in IUGR piglets (Liu et al. 2011). Moreover, evidence suggests that folic acid supplementation may exert its effects

through epigenetic mechanisms, involving reduced methylation of gene promoters such as GR and PPAR α (Lillicrop et al. 2005; Burdge et al. 2008; Lillicrop et al. 2008). A further study done by Burdge et al. (2009), in which folic acid supplementation was given to juvenile-pubertal (post-natal day 56) rats of maternal protein restricted mothers lead to favourable epigenetic changes in the promoters of hepatic GR and PPAR α (Burdge et al. 2009). Further studies are warranted to determine whether intervention in the offspring diet is comparable to dietary intervention in the mother and whether plasticity lasts longer than we have anticipated. For a review of intervention strategies in the rat, please see Fig. 2.

Conclusion

This review presents only a small piece of the puzzle with regards to the molecular mechanisms underlying the fetal programming of chronic adult diseases. Undoubtedly, there exist many other key players involved that we have not discussed and many mechanism that have yet to be elucidated. However, we have come a long way in terms of understanding the hypothesis that Barker postulated decades ago. As we come to understand the molecular mechanisms behind the programming of adult disease, we come closer to not only understanding the development of these diseases but preventing their onset as well.

Elucidating the processes of integral, multifaceted proteins provides significant promise in treating complex,

systemic diseases such as the Metabolic Syndrome. The most promising targets for the investigation of fetal programming in the near future will likely be transcription factors and nuclear receptors involved in multiple pathways. These include LXR α and the PPARs, among others. The role of LXR in cholesterol metabolism, lipid metabolism and glucose metabolism make it an auspicious candidate for linking the three pathways together.

Our recent understanding of epigenetics has shed light on many of the possible mechanisms that may be responsible for the long-term programming of disease. Though traditional genetic inheritance increases the disposition for developing disease, it is ultimately epigenetics and the environment that will determine the outcome of that disease. This provides great hope for therapeutic intervention, as one's future is no longer determined only by what is passed on by their parents. By understanding how our diet, lifestyle, and external environment can affect our DNA methylation profiles, chromatin remodeling, and such, we can learn to take control of our internal environment. This holds true especially for expectant mothers and what they can do for their future children. The evidence underlying the importance of a proper diet during pregnancy has never been so strong. Furthermore, investigation using pharmaceutical therapies provides more support for how the early life environment can be manipulated. Current studies investigating the efficacy of early life intervention are encouraging. For example, work by Pinney et al. (2011) remarkably found that intervention during neonatal life with a transcription factor agonist (*Exendin-4*) lead to permanent changes in later

Fig. 2 Windows of opportunity for therapeutic intervention to prevent IUGR related development of chronic disease in rats and possible mechanisms of action. **a** The period of gestation, beginning at conception is an opportune moment for the prevention and intervention of IUGR. Dietary supplementation and proper nutritional intake are likely to be the most effective forms of prevention and intervention of IUGR during this period. **b** The period right after birth until weaning is a

period of rapid development and high plasticity. At this moment IUGR has already occurred and programming may have taken its toll, however organ plasticity still allows for intervention—especially with pharmaceuticals that modulate transcription factor activity. Dietary intervention in the offspring may reverse the effects of IUGR. Dietary intervention in the mother may also be effective as the offspring still rely on the mother for milk as well

life and ultimately a reversal of the programmed phenotype, likely mediated through epigenetic mechanisms such as histone acetylation and DNA methylation. Currently, *Exendin-4* is already used to treat diabetic patients. However, if we can determine its efficacy and safety during the neonatal period, we can develop therapies to prevent the development of disease in IUGR infants. As well, these studies warrant further investigation with other transcription factor agonists and antagonists.

References

- Altmann S, Murani E, Schwerin M, Metges CC, Wimmers K, Ponsuksili S (2012) Maternal dietary protein restriction and excess affects offspring gene expression and methylation of non-SMC subunits of condensin I in liver and skeletal muscle. *Epigenetics Off J DNA Methylation Soc* 7(3):239–252
- Angione AR, Jiang C, Pan D, Wang YX, Kuang S (2011) PPARdelta regulates satellite cell proliferation and skeletal muscle regeneration. *Skelet muscle* 1(1):33
- Apfel R, Benbrook D, Lernhardt E, Ortiz MA, Salbert G, Pfahl M (1994) A novel orphan receptor specific for a subset of thyroid hormone-responsive elements and its interaction with the retinoid/thyroid hormone receptor subfamily. *Mol Cell Biol* 14(10):7025–7035
- Barak Y, Liao D, He W, Ong ES, Nelson MC, Olefsky JM, Boland R, Evans RM (2002) Effects of peroxisome proliferator-activated receptor delta on placentation, adiposity, and colorectal cancer. *Proc Natl Acad Sci U S A* 99(1):303–308
- Barker DJ (1990) The fetal and infant origins of adult disease. *BMJ (Clin Res Ed)* 301(6761):1111
- Barker DJ, Eriksson JG, Forsen T, Osmond C (2002) Fetal origins of adult disease: strength of effects and biological basis. *Int J Epidemiol* 31(6):1235–1239
- Barnes PJ (1998) Anti-inflammatory actions of glucocorticoids: molecular mechanisms. *Clin Sci (Lond)* 94(6):557–572
- Barth TK, Imhof A (2010) Fast signals and slow marks: the dynamics of histone modifications. *Trends Biochem Sci* 35(11):618–626
- Bateson P, Barker D, Clutton-Brock T, Deb D, D'Udine B, Foley RA, Gluckman P, Godfrey K, Kirkwood T, Lahr MM, McNamara J, Metcalfe NB, Monaghan P, Spencer HG, Sultan SE (2004) Developmental plasticity and human health. *Nature* 430(6998):419–421
- Bayol SA, Simbi BH, Stickland NC (2005) A maternal cafeteria diet during gestation and lactation promotes adiposity and impairs skeletal muscle development and metabolism in rat offspring at weaning. *J Physiol* 567(Pt 3):951–961
- Begum G, Stevens A, Smith EB, Connor K, Challis JR, Bloomfield F, White A (2012) Epigenetic changes in fetal hypothalamic energy regulating pathways are associated with maternal undernutrition and twinning. *The FASEB J* 26(4):1694–1703.
- Benediktsson R, Lindsay RS, Noble J, Seckl JR, Edwards CR (1993) Glucocorticoid exposure in utero: new model for adult hypertension. *Lancet* 341(8841):339–341
- Bertram C, Trower AR, Copin N, Jackson AA, Whorwood CB (2001) The maternal diet during pregnancy programs altered expression of the glucocorticoid receptor and type 2 11beta-hydroxysteroid dehydrogenase: potential molecular mechanisms underlying the programming of hypertension in utero. *Endocrinology* 142(7):2841–2853
- Bispham J, Gardner DS, Gnanalingham MG, Stephenson T, Symonds ME, Budge H (2005) Maternal nutritional programming of fetal adipose tissue development: differential effects on messenger ribonucleic acid abundance for uncoupling proteins and peroxisome proliferator-activated and prolactin receptors. *Endocrinology* 146(9):3943–3949
- Blondeau B, Lesage J, Czernichow P, Dupouy JP, Breant B (2001) Glucocorticoids impair fetal beta-cell development in rats. *Am J Physiol Endocrinol Metab* 281(3):E592–E599
- Brennecke J, Stark A, Russell RB, Cohen SM (2005) Principles of microRNA-target recognition. *PLoS Biol* 3(3):e85
- Burdge GC, Hanson MA, Slater-Jefferies JL, Lillycrop KA (2007a) Epigenetic regulation of transcription: a mechanism for inducing variations in phenotype (fetal programming) by differences in nutrition during early life? *Br J Nutr* 97(6):1036–1046
- Burdge GC, Slater-Jefferies J, Torrens C, Phillips ES, Hanson MA, Lillycrop KA (2007b) Dietary protein restriction of pregnant rats in the F0 generation induces altered methylation of hepatic gene promoters in the adult male offspring in the F1 and F2 generations. *Br J Nutr* 97(3):435–439
- Burdge GC, Lillycrop KA, Jackson AA, Gluckman PD, Hanson MA (2008) The nature of the growth pattern and of the metabolic response to fasting in the rat are dependent upon the dietary protein and folic acid intakes of their pregnant dams and post-weaning fat consumption. *Br J Nutr* 99(3):540–549
- Burdge GC, Lillycrop KA, Phillips ES, Slater-Jefferies JL, Jackson AA, Hanson MA (2009) Folic acid supplementation during the juvenile-pubertal period in rats modifies the phenotype and epigenotype induced by prenatal nutrition. *J Nutr* 139(6):1054–1060
- Carter AM (2000) Placental oxygen consumption. Part I: in vivo studies—a review. *Placenta* 21(Suppl A):S31–S37
- Cascio S, Zaret KS (1991) Hepatocyte differentiation initiates during endodermal-mesenchymal interactions prior to liver formation. *Development* 113(1):217–225
- Chamson-Reig A, Thyssen SM, Hill DJ, Arany E (2009) Exposure of the pregnant rat to low protein diet causes impaired glucose homeostasis in the young adult offspring by different mechanisms in males and females. *Exp Biol Med* (Maywood, NJ) 234(12):1425–1436
- Chinetti G, Fruchart JC, Staels B (2000) Peroxisome proliferator-activated receptors (PPARs): nuclear receptors at the crossroads between lipid metabolism and inflammation. *Inflammation research* 49(10):497–505
- Cooper WN, Khulan B, Owens S, Elks CE, Seidel V, Prentice AM, Belteki G, Ong KK, Affara NA, Constancia M, Dunger DB (2012) DNA methylation profiling at imprinted loci after periconceptional micronutrient supplementation in humans: results of a pilot randomized controlled trial. *The FASEB J* 26(5):1782–90
- Covarrubias L, Hernandez-Garcia D, Schnabel D, Salas-Vidal E, Castro-Obregon S (2008) Function of reactive oxygen species during animal development: passive or active? *Dev Biol* 320(1):1–11
- Curhan GC, Chertow GM, Willett WC, Spiegelman D, Colditz GA, Manson JE, Speizer FE, Stampfer MJ (1996a) Birth weight and adult hypertension and obesity in women. *Circulation* 94(6):1310–1315
- Curhan GC, Willett WC, Rimm EB, Spiegelman D, Ascherio AL, Stampfer MJ (1996b) Birth weight and adult hypertension, diabetes mellitus, and obesity in US men. *Circulation* 94(12):3246–3250
- Dahri S, Snoeck A, Reusens-Billen B, Remacle C, Hoet JJ (1991) Islet function in offspring of mothers on low-protein diet during gestation. *Diabetes* 40(Suppl 2):115–120
- Desai M, Hales CN (1997) Role of fetal and infant growth in programming metabolism in later life. *Biol Rev Camb Philos Soc* 72(2):329–348
- Duffield JA, Vuocolo T, Tellam R, McFarlane JR, Kauter KG, Muhlhausler BS, McMillen IC (2009) Intrauterine growth restriction and the sex specific programming of leptin and peroxisome proliferator-activated

- receptor gamma (PPARGgamma) mRNA expression in visceral fat in the lamb. *Pediatr Res* 66(1):59–65
- Dumortier O, Theys N, Ahn MT, Remacle C, Reusens B (2011) Impairment of rat fetal beta-cell development by maternal exposure to dexamethasone during different time-windows. *PLoS One* 6(10):e25576
- Einstein F, Thompson RF, Bhagat TD, Fazzari MJ, Verma A, Barzilai N, Grealley JM (2010) Cytosine methylation dysregulation in neonates following intrauterine growth restriction. *PLoS one* 5(1):e8887
- Farmer SR (2006) Transcriptional control of adipocyte formation. *Cell Metab* 4(4):263–273
- Fluhr JW, Crumrine D, Mao-Qiang M, Moskowitz DG, Elias PM, Feingold KR (2005) Topical liver x receptor activators accelerate postnatal acidification of stratum corneum and improve function in the neonate. *J Invest Dermatol* 125(6):1206–1214
- Ford ES, Giles WH, Mokdad AH (2004) Increasing prevalence of the metabolic syndrome among u.s. Adults. *Diabetes Care* 27(10):2444–2449
- Franco Mdo C, Dantas AP, Akamine EH, Kawamoto EM, Fortes ZB, Scavone C, Tostes RC, Carvalho MH, Nigro D (2002) Enhanced oxidative stress as a potential mechanism underlying the programming of hypertension in utero. *J Cardiovasc Pharmacol* 40(4):501–509
- Gesina E, Tronche F, Herrera P, Duchene B, Tales W, Czernichow P, Breant B (2004) Dissecting the role of glucocorticoids on pancreas development. *Diabetes* 53(9):2322–2329
- Giussani DA, Phillips PS, Anstee S, Barker DJ (2001) Effects of altitude versus economic status on birth weight and body shape at birth. *Pediatr Res* 49(4):490–494
- Giussani DA, Camm EJ, Niu Y, Richter HG, Blanco CE, Gottschalk R, Blake EZ, Horder KA, Thakor AS, Hansell JA, Kane AD, Wooding FB, Cross CM, Herrera EA (2012) Developmental programming of cardiovascular dysfunction by prenatal hypoxia and oxidative stress. *PLoS One* 7(2):e31017
- Gomez-Valades AG, Mendez-Lucas A, Vidal-Alabro A, Blasco FX, Chillon M, Bartrons R, Bermudez J, Perales JC (2008) Pck1 gene silencing in the liver improves glycemia control, insulin sensitivity, and dyslipidemia in db/db mice. *Diabetes* 57(8):2199–2210
- Greengard O, Federman M, Knox WE (1972) Cytomorphometry of developing rat liver and its application to enzymic differentiation. *J Cell Biol* 52(2):261–272
- Gregor MF, Yang L, Fabbri E, Mohammed BS, Eagon JC, Hotamisligil GS, Klein S (2009) Endoplasmic reticulum stress is reduced in tissues of obese subjects after weight loss. *Diabetes* 58(3):693–700
- Grundy SM (2008) Metabolic syndrome pandemic. *Arterioscler Thromb Vasc Biol* 28(4):629–636
- Gruppiso PA, Awad M, Bienieki TC, Boylan JM, Fernando S, Faris RA (1997) Modulation of mitogen-independent hepatocyte proliferation during the perinatal period in the rat. *In Vitro Cell Dev Biol Anim* 33(7):562–568
- Gualdi R, Bossard P, Zheng M, Hamada Y, Coleman JR, Zaret KS (1996) Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control. *Genes Dev* 10(13):1670–1682
- Guan H, Arany E, van Beek JP, Chamson-Reig A, Thyssen S, Hill DJ, Yang K (2005) Adipose tissue gene expression profiling reveals distinct molecular pathways that define visceral adiposity in offspring of maternal protein-restricted rats. *Am J Physiol Endocrinol Metab* 288(4):E663–E673
- Habib S, Gattineni J, Twombly K, Baum M (2011) Evidence that prenatal programming of hypertension by dietary protein deprivation is mediated by fetal glucocorticoid exposure. *Am J Hypertens* 24(1):96–101
- Hales CN, Barker DJ, Clark PM, Cox LJ, Fall C, Osmond C, Winter PD (1991) Fetal and infant growth and impaired glucose tolerance at age 64. *BMJ (Clin Res Ed)* 303(6809):1019–1022
- Hewitt KN, Boon WC, Murata Y, Jones ME, Simpson ER (2003) The aromatase knockout mouse presents with a sexually dimorphic disruption to cholesterol homeostasis. *Endocrinology* 144(9):3895–3903
- Hewitt KN, Pratis K, Jones ME, Simpson ER (2004) Estrogen replacement reverses the hepatic steatosis phenotype in the male aromatase knockout mouse. *Endocrinology* 145(4):1842–1848
- Hockaday TD, Yajnik CS (2003) –to: Hales CN, Barker DJP (1992) Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* 35:595–601. *Diabetologia* 46(2):303–304
- Huxley R, Owen CG, Whincup PH, Cook DG, Rich-Edwards J, Smith GD, Collins R (2007) Is birth weight a risk factor for ischemic heart disease in later life? *Am J Clin Nutr* 85(5):1244–1250
- Imai T, Takakuwa R, Marchand S, Dentz E, Bornert JM, Messaddeq N, Wendling O, Mark M, Desvergne B, Wahli W, Chambon P, Metzger D (2004) Peroxisome proliferator-activated receptor gamma is required in mature white and brown adipocytes for their survival in the mouse. *Proc Natl Acad Sci U S A* 101(13):4543–4547
- Isseman I, Green S (1990) Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347(6294):645–650
- Jackson AA, Dunn RL, Marchand MC, Langley-Evans SC (2002) Increased systolic blood pressure in rats induced by a maternal low-protein diet is reversed by dietary supplementation with glycine. *Clin Sci (Lond)* 103(6):633–639
- Janowski BA, Willy PJ, Devi TR, Falck JR, Mangelsdorf DJ (1996) An oxysterol signalling pathway mediated by the nuclear receptor LXR alpha. *Nature* 383(6602):728–731
- Jaquet D, Gaboriau A, Czernichow P, Levy-Marchal C (2000) Insulin resistance early in adulthood in subjects born with intrauterine growth retardation. *J Clin Endocrinol Metab* 85(4):1401–1406
- Jenuwein T, Allis CD (2001) Translating the histone code. *Science (New York, NY)* 293(5532):1074–1080
- Karowicz-Bilinska A, Suzin J, Sieroszewski P (2002) Evaluation of oxidative stress indices during treatment in pregnant women with intrauterine growth retardation. *Med Sci Monit Int Med J Exp Clin Res* 8(3):CR211–CR216
- Ke X, Lei Q, James SJ, Kelleher SL, Melnyk S, Jernigan S, Yu X, Wang L, Callaway CW, Gill G, Chan GM, Albertine KH, McKnight RA, Lane RH (2006) Uteroplacental insufficiency affects epigenetic determinants of chromatin structure in brains of neonatal and juvenile IUGR rats. *Physiol Genomics* 25(1):16–28
- Keller H, Dreyer C, Medin J, Mahfoudi A, Ozato K, Wahli W (1993) Fatty acids and retinoids control lipid metabolism through activation of peroxisome proliferator-activated receptor-retinoid X receptor heterodimers. *Proc Natl Acad Sci U S A* 90(6):2160–2164
- Keyes LE, Armaza JF, Niermeyer S, Vargas E, Young DA, Moore LG (2003) Intrauterine growth restriction, preeclampsia, and intrauterine mortality at high altitude in Bolivia. *Pediatr Res* 54(1):20–25
- Khorram O, Han G, Bagherpour R, Magee TR, Desai M, Ross MG, Chaudhri AA, Toloubeydokhti T, Pearce WJ (2010) Effect of maternal undernutrition on vascular expression of micro and messenger RNA in newborn and aging offspring. *Am J Physiol Regul Integr Comp Physiol* 298(5):R1366–R1374
- Kim YI, Pogribny IP, Basnakian AG, Miller JW, Selhub J, James SJ, Mason JB (1997) Folate deficiency in rats induces DNA strand breaks and hypomethylation within the p53 tumor suppressor gene. *The American Journal of Clinical Nutrition* 65(1):46–52
- Kliwer SA, Umesono K, Noonan DJ, Heyman RA, Evans RM (1992) Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature* 358(6389):771–774

- Knight DB, Liggins GC, Wealthall SR (1994) A randomized, controlled trial of antepartum thyrotropin-releasing hormone and betamethasone in the prevention of respiratory disease in preterm infants. *Am J Obstet Gynecol* 171(1):11–16
- Kung JW, Currie IS, Forbes SJ, Ross JA (2010) Liver development, regeneration, and carcinogenesis. *J Biomed Biotechnol* 2010:984248
- Langley SC, Browne RF, Jackson AA (1994) Altered glucose tolerance in rats exposed to maternal low protein diets in utero. *Comp Biochem Physiol Physiol* 109(2):223–229
- Langley-Evans SC (1997) Hypertension induced by foetal exposure to a maternal low-protein diet, in the rat, is prevented by pharmacological blockade of maternal glucocorticoid synthesis. *J Hypertens* 15(5):537–544
- Law MR, Wald NJ, Rudnicka AR (2003) Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis. *BMJ (Clin Res Ed)* 326(7404):1423
- Leader JE, Wang C, Popov VM, Fu M, Pestell RG (2006) Epigenetics and the estrogen receptor. *Ann N Y Acad Sci* 1089: 73–87
- Lefebvre P, Chinetti G, Fruchart JC, Staels B (2006) Sorting out the roles of PPAR alpha in energy metabolism and vascular homeostasis. *J Clin Invest* 116(3):571–580
- Lehmann JM, Kliewer SA, Moore LB, Smith-Oliver TA, Oliver BB, Su JL, Sundseth SS, Winegar DA, Blanchard DE, Spencer TA, Willson TM (1997) Activation of the nuclear receptor LXR by oxysterols defines a new hormone response pathway. *J Biol Chem* 272(6):3137–3140
- Lenzen S, Drinkgern J, Tiedge M (1996) Low antioxidant enzyme gene expression in pancreatic islets compared with various other mouse tissues. *Free Radic Biol Med* 20(3):463–466
- Lesage J, Blondeau B, Grino M, Breant B, Dupouy JP (2001) Maternal undernutrition during late gestation induces fetal overexposure to glucocorticoids and intrauterine growth retardation, and disturbs the hypothalamo-pituitary adrenal axis in the newborn rat. *Endocrinology* 142(5):1692–1702
- Levitt NS, Lindsay RS, Holmes MC, Seckl JR (1996) Dexamethasone in the last week of pregnancy attenuates hippocampal glucocorticoid receptor gene expression and elevates blood pressure in the adult offspring in the rat. *Neuroendocrinology* 64(6):412–418
- Lewis RM, James LA, Zhang J, Byrne CD, Hales CN (2001) Effects of maternal iron restriction in the rat on hypoxia-induced gene expression and fetal metabolite levels. *Br J Nutr* 85(2):193–201
- Li T, Ma H, Chiang JY (2008) TGFbeta1, TNFalpha, and insulin signaling crosstalk in regulation of the rat cholesterol 7alpha-hydroxylase gene expression. *J Lipid Res* 49(9):1981–1989
- Liggins GC, Howie RN (1972) A controlled trial of antepartum glucocorticoid treatment for prevention of the respiratory distress syndrome in premature infants. *Pediatrics* 50(4):515–525
- Lillicrop KA, Phillips ES, Jackson AA, Hanson MA, Burdge GC (2005) Dietary protein restriction of pregnant rats induces and folic acid supplementation prevents epigenetic modification of hepatic gene expression in the offspring. *J Nutr* 135(6):1382–1386
- Lillicrop KA, Slater-Jefferies JL, Hanson MA, Godfrey KM, Jackson AA, Burdge GC (2007) Induction of altered epigenetic regulation of the hepatic glucocorticoid receptor in the offspring of rats fed a protein-restricted diet during pregnancy suggests that reduced DNA methyltransferase-1 expression is involved in impaired DNA methylation and changes in histone modifications. *Br J Nutr* 97(6):1064–1073
- Lillicrop KA, Phillips ES, Torrens C, Hanson MA, Jackson AA, Burdge GC (2008) Feeding pregnant rats a protein-restricted diet persistently alters the methylation of specific cytosines in the hepatic PPAR alpha promoter of the offspring. *Br J Nutr* 100(2):278–282
- Liu JB, Chen DW, Yu B, Mao XB (2011) Effect of maternal folic acid supplementation on hepatic one-carbon unit associated gene expressions in newborn piglets. *Mol Biol Rep* 38(6):3849–3856
- Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ (2006) Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 367(9524):1747–1757
- Lu NZ, Cidlowski JA (2005) Translational regulatory mechanisms generate N-terminal glucocorticoid receptor isoforms with unique transcriptional target genes. *Mol Cell* 18(3):331–342
- Lucas A, Baker BA, Desai M, Hales CN (1996) Nutrition in pregnant or lactating rats programs lipid metabolism in the offspring. *Br J Nutr* 76(4):605–612
- Lund EG, Peterson LB, Adams AD, Lam MH, Burton CA, Chin J, Guo Q, Huang S, Latham M, Lopez JC, Menke JG, Milot DP, Mitnaul LJ, Rex-Rabe SE, Rosa RL, Tian JY, Wright SD, Sparrow CP (2006) Different roles of liver X receptor alpha and beta in lipid metabolism: effects of an alpha-selective and a dual agonist in mice deficient in each subtype. *Biochem Pharmacol* 71(4): 453–463
- Luquet S, Lopez-Soriano J, Holst D, Fredenrich A, Melki J, Rassoulzadegan M, Grimaldi PA (2003) Peroxisome proliferator-activated receptor delta controls muscle development and oxidative capability. *FASEB J Off Publ Fed Am Soc Exp Biol* 17(15):2299–2301
- Mangelsdorf DJ, Thummel C, Beato M, Herrlich P, Schutz G, Umesono K, Blumberg B, Kastner P, Mark M, Chambon P, Evans RM (1995) The nuclear receptor superfamily: the second decade. *Cell* 83(6):835–839
- Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorf LA, Hunter DJ, McCarthy MI, Ramos EM, Cardon LR, Chakravarti A, Cho JH, Guttacher AE, Kong A, Kruglyak L, Mardis E, Rotimi CN, Slatkin M, Valle D, Whittemore AS, Boehnke M, Clark AG, Eichler EE, Gibson G, Haines JL, Mackay TF, McCarroll SA, Visscher PM (2009) Finding the missing heritability of complex diseases. *Nature* 461(7265):747–753
- Marmorstein R, Trievel RC (2009) Histone modifying enzymes: structures, mechanisms, and specificities. *Biochim Biophys Acta* 1789(1):58–68
- McGuire S (2011) Shields M., Carroll M.D., Ogden C.L. Adult Obesity Prevalence in Canada and the United States. NCHS Data Brief no. 56, Hyattsville, MD: National Center for Health Statistics, 2011. *Adv Nutr (Bethesda, MD)* 2(4):368–369
- Michalik L, Auwerx J, Berger JP, Chatterjee VK, Glass CK, Gonzalez FJ, Grimaldi PA, Kadowaki T, Lazar MA, O'Rahilly S, Palmer CN, Plutzky J, Reddy JK, Spiegelman BM, Staels B, Wahli W (2006) International Union of Pharmacology. LXI. Peroxisome proliferator-activated receptors. *Pharmacol Rev* 58(4):726–741
- Mitro N, Godio C, De Fabiani E, Scotti E, Galmozzi A, Gilardi F, Caruso D, Vigil Chacon AB, Crestani M (2007a) Insights in the regulation of cholesterol 7alpha-hydroxylase gene reveal a target for modulating bile acid synthesis. *Hepatology (Baltimore, MD)* 46(3):885–897
- Mitro N, Mak PA, Vargas L, Godio C, Hampton E, Molteni V, Kreusch A, Saez E (2007b) The nuclear receptor LXR is a glucose sensor. *Nature* 445(7124):219–223
- Moore LG, Niermeyer S, Zamudio S (1998) Human adaptation to high altitude: regional and life-cycle perspectives. *Am J Phys Anthropol (Suppl 27)*: 25–64
- Mouillet JF, Chu T, Hubel CA, Nelson DM, Parks WT, Sadovsky Y (2010) The levels of hypoxia-regulated microRNAs in plasma of pregnant women with fetal growth restriction. *Placenta* 31(9):781–784

- Muhlhausler BS, Duffield JA, McMillen IC (2007) Increased maternal nutrition stimulates peroxisome proliferator activated receptor- γ , adiponectin, and leptin messenger ribonucleic acid expression in adipose tissue before birth. *Endocrinology* 148(2):878–885
- Murrell A, Heeson S, Bowden L, Constancia M, Dean W, Kelsey G, Reik W (2001) An intragenic methylated region in the imprinted *Igf2* gene augments transcription. *EMBO Rep* 2(12):1101–1106
- Nakazono K, Watanabe N, Matsuno K, Sasaki J, Sato T, Inoue M (1991) Does superoxide underlie the pathogenesis of hypertension? *Proc Natl Acad Sci U S A* 88(22):10045–10048
- Nestel P, Lyu R, Low LP, Sheu WH, Nitiyanant W, Saito I, Tan CE (2007) Metabolic syndrome: recent prevalence in East and Southeast Asian populations. *Asia Pac J Clin Nutr* 16(2):362–367
- Nijland MJ, Mitsuya K, Li C, Ford S, McDonald TJ, Nathanielsz PW, Cox LA (2010) Epigenetic modification of fetal baboon hepatic phosphoenolpyruvate carboxykinase following exposure to moderately reduced nutrient availability. *J Physiol* 588(Pt 8):1349–1359
- Nilsson PM, Ostergren PO, Nyberg P, Soderstrom M, Allebeck P (1997) Low birth weight is associated with elevated systolic blood pressure in adolescence: a prospective study of a birth cohort of 149378 Swedish boys. *J Hypertens* 15(12 Pt 2):1627–1631
- Nyirenda MJ, Lindsay RS, Kenyon CJ, Burchell A, Seckl JR (1998) Glucocorticoid exposure in late gestation permanently programs rat hepatic phosphoenolpyruvate carboxykinase and glucocorticoid receptor expression and causes glucose intolerance in adult offspring. *J Clin Invest* 101(10):2174–2181
- Oakley RH, Sar M, Cidlowski JA (1996) The human glucocorticoid receptor beta isoform. Expression, biochemical properties, and putative function. *J Biol Chem* 271(16):9550–9559
- Ogden CL, Carroll MD, Kit BK, Flegal KM (2012) Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA* 307(5):483–490
- Ozaki T, Nishina H, Hanson MA, Poston L (2001) Dietary restriction in pregnant rats causes gender-related hypertension and vascular dysfunction in offspring. *J Physiol* 530(Pt 1):141–152
- Ozcan U, Cao Q, Yilmaz E, Lee AH, Iwakoshi NN, Ozdelen E, Tuncman G, Gorgun C, Glimcher LH, Hotamisligil GS (2004) Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. *Science (New York, NY)* 306(5695):457–461
- Parimi PS, Cripe-Mamie C, Kalhan SC (2004) Metabolic responses to protein restriction during pregnancy in rat and translation initiation factors in the mother and fetus. *Pediatr Res* 56(3):423–431
- Peters JM, Lee SS, Li W, Ward JM, Gavrilova O, Everett C, Reitman ML, Hudson LD, Gonzalez FJ (2000) Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(δ). *Mol Cell Biol* 20(14):5119–5128
- Peterside IE, Selak MA, Simmons RA (2003) Impaired oxidative phosphorylation in hepatic mitochondria in growth-retarded rats. *Am J Physiol Endocrinol Metab* 285(6):E1258–E1266
- Petrik J, Reusens B, Arany E, Remacle C, Coelho C, Hoet JJ, Hill DJ (1999) A low protein diet alters the balance of islet cell replication and apoptosis in the fetal and neonatal rat and is associated with a reduced pancreatic expression of insulin-like growth factor-II. *Endocrinology* 140(10):4861–4873
- Pinney SE, Jaecle Santos LJ, Han Y, Stoffers DA, Simmons RA (2011) Exendin-4 increases histone acetylase activity and reverses epigenetic modifications that silence *Pdx1* in the intrauterine growth retarded rat. *Diabetologia* 54(10):2606–2614
- Prufer K, Boudreaux J (2007) Nuclear localization of liver X receptor alpha and beta is differentially regulated. *J Cell Biochem* 100(1):69–85
- Raab EL, Vuguin PM, Stoffers DA, Simmons RA (2009) Neonatal exendin-4 treatment reduces oxidative stress and prevents hepatic insulin resistance in intrauterine growth-retarded rats. *Am J Physiol Regul Integr Comp Physiol* 297(6):R1785–R1794
- Ray A, Prefontaine KE (1994) Physical association and functional antagonism between the p65 subunit of transcription factor NF- κ B and the glucocorticoid receptor. *Proc Natl Acad Sci U S A* 91(2):752–756
- Reckelhoff JF (2001) Gender differences in the regulation of blood pressure. *Hypertension* 37(5):1199–1208
- Rees WD, Hay SM, Brown DS, Antipatis C, Palmer RM (2000) Maternal protein deficiency causes hypermethylation of DNA in the livers of rat fetuses. *J Nutr* 130(7):1821–1826
- Regnault TR, Zhao L, Chiu JS, Gotthel SK, Foran A, Yee SP (2010) Peroxisome Proliferator-Activated Receptor -beta/delta, -gamma Agonists and Resveratrol Modulate Hypoxia Induced Changes in Nuclear Receptor Activators of Muscle Oxidative Metabolism. *PPAR Res* 2010:129173
- Repa JJ, Mangelsdorf DJ (2000) The role of orphan nuclear receptors in the regulation of cholesterol homeostasis. *Annu Rev Cell Dev Biol* 16:459–481
- Reynolds RM (2010) Corticosteroid-mediated programming and the pathogenesis of obesity and diabetes. *J Steroid Biochem Mol Biol* 122(1–3):3–9
- Rhen T, Cidlowski JA (2005) Antiinflammatory action of glucocorticoids—new mechanisms for old drugs. *N Engl J Med* 353(16):1711–1723
- Robertson RP, Harmon J, Tran PO, Tanaka Y, Takahashi H (2003) Glucose toxicity in beta-cells: type 2 diabetes, good radicals gone bad, and the glutathione connection. *Diabetes* 52(3):581–587
- Samuelsson AM, Matthews PA, Argenton M, Christie MR, McConnell JM, Jansen EH, Piersma AH, Ozanne SE, Twinn DF, Remacle C, Rowlerson A, Poston L, Taylor PD (2008) Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: a novel murine model of developmental programming. *Hypertension* 51(2):383–392
- Sapolsky RM, Romero LM, Munck AU (2000) How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 21(1):55–89
- Schaiff WT, Knapp FF Jr, Barak Y, Biron-Shental T, Nelson DM, Sadovsky Y (2007) Ligand-activated peroxisome proliferator activated receptor gamma alters placental morphology and placental fatty acid uptake in mice. *Endocrinology* 148(8):3625–3634
- Schuler M, Ali F, Chambon C, Duteil D, Bornert JM, Tardivel A, Desvergne B, Wahli W, Chambon P, Metzger D (2006) PGC1 α expression is controlled in skeletal muscles by PPARbeta, whose ablation results in fiber-type switching, obesity, and type 2 diabetes. *Cell Metab* 4(5):407–414
- Seckl JR (2004) Prenatal glucocorticoids and long-term programming. *European J Endocrinol/Eur Fed Endocr Soc* 151(Suppl 3):U49–U62
- Simmons R (2005) Developmental origins of adult metabolic disease: concepts and controversies. *Trends Endocrinol Metab* 16(8):390–394
- Simmons RA, Suponitsky-Kroyter I, Selak MA (2005) Progressive accumulation of mitochondrial DNA mutations and decline in mitochondrial function lead to beta-cell failure. *J Biol Chem* 280(31):28785–28791
- Sohi G, Marchand K, Revesz A, Arany E, Hardy DB (2011a) Maternal protein restriction elevates cholesterol in adult rat offspring due to repressive changes in histone modifications at the cholesterol 7 α -hydroxylase promoter. *Mol Endocrinol (Baltimore, MD)* 25(5):785–798
- Sohi G, Revesz A, Arany E, Hardy DB (2011b) The liver X receptor mediates the impaired cholesterol metabolism exhibited in the offspring of maternal protein restricted rats. *Reprod Sci* 18(4):F163

- Sohi G, Revesz A, Hardy DB (2012) Maternal protein restriction leads to elevated hepatic endoplasmic reticulum stress and insulin resistance in the adult rat offspring. *Reprod Sci* 19(3):91A
- Song C, Kokontis JM, Hiipakka RA, Liao S (1994) Ubiquitous receptor: a receptor that modulates gene activation by retinoic acid and thyroid hormone receptors. *Proc Natl Acad Sci U S A* 91(23):10809–10813
- Stevens A, Begum G, Cook A, Connor K, Rumball C, Oliver M, Challis J, Bloomfield F, White A (2010) Epigenetic changes in the hypothalamic proopiomelanocortin and glucocorticoid receptor genes in the ovine fetus after periconceptional undernutrition. *Endocrinology* 151(8):3652–3664
- Stoffers DA, Desai BM, DeLeon DD, Simmons RA (2003) Neonatal extendin-4 prevents the development of diabetes in the intrauterine growth retarded rat. *Diabetes* 52(3):734–740
- Talens RP, Boomsma DI, Tobi EW, Kremer D, Jukema JW, Willemsen G, Putter H, Slagboom PE, Heijmans BT (2010) Variation, patterns, and temporal stability of DNA methylation: considerations for epigenetic epidemiology. *FASEB J Off Publ Fed Am Soc Exp Biol* 24(9):3135–3144
- Talukdar S, Hillgartner FB (2006) The mechanism mediating the activation of acetyl-coenzyme A carboxylase- α gene transcription by the liver X receptor agonist T0-901317. *J Lipid Res* 47(11):2451–2461
- Thomassin H, Flavin M, Espinas ML, Grange T (2001) Glucocorticoid-induced DNA demethylation and gene memory during development. *EMBO J* 20(8):1974–1983
- Thorn SR, Regnault TR, Brown LD, Rozance PJ, Keng J, Roper M, Wilkening RB, Hay WW Jr, Friedman JE (2009) Intrauterine growth restriction increases fetal hepatic gluconeogenic capacity and reduces messenger ribonucleic acid translation initiation and nutrient sensing in fetal liver and skeletal muscle. *Endocrinology* 150(7):3021–3030
- Tiedge M, Lortz S, Drinkgern J, Lenzen S (1997) Relation between antioxidant enzyme gene expression and antioxidative defense status of insulin-producing cells. *Diabetes* 46(11):1733–1742
- Tontonoz P, Hu E, Graves RA, Budavari AI, Spiegelman BM (1994) mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes Dev* 8(10):1224–1234
- Torrens C, Brawley L, Anthony FW, Dance CS, Dunn R, Jackson AA, Poston L, Hanson MA (2006) Folate supplementation during pregnancy improves offspring cardiovascular dysfunction induced by protein restriction. *Hypertension* 47(5):982–987
- Unterberger A, Szyf M, Nathanielsz PW, Cox LA (2009) Organ and gestational age effects of maternal nutrient restriction on global methylation in fetal baboons. *Journal of medical primatology* 38(4):219–227
- Valera A, Pujol A, Pelegrin M, Bosch F (1994) Transgenic mice overexpressing phosphoenolpyruvate carboxykinase develop non-insulin-dependent diabetes mellitus. *Proc Natl Acad Sci U S A* 91(19):9151–9154
- Valtat B, Dupuis C, Zenaty D, Singh-Estivalet A, Tronche F, Breant B, Blondeau B (2011) Genetic evidence of the programming of beta cell mass and function by glucocorticoids in mice. *Diabetologia* 54(2):350–359
- van Straten EM, Bloks VW, Huijckman NC, Baller JF, Meer H, Lutjohann D, Kuipers F, Plosch T (2010) The liver X-receptor gene promoter is hypermethylated in a mouse model of prenatal protein restriction. *Am J Physiol Regul Integr Comp Physiol* 298(2):R275–R282
- Vo T, Sohi G, Revesz A, Hardy DB (2012) Maternal protein restriction results in altered transcriptional and epigenetic regulation of hepatic liver X receptor (LXR) target genes leading to impaired glucose homeostasis in adult rat offspring. *Reprod Sci* 19(3):118A
- von Bergen NH, Koppenhafer SL, Spitz DR, Volk KA, Patel SS, Roghair RD, Lamb FS, Segar JL, Scholz TD (2009) Fetal programming alters reactive oxygen species production in sheep cardiac mitochondria. *Clin Sci (Lond)* 116(8):659–668
- Wang YX, Zhang CL, Yu RT, Cho HK, Nelson MC, Bayuga-Ocampo CR, Ham J, Kang H, Evans RM (2004) Regulation of muscle fiber type and running endurance by PPARdelta. *PLoS Biol* 2(10):e294
- Wang Z, Huang Z, Lu G, Lin L, Ferrari M (2009) Hypoxia during pregnancy in rats leads to early morphological changes of atherosclerosis in adult offspring. *Am J Physiol Heart Circ Physiol* 296(5):H1321–H1328
- Waterland RA (2006) Assessing the effects of high methionine intake on DNA methylation. *The Journal of nutrition* 136(6):1706S–1710S
- Whorwood CB, Firth KM, Budge H, Symonds ME (2001) Maternal undernutrition during early to midgestation programs tissue-specific alterations in the expression of the glucocorticoid receptor, 11beta-hydroxysteroid dehydrogenase isoforms, and type 1 angiotensin ii receptor in neonatal sheep. *Endocrinology* 142(7):2854–2864
- Wilson MJ, Shivapurkar N, Poirier LA (1984) Hypomethylation of hepatic nuclear DNA in rats fed with a carcinogenic methyl-deficient diet. *The Biochemical journal* 218(3):987–990
- Willy PJ, Umesono K, Ong ES, Evans RM, Heyman RA, Mangelsdorf DJ (1995) LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes Dev* 9(9):1033–1045
- Woodall SM, Johnston BM, Breier BH, Gluckman PD (1996) Chronic maternal undernutrition in the rat leads to delayed postnatal growth and elevated blood pressure of offspring. *Pediatr Res* 40(3):438–443
- Woods LL, Weeks DA, Rasch R (2004) Programming of adult blood pressure by maternal protein restriction: role of nephrogenesis. *Kidney Int* 65(4):1339–1348
- WRITING GROUP MEMBERS, Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N, Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A, McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W, Sacco R, Sorlie P, Roger VL, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J, American Heart Association Statistics Committee and Stroke Statistics Subcommittee (2010) Heart disease and stroke statistics—2010 update: a report from the American Heart Association. *Circulation* 121(7):e46–e215
- Xu C, Liu S, Fu H, Li S, Tie Y, Zhu J, Xing R, Jin Y, Sun Z, Zheng X (2010) MicroRNA-193b regulates proliferation, migration and invasion in human hepatocellular carcinoma cells. *Eur J Cancer (Oxford, England: 1990)* 46(15):2828–2836
- Yamagata K, Furuta H, Oda N, Kaisaki PJ, Menzel S, Cox NJ, Fajans SS, Signorini S, Stoffel M, Bell, GI (1996) Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). *Nature* 384(6608):458–460
- Yan X, Huang Y, Zhao JX, Long NM, Uthlaut AB, Zhu MJ, Ford SP, Nathanielsz PW, Du M (2011) Maternal obesity-impaired insulin signaling in sheep and induced lipid accumulation and fibrosis in skeletal muscle of offspring. *Biol Reprod* 85(1):172–178
- Young BK, Klein SA, Katz M, Wilson SJ, Douglas GW (1980) Intravenous dexamethasone for prevention of neonatal respiratory distress: A prospective controlled study. *Am J Obstet Gynecol* 138(2):203–209
- Yung HW, Calabrese S, Hynx D, Hemmings BA, Cetin I, Charnock-Jones DS, Burton GJ (2008) Evidence of placental translation inhibition and endoplasmic reticulum stress in the etiology of human intrauterine growth restriction. *Am J Pathol* 173(2):451–462
- Yung HW, Cox M, Tissot van Patot M, Burton GJ (2012) Evidence of endoplasmic reticulum stress and protein synthesis inhibition in the placenta of non-native women at high altitude. *The FASEB J* 26(5):1970–1981

- Zhang Z, Burch PE, Cooney AJ, Lanz RB, Pereira FA, Wu J, Gibbs RA, Weinstock G, Wheeler DA (2004) Genomic analysis of the nuclear receptor family: new insights into structure, regulation, and evolution from the rat genome. *Genome Res* 14(4):580–590
- Zhang J, Lewis RM, Wang C, Hales N, Byrne CD (2005) Maternal dietary iron restriction modulates hepatic lipid metabolism in the fetuses. *Am J Physiol Regul Integr Comp Physiol* 288(1):R104–R111
- Zhang J, Zhang F, Didelot X, Bruce KD, Cagampang FR, Vatish M, Hanson M, Lehnert H, Ceriello A, Byrne CD (2009) Maternal high fat diet during pregnancy and lactation alters hepatic expression of insulin like growth factor-2 and key microRNAs in the adult offspring. *BMC Genomics* 10:478