Published in final edited form as:

J Med Primatol. 2009 June ; 38(3): 151–155. doi:10.1111/j.1600-0684.2008.00334.x.

Fetal Blood Sampling During the Second and Third Trimester in Baboons

Saju D. Joy, MD^{1,2}, Richard O'Shaughnessy, MD², Natalia Schlabritz-Loutsevitch, MD, PhD^{3,4}, M Michelle Leland, DVM⁴, Patrice Frost, DVM³, and Patty Fan-Havard, PharmD^{2,5}

¹Department of Obstetrics & Gynecology, Wake Forest University School of Medicine

²Department of Obstetrics & Gynecology, The Ohio State University

³Southwest National Primate Research Center at the Southwest Foundation for Biomedical Research

⁴Department of Obstetrics and Gynecology, University of Texas Health Science Center at San Antonio

⁵Department of Pharmacy Practice, University at Buffalo

Abstract

Purpose—The baboons (*Papio cynocephalus*) have similarities with human placentation and fetal development. Fetal blood sampling allows investigators to assess fetal condition at a specific point in gestation as well as transplacental transfer of medications. Unfortunately, assessing fetal status during gestation has been difficult and fetal instrumentation associated with high rate of pregnancy loss. Our objectives are to describe the technique of ultrasound guided cordocentesis (UGC) in baboons, report post-procedural outcomes, and review existing publications.

Research design—This is a procedural paper describing the technique of UGC in baboons. After confirming pregnancy and gestational age via ultrasound, animals participating in approved research protocols that required fetal assessment underwent UGC.

Results—We successfully performed UGC in four animals (5 samples) using this technique. Animals were sampled in the second and third trimesters with fetal blood sampling achieved by sampling a free cord loop, placental cord insertion site or the intrahepatic umbilical vein. All procedures were without complication and these animals delivered at term.

Conclusions—Ultrasound guided fetal umbilical cord venipuncture is a useful and safe technique to sample the fetal circulation with minimal risk to the fetus or mother. We believe this technique could be used for repeated fetal venous blood sampling in the baboons.

Keywords

Fetal blood sampling; Baboon; *Papio cynocephalus*; PUBS; cordocentesis

The primate model continues to be an essential component of clinical research. The baboons (*Papio cynocephalus*) have been of particular interest due to their similarities with human placentation and fetal development. Fetal blood sampling allows the investigator to assess fetal condition at a specific point in gestation as well as transplacental transfer of

Address reprint requests to: Patty Fan-Havard, Pharm.D., Division of Pharmacy Practice and Administration, College of Pharmacy, 500 West 12th Avenue, Columbus, OH 43210-1291, havard.1@osu.edu.
At the time of writing, Dr. Saju D. Joy was Maternal Fetal Medicine Fellow, Department of Obstetrics and Gynecology, The Ohio State University.

medications or markers of interest. A method for fetal sampling in the pregnant macaque using fetoscopy was described by Davidson et al.² in 1978 and by Morris et al.³ in 1980 for baboons. Hankins et al.⁴ described arterial blood sampling at 170 days gestation in pregnant baboons. Since fetal instrumentation in this species is associated with high rate of pregnancy loss², it is essential for pregnancy related studies conducted in the baboon model to be able to develop the techniques of multiple sampling from fetal circulation. In humans, the procedure of cordocentesis was established in 1983⁵ and allows for fetal blood sampling as early as 17 weeks gestation. It is often performed during the 18th–20th weeks of gestation with a complication rate of less than 1% with experienced surgeons. The objectives of this paper are to describe the detailed technique of ultrasound guided cordocentesis (UGC) in baboons at different gestational ages, report the maternal and newborn outcomes post-procedure, and review the existing publications and discuss advantages and disadvantages of each of them.

Material and Methods

Animals studied

This study involved five pregnant baboons (*Papio cynocephalus*) from the colony maintained at the Southwest National Primate Research Center (SNPRC), Southwest Foundation for Biomedical Research (SFBR). All baboons were housed in outdoor metal and concrete gang cages, each containing 10–15 females and one breeding male. Animals were fed Purina Monkey Diet 5038 and 5 LEO (Purina, St. Louis, Missouri, USA) and water *ad libitum*. All animal procedures were performed in accordance with protocols approved by the SFBR Institutional Animal Care and Use Committee (IACUC). Animal characteristics are presented in Table 1.

Study design and blood sampling

Normal baboon gestation lasts 180 days. At 90 days of gestation, (0.5 gestation, or 0.5 G) pregnant baboons maintained in group caging were weighed and underwent a detailed ultrasound examination (GE LOGIQ 5 (GE Healthcare Technologies, Waukesha, WI, USA) ultrasound system,) following tranquilization with ketamine hydrochloride (Ketaset, Fort Dodge, Iowa, USA), 10 mg/kg administered intramuscularly, intubation and inhalation of isoflurane (2 l/min). If necessary, Doppler flow was used to enhance the identification of the cord location and cord insertion sites. The ideal target was the umbilical vein at the placental insertion site where the cord remains anchored. Alternative locations included the umbilical cord near the fetal insertion site, into a free loop of cord, or fetal intrahepatic umbilical vein, however, each of these locations carries an increased risk of trauma secondary to fetal movement.

After identifying an approximate point of needle insertion, the maternal abdomen was prepped and draped to establish a sterile field. Using sterile technique, an ultrasound evaluation was repeated to reassess fetal position and correct for any fetal movements. Throughout the procedure, evaluation for fetal well-being was accomplished with either acoustic monitoring of fetal heart rate or visualization of fetal cardiac activity and/or umbilical blood flow. The greatest amount of preparation was in the ultrasound evaluation to identify an ideal target (i.e. best needle approach with the clearest ultrasound image).

Using ultrasound guidance, a 22 gauge spinal needle was advanced through the maternal abdomen. As the needle approaches the target umbilical vein, the ultrasound image was magnified. This allowed the surgeon to rest the needle tip against the vessel wall and reconfirm midline needle placement. After confirming correct location, a sharp thrust was applied to overcome the resistance of the vascular wall. After confirming intraluminal

location of the needle, the stylette was removed and backflow observed. Images I and II show sampling from a free loop of cord and Image III, a site for intrahepatic sampling.

Fetal blood samples were aspirated. If free fetal blood was not obtained, successful venipuncture may be obtained by needle withdrawal in very small increments with gentle suction and needle rotation. Umbilical vein identity was confirmed by infusing 0.3 mL of normal saline solution with clear ultrasound visualization of venous turbulence. We prefer to approach a length of vein in sagittal section with needle advancement from the superior to inferior direction. This permits visualization of the needle throughout its course. Hematologic measurements require pure fetal blood obtained under free-flowing conditions. Therefore, heparin flush of the syringe was used to prevent clotting in the tube of the fetal blood delivered immediately for evaluation.

After obtaining the sample, the needle was removed at the same angle of insertion. Transient bleeding from the needle puncture site is a common finding often lasting less than one minute. This bleeding is visible with all transamniotic procedures and is always monitored to cessation. After completing the procedure, fetal well-being was confirmed with cardiotocography and an ultrasound evaluation to reconfirm hemostasis.

Results

Maternal and fetal baboon outcomes were evaluated in five baboons. Serial fetal blood collection by cordocentesis was successfully obtained during second (GD 120) and third (GD 171) trimesters in one baboon. An additional three baboons had UGC performed during the second or third trimester. There were no postsurgical, maternal or fetal complications observed in these animals. Unsuccessful fetal blood collection occurred in only one baboon and this was due to fetal position (spine located anteriorly) and placental location (posterior placenta obstructed by the fetus). This fetus had no post-procedural complications and was delivered at term. However, this infant died six days following delivery, suspected to be due to pneumonia.

Low amniotic fluid volume was noted in third trimester baboons, which may result in a more difficult route of sampling by cordocentesis. However, the intrahepatic vein provides an alternative site for fetal blood collection especially if low amniotic fluid or fetal orientation limits umbilical vein access. Fetal blood sampling was successfully obtained from the hepatic vein in a separate third trimester dam (GD 156). No maternal and fetal complications were observed and the dam delivered a healthy pup at term. The deliveries for these two baboons occurred at term via cesarean delivery.

Discussion

Morris et al. used a 27 gauge needle for fetal sampling in midtrimester pregnant baboons.³ We prefer the 22 gauge needle at all gestational ages. We are concerned that a smaller gauge needle is more difficult to guide, can easily occlude when suction is applied to the syringe and may be associated with more hemolysis.^{3, 6} Table 2 lists the umbilical cord diameters during human^{7, 8} and baboon gestation. Although the umbilical cord diameter increases in size during the baboons second trimester, it is known to be generally 4–5 mm at 91–98 days gestation. Additionally, the subhuman primate uterus, in contrast to the human organ, is a very thin, ballotable, fluid-filled sac with walls which tend to invaginate rather than be perforated.³

If there is excessive fetal movement compromising access to the ideal cord site or placing increased risk to the fetus, paralyzing the fetus for a short interval is an option. Intravenous or intramuscular vecuronium (0.1mg/kg) is an available and well-accepted technique.⁹ In

this study, all animals received general anesthesia and the fetus remained immobile for the UGC. The clinician must verify that the blood sample is fetal in origin. This is most commonly achieved by comparing the mean corpuscular volume (MCV) of the retrieved cells with that of maternal blood. Fetal red blood cells are considerably larger than maternal cells. Additionally, Anti-I studies could be performed on the aspirated blood to confirm fetal sampling.

The estimated fetal loss rate with cordocentesis is approximately one percent per procedure. This fetal loss rate varies from one to 6.7 percent depending on the operator experience and the differences in patient selection at any particular center.⁸ Complications can include bradycardia (52%), fetal bleeding (40–60%), where as, cord hematoma, infection, maternal alloimmunization, etc. are less frequent.

Other studies have sampled the umbilical artery⁴ as well as introducing a catheter into the fetal vessel for repetitive samplings.^{3, 11, 12} Some of these catheter trials were complicated by excessive fetal bleeding and fetal loss when the catheter was pulled out of the vessel.³ Another approach was via fetoscopy where by, a variety of procedures could be carried out under direct visualization including fetal blood sampling, laser ablation of placental communicating vessels, etc.⁴ The obvious disadvantage was in entering the intrauterine cavity with a larger diameter instrument (trocar and fetoscopic sleeve). However, an advantage of fetoscopy over ultrasound guided cordocentesis was the direct visualization of placental and umbilical cord vessels. Thus, using guides attached to the fetoscope, directed fetal blood sampling was easily accomplished. After completing the procedure, confirmation of hemostasis was visualized.

Although other studies have shown different techniques for fetal blood sampling in the baboon, the procedure we present has proven successful in human subjects. Although, fetoscopy is gaining popularity with the success of placental vessel laser ablation in the setting of twin-twin transfusion syndrome, there remain few centers and experienced operators to successfully complete the procedure. Finally, two reasons that an umbilical venipuncture is preferred to arterial sampling is that 1) the vein represents a measure of placental transfer without fetal metabolism and 2) the artery is more prone to spasm and reflex bradycardia. The risk for fetal loss with this procedure is inversely proportional to user experience.

Conclusion

The procedure we describe for fetal sampling in baboons is similar to that in humans. We successfully performed UGC in four animals (5 samples) using this technique. Three of the animals were in the second trimester and fetal blood sampling achieved by sampling a free cord loop or placental cord insertion site. One of the animals was in the third trimester, and due to fetal position and the observation of decreased amniotic fluid volume, we proceeded with sampling the intrahepatic umbilical vein. All procedures were without complication and these animals delivered at term. Ultrasound guided fetal umbilical cord venipuncture is a useful tool to sample the fetal circulation with minimal risk to the fetus or mother. We feel that this technique could be used for repeated fetal venous blood sampling in the baboons.

Acknowledgments

This study was funded by the pilot study grant mechanisms from Southwest National Primate Center (NIH/NCRR GRANT P51 RR013986) to Dr. P. Havard and to Dr. N. Schlambitz-Loutsevitch. Authors are thankful for the financial support of Department of Obstetrics and Gynecology University of Texas Health Science Center at San Antonio.

References

1. Fazleabas AT. A baboon model for simulating pregnancy. *Methods Mol Med*. 2006; 121:101–110. [PubMed: 16251738]
2. Davidson, Jr, EC; Morris, JA.; O'Grady, JP.; Hendrickx, AG.; Anderson, J.; Kaback, M.; Frazer, R. Sampling the fetoplacental circulation. III. Combined laparoscopy-fetoscopy in the pregnant macaque for hemoglobin identification. *Am J Obstet Gynecol*. 1978 Dec 15; 132(8):833–844. [PubMed: 153713]
3. Morris JA, Davidson EC, Makabali G, O'Grady JP, Anderson J, Hendrickx AG. Sampling the fetoplacental circulation. Preliminary experience in the pregnant baboon (*papio cynocephalus*). *Am J Obstet Gynecol*. 1980 Feb; 136(4):531–536. [PubMed: 6153507]
4. Hankins GD, Lowery CL Jr, Scott RT, Morrow WR, Carey KD, Leland MM, Colvin EV. Transplacental transfer of zidovudine in the near-term pregnant baboon. *Am J Obstet Gynecol*. 1990 Sep; 163(3):728–732. [PubMed: 2403153]
5. Daffos F, Capella-Pavlovsky M, Forestier F. A new procedure for fetal blood sampling in utero: preliminary results of fifty-three cases. *Am J Obstet Gynecol*. 1983 Aug 15; 146(8):985–987. [PubMed: 6881235]
6. Kennedy C, Angermuller S, King R, Noviello S, Walker J, Warden J, Vang S. A comparison of hemolysis rates using intravenous catheters versus venipuncture tubes for obtaining blood samples. *J Emerg Nurs*. 1996; 22(6):566–569. [PubMed: 9060320]
7. Weissman A, Jakobi P, Bronshtein M, Goldstein I. Sonographic measurements of the umbilical cord and vessels during normal pregnancies. *J Ultrasound Med*. 1994; 13:11–14. [PubMed: 7636947]
8. Togni FA, Araujo JA, Moron AF, et al. Reference intervals for the cross sectional area of the umbilical cord during gestation. *J. Perinat. Med*. 2007; 35:130–134. [PubMed: 17302517]
9. Mouw RJC, Klumper F, Hermans J, et al. Effect of atracurium or pancuronium on the anemic fetus during and directly after intravascular intrauterine transfusion: a double blind randomized study. *Acta Obstet Gynecol Scand*. 1999; 78:763–766.
10. Van Camp IL, Klumper FJ, Oepkes D, et al. Complications of intrauterine intravascular transfusion for fetal anemia due to maternal red-cell alloimmunization. *Am J Obstet Gynecol*. 2005; 192(1): 171–177. [PubMed: 15672021]
11. Stark RI, Daniel SS, James LS, MacCarter G, Morishima HO, Niemann WH, Rey H, Tropper PJ, Yeh MN. Chronic instrumentation and longterm investigation in the fetal and maternal baboon: tether system, conditioning procedures and surgical techniques. *Lab Anim Sci*. 1989 Jan; 39(1): 25–32. [PubMed: 2918681]
12. Garland M, Szeto HH, Daniel SS, Tropper PJ, Myers MM, Stark RI. Placental transfer and fetal metabolism of zidovudine in the baboon. *Pediatr Res*. 1998 Jul; 44(1):47–53. [PubMed: 9667370]

Image I.
Free loop of umbilical cord.

Image II.
Free loop of umbilical cord.

Image III.
Intrahepatic site for sampling.

Table 1

Characteristics of the animals on this study, G- gestation

Animal number	Number of cordocentesis procedures performed	Gestational age at procedure	Age (years)	Weight (kg)	Number of the previous offsprings	Number of previous c. sections	Outcome
#1	2	0.67 & 0.75 G	15	21	10	3	Delivered healthy infant
#2	1	0.87 G	17	20	10	3	Delivered healthy infant
#3	1	0.69 G	7	14	0	0	Delivered healthy infant at term
#4	1	0.66 G	11	19	3	1	Delivered healthy infant at term
#5	1- Attempt unsuccessful	0.66 G	11	18	4	1	Infant was born at term and died at 6 days of age from pneumonia

Table 2

Comparative Umbilical Cord Characteristics Between Humans and Baboons

Human GD	Baboon GD Please check!!!	Human Umbilical Cord Diameter		Human Umbilical Vein Diameter		Baboon Umbilical Cord Diameter#
		Mean (cm)	SD (cm)	Mean (cm)	SD (cm)	
140-152	91 - 98 (0.51 - 0.54 G)	0.44	NR ¹	NR	NR	0.4 - 0.5
153-162	98 - 105 (0.54 - 0.58 G)	0.61	0.1	0.2	NR	--
174-184	112 - 119 (0.62 - 0.66 G)	0.7	0.1	0.24	0.06	--
196-207	126 - 133 (0.70 - 0.74 G)	1.01	0.14	0.36	0.06	--
218-230	140 - 147 (0.78 - 0.82 G)	1.19	0.16	0.41	0.07	--
241-249	154 - 161 (0.86 - 0.89G)	1.29	0.12	0.47	0.07	--
260-272	168 - 175 (0.93 - 0.97 G)	1.39	0.08	0.54	0.06	1.0

Values for the human data is from Weissman et al.⁷ More recent human data provides a regression function for the average of the cord sectional area by gestational age (GA): **Cordaverage = -673.128 + 51.279*GA - 0.732*GA²**; and the standard deviation (SD): **CordSD = 4.24 + 1.28*GA**. (Togni et al)⁸

GD = Gestational days; SD = Standard deviation