Endoscopic Endonasal Management of Recurrent Petrous Apex Cholesterol Granuloma

Nancy McLaughlin¹ Daniel F. Kelly¹ Daniel M. Prevedello² Kiarash Shahlaie¹ Ricardo L. Carrau³
Amin B. Kassam⁴

¹Brain Tumor Center, John Wayne Cancer Institute of Saint John's Health Center, Santa Monica, California, United States

²Department of Neurological Surgery, The Ohio State University, Columbus, Ohio, United States

³Department of Head and Neck Surgery, The Ohio State University, Columbus, Ohio, United States

⁴Department of Surgery, Division of Neurosurgery, University of Ottawa, Ontario, Canada

Address for correspondence and reprint requests Amin B. Kassam, M.D., F.R.C.S.C., Neurosurgery Division, The Ottawa Hospital - Civic Campus, 1053 Carling Avenue, Ottawa, Canada K1Y 4E9 (e-mail: kassamab@gmail.com).

J Neurol Surg B 2012;73:190–196.

Abstract

Keywords

- ▶ cholesterol granuloma
- ▶ petrous apex
- ▶ endoscopy
- ▶ transsphenoidal approach
- ▶ expanded endonasal approach

Introduction Petrous apex cholesterol granulomas (PACGs) are uncommon lesions. Recurrence following transcranial or endonasal approaches to aerate the cyst occurs in up to 60% of cases. We describe the technical nuances pertinent to the endonasal endoscopic management of a recurrent symptomatic PACG and review the literature.

Results A 19-year-old woman presented with a recurrent abducens nerve paresis. Four months prior, she underwent an endonasal transsphenoidal surgery (TSS) for drainage of a symptomatic PACG. Current imaging documented recurrence of the right PACG. Transsphenoidal and infrapetrous approaches were performed to obtain a wider bony opening along the petrous apex and drain the cyst. A Doyle splint was inserted into the cyst's cavity and extended out into the sphenoid sinus, maintaining patency during the healing process. Three months after surgery, the splint was removed endoscopically, allowing visualization of a patent cylindrical communication between both aerated cavities. The patient remains symptom-free and recurrence-free.

Conclusion Endoscopic endonasal surgery must be adapted to manage a recurrent PACG. A TSS may not be sufficient. An infrapetrous approach with wider bony opening, extensive removal of the cyst's anterior wall, and the use of a stent are indicated for the treatment of recurrent PACG and to prevent recurrences.

Introduction

Cholesterol granulomas (CGs) are inflammatory lesions that occur rarely in the petrous apex.¹ Initially, these lesions tend to remain asymptomatic. As they expand, headaches and cranial neuropathies may arise.^{1,2} The treatment of symptomatic petrous apex CG (PACG) is surgical drainage of the cyst and permanent aeration of petrous apex air cells to prevent recurrence.^{1,3} Several transcranial approaches have been used to treat these lesions, the choice of which depends on the patient's preoperative hearing status and the lesion's

location in relation to critical neurovascular structures.^{2–5} These traditional lateral skull base approaches expose to potential structural damage to the inner ear and facial nerve as well as added morbidities of any intracranial procedure.⁴ They also have the shortcoming of often producing a relatively narrow bony opening, potentially compromising long-term drainage, and preventing the use of stents between the aerated cavity and the sphenoid sinus. In addition to having recurrence rates as high as 60%, repeat transcranial surgeries for recurrent lesions may be associated with significant

received

August 7, 2010

accepted

November 15, 2010

published online

May 17, 2012

Copyright © 2012 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 584-4662.

DOI <http://dx.doi.org/10.1055/s-0032-1312706>.
ISSN 2193-6331.

Figure 1 Preoperative (top row) and postoperative (bottom row) magnetic resonance imaging in T2-weighted image axial (A, D), T1-weighted image sagittal (B, E), and T1-weighted image postgadolinium coronal (C, F) views.

morbidities.^{2,3} Recently, the endoscopic endonasal approach has gained popularity in the neurosurgical community. This less invasive approach is attractive for the initial surgical management of symptomatic PACG and possibly even more so for symptomatic recurrent PACG.

In this study, we describe a case of recurrent symptomatic PACG treated by an expanded endonasal approach. We review all reported cases treated by an endoscopic endonasal approach and attempt to establish management pearls regarding the endoscopic endonasal management of PACG.

Case Report and Technique Description

A 19-year-old woman presented with an ongoing severe headache of subacute onset and 3-day history of diplopia. Physical examination was unremarkable except for right abducens nerve palsy. Head computed tomography scan demonstrated expansion of the right petrous apex. Magnetic resonance imaging showed a right petrous apex mass measuring $25 \times 14 \times 14$ mm abutting the posterior wall of the sphenoid sinus. The lesion was hyperintense on both T1- and T2-weighted images and did not enhance following gadolinium administration (**Fig. 1**). Given her rapid onset of symptoms, surgical drainage was recommended.

The patient underwent an endonasal transsphenoidal approach using intermittently the operating microscope and the endoscope for visualization. The location of the lesion

was confirmed in the lower inferior right aspect of the sphenoid sinus with intraoperative neuronavigation. After creating a wide bony opening to the face of the CG, its capsule was incised. Characteristic greenish fluid with cholesterol crystals poured forth. After satisfactory drainage, the bony opening was widened and multiple fragments of the capsule were removed. Final inspection of the cyst's cavity showed complete drainage of the lesion. Postoperatively, symptoms progressively resolved.

Four months after initial surgery, clinicoradiological recurrence occurred and mandated a repeat transsphenoidal and infrapetrous approaches (**Fig. 2**). After removing scar tissue from the prior bony opening in the sphenoid floor, care was taken to identify the vertical and horizontal segments of the internal carotid artery (ICA). The high speed drill was used to widen the bony opening inferior to the petrous carotid into the petrous apex in a medial and inferior direction. Once exposed on a large surface, the anterior wall of the cyst was removed, the content of the cyst was drained, and its lining partly removed. A 45-degree endoscope allowed visualization of the empty cavity (**Fig. 3**). A Doyle splint was inserted into the cyst's cavity on ~ 1 cm and extended out into the sphenoid sinus, maintaining patency during the healing process. At 3 months after surgery, the Doyle splint was removed under endoscopic visualization. A healed cylindrical communication between the involved petrous apex cells and the sphenoid sinus was appreciated (**Fig. 4**). A rubber

Figure 2 Magnetic resonance imaging at clinical recurrence in T2-weighted image axial (A), T1-weighted image sagittal (B), and T1-weighted image postgadolinium coronal (C) views.

Figure 3 Intraoperative view of the repeat transsphenoidal and intrapetrous approaches. (A) Exposure of the petrous apex and cyst on a large surface. (B) Widening of the bony opening inferior to the petrous carotid into the petrous apex in a medial and inferior direction. (C) Visualization of the emptied cavity with a 45-degree endoscope. (D) Insertion of a Doyle splint into the cyst's cavity.

Figure 4 Intraoperative view of the removal of the Doyle splint with irrigation of the cavity.

Table 1 Summary of Reported Cases of Petrous Apex Cholesterol Granuloma Treated Via the Endonasal Endoscopic Approach

Author	Age, Sex	Preoperative Symptoms	Location of CG in Relation to SS	Symptoms on FU (Length of FU, in Months)	Procedure	Complication	Symptomatic Recurrence
Griffith and Terrell (1996) ³	34, M	Dysequilibrium	Protruding into the posterior SS, bony erosion	Asymptomatic (18)	TS drainage of cyst; wide opening of the cyst cavity	Transient epistaxis	None
	24, M	Hearing loss, V3 hypesthesia	Protruding into SS	Asymptomatic after third OR (12)	First: Subtemporal transzygomatic approach Second: TS drainage of cyst Third: TS drainage of cyst and wide opening of the cyst cavity	None	Recurrence 2 months after first OR Recurrence 2 months FU after second OR No recurrence after third OR
Michaelson et al (2001) ⁸	13, F	H/A, sixth nerve palsy	Abutting the posterior SS	Asymptomatic (6)	TS drainage of cyst; wide opening of the cyst cavity; marsupialization	None	None
DiNardo et al (2003) ⁹	62, F	Dysequilibrium	Abutting the posterior SS, separated by a thin layer of bone	Asymptomatic (12)	TS drainage of cyst; wide opening of the cyst cavity	None	None
Presutti et al (2006) ¹⁰	38, M	H/A, sixth nerve palsy, increasing vertigo	Abutting the posterolateral wall of the SS	Asymptomatic (3)	TS drainage of cyst; wide opening of the cyst cavity; placement of T-shaped stent	None	None
Oyama et al (2007) ¹¹	28, F	Hemifacial pain	Protruding in the posterior SS, separated by a thin layer of bone	Asymptomatic (24)	TS drainage of cyst; wide opening of the cyst cavity	None	None
Georgalas et al (2008) ⁷	30, M	Facial palsy, sensorineural hearing loss, vertigo	Abutting the posterolateral SS	Asymptomatic (120)	TS drainage of cyst; wide opening of the cyst cavity; marsupialization	None	None
	13, M	H/A, sixth nerve palsy	Sphenopetrous lesion, almost completely obliterating SS	Asymptomatic (72)	First: Middle fossa approach Second: TS drainage of cyst; wide opening of the cyst cavity; marsupialization	None	Recurrence after first OR (time to recurrence not specified) No recurrence after second OR
	44, F	H/A	Protruding into the posterior SS	Asymptomatic (24)	TS drainage of cyst; wide opening of the cyst cavity; marsupialization	None	None

(Continued)

Table 1 (Continued)

Author	Age, Sex	Preoperative Symptoms	Location of CG in Relation to SS	Symptoms on FU (Length of FU, in Months)	Procedure	Complication	Symptomatic Recurrence
Samadian et al (2009) ¹²	28, F	H/A, sixth nerve palsy, hearing loss	Protruding in the posterior SS, separated by a thin layer of bone	Asymptomatic (48)	TS drainage of cyst; wide opening of the cyst cavity; marsupialization; placement of a silicone tube drain	None	None
Zanation et al (2009) ¹³	8 patients ^a	All had H/A Two had occasional vertigo	No detail	No detail	TS: 2 TS with carotid lateralization: 3 IP approach: 3 Drainage of cyst and stenting	None	One recurrence at 2.5 years after first OR
Present case	19, F	H/A, sixth nerve palsy	Abutting the posterior SS	Asymptomatic to date	First: TS drainage of cyst; wide opening of the cyst cavity; partial marsupialization Second: TS and IP for drainage of cyst; very wide opening of the cyst cavity; marsupialization; placement of a Doyle splint	After first OR: delayed epistaxis	Recurrence 4 months after first OR No recurrence yet after second OR

CG, cholesterol granuloma; FU, follow-up; H/A, headache; IP, infrapetrous; OR, operation; SS, sphenoid sinus; TS, transsphenoidal.

^aSpecific demographic data could not be extracted from the original paper.

catheter was used to irrigate the aerated cavity. The patient currently remains symptom-free and recurrence-free.

Discussion

CGs rarely occur in the petrous apex because the petrous apex is pneumatized in only 30% of temporal bones.⁶ They result from obstruction of the normal aeration of petrous apex air cells. This obstruction is believed to lead to negative pressure within the air cells, eventually causing mucosal edema, tissue breakdown, and hemorrhage and with accumulation of breakdown products such as cholesterol.^{1,2} The cholesterol crystals induce a granulomatous inflammatory reaction resulting in the typical CG.

For symptomatic patients, surgery is presently the only treatment option available. Surgical treatment implies cyst drainage, with or without removal of its lining, and restoration of air cell aeration. Some authors have also proposed complete excision of the lesion followed by obliteration of the cavity with vascularized tissue.² Numerous transcranial approaches have been performed to treat PACG such as middle cranial fossa approaches and lateral skull base approaches including the translabyrinthine, infratrochlear, infralabyrinthine, transcochlear approaches.^{2–5} The specific approach recommended depended on the patient's hearing status and the cyst's location. Given the lateral trajectory of transcranial skull base approaches, they inevitably expose to potential structural damage to the inner ear and facial nerve as well as added morbidities of any intracranial procedure.^{4,7}

In addition, although satisfactory cyst drainage may be accomplished through these transcranial routes, bony opening obtained is often relatively narrow, rendering difficult intraoperative splinting techniques and potentially putting at risk chronic drainage. Indeed, recurrences have been reported to be as high as 60% irrespective of the used transcranial approach and degree of completeness of removal.^{2,3} Stenosis of the opening used to exteriorize the cyst may result in reaccumulation of its content and recurrence of symptoms. Just as for initial procedures, revision transcranial surgeries for recurrent lesions may also be associated with significant morbidities.

Contemporary progresses in endoscopic surgery as well as in image-guidance technology have allowed treatment of PACG through a less invasive route. Petrous lesions located medial to the ICA and that abut, protrude, or invade the sphenoid sinus may be safely approached through a transsphenoidal approach^{3,7–13} (►Table 1). The use of the endoscope as visualization technique has enabled a more complete drainage of the cyst.¹⁴ The cyst cavity may be closely inspected under endoscopic visualization to remove septations between separate fluid loculations or any remote debris.¹⁴

Including the present case, 19 cases of PACG have been treated through an endoscopic endonasal approach (►Table 1).^{3,7–13} Of the 19 patients, 16 have been drained initially through a medial transsphenoidal approach (with or without lateralization of the ICA). Of these 16 patients, 3 developed a symptomatic recurrence that required repeat surgery. The remaining three patients had been drained

initially via an infrapetrous approach, and did not develop a recurrence.¹³ Accordingly, 3 of the 19 patients (15.7%) with PACG treated via an endonasal route presented a recurrence at 2, 4, and 30 months postoperative. This recurrence rate (15.7%) is significantly lower than that reported following other transcranial approaches (up to 60%). Several factors may contribute to this lower recurrence rate including a wider bony opening and, consequently, a more extensive removal of the cyst's anterior or medial wall, afforded by the endonasal endoscopic approaches (especially with the infrapetrous approach). In addition, the superior visualization of the cavity using angled endoscopes enables to assess and address remaining septations and/or debris; therefore, optimizing the drainage.¹⁴ Furthermore, the use of a stent to preserve a drainage pathway throughout the healing process may contribute to reducing recurrences. This is especially important with the infrapetrous approach since the path of dissection is deeper than that performed in the medial transsphenoidal approach, and therefore may be more at risk of scarring.¹³ All of these considerations are of utmost importance when treating a recurrent PACG. We favor the endonasal endoscopic infrapetrous approach for recurrent PACG regardless of the initial surgical approach. In comparison to a transcranial surgery which may be associated with significant morbidity, no complications have yet occurred following endonasal endoscopic treatment for recurrent PACG.

Conclusion

Overall, the endoscopic endonasal route enables treatment of PACG through an anterior trajectory, preserving hearing, vestibular, and facial function. This approach is significantly less invasive than traditional transcranial procedures with overall reduced morbidity, shortened procedure time, reduced hospital stay, and better cosmetic and lower recurrence rates. It should be considered as the preferred surgical alternative for initial treatment of symptomatic PACG but also for the treatment of symptomatic recurrence, independently of the initial surgical route. Repeat endoscopic endonasal surgery must be adapted to recurrent PACG. The infrapetrous approach with wider bony opening, extensive anterior cyst wall removal, and the use of a stent should be the treatment of choice for recurrent PACG.

References

- 1 Sincoff EH, Liu JK, Matsen L, et al. A novel treatment approach to cholesterol granulomas. Technical note. *J Neurosurg* 2007;107(2):446–450
- 2 Eisenberg MB, Haddad G, Al-Mefty O. Petrous apex cholesterol granulomas: evolution and management. *J Neurosurg* 1997;86(5):822–829
- 3 Griffith AJ, Terrell JE. Transsphenoid endoscopic management of petrous apex cholesterol granuloma. *Otolaryngol Head Neck Surg* 1996;114(1):91–94
- 4 Brackmann DE, Toh EH. Surgical management of petrous apex cholesterol granulomas. *Otol Neurotol* 2002;23(4):529–533

- 5 Terao T, Onoue H, Hashimoto T, Ishibashi T, Kogure T, Abe T. Cholesterol granuloma in the petrous apex: case report and review. *Acta Neurochir (Wien)* 2001;143(9):947–952
- 6 Thedinger BA, Nadol JB Jr, Montgomery WW, Thedinger BS, Greenberg JJ. Radiographic diagnosis, surgical treatment, and long-term follow-up of cholesterol granulomas of the petrous apex. *Laryngoscope* 1989;99(9):896–907
- 7 Georgalas C, Kania R, Guichard JP, Sauvaget E, Tran Ba Huy P, Herman P. Endoscopic transsphenoidal surgery for cholesterol granulomas involving the petrous apex. *Clin Otolaryngol* 2008;33(1):38–42
- 8 Michaelson PG, Cable BB, Mair EA. Image-guided transphenoidal drainage of a cholesterol granuloma of the petrous apex in a child. *Int J Pediatr Otorhinolaryngol* 2001;57(2):165–169
- 9 DiNardo LJ, Pippin GW, Sismanis A. Image-guided endoscopic transsphenoidal drainage of select petrous apex cholesterol granulomas. *Otol Neurotol* 2003;24(6):939–941
- 10 Presutti L, Villari D, Marchioni D. Petrous apex cholesterol granuloma: transsphenoid endoscopic approach. *J Laryngol Otol* 2006;120(6):e20
- 11 Oyama K, Ikezono T, Tahara S, Shindo S, Kitamura T, Teramoto A. Petrous apex cholesterol granuloma treated via the endoscopic transsphenoidal approach. *Acta Neurochir (Wien)* 2007;149(3):299–302, discussion 302
- 12 Samadian M, Vazirnezami M, Moqaddasi H, Rakhshan M, Khormaee F, Ashraf H. Endoscopic transrostral-transsphenoidal approach to petrous apex cholesterol granuloma: case report. *Turk Neurosurg* 2009;19(1):106–111
- 13 Zanation AM, Snyderman CH, Carrau RL, Gardner PA, Prevedello DM, Kassam AB. Endoscopic endonasal surgery for petrous apex lesions. *Laryngoscope* 2009;119(1):19–25
- 14 Mattox DE. Endoscopy-assisted surgery of the petrous apex. *Otolaryngol Head Neck Surg* 2004;130(2):229–241