Survival in Sinonasal Melanoma: A Meta-analysis

Mitchell R. Gore, M.D., Ph.D.¹ Adam M. Zanation, M.D.¹

¹ Division of Skull Base Surgery/Rhinology, Department of Otolaryngology–Head and Neck Surgery, UNC-Chapel Hill School of Medicine, Chapel Hill, North Carolina

J Neurol Surg B 2012;73:157–162.

Address for correspondence and reprint requests Adam M. Zanation, M.D., Assistant Professor, Division of Skull Base Surgery/Rhinology, Department of Otolaryngology/Head and Neck Surgery, CB#7070, UNC-Chapel Hill School of Medicine, Chapel Hill, NC 27599-7070 (e-mail: adam_zanation@med.unc.edu).

Abstract

Sinonasal melanoma is an uncommon tumor which carries a poor prognosis and high rates of local and regional recurrence and distant metastasis. While surgical resection is the mainstay of treatment, the utility of multimodality therapy has not been well studied or established. We sought to better evaluate the optimal treatment modality for sinonasal melanoma. We reviewed 39 case reports involving 423 patients with sinonasal melanoma and present a meta-analysis comparing survival by treatment modality. The two-tailed *p*-value for survival by treatment modality was determined. The number of primary site/local, regional, and distant recurrences was determined where data was available. There was a nonsignificant increase in survival for patients treated with surgery + radiotherapy versus surgery alone. There was a statistically significant increase in survival for surgery + chemotherapy versus chemotherapy alone and versus surgery alone. Patients treated with combined surgery, radiation, and chemotherapy had a statistically shorter survival interval than patients treated with surgery + chemotherapy, which may reflect more advanced disease in patients treated with triple therapy. There was no statistically significant increase in survival found for the addition of radiation to surgery. This meta-analysis demonstrates that multimodality therapy, particularly the addition of chemo-or immunotherapy to surgery, may increase survival in a subset of patients. Radiation therapy did not appear to increase survival. There may be a significant increase in overall survival with combined modality therapy with surgery and chemo/immunotherapy versus single modality therapy. Level of evidence: III. Grade of recommendation: C.

Keywords

- ▶ sinonasal
- ▶ melanoma
- ▶ skull base
- ▶ endoscopic

Mucosal melanoma of the head and neck accounts for only 25 to 30% of total melanomas with sinonasal melanoma accounting for less than 5% of these. Sinonasal melanoma carries a relatively poor prognosis, owing to the tendency of sinonasal melanoma to present at an advanced stage. A standardized treatment regimen has been difficult to develop, owing to the low incidence of the disease. Typically sinonasal melanoma has been thought to be a radiation-resistant disease, and thus surgical resection with clear margins has been the key component of therapy with curative intent, while chemotherapy and immunotherapy have

typically been reserved for unresectable or metastatic disease. Recent literature has suggested, however, that there may be a 50 to 75% initial response rate when radiotherapy alone is used.¹ It has not been shown definitively that there is a survival advantage with the addition of radio- or chemotherapy to surgery. One of our primary aims was to determine whether there is a statistically significant increase in survival with combined therapy versus single modality therapy. In addition, we sought to characterize the most frequent primary site location, as well as the frequency of metastasis to the neck.

received

July 9, 2011

accepted

October 18, 2011

published online

February 22, 2012

Copyright © 2012 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 584-4662.

DOI <http://dx.doi.org/10.1055/s-0032-1301400>.
ISSN 2193-6331.

Table 1 Average Survival (Months) by Treatment Modality

Modality	n	Average Survival (Months)	Standard Deviation
Surgery	53	24.15	19.78
Surgery + Radiotherapy	91	30.12	29.27
Surgery + Radiotherapy + Chemotherapy	26	23.45	24.56
Radiotherapy	15	18.73	17.66
Chemotherapy	3	12.7	5.8
Surgery + Chemotherapy	15	38.8	17.1
Chemotherapy + Radiotherapy	3	38.7	40.3

Methods

Average survival was calculated as the average of the longest survival reported for each patient in each treatment group. Chemotherapy and immunotherapy were grouped together in the chemotherapy group. Two-sided *p*-values for survival were calculated using the Student's *t*-test with values <0.05 considered significant.

Results

Thirty-nine reviews, case series, and case reports were studied totaling 423 patients with sinonasal melanoma.²⁻⁴⁰ Of the 206 patients for whom treatment and survival data was provided, 53 patients were treated with surgery alone, while 91 were treated with surgery and radiotherapy and 26 patients were treated with surgery along with chemoradiation. Fifteen patients were treated with radiotherapy alone whereas 3 patients were treated with chemotherapy alone, 15 patients were treated with surgery plus chemotherapy, and 3 patients were treated with chemoradiation. ►**Table 1** shows average survival by treatment modality, whereas ►**Table 2** shows *p*-values for comparisons between

treatment modalities. The average survival was 24.15 months for surgery alone, 30.12 months for surgery and radiotherapy, and 23.45 months for surgery and chemoradiation. The two-tailed *p*-value was not significant for any of these differences in survival. The average survival for radiotherapy, chemotherapy, surgery + chemotherapy, and chemoradiotherapy was 18.73, 12.67, 38.83, and 38.67 months respectively. Of these, the increases in survival interval between surgery + chemotherapy and both chemotherapy alone and surgery alone were statistically significant, with *p*-values of 0.0205 and 0.0112, respectively. There was also a statistically significant decrease in survival for patients treated with surgery, radiation, and chemotherapy versus patients treated with surgery and chemotherapy, with a *p*-value of 0.0387.

Of the 423 patients, there were 17 patients specifically reported to have metastases to the neck, but 36 reported regional metastases. Two hundred and six recurrences were reported, for a recurrence rate of 54.6%. The most common primary site was the nasal cavity/nasopharynx with 165 cases. Unspecified sinus was next most frequent with 54 cases, followed by turbinate unspecified with 34 and the nasal septum with 33 cases.

Table 2 Two-Tailed *p*-Value by Treatment Modality

Modality	Two-Tailed <i>p</i> -Value for Overall Survival
Surgery survival vs. surgery + radiotherapy survival	0.1895
Surgery survival vs. triple therapy survival	0.8925
Surgery + radiotherapy survival vs. triple therapy survival	0.2921
Surgery survival vs. radiotherapy survival	0.3421
Surgery + radiotherapy survival vs. radiotherapy survival	0.1474
Triple therapy survival vs. radiotherapy survival	0.5182
Chemotherapy + radiotherapy survival vs. chemotherapy alone survival	0.3305
Surgery + chemotherapy survival vs. chemotherapy alone survival	0.0205 ^a
Surgery survival vs. chemotherapy survival	0.3242
Triple therapy survival vs. chemotherapy survival	0.4615
Surgery + chemotherapy survival vs. surgery alone survival	0.0112 ^a
Triple therapy survival vs. surgery + chemotherapy survival	0.0387 ^a
Chemotherapy + surgery vs. chemotherapy + radiation survival	0.9907

^aStatistically significant.

Discussion

The question of whether to add radiation and chemotherapy to surgery in treatment of nasal melanoma has been a difficult question to answer. Given the low incidence of melanoma of the sinonasal cavity it is difficult to assemble randomized control trials or even large case-control studies. Sinonasal melanoma has been thought to be a radioresistant disease, with surgery with negative margins being the primary modality for attempted cure. We sought to elucidate whether combined therapy increased survival in primary sinonasal melanoma over single modality therapy. We performed a meta-analysis of 423 patients with histologically confirmed sinonasal melanoma. The majority of patients (135/206) for whom data on treatment modality was provided were treated with some form of combined modality therapy. The majority were treated with surgery + radiation or surgery alone. No significant difference was seen between surgery or combined surgery + radiation. Interestingly, survival was actually decreased for patients treated with surgery, radiation, and chemotherapy, although this difference was nonsignificant versus surgery alone or surgery + radiation, but was significant versus surgery + chemotherapy (23.45 months vs. 38.83 months, $p = 0.0387$). This may reflect a higher stage of disease in patients treated with triple therapy. This hypothesis is difficult to confirm as only two studies reported the specific stage for patients in these two treatment groups, with three patients in both the triple therapy and surgery + chemotherapy group noted to have metastatic disease and no specific data on stage available for the remainder of the patients in these two treatment groups. As might be expected, the shortest survival was found for chemotherapy alone at 12.7 months. This difference was not significant for chemotherapy alone versus surgery alone or radiation alone. There was, however, a statistically significant ($p = 0.0205$) increase in survival for surgical treatment + chemotherapy (38.82 ± 17.05 months) versus chemotherapy alone (12.7 ± 5.8 months). There was also a statistically significant increase in survival for surgery + chemotherapy versus surgery alone (24.15 ± 19.78 , p -value = 0.0112). Other than the statistically significant increase in survival for surgery + chemotherapy versus chemotherapy alone and surgery alone, and the decreased survival in patients treated with triple modality therapy versus surgery + chemotherapy, no significant increase in survival was seen for any other single- or multimodality therapy. This is consistent with previous data showing that radiotherapy adds little to overall survival in sinonasal melanoma.

The vast majority of primary sinonasal melanomas occurred in the nasal cavity or nasopharynx. The most common sinus primaries were maxillary and ethmoid, with primary sphenoid sinonasal melanoma being quite rare, with only five cases seen. Average survival for all modalities was 27.41 ± 24.32 months. This number is consistent with other studies showing an average survival of ~ 2 years.⁴¹

Distant metastases were by far the most common site of recurrence (90, 44%), with locoregional (45, 22%), regional (36, 17%), and local (35, 17%) recurrence much less prevalent

than distant metastases. The rate of regional metastasis in this meta-analysis is consistent with the 5 to 25% range found in most series. Given this relatively low rate in sinonasal melanoma, elective neck dissection has not been widely advocated, in contrast to oral cavity mucosal melanoma, which carries a much higher rate of nodal metastasis.^{10,42}

This seems to indicate that even with good local or regional control, sinonasal melanoma is difficult to control as distant spread is likely to occur even in the presence of good locoregional control. This is consistent with other reports^{41,43} indicating that even after extensive surgical resection, local control has been reported to be below 50%, and that distant and regional metastases are relatively common, occurring in 25% or more patients in most series.^{41,43} This may at least partially explain the lack of survival benefit with combined modality therapy versus single modality therapy. The propensity for recurrence is so high that significant improvements in local or locoregional control with combined therapy are likely lost due to the high frequency of recurrence, particular of distant recurrence/metastasis.

Ninety percent of the patients in this meta-analysis (185/206 patients) for whom treatment data was reported received some form of surgical therapy. Our data and previous reports indicate that while surgery is the primary modality for treatment of sinonasal melanoma, survival remains poor even in patients with gross local control, likely secondary to both the high rate of distant metastases (44% in our data) along with the high rate of microscopic disease.⁴³ Patients with positive surgical margins have a greatly increased risk of dying from mucosal melanoma, and lack of local control has been shown to increase the rate of distant metastases.⁴⁴ The poor survival in sinonasal melanoma even with aggressive surgical management in the vast majority of patients may result from the presence of microscopic disease as well as occult distant metastases.

Fifteen of 206 patients (7%) were treated with radiotherapy alone, while 91 of 206 (44%) were treated with adjuvant radiotherapy with surgery, and 26/206 (13%) with triple therapy and 3/206 (0.01%) with chemoradiation. Adjuvant radiotherapy has become widely used in sinonasal melanoma, despite the lack of clear evidence of a survival benefit. Temam et al⁴⁵ found that while adjuvant radiotherapy produced a statistically significant increase in local control in head and neck melanoma, the rate of distant metastasis and disease-free survival was not affected in a statistically significant fashion. Krengli et al⁴⁶ and Owens et al⁴⁷ found similar results, with radiation improving local and locoregional control, but not affecting overall survival. We similarly found no statistically significant increase in overall survival for the addition of radiation to any other modality, or as primary therapy. Sufficient data was not available on our meta-analysis population to examine the effect of either adjuvant or primary radiation on local or locoregional control. Owens et al felt that the lack of improvement in overall survival with either primary radiation or adjuvant radiotherapy was due to the high rate of metastatic disease in these series. This seems to be a likely explanation, as in our meta-analysis we noted that the majority of recurrences (90) were distant

recurrences, which is consistent with the high rate of metastasis seen in these prior studies.

In our meta-analysis 3 of 206 patients (0.01%) were treated with chemotherapy or immunotherapy alone, 3 of 206 (0.01%) with chemotherapy with radiotherapy, 15/206 with surgery + chemotherapy (0.07%), and 26/206 (0.12%) with triple therapy. We observed a statistically significant increase in survival for surgery + chemotherapy versus both chemotherapy alone and surgery alone. We also noted a statistically significant decrease in survival for patients treated with triple therapy versus surgery + chemotherapy only. Due to the lack of specific data given on the immunotherapy or chemotherapy regimen given on each patient, we grouped chemotherapy and bio- or immunotherapy together. There is no standardized systemic therapy for sinonasal melanoma, and many patients are treated with a chemotherapeutic agent in conjunction with a biologic or immunotherapeutic agent such as Bacillus Calmette-Guérin, interferon- α , or interleukin-2. Ives et al examined the effectiveness of chemotherapy versus immunotherapy in patients with metastatic cutaneous melanoma.⁴⁸ They specifically examined the effect of adding interferon- α (IFN) $\pm$ interleukin-2 (IL-2) to chemotherapy in patients with metastatic melanoma. Their meta-analysis of trials of biochemotherapy versus chemotherapy evaluated rates of partial response, complete response and overall (partial + complete) response; response duration; progression-free survival, overall survival, and toxicity. A subgroup analysis was performed by type of immunotherapy, according to whether IFN only or IFN and IL-2 were administered in the biochemotherapy arm. They found a clear benefit for biochemotherapy for partial, complete, and overall response for both the IFN and IFN + IL-2 groups. They found no benefit in overall survival for biochemotherapy versus chemotherapy. There was a significant heterogeneity in the types of chemotherapy utilized. Most trials used a single-agent chemotherapy regimen, DTIC, temozolomide, or vindesine. One trial used combination aranoza and cisplatin. The trials of chemotherapy $\pm$ IFN and IL-2 used combination DTIC and cisplatin or a triple chemotherapy regimen of DTIC and cisplatin combined with carmustine, vinblastine, or vindesine. The studies included in our meta-analysis did not provide consistent data on the specific regimen used.

Data on chemotherapy and biochemotherapy in sinonasal melanoma is limited. Bartell et al⁴⁹ reported on 15 patients who received various biochemotherapy regimens for advanced head and neck mucosal melanoma. After a median follow-up of 13 months, 3 patients (20%) had partial response, and 4 patients (27%) had complete response. The median time to disease progression for all 15 patients was 10 months and the median overall survival duration for all patients was 22 months. They concluded that biochemotherapy for advanced head and neck mucosal melanoma should be considered as a systemic treatment option both as adjuvant therapy for metastatic disease as well as neoadjuvant therapy for patients with aggressive or extensive local disease.

The decrease in survival for triple therapy versus surgery + chemotherapy in our data may well be due to a higher stage in the triple therapy group. It is difficult to definitively answer

this due to the limited patient-specific data available in the manuscripts analyzed. The statistically significant increase in overall survival for surgery + chemo- or biochemotherapy versus surgery alone and chemotherapy alone is, to our knowledge, the first demonstrated increase in overall survival in sinonasal melanoma for combined therapy. It seems likely that this is due to response to chemo- and biochemotherapy of metastatic disease, as several studies have demonstrated the lack of improvement in overall survival with improvement in local control. Again, the limited patient-specific data provided makes this difficult to establish, but it seems clear that systemic therapy improves overall survival while the addition of radiotherapy does not significantly improve overall survival, but may improve local control.

There are several weaknesses of this study. First, the retrospective data makes selection bias a possibility. However, it is extremely difficult to assemble single- or multicenter prospective trials for uncommon tumors such as sinonasal melanoma. A randomized prospective multicenter trial for sinonasal melanoma would potentially allow better standardization of treatment regimens as well as more accurate comparison of local control, disease-free survival, and overall survival between treatment modalities. The varying lengths of follow-up is another issue, however most studies followed patients for several years, and in many cases the case series included patients over the course of a decade. Given the relatively constant average survival for sinonasal melanoma, $\sim$ 2 years, we believe this gives fairly accurate longitudinal follow-up in the data included in our meta-analysis. Another weakness is the inclusion of chemotherapy with biochemotherapy and immunotherapy as a single group. A more specific subgroup analysis by specific treatment regimen would be ideal, however the limited number of patients with sinonasal melanoma and the lack of individual data on each patient makes this difficult. As other large series such as the Ives meta-analysis⁴⁸ grouped different regimens together and found significant differences which were maintained whether the chemo- and biochemotherapy groups were analyzed separately or as subgroups, we believe this does not significantly affect the overall improvement on survival found in this meta-analysis.

Conclusion

To our knowledge, this is the largest meta-analysis of patients with primary sinonasal melanoma. The data demonstrates that there is no survival advantage for combined radiotherapy + surgery or chemoradiotherapy + surgery versus surgery alone. We did find a significant overall survival advantage for surgery + chemotherapy versus surgery alone and versus chemotherapy alone. The average overall survival was 27.41 months for all therapies, confirming past studies pointing to a fairly dismal prognosis for this rare malignancy. We therefore recommend aggressive treatment primarily with surgery for resectable disease with the goal of achieving local control, with the caveat that a significant number of patients will have distant recurrence (44% in our dataset) which is not significantly impacted by adjuvant treatment such as

radiotherapy. It appears that improvements in overall survival must result from improvements in response of metastatic disease and distant recurrences with chemotherapy, immunotherapy, or biochemotherapy. We recommend aggressive surgical resection as the primary modality, combined with radiotherapy as necessary for local control. We recommend chemotherapy and biochemotherapy as both adjuvant treatment for distant disease as well as neoadjuvant chemotherapy as this appears to improve overall survival. In conclusion, this meta-analysis is, to our knowledge, the largest meta-analysis of patients with mucosal melanoma of the nasal cavity/paranasal sinuses. It confirms earlier retrospective studies demonstrating that sinonasal melanoma has a poor prognosis, with survival on the order of 24 months. It also demonstrates that the addition of radiotherapy to surgical resection does not significantly improve overall survival, but that overall survival is improved with combined surgery + chemotherapy.

Note

Presented at the 20th Annual North American Skull Base Society Meeting, New Orleans, LA, October 14th to 18th, 2009.

References

- Dwivedi R, Dwivedi R, Kazi R, Kumar S, Agarwal SP. Mucosal melanoma of nasal cavity and paranasal sinus. *J Cancer Res Ther* 2008;4(4):200–202
- Kuo CT, Hoon DS, Takeuchi H, et al. Prediction of disease outcome in melanoma patients by molecular analysis of paraffin-embedded sentinel lymph nodes. *J Clin Oncol* 2003; 21(19, Issue 19):3566–3572
- Wagner M, Morris CG, Werning JW, Mendenhall WM. Mucosal melanoma of the head and neck. *Am J Clin Oncol* 2008;31(1):43–48
- Choi YS, Han GS, Choi JS, Jang TY. Two cases of malignant melanoma in turbinate and maxillary sinus. *Eur Arch Otorhinolaryngol* 2006;263(11):996–1000
- Narasimhan K, Kucuk O, Lin H-S, et al. Sinonasal mucosal melanoma: a 13-year experience at a single institution. *J Neurol Surg B* 2009;19(4):255–262
- Cheng YF, Lai CC, Ho CY, Shu CH, Lin CZ. Toward a better understanding of sinonasal mucosal melanoma: clinical review of 23 cases. *J Chin Med Assoc* 2007; 70(1, Issue 1):24–29
- Thompson LD, Wieneke JA, Miettinen M. Sinonasal tract and nasopharyngeal melanomas: a clinicopathologic study of 115 cases with a proposed staging system. *Am J Surg Pathol* 2003;27(5):594–611
- Kardon DE, Thompson LD. Sinonasal mucosal malignant melanoma: report of an unusual case mimicking schwannoma. *Ann Diagn Pathol* 2000;4(5):303–307
- Dauer EH, Lewis JE, Rohlinger AL, Weaver AL, Olsen KD. Sinonasal melanoma: a clinicopathologic review of 61 cases. *Otolaryngol Head Neck Surg* 2008;138(3):347–352
- Moreno MA, Hanna EY. Management of mucosal melanomas of the head and neck: did we make any progress? *Curr Opin Otolaryngol Head Neck Surg* 2010;18(2):101–106
- Slavik T, Hannah M, Schroder RA. Primary chondroid melanoma of the nasal skin: a rare melanoma variant at a previously undocumented site. *J Cutan Pathol* 2007;34(5):427–430
- McLean N, Tighiouart M, Muller S. Primary mucosal melanoma of the head and neck. Comparison of clinical presentation and histopathologic features of oral and sinonasal melanoma. *Oral Oncol* 2008;44(11):1039–1046
- Racic G, Kurtovic D, Roje Z, Tomic S, Dogas Z. Primary mucosal melanoma of the eustachian tube. *Eur Arch Otorhinolaryngol* 2004;261(3):139–142
- McLean N, Tighiouart M, Muller S. Primary mucosal melanoma of the head and neck. Comparison of clinical presentation and histopathologic features of oral and sinonasal melanoma. *Oral Oncol* 2008;44(11):1039–1046
- Dauer EH, Lewis JE, Rohlinger AL, Weaver AL, Olsen KD. Sinonasal melanoma: a clinicopathologic review of 61 cases. *Otolaryngol Head Neck Surg* 2008;138(3):347–352
- Rinaldo A, Shaha AR, Patel SG, Ferlito A. Primary mucosal melanoma of the nasal cavity and paranasal sinuses. *Acta Otolaryngol* 2001;121(8):979–982
- Lynch SC, Lee AG, Graham SM, Kirby PA. Primary melanoma of the sphenoid sinus presenting with a third cranial nerve palsy. *J Neuroophthalmol* 2005;25(4):289–292
- Carter TR. Pathologic quiz case 1. Malignant melanoma. *Arch Otolaryngol Head Neck Surg* 1986;112(4):450–453
- Shinbori T, Uyama E, Eto K, Kohroggi H, Araki S. [An autopsy case of malignant melanoma possibly originating in the sphenoid sinus]. *Rinsho Shinkeigaku* 1988;28(6):636–642
- Pino Rivero V, Keituaqwa Yáñez T, Marcos García M, Trinidad Ruíz G, Pardo Romero G, Blasco Huelva A. [Melanoma of sphenoid sinus. Case presentation and review of literature]. *Acta Otorrinolaringol Esp* 2004;55(1):45–48
- Kung B, Deschenes GR, Keane W, Cunnane M, Jacob-Ampuero MP, Rosen M. Paranasal sinus melanoma masquerading as chronic sinusitis and nasal polyposis. *Ear Nose Throat J* 2007;86(9):561–564
- Martin JM, Porceddu S, Weih L, Corry J, Peters LJ. Outcomes in sinonasal mucosal melanoma. *ANZ J Surg* 2004;74(10):838–842
- Ferraro RE, Schweinfurth JM, Highfill GR. Mucosal melanoma of the sinonasal tract. *Am J Otolaryngol* 2002;23(5):321–323
- Wagner M, Morris CG, Werning JW, Mendenhall WM. Mucosal melanoma of the head and neck. *Am J Clin Oncol* 2008;31(1):43–48
- Rinaggio J, Hameed M, Baredes S. Melanoma with cartilaginous differentiation originating within the mucosa of the nasal cavity. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;106(6):861–865
- Eyden B, Pandit D, Banerjee SS. Malignant melanoma with neuroendocrine differentiation: clinical, histological, immunohistochemical and ultrastructural features of three cases. *Histopathology* 2005;47(4):402–409
- Sanderson AR, Gaylis B. Malignant melanoma of the sinonasal mucosa: two case reports and a review. *Ear Nose Throat J* 2007;86(5):287–289, 294
- Dass A, Virk RS, Hundal H, Mohan H. Malignant melanoma of the mucous membranes of the head and neck: three case reports. *Ear Nose Throat J* 2006;85(4):268–270
- Malaguarnera M, Vinci M, Pistone G. Malignant melanoma of nasal cavity: case report and review of the literature. *Cancer Biother Radiopharm* 2002;17(1):29–34
- Combs SE, Konkel S, Schulz-Ertner D, et al. Intensity modulated radiotherapy (IMRT) in patients with carcinomas of the paranasal sinuses: clinical benefit for complex shaped target volumes. *Radiat Oncol* 2006;1:23
- Hofbauer GF, Böni R, Simmen D, et al. Histological, immunological and molecular features of a nasal mucosa primary melanoma associated with nasal melanosis. *Melanoma Res* 2002;12(1):77–82
- Helsel JC, Bardales RH, Mukunyadzi P. Fine-needle aspiration biopsy cytology of malignant neoplasms of the sinonasal tract. *Cancer* 2003;99(2):105–112

- 33 Bridger AG, Smee D, Baldwin MA, Kwok B, Bridger GP. Experience with mucosal melanoma of the nose and paranasal sinuses. *ANZ J Surg* 2005;75(4):192–197
- 34 Podboj J, Smid L. Endoscopic surgery with curative intent for malignant tumors of the nose and paranasal sinuses. *Eur J Surg Oncol* 2007;33(9):1081–1086
- 35 Poetker DM, Toohill RJ, Loehrl TA, Smith TL. Endoscopic management of sinonasal tumors: a preliminary report. *Am J Rhinol* 2005;19(3):307–315
- 36 Feeley C, Theaker J. Epithelial markers in primary sinonasal mucosal melanoma. *Histopathology* 2004;45(1):96–98
- 37 Lund VJ, Howard DJ, Harding L, Wei WJ. Management options and survival in malignant melanoma of the sinonasal mucosa. *Laryngoscope* 1999;109(2 Pt 1):208–211
- 38 Díaz Molina JP, Rodrigo Tapia JP, Llorente Pendas JL, Suárez Nieto C. [Sinonasal mucosal melanomas. Review of 17 case]. *Acta Otorrinolaringol Esp* 2008;59(10):489–493
- 39 Tabaee A, Nyquist G, Anand VK, Singh A, Kacker A, Schwartz TH. Palliative endoscopic surgery in advanced sinonasal and anterior skull base neoplasms. *Otolaryngol Head Neck Surg* 2010;142(1):126–128
- 40 Roth TN, Gengler C, Huber GF, Holzmann D. Outcome of sinonasal melanoma: clinical experience and review of the literature. *Head Neck* 2010;32(10):1385–1392
- 41 McLean N, Tighiouart M, Muller S. Primary mucosal melanoma of the head and neck. Comparison of clinical presentation and histopathologic features of oral and sinonasal melanoma. *Oral Oncol* 2008;44(11):1039–1046
- 42 Patel SG, Prasad ML, Escrig M, et al. Primary mucosal malignant melanoma of the head and neck. *Head Neck* 2002;24(3):247–257
- 43 Medina JE, Ferlito A, Pellitteri PK, et al. Current management of mucosal melanoma of the head and neck. *J Surg Oncol* 2003;83(2):116–122
- 44 Penel N, Mallet Y, Mirabel X, Van JT, Lefebvre JL. Primary mucosal melanoma of head and neck: prognostic value of clear margins. *Laryngoscope* 2006;116(6):993–995
- 45 Temam S, Mamelle G, Marandas P, et al. Postoperative radiotherapy for primary mucosal melanoma of the head and neck. *Cancer* 2005;103(2, Issue 2):313–319
- 46 Krengli M, Masini L, Kaanders JH, et al. Radiotherapy in the treatment of mucosal melanoma of the upper aerodigestive tract: analysis of 74 cases. A Rare Cancer Network study. *Int J Radiat Oncol Biol Phys* 2006;65(3):751–759
- 47 Owens JM, Roberts DB, Myers JN. The role of postoperative adjuvant radiation therapy in the treatment of mucosal melanomas of the head and neck region. *Arch Otolaryngol Head Neck Surg* 2003;129(8):864–868
- 48 Ives NJ, Stowe RL, Lorigan P, Wheatley K. Chemotherapy compared with biochemotherapy for the treatment of metastatic melanoma: a meta-analysis of 18 trials involving 2,621 patients. *J Clin Oncol* 2007;25(34):5426–5434
- 49 Bartell HL, Bedikian AY, Papadopoulos NE, et al. Biochemotherapy in patients with advanced head and neck mucosal melanoma. *Head Neck* 2008;30(12):1592–1598