The nucleotide sequence surrounding the origin of DNA replication of Col E₁

Deepak Bastia *

Department of Human Genetics, Yale University School of Medicine, New Haven, CT 06510, USA

Received 23 June 1977

SUMMARY

The DNA of Col E₁ replicates from a unique origin located at a distance of 17-19% of the genome length from the single Eco R₁ cleavage site. The nucleotide sequence about this site has been determined by a combination of RNA and DNA sequencing techniques. The principal features of the sequence are two palindromes, one of which resembles a palindrome located in the intercistronic region of ØX174. The sequence also contains stretches of purine and pyrimidine clusters of the following compositions: pAT₅G, pC₂T₅G, pGT₅G. The origin sequence demonstrates that initiation of DNA replication takes place in an intercistronic region of Col E₁ DNA, although the possibility that this region makes small polypeptides 30-40 residues long cannot be strictly eliminated at this time.

INTRODUCTION

The colicinogenic plasmid Col E₁ not only has become a useful vector in molecular cloning experiments (1), but also a favorable system for the study of DNA replication in vivo (2) and in vitro (3). The DNA of Col E₁ replicates from a unique origin located at a distance of 17-19% of the genome length from the single Eco R₁ cleavage site (3, 4, 5). The replication of this DNA is unidirectional and therefore the terminus of replication is located at or near the origin. The replication intermediates are partially supercoiled "Cairns forms" or θ shaped structures (3, 5, 6). The replication in vivo requires DNA polymerase I and the initiation is rifampicin sensitive (2). Tomizawa (7) has shown that only the initiation in vitro of a new DNA chain complementary to the heavy strand at the origin is rifampicin sensitive, thus suggesting the involvement of an RNA primer at this step of initiation. The on-going replication does not require RNA synthesis or protein synthesis (2). Inhibition of host protein synthesis by chloramphenicol leads to continuation of plasmid replication in the

absence of host DNA replication and thus results in up to 100-fold amplification of the number of copies of plasmids per cell (8).

Hitherto, the structure of only one replication origin; namely, that of SV40 is known (9). Although the entire sequence of ØX174 chromosome has been deduced, the exact location of the replication origin is not known (10). In view of the lack of information regarding the replication origin of plasmid chromosomes, we have sequenced approximately 260 nucleotides of Col E₁ spanning the chromosome from 16.5-20% of the genome length from the Eco R₁ cleavage site. Electron microscopic data from three different laboratories have located the replication origin in this part of Col E₁ chromosome (3, 4, 5). Analysis of Col E₁ DNA carrying large deletions also supports this localization (Inselburg, personal communication) of the replication origin.

MATERIALS AND METHODS

DNA was prepared from JC411 (Col E₁) (kindly supplied by Don Helinski) as previously described (8). Preparation of restriction fragments (19), RNA sequencing (12, 13), DNA sequencing by partial snake venom digestions (14) and by chemical cleavage method (15) are published.

RESULTS

Bastia and Weissman (submitted for publication) have reported a detailed restriction map of the part of Col E₁ DNA containing the replication origin and the relaxation site which is shown in Figure 1. Oka and Takanami (11) have previously reported the Hae III and Hae II cleavage map of Col E₁. The electron microscopic localization of replication origin, translated into the restriction map, places the replication origin in the fragment Hae II-E (see Figure 1) near the Hae III-N - Hae III-H junction. In this paper I present the sequence of the fragments F₁, F₄, and approximately half of F₉ (see Figure 1). These fragments cover approximately 16.5-20% of the genome length from the Eco R₁ cleavage site.

Sequence of F₁:

The fragment of F₁ is 160 base pairs long and contains three Hinf sites and a single Hha I/Hae II site. The strategy for sequencing F₁ was to trans-

Fig. 2 T₁ fingerprint of αP^{32} GTP labeled RNA transcript of F₁

Partial snake venom digestions of 5' end labeled subfragments of F₁:

The subfragments of F₁ were labeled at one of the two 5' phosphoryl ends (with γ -³²P ATP and polynucleotide kinase) by labeling the appropriate parental fragment containing the subfragment at both the 5' ends followed by recleaving with an appropriate restriction enzyme or by separating the two strands of the subfragment labeled at both 5' ends, by two-dimensional homochromatography as previously described (14). The fragments asymmetrically labeled at one of the two 5' ends were subjected to partial hydrolysis with snake venom phosphodiesterase and the partial di-

TABLE 1

Oligonucleotides in extensive T₁ RNase digest of transcripts of the DNA fragment F₁^(a)
 Products of extensive pancreatic RNase digestion^(c)

Oligo- nucleotide number (b)	[α- ³² P] labeled precursor				Sequence of product (d)
	CTP	ATP	GTP	UTP	
1	AAAAAC <u>AC</u> U	C, <u>AC</u> , AAAAAC	AAAAAC, G	AC	ACUCACACAAAAACG (G)
2	<u>AC</u> C, U	<u>C</u> , AC	C, G	<u>AU</u> U	CAUUUCACACCCGG
3	AC, G, C, <u>U</u>	C, U	AC	AU, C, <u>U</u>	CAUUUUUCCUCCUACG (C)
4	AU, <u>C</u> =	U, G	U	AU, C, U	CCUUUAUCCCGU (A)
5	AU, AC, <u>C</u>	U, C, AC, AG	AG	AU, U	UUAUCCACAG (A)
6	AAC, AC	AC, AAC, U	AG	AG	UAACACAG (U)
8	G, C, U	-	C	C, <u>U</u>	CUUUUCCG (C)
9	C, <u>U</u> , G	-	C	<u>C</u> , U	CUUCCUCG (C)
10	AC, U	C, G	U	AC, C	CUCACUG (A) AUAACUG (U)
11	AAU	AAU, C	AG	AAU	AAUCAG (G)
13	U	-	U	AU, C, G	AUUCUG (U)
15	AU, <u>AC</u>	<u>C</u> , U, G	<u>U</u> , C	<u>AU</u> , <u>C</u> , U, G =	CAUCUG (U) UCACUG (A) AC UUACCG (U)
16	-	AU, AAG	AAG	AU	AUAAG (G)
17	AAAAG	AAAAG	AAAAG	-	AAAAG (C)
18	-	U, AAG	AAG (G)	-	UAAG (G)
131	-	-	U	<u>U</u> , G	UUUUG (U)
181	AC	U, G	U	<u>AC</u>	UACUG (A)
201	AC	AC, C, AG	AG	-	CACAG (A)
19	G, U, C AC	C, U, G	AC, C, AG	AC, C	UACCG (C) ACUCGC UCCAGC CUACGC
23	U, G		G, <u>C</u> , U	C	CUCG (G) (C)
24	AC -		AC, C	-	ACCG (A) CACG (U)
25	G, AG	U, G	AU, AG	AU	AUG (C) UAG (A) (C)
26	G	G	<u>U</u> , G	G	UG (G) (U) (A) (C)
27	U, G	-	C, U, G	C	CUG (C) UCG (G)
30	G	G	<u>C</u> , G	G	CG (A) (U) (C) (G)
124	C, G	-	C	-	CCG (C)
231	AC G, U	C,	<u>U</u> , AG	<u>AG</u> , AC	UCAG (U) ACUG (C)
241	AAG	AAG	AAG	-	AAG (C)
261	AG	AG	AG	-	AG (C) (A) (G)
281	AC, AG	C, G	AC, AG	G	CAG (C) ACG (A) (U)

a) See Figure 1.

b) Number refers to the oligonucleotide in Figure 2.

c) The underlined products are present in more than one mole. Where appropriate, the products were analyzed by alkaline hydrolysis to determine the nearest neighbors.

d) Letters in parenthesis indicate nearest neighbors. Sequences were deduced from results shown in Tables 1 and 2 and in Figures.

TABLE 2

Products of extensive U2 RNase digest of oligonucleotides listed in Table 1

Oligo-nucleotide number (a)	Product (b) [α - 32 P] labeled precursor			Sequence of product (c)
	CTP	GTP	UTP	
1	-	1.0	-	CG
	.2	-	0.66	CUCA (C)
2	.19	.22	-	CCCG
	.05	-	0	UUUCA (C)
	CA	0	-	CA (G)
4	0	0	0	UCCCCUG
	.18	-	.12	CCUUA
5	.22	1.1	-	CA
	-	-	.23	UCCA (C)
	-	-	.41	UUA
6	CA	.82	.8	UG
10	-	.45	.45	CUG
11	.48	.49	-	UCA
13	-	.05	-	UUCUG
15	.12	.14	.13	UCUG
	.23	-	.25	UUA (C)
	.5	.5	-	UCA+CCG

a) See footnote (b) of Table 1.

b) Numbers refer to the relative mobility with respect to the xylene cyanol tracking dye.

c) Letters in parenthesis indicate nearest neighbors. Products were analyzed by (i) digestion with alkali; (ii) re-electrophoresis on DEAE paper, pH 1.7, where appropriate.

gests were resolved into a trail of spots by electrophoresis on Cellogel in the first dimension followed by homochromatography in the second dimension. The trails generated by partial digestion (referred to hereafter as SVP trails) of various subfragments of F_1 were used to overlap the T_1 products (representative SVP trails are shown in Figure 3).

The subfragments of F_1 were also asymmetrically labeled at one of the two 5' phosphoryl ends as described above and subjected to base specific chemical degradation procedure described by Maxam and Gilbert (15). The products were fractionated in 20% acrylamide-urea slab gels and representative gel patterns are shown in Figures 4 and 5.

The T_1 oligonucleotides of F_1 were ordered and overlapped using the sequence derived from snake venom trails and the Maxam-Gilbert patterns.

Sequence of F_4 :

The fragment F_4 (Hae III-N, Figure 1) is 43 base pairs long and its sequence was derived entirely by DNA sequencing methods: (a) by examining

Fig. 3 (a - j) Snake venom trails of the fragments P_1 (a-g), P_4 (i), and P_9 (h).

Fig. 3a

Fig. 3b

Fig. 3d

Fig. 3c

Fig. 3j

Fig. 3i

Fig. 4(a,b) Maxam - Gilbert pattern of Hpa II J labeled at the 5' end of its light strand.

the SVP trails of both strands of F_4 after strand separation by homochromatography, (Representative SVP trails are shown in Figure 3.); and (b) the sequence was confirmed by performing the base specific chemical degradation of the fragment Hpa II-J, which includes Hae III-N (Figure 1) labeled at its right 5' phosphoryl end and of Hae III-N labeled at its right 5' end.

The Maxam-Gilbert pattern of the Hae III-N was identical to that of the Hpa II-J (Figure 4a,b) up to the Hae III recognition sequence of GGCC

Both procedures described above yielded identical deductions of the sequence of Hae III-N (F_4).

Sequence of right half of F_9 :

The fragment F_9 is approximately 134 base pairs long and contains no cleavage site for the restriction enzymes available to us. The sequence of up to 16 residues from the right 5' end of F_9 was determined by snake venom partial digestion procedure (Figure 3c).

Maxam-Gilbert gel patterns of F_9 labeled at its Hae III 5' phosphoryl end were obtained and are presented in Figure 5a,b,c. The data from the gels used in the construction of the sequence presented in Figure 6A, were obtained from six independent sets of experiments. In the absence of other lines of evidence, these repeat gel runs were essential, especially in cases where certain T residues showed up in some gels as C residues. For example, the residues at positions 12 and 31 on the heavy strand (bottom line, Figure 6A) in some gels could be misread as C residues. However, with repeated runs, this potential misinterpretation of gel pattern was avoided. This is particularly crucial for the interpretation of the reading frame of the sequence as described in the Discussion section. I have also obtained chemical cleavage patterns of A and G residues of the light strand of F_9 (not shown) labeled at the Hha I end. The part of the light strand that is expected to overlap the heavy strand sequence of F_9 (Figure 6A) between the residues 1 to 16. Although one can determine the number of pyrimidine residues only approximately by examining the gaps between two purine

Fig. 5 (a, b, c) Maxam-Gilbert gel pattern of F₉

```

                                10
                                CTGACTTGAGCGT
                                GACTGAACTCGCA
                                40
          20          30          40
CTATTTTGTGATGCTCGTCAGGGGGCGGAGCCTATGG
GATAAAACACTACGAGCAGTCCCCCGCCTCGGATACC

          60          70          80
AAAAACGCCCTGCTACGTGGCCTTCTTCTGTTCTAG
TTTTTGGGACGATGCACCGGAAGAAGGACAAGGATC

90          100          110          120
TCTTTTGCTCÁCATGTTCTTĠCCGGCCTTAĠCCCTGA
AGAAAACGAGTGTACAAGAACGGCCGGAATAGGGGACT

130          140          150          160
TTCTGTGGATAACTGTGTACCĠTTTTTGTGTĠAGTCAG
AAGACACCTATTGACACAATGGCAAAACACACTCAGTC

170          180          190          200
TACĠGCTCGCCGĠAGTCĠAACĠACĠGAGCGTAGĠGAGTCAG
ATGGCGAGCGGCGTCAGCTTĠCTGGCTCGCATCGCTCAGTC

210          220          230          240
TGAGCGAGGAAGĠGGAAAAGCGCTGGACGTGCAĠTTTCTC
ACTCGCTCCTTCGCTTTTCGCGACCTĠGACGTAAGAAG

250          260          270
CTTACGCATCTGTĠCGGCATTĠĠACACC
GAATGCGTAGACACGCCGTAAAGTGTGG

```

Fig. 6A The nucleotide sequence surrounding the origin of replication of Col E. Top and bottom lines represent the heavy and light strands, respectively.

Fig. 6B Palindromes in the sequence presented in 6A.
a, b Col E₁
c Palindrome present in the intercistronic region of ØX174 (16)

residues, the pattern is not inconsistent with the deduced sequence of F_9 as shown below:

	light strand	
	1	16
pGGGPyPyPyPyGPyPyAPyPyPyPyPyGAPyPyPyGAGPyPyPyA		
CCCA A A GCG G TG G A G A <u>CTG</u> A A <u>CTCA</u> G A <u>T</u>		
	heavy strand	

Although the sequence (Figure 6A) from residue 1 to 55 was derived exclusively from repeated runs of Maxam-Gilbert gels, the rest of the 222 residues were either confirmed by partial snake venom phosphodiesterase digestions or by examination of T_1 products of the RNA transcripts or by both these procedures.

DISCUSSION

Two lines of evidence place the replication origin of Col E_1 in the fragment Hae II-E and near the Hae III-N - Hae III-H junction (Figure 1): (a) electron microscopy of replicating intermediates of Col E_1 cut with Eco R_1 (3, 4 5); (b) even more persuasive evidence comes from the observation that micro Col E_1 DNA which is only 800 base pairs long contains this stretch of Col E_1 , but lacks the region spanning 0-14% of DNA genome length from Eco R_1 site and is able to replicate itself (Inselburg, personal communication). Although electron microscopic localization of replication origin may be subject to a certain degree of imprecision, at least, in the case of SV40, it has proven to be remarkably accurate (see discussion 9). Recently, Tomizawa and co-workers have located the 3' end of the presumptive primer RNA near the F_4 - F_9 junction (personal communication from Dr. Tomizawa).

The salient features of the sequence presented in this report are two palindromes (Figure 6B a, b), one of which bears a striking resemblance to the largest palindrome discovered by Fiddes (16) in the intercistronic region between gene G and F of ϕ X174 (Figure 6B, c).

The resemblance of the two palindromes may be more than a coincidence and may suggest a common recognition site for a protein.

The other striking features of the sequence are clusters of 5As and 5Ts on both strands of the DNA at the positions 17, 53 and 151 (Figure 5A). There

is a stretch of 6 Gs in a row on the light strand at the position 35. The sequence of F_4 is very pyrimidine rich on the light strand.

Examination of the reading frames of the sequence shows that starting from the residues 1, 2 and 3 on the light strand (top line, Figure 6A), one encounters chain terminators: at the residues 7 (UGA), 2 (UGA) and 87 (UAG), respectively. Similarly, on the heavy strand (bottom line, Figure 5A) starting from the residues 275, 274 and 273, one encounters non-sense codons at positions 251 (UAA), 271 (UGA) and 165 (UGA). This observation is meaningful only if the sequence is completely free of errors. We believe this to be the case. Thus, the replication of origin of Col E_1 is located in an intercistronic area, a situation analogous to that of M 13 (20) and SV40 (9), but unlike that of ϕ X174, which has the replication origin somewhere within cistron A (10, 11).

I have sequenced approximately 100 nucleotides beyond (to the left of) the residue using the Maxam-Gilbert method (data not shown). This sequence is tentative in nature, nevertheless does not show any palindromes or repeated sequences. The purine stretch AGGGGGAAA, however, was found upstream in the direction of replication from position 1 of the sequence presented in Figure 6A.

We have also sequenced approximately 300 nucleotides downstream in the direction of replication beyond position 262 and this sequence which contains the relaxation site of Col E_1 (Bastia and Weissman, 1977, submitted for publication) also contains amber, ochre and opal codons in all three reading frames of both strands and thus is a continuation of the untranslated segment containing the replication origin sequenced in the present report. The possibility that the presumptive intercistronic region codes for very small peptides up to 30 residues long, however, cannot be eliminated at this time. With the exception of the two palindromes (Figure 6B a, b), the replication origin of Col E_1 does not feature the extensive two-fold rotational symmetries present in the origin of replication of SV40 (9) and fd (Schaller, personal communication).

The sequence PuU_6Pu has been identified as a strong terminator of transcription in prokaryotes (13, 17). This sequence was not found in the replication origin sequence presented in this report. However, 5 Ts in a

row were found in 3 places. Runs of 4 or 5 Ts in a row with or without interruptions by other pyrimidines are also found at the sites where transcription terminates (18).

How long is the *cis* active site constituting the replication origin? Which are the binding sites for various initiator proteins? Hopefully, these questions will be answered in future experiments.

ACKNOWLEDGMENTS

This work was supported by a grant from the National Institutes of Health to Dr. Sherman M. Weissman. I thank Rosemarie Cavalier, Ann Goss, Regina Ezekiel, Nina Lepak and Richard Wang for technical assistance. I thank Drs. Allan Maxam and Walter Gilbert for making their DNA sequencing procedure available to me before publication. I also thank Drs. J. Inselburg and J. Tomizawa for communicating their results prior to publication.

*Present address: Department of Microbiology, University of Alabama Medical Center, Birmingham, Alabama

REFERENCES

- 1 Herschfield, V., Boyer, H.T., Yanofsky, C., Lovett, M. and Helinski, D. (1974) *Proc. Nat. Acad. Sci. USA* 71, 3455-3459
- 2 Helinski, D. (1976) *Fed. Proc.* 35, 2026
- 3 Tomizawa, J., Sakakibara, Y. and Kakefuda, P. (1974) *Proc. Nat. Acad. Sci. USA* 71, 2260-2264
- 4 Inselburg, J. (1974) *Proc. Nat. Acad. Sci. USA* 71, 2256-2259
- 5 Lovett, M., Katz, L. and Helinski, D. (1974) *Nature* 251, 337-340
- 6 Fuke, M. and Inselburg, J. (1972) *Proc. Nat. Acad. Sci. USA* 69, 89-92
- 7 Tomizawa, J. (1975) *Nature* 257, 253-254
- 8 Clewell, D. (1972) *J. Bacteriol.* 110, 667-676
- 9 Subramanian, K.N., Dhar, R. and Weissman, S.M. (1977) *J. Biol. Chem.* 252, 355-367.
- 10 Sanger, F., Air, G., Barrell, B., Brown, N., Coulson, A., Fiddes, J., Hutchison, C., Slocumbe, P. and Smith, M. (1977) *Nature* 265, 687-695
- 11 Oka, A. and Takanami, M. (1976) *Nature* 264, 193-195
- 12 Brownlee, G.C. and Sanger, F. (1969) *Eur. J. Biochem.* 11, 395-408
- 13 Lebowitz, P., Weissman, S.M. and Radding, C. (1971) *J. Biol. Chem.* 246, 5120-5139
- 14 Maniatis, T., Jeffrey, A. and Kleid, D. (1975) *Proc. Nat. Acad. Sci. USA* 72, 1184-1188
- 15 Maxam, A. and Gilbert, W. (1977) *Proc. Nat. Acad. Sci. USA* 74, 560-564
- 16 Fiddes, J.C. (1976) *J. Mol. Biol.* 107, 1-24

- 17 Squires, C., Lee, F., Bertrand, K., Bronson, M. and Yanofsky, C. (1976) J. Mol. Biol. 103, 351-381
- 18 Ohe, K. and Weissman, S.M. (1971) J. Biol. Chem. 246, 6991-7002
- 19 Subramanian, K.N., Pan, J. Zain, B.S. and Weissman, S.M. (1977) Nucl. Acid Res. 1, 727-752
- 20 Suggs, S.V. and Ray, D.S. (1977) J. Mol. Biol. 110, 147-153

ABBREVIATIONS

γ NTP	ribonucleotide triphosphate
Hae II	endonuclease II from <u>Hemophilus aegyptius</u>
Hae III	endonuclease III from <u>Hemophilus aegyptius</u>
Hha I	endonuclease I from <u>Hemophilus hemolyticus</u>
Hpa II	endonuclease II from <u>Hemophilus parahemolyticus</u>
Hinf I	endonuclease I from <u>Hemophilus influenzae</u> serotype f