Published in final edited form as:

Chem Res Toxicol. 2012 August 20; 25(8): 1581–1597. doi:10.1021/tx200526y.

Reactive Nitrogen Species Reactivities with Nitrones: Theoretical and Experimental Studies

Kevin M. Nash[†], Antal Rockenbauer[§], and Frederick A. Villamena^{†,‡,*}

[†]Department of Pharmacology, College of Medicine, The Ohio State University, Columbus, OH, 43210, United States

[‡] Davis Heart and Lung Research Institute, College of Medicine, The Ohio State University, Columbus, OH, 43210, United States

[§]Molecular Pharmacology, Research Center for Natural Sciences, H-1025 Budapest, Pusztaszeri 59, Hungary

Abstract

Reactive nitrogen species (RNS) such as nitrogen dioxide ($\bullet\text{NO}_2$), peroxyxynitrite (ONOO^-), and nitrosoperoxycarbonate (ONOOCO_2^-) are among the most damaging species present in biological systems due to their ability to cause modification of key biomolecular systems through oxidation, nitrosylation and nitration. Nitrone spin traps are known to react with free radicals and non-radicals *via* electrophilic and nucleophilic addition reactions, and have been employed as reagents to detect radicals using electron paramagnetic resonance (EPR) spectroscopy, and as pharmacological agents against oxidative stress-mediated injury. This study examines the reactivity of cyclic nitrones such as 5,5-dimethylpyrroline *N*-oxide (DMPO) with $\bullet\text{NO}_2$, ONOO^- , ONOOCO_2^- , SNAP and SIN-1 using EPR. The thermochemistries of nitrone reactivity with RNS, and isotropic hfsc's of the addition products were also calculated at the PCM(water)/B3LYP/6-31+G**//B3LYP/6-31G* level of theory with and without explicit water molecules in order to rationalize the nature of the observed EPR spectra. Spin trapping of other RNS such as azide ($\bullet\text{N}_3$), nitrogen trioxide ($\bullet\text{NO}_3$), amino ($\bullet\text{NH}_2$) radicals, and nitroxyl (HNO) were also theoretically and experimentally investigated by EPR spin trapping and mass spectrometry. This study also shows other spin traps such as AMPO, EMPO and DEPMPO can react with radical and non-radical RNS, thus, making spin traps suitable probes as well as antioxidants against RNS mediated oxidative damage.

Introduction

It has become clear that reactive nitrogen species (RNS) along with reactive oxygen species (ROS) have been implicated in the pathogenesis of various diseases¹⁻⁶ or as key mediators in cell signaling⁷ and immune response.⁸ Nitrosative stress[†] results from a myriad of biomolecular modifications⁹⁻¹⁵ caused by RNS such as the paramagnetic nitric oxide ($\bullet\text{NO}$), nitrogen dioxide ($\bullet\text{NO}_2$), as well as the diamagnetic peroxyxynitrite (ONOO^-). Nitric oxide is known to be formed *in vivo* by nitric oxide synthase (NOS) from L-arginine and O_2 ,¹⁶ or

*Corresponding Author 390 Biomedical Research Tower, 460 W. 12th Ave. Columbus, OH 43210, USA. Phone: 614-292-8215. Fax: 614-699-0999. villamena.1@osu.edu.

Supporting Information

Additional EPR spectra, tables, thermodynamic data of all RNS, ROS, nitrones and their respective spin adducts, and complete reference 67. This material is available free of charge *via* internet www.pubs.acs.org.

[†]The term "nitrosative stress" is a misnomer as commonly used in the literature because most manifestations of which are not due to nitrosation reaction but will be referred to in general as stress caused by RNS.

can be released directly by synthetic molecules such as *S*-nitroso-*N*-acetyl-DL-penicillamine (SNAP).¹⁷ The NO-generating function of hemoglobin as an *S*-nitrosothiol synthase using nitrite as substrate has been proposed.¹⁸ Nitric oxide is a free radical that serves as an intracellular messenger and vasodilator, and its toxicity is generally limited to its reaction or oxidation to more highly reactive species such as ONOO⁻ and [•]NO₂ (Scheme 1).¹⁹

Peroxynitrite is formed from the addition reaction of [•]NO with superoxide (O₂^{•-}) at a diffusion-controlled rate.^{20,21} Peroxynitrite is known to exist in the relatively stable cis-conformation, or gain a proton to form peroxynitrous acid (ONOOH, pK_a 6.8). Like O₂^{•-}, ONOO⁻ is capable of reacting with protein active sites containing Cu, Zn, sulfhydryl and Fe-S clusters to cause nitration and protein cleavage resulting in enzyme deactivation.²²⁻²⁵ Currently, indirect methods of ONOO⁻ detection are limited by their sensitivity and specificity, which include fluorescence,²⁶⁻²⁸ electrochemistry,²⁹ tyrosine, tryptophan and DNA nitration,^{30,31} and indirect radical detection *via* electron paramagnetic resonance (EPR) spectroscopy.³² It should be noted however that detection of nitration is not exclusively an indication of peroxynitrite production and that the nitrating agent, NO₂, can originate from other routes.³³ Attempts to trap [•]NO and O₂^{•-} concurrently using Fe(DTCS)₂ and various nitrones, respectively, only yielded individual spectrum of [•]NO, O₂^{•-} and HO[•] adducts due to the varying rates of reaction of these radicals with the spin traps.³⁴ One emerging method of detection is the selective reaction of ONOO⁻ with boronates coupled with fluorescence spectroscopy.³⁵

One relevant mechanism for ONOO⁻/ONOOH decay is its homolytic cleavage through [•]ONO⁻ and [•]ONO⁻OH intermediates (Scheme 2).³⁶ For ONOOH, the rate of radical cleavage has been reported to be $0.35 \pm 0.03 \text{ s}^{-1}$, with about 30% [•]OH and [•]NO₂ release at pH < 5 *via* escape from the solvent cage. The rate constant for ONOOH isomerization to nitric acid (HNO₃) was found to be $1.1 \pm 0.1 \text{ s}^{-1}$.³⁷ Hydrolysis of [•]NO₂ results in the formation of NO₂⁻ and NO₃⁻.^{37,38} For ONOO⁻, the rate of radical cleavage has been reported at $\approx 10^{-6} \text{ s}^{-1}$, with negligible [•]NO₂ and O^{•-} release.³⁹ While the low rate of homolytic cleavage of ONOO⁻ makes the reaction trivial, ONOO⁻ is known to react with dissolved CO₂ to form nitrosoperoxycarbonate (ONOOCO₂⁻), an oxidizing species that undergoes homolytic cleavage to form 30% CO₃^{•-} and [•]NO₂.^{40,41} The decay of ONOOCO₂⁻ and ONOOH has been shown to vary depending on the ability of the solvent to hold the intermediate species in the solvent cage, and is therefore, dependent on the viscosity of solvent.⁴² Peroxynitrite is a strong nucleophile, and has been shown to cause β-scission of carbonyl groups,^{43,44} where acyl- and H-spin adducts have been observed using EPR spin trapping.^{45,46} Peroxynitrite has recently been shown to form peroxynitrate (O₂NOO⁻) at neutral pH through combination of ONOO⁻ and ONOOH to form O₂NOOH and nitrite (NO₂⁻).⁴⁷ [•]NO₂ is also known to dimerize into N₂O₄ at moderate concentrations which effectively hides its radical character.⁴⁸

Nitrogen trioxide radical ([•]NO₃) is an important oxidant in the troposphere and has been considered as a major oxidant for a variety of unsaturated gas-phase organic species. The reactivity of [•]NO₃ radical with organic compounds occurs *via* hydrogen atom abstraction or addition to double bond with a rate constant range of 10^5 - $10^7 \text{ M}^{-1}\text{s}^{-1}$ and $5 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$, respectively.⁴⁹ Azide anion binds to many heme proteins and metalloenzymes inhibiting their activities. It is also a potent inhibitor of mitochondrial electron transport chain by blocking cytochrome *c* oxidase and preventing ATP hydrolysis.⁵⁰ Azidyl radical ([•]N₃) has been generated in solution and their formation was observed using spin trapping technique *via* oxidation of azide anion by peroxidases in the presence of H₂O₂,⁵¹ succinate-driven respiration in azide-inhibited rat brain submitochondrial particles,⁵² or *via* oxidation of azide anion in cadmium sulfide and zinc oxide suspensions.⁵³ Ammonia is one of the most important trace gases in the atmosphere. Activation of NH₃ in aqueous solution leads to the

formation of a reactive amidogen ($\cdot\text{NH}_2$) radical which can later react with a variety of molecular species such as O_2 , amino acids and melanins.⁵⁴ Nitroxyl (HNO) ($\text{pK}_\text{a} = 11.4$) is the one-electron reduction product of $\cdot\text{NO}$ and has been shown to regulate cellular function with unique pharmacological properties, specifically to cardiovascular diseases.⁵⁵

Although EPR has been used to detect RNS in biological systems with hydroxylamine redox probes (*e.g.*, TEMPONE-H),⁵⁶ the interpretation of the observed signal due to the formation of nitroxide is complicated by the fact that the nitroxide can also be formed from a variety of other reactions.⁵⁷ In this study we explored the reactivities of various RNS using EPR spin trapping. The spin trapping technique unambiguously identifies free radicals *via* formation of a persistent radical adduct from the addition reaction of a radical to a nitron spin trap.^{58,59} Furthermore, spin adducts have also been shown to be formed *via* nucleophilic addition reaction to nitrones and subsequent oxidation to the paramagnetic adduct (Forrester-Hepburn mechanism), as demonstrated by others,^{60,61} making this technique appropriate to study reactive non-radical species as well. Another path is that of inverted spin trapping as proposed by Eberson⁶² where one-electron oxidation of nitron yields the nitron radical cation [nitron] $^{\cdot+}$ and its subsequent addition to a nucleophile (Nu^-) forms the nitron-Nu spin adduct. Nitrones have been used for the detection of RNS,⁶³ and have been shown to trap decomposition products and tertiary radicals formed from ONOO^- .⁶⁴ In this study, we also computationally investigated the thermodynamics of RNS reaction to nitrones with the aim to rationalize the nature of the EPR spectral data obtained from the reactions. Studies on the reaction of RNS with nitrones is important, not only for the purpose of RNS detection, but also to partly provide a rationale for the protective properties of nitrones against RNS-mediated cellular toxicity.⁶⁵

Experimental Procedures

Computational Study

Initial conformational search of the RNS, ROS and spin traps and their respective adducts was carried out using Spartan 04 at the MMFF level. Density functional theory (DFT),⁶⁶ at the B3LYP/6-31G* level of theory was employed in this study to determine the optimized geometry and each yielded no imaginary vibrational frequency. All calculations were performed with Gaussian 03,⁶⁷ at the Ohio Supercomputer Center. A scaling factor of 0.9806,⁶⁸ was used for the zero-point vibrational energy (ZPE) corrections for all the B3LYP/6-31G* geometries. The effects of solvation on the gas-phase calculations were also investigated by using the PCM,^{69,70} and the spin and charge densities were obtained from natural population analysis (NPA) approach,⁷¹ at the PCM/B3LYP/6-31+G** level of theory. Optimization was also performed for adducts with two explicit water molecules for prediction of hyperfine splitting constants at the PCM(water)/B3LYP/6-31+G**//B3LYP/6-31G*. Negligible spin contamination of $0.75 < \langle S^2 \rangle < 0.76$ was obtained for all the minima. Using the B3LYP/6-31G* optimized structures, additional calculation was performed for RNS/ROS shown in Table 1 at the G3(MP2) level of theory which is known to perform better than DFT for calculations involving enthalpies of formation, ionization potentials, electron affinities, proton affinities^{72,73}

General EPR Experiments

EPR measurements were taken at room temperature with microwave power 10 mW, modulation amplitude 1 G, receiver gain 1.0×10^5 , scan time 21.4 s, time constant 42.0 s, and sweep width 120 G. Solvents used include DMSO or a 10 mM potassium phosphate buffer (pH 7.0) containing 100 μM diethylenetriaminepentaacetic acid (DTPA). All solutions were bubbled with nitrogen gas prior to irradiation. Sample cell used were 50 μL quartz or glass capillary tubes for UV or non-UV irradiation experiments, respectively. For UV

irradiation experiments, the sample cell was irradiated with a Spectroline low-pressure mercury vapor lamp with 0.64 cm × 5.4 cm dimensions and a 254-nm wavelength. The spectrum simulation was carried out by an automatic fitting program.⁷⁴

Peroxynitrite Trapping

Peroxynitrite (90% in 0.3 M NaOH) was commercially obtained with nitrite/nitrate as impurities and was stored at -86 °C. Peroxynitrite concentration was determined spectrophotometrically at 302 nm. In a typical experiment, solutions contain 20 mM DMPO and 8-12 mM ONOO⁻ in 70% (v/v) DMSO in PBS. For inert atmosphere conditions, solvents (*e.g.*, 70% (v/v) DMSO in PBS) were first purged with argon before making the DMPO and ONOO⁻ solutions. For ONOO⁻ in 100% water, ONOO⁻ was prepared according to the method previously described.⁷⁵ One milliliter of aqueous solutions of 0.7 M HCl and 0.6 M H₂O₂ was extruded simultaneously with 1 mL 0.6 M NaNO₂ using 1 mL plastic syringes into a stirred ice-cold 0.3 mL solution of 3 M NaOH. The excess H₂O₂ was removed with MnO₂. The concentration of ONOO⁻ solution was calculated to be 15 mM based on an extinction coefficient of 1700 M⁻¹ cm⁻¹ at 302 nm. For ONOOCO₂⁻ generation, a solution of DMPO in DMSO was purged with argon and subsequently bubbled with CO₂. Stock argon-purged DMSO solution of ONOO⁻ was then added to the DMPO solution to make 70% (v/v) DMSO-30% PBS final solution.

[•]NO₂ Generation

a) A solution containing 20 mM nitron and 300 mM NaNO₂ in PBS was transferred to a quartz capillary tube and was UV-photolyzed in the EPR cavity according to previous procedure.⁷⁶ b) DMSO was bubbled with argon for 15 min, then [•]NO gas was bubbled through the solvent for 15 min. A 25 μL of [•]NO-saturated DMSO solution was then added to a 25 μL solution of 50 mM nitron in DMSO.

Spin Trapping using SNAP

Final aqueous solution contains 100 mM nitron and 65 mM SNAP. For inert atmosphere conditions, solutions of nitron and SNAP were argon purged prior to their addition to prevent [•]NO₂ formation. An aliquot of 70 μL was transferred to capillary tube and sample was irradiated in the cavity using visible light while acquiring EPR spectra.

Generation of [•]N₃, [•]NH₂, [•]NO₃ and HNO

PBS (pH 7.4) solutions of NaN₃, NH₂OH, or HNO₃ (100 mM) in the presence of 0.2% H₂O₂ and 50 mM DMPO were UV irradiated to afford the corresponding radical adducts. Spin adducts from HNO was carried out using Ar purged Angeli's salt solution (120 mM) in distilled water in the presence of DMPO (100 mM) and subsequent acidification with HNO₃ to give a final acid concentration of 20 mM.

¹⁷O-Isotopic Labeling Studies

O-17 labeled nitrogen dioxide, [•]N¹⁷O₂, was prepared according to the procedure shown above by mixing saturated DMSO solutions of [•]NO and O₂ (28% ¹⁷O atom) in DMSO.

Mass Spectral Studies

a) GC-MS analysis was carried out using a positive ion electron impact ionization (EI) detection. Ten microliter of the CH₃Cl fraction was injected to the column at an initial temperature of 40 °C with ramp of 20 °C min⁻¹ up to a maximum of 250 °C. MS detection was conducted at 200 °C ion source temperature, electron energy of 70 eV and scan speed of 1.6584 scans s⁻¹. In a typical experiment for DMPO-NO₂ adduct formation, 1 μL of pure

DMPO (~10 M) was added to a 25 μL O_2 -saturated DMSO solution followed by 25 μL NO-saturated DMSO solution. Seventy microliter of CH_3Cl was then added and the resulting mixture was vigorously mixed and 10 μL of the bottom CH_3Cl fraction was injected into the column. The procedure was repeated for DMPO alone and O_2/NO in the absence of DMPO. For DMPO-ONOO, ONOO $^-$ was prepared according to the method previously described.⁷⁵ A 70 μL aliquot solution of ONOO $^-$ (15 mM) was added to a 1 μL of pure DMPO followed by 70 μL of CH_3Cl . The resulting mixture was vigorously mixed and 10 μL of the bottom CH_3Cl fraction was injected into the column. The procedure was repeated for DMPO alone and ONOO $^-$ solution in the absence of DMPO. For GC-MS analysis of DMPO-adducts generated from ONOOCO $_2^-$, SNAP, $\cdot\text{NO}_3$, $\cdot\text{NH}_2$, $\cdot\text{N}_3$, and HNO, procedures for their generation as mentioned above were followed except that distilled water was used instead of PBS. **b)** Time-of-flight (TOM) mass spectral analysis was carried out on DMPO-ONOO system only since solution containing DMSO from the generation of DMPO- NO_2 is not compatible for this type of analysis. To a 70 μL of freshly prepared aqueous solution of ONOO $^-$ (15 mM) was added 1 μL of pure DMPO. The resulting solution (70 μL) was directly infused into the spectrometer. All measurements were done in triplicate.

Results and Discussion

Redox Properties

Free energies ($\Delta G_{298\text{K},\text{red}}$) of the various RNS formation and decomposition pathways were computationally studied at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory. Reduction free energies ($\Delta G_{298\text{K},\text{red}}$) for each of the species were calculated and are shown in Table 1. Three oxidizing ROS, $\text{O}_2^{\cdot-}$, $\text{HO}_2^{\cdot}$, and $\cdot\text{OH}$, were also included for comparison. For one-electron reduction reactions, the $\Delta G_{298\text{K},\text{red}}$ of $\cdot\text{NO}_3$ to NO_3^- was found to be the most favorable, even more so than the $\Delta G_{298\text{K},\text{red}}$ of the highly reactive $\cdot\text{OH}$. Reduction of N_2O_4 resulted in an 18.4 kcal mol $^{-1}$ decrease in $\Delta G_{298\text{K},\text{red}}$ compared to its monomeric form $\cdot\text{NO}_2$. *Cis*-peroxynitrite was found to be the least oxidizing of the RNS's studied ($\Delta G_{298\text{K},\text{red}} = -55.7$ kcal/mol), and has free energy of reduction close to $\text{O}_2^{\cdot-}$ ($\Delta G_{298\text{K},\text{red}} = -53.7$ kcal/mol). Previous studies⁷⁷ support our theoretical data showing that *trans*-ONOO $^-$ is more oxidizing than *cis*-ONOO $^-$, by 7.1 kcal mol $^{-1}$. In contrast to ONOO $^-$, the protonated form, *cis*-ONOOH, is more oxidizing by 3.2 kcal mol $^{-1}$ than its *trans* analogue but in general, *cis*- or *trans*-ONOOH are more oxidizing than the *cis*- and *trans*-ONOO $^-$.

The $\Delta G_{298\text{K}}$ of oxidation of DMPO to DMPO $^{*+}$ was found to be 139.7 kcal mol $^{-1}$,⁷⁸ and therefore, $\cdot\text{NO}_3$ is the only RNS in this study that can spontaneously oxidize DMPO with an overall $\Delta G_{298\text{K},\text{rxn}} = -12.6$ kcal mol $^{-1}$. Thermodynamics of oxidation of DMPO by ONOO $^-$ is also unlikely due to the highly endoergic free energy for this process of 84 kcal/mol as had been previously been ruled out by other.⁶⁴ Due to the high oxidation potential of DMPO ($E_{\text{DMPO}^{*+}/\text{DMPO}} = 1.63$ V), and by using [^{17}O] labeling and EPR spin trapping, the formation of DMPO $^{*+}$ from $\text{HO}^{\cdot}$ and $\text{SO}_4^{\cdot-}$ was discarded.⁶⁰ Instead, nucleophilic addition to DMPO and its subsequent oxidation to give an EPR detectable species is the more plausible mechanism for radical adduct formation in sulfite/HRP/ H_2O_2 system. RNS reactions with nitrones may not be limited to simple electron transfer reactions and therefore, nucleophilic addition reactions of RNS to nitrones will be explored in the subsequent sections.

We performed additional calculations at the G3(MP2)//B3LYP/6-31G* level of theory in aqueous phase. G3(MP2) values have been shown to give accurate gas-phase organic thermochemistries for small molecules.⁷⁹ Using the G3(MP2) approach, more exoergic free energy of reduction was observed for $\text{HO}^{\cdot}$ compared to that of $\cdot\text{NO}_2$ and is more consistent with the experimental thermochemical data of $E^{\circ}(\text{HO}^{\cdot}_{\text{g}}/\text{HO}^-_{\text{aq}}) = 1.98$ V⁸⁰ and $E^{\circ}(\cdot\text{NO}_2/\text{NO}_2^-_{\text{aq}}) = 1.04$ V.⁸¹

Nitrogen dioxide radical ($\bullet\text{NO}_2$)

Figure 1 shows the optimized geometries of $\bullet\text{NO}_2$, and the O- and N-centered radical adducts, DMPO- NO_2 . NBO calculation of $\bullet\text{NO}_2$ shows negative charges on the two O-atoms and a positive charge on the N-atom with 44% of the spins residing on the N-atom, and 25% on each of the O-atoms. The N-centered radical adduct formation was exoergic with $\Delta G_{298\text{K},\text{rxn}} = -1.4$ kcal/mol while the O-centered radical adduct is slightly endoergic with $\Delta G_{298\text{K},\text{rxn}} = 1.2$ kcal/mol. Unlike $\text{O}_2^{\bullet-}$, the more favored addition of an electropositive center (*i.e.*, the N-atom) to the nitronyl-C indicates that the $\bullet\text{NO}_2$ addition to DMPO is electrophilic in nature which may be due to the high spin density distribution on the N-atom. The small energy difference between the N-centered and O-centered addition reaction could also indicate that the latter mechanism can occur as will be discussed below.

Attempts to generate $\bullet\text{NO}_2$ by UV irradiation of NaNO_2 with DMPO in PBS only yielded an acyclic adduct with $a_{\text{N}} = 7.3$, $a_{\text{H}} = 4.1$, $a_{\text{H}} = 4.1$ (Figure not shown). Spin trapping using $\bullet\text{NO}/\text{O}_2$ solution in DMSO, however, gave an unidentified O-centered radical adduct which will be called DMPO-OX with $a_{\text{N}} = 13.8$, $a_{\text{H}} = 11.7$ (see Figure 2a). Table 2 and Table S1 list the predicted hfsc's for various $\bullet\text{NO}_2$ adducts (in the presence of bulk dielectric effect of water as well as in the presence of two explicit water molecules) and shows that N-centered radical adduct must exhibit a significant nitro-N hfsc, but experimental evidence shows otherwise, indicating that the O-centered radical adduct could be the major product in DMSO solution. Attempt to further prove the formation of O-centered DMPO adduct with NO_2 , EPR spectrum was obtained from mixing ^{17}O -labeled O_2 (28% O-17 atom) and $\bullet\text{NO}$ saturated DMSO solutions in the presence of DMPO. Solution only yielded the same spectrum as above with $a_{\text{N}} = 13.8$ G and $a_{\text{H}} = 11.8$ G, and did not yield hfsc originating from ^{17}O due perhaps to the low concentration of N^{17}O_2 formed from the complex mechanism of its formation from $\bullet\text{NO}$ and O_2 as well as the low final yield of DMPO- N^{17}O_2 . The non-observance of the ^{17}O was exacerbated by its nuclear spin of $I = 5/2$ that can drastically reduce the amplitude of spectrum by a factor of six. GC-MS analysis, however, showed a distinctive GC peak at ~ 4.4 min with significantly intense base mass peak of 158.98 m/z corresponding to DMPO- NO_2 adduct (see Figure S7a of the SI). Considering that the energy difference for the formation of N- and O-centered radical adducts is only 2.6 kcal/mol, the formation of O-centered radical adduct is not improbable. Moreover, addition of the electronegative center (*i.e.*, the O atom) to the nitronyl follows that of the nucleophilic addition reaction of $\text{O}_2^{\bullet-}$ to nitrones. Based on the thermodynamics of O-centered radical adduct formation from NO_2 which is only 2.6 kcal/mol difference compared to the formation of the N-centered adduct, the fact that the reaction of NO_2 with DMPO yielded an adduct with no additional hfsc due to N, and that the mass spectral analysis is consistent with DMPO- NO_2 formula, it is reasonable to assume that the adduct formed has a general structure of DMPO-ONO.

Spin trapping using $\bullet\text{NO}/\text{O}_2$ in the presence of AMPO, EMPO, or DEPMPO were also carried out (Figure 2b-2d). Similar to that observed for DMPO, significant amounts of nitronyl-ONO adducts were evident. Previously,⁸² it has been shown that olefins exhibited fast reactions with radicals generated from $\bullet\text{NO}/\text{O}_2$ systems yielding $\bullet\text{NO}_2$ addition product *via* O-atom attack. Figure 1 shows the thermodynamics of nitronyl- NO_2 formation which generally shows slightly higher thermodynamic favorability for the formation of the trans-isomers compared to the corresponding cis-isomers, as well as $\bullet\text{NO}_2$ addition *via* the O-atom rather than *via* the N-atom addition. Figure 2 shows the calculated hfsc's for the more preferred nitronyl-ONO adducts, and qualitative trends can be made from the relative magnitudes of a_{N} and $a_{\beta\text{-H}}$, as well as the presence of additional a_{N} for the AMPO-ONO cis and trans adducts as experimentally observed (Figure 2b).

The formation of NO_2 from NO and O_2 is quite complex and may form intermediate species that can react with nitrones. The formation of intermediate species such as N_2O_4 or NO_3 should yield nitron-ONO and nitron- ONO_2 (to ultimately give DMPO-OH), respectively. Since formation of nitron-OH adduct was not evident any of the spectra in Figure 2, therefore, one cannot discount the formation of nitron-ONO from N_2O_4 since the free energy of N-N homolytic cleavage for N_2O_4 to give two molecules of NO_2 is only endoergic by 0.5 kcal/mol.

Peroxynitrite (ONOO^-) and Peroxynitrous Acid (ONOOH)

The electronic property of ONOO^- as shown in Figure 3, is characterized by high negative charge density on the terminal peroxy-O, O(3), for both *cis* and *trans* isomers (-0.68e and -0.72e, respectively). The charge density on the nitroso-O, O(1), is less negative with a charge of -0.21e. The *cis*- ONOO^- conformation is 4.2 kcal/mol more favored than the *trans* isomer. Scheme 3 shows the various modes of addition of ONOO^- to DMPO. Theoretical studies show that nucleophilic addition of ONOO^- to DMPO results in ring opening to form the nitroso-aldehyde and nitrite with $\Delta G_{298\text{K},\text{rxn}}$ of -34.4 kcal/mol. Formation of a triplet adduct from ONOO^- and DMPO gave a highly endoergic product for the O(1)-addition (where O(1) corresponds to the doubly-bonded O) with $\Delta G_{298\text{K},\text{rxn}} = 54.1$ kcal/mol (Figure not shown). Addition of the terminal peroxy-O, that is O(3), to DMPO gave exoergic free energy of -9.0 kcal/mol yielding $\cdot\text{NO}_2$ and $\text{DMPO-O}^{\cdot-}$ radicals, but formation of DMPO-NO and $\text{O}_2^{\cdot-}$ *via* addition through the N atom gave endoergic $\Delta G_{298\text{K},\text{rxn}}$ of 26.2 kcal/mol. It has been previously shown that the reaction of ONOO^- with DMPO in ^{17}O -labeled water yielded DMPO-OH spectra under low oxygen tensions, as well as a methyl radical adduct to DBNBS in the presence of ONOO^- in DMSO.⁶⁴ This experimental evidence support that nucleophilic addition of ONOO^- to DMPO can result in the formation of DMPO-OH adduct from $\text{DMPO-O}^{\cdot-}$ with slightly endoergic $\Delta G_{298\text{K},\text{rxn}}$ of 3.8 kcal/mol (Scheme 3). The formation of doublet adducts from ONOO^- *via* its addition to $[\text{DMPO}]^{*+}$ is highly exoergic with O(3) addition being the most favorable with $\Delta G_{298\text{K},\text{rxn}} = -26.3$ kcal/mol.

EPR spin trapping of ONOO^- in the presence of DMPO was initially performed in DMSO/ H_2O . Figure 4a shows that the spectrum obtained under normal atmospheric conditions was markedly different than those obtained when the solution was purged with argon (Figure 4b). EPR simulation of the resulting spectra under normal atmospheric conditions revealed hfsc's corresponding to an unknown O-centered radical adduct, DMPO-OX (80%; $a_{\text{N}} = 14.0$, $a_{\text{H}} = 9.8$, $a_{\text{H}\gamma} = 1.8$) and a C-centered radical adduct, DMPO-R (20%; $a_{\text{N}} = 15.5$, $a_{\text{H}} = 21.5$). The nature of the DMPO-OX will be further discussed in the succeeding section. Using the same spin trapping conditions with an argon purge yielded a different mixture of radical adducts; a C-centered radical adduct DMPO-R (80%; $a_{\text{N}} = 15.3$, $a_{\text{H}} = 21.5$), DMPOOH (16%; $a_{\text{N}} = 15.4$, $a_{\text{H}} = 15.0$) and an adduct assigned to DMPO-ONX (4%; $a_{\text{N}} = 9.4$, $a_{\text{H}} = 19.9$). The DMPO-R adduct can be assigned to the formation of DMPO-CH_3 , perhaps due to the formation of $\text{HO}^{\cdot}$ from the O-O homolytic cleavage of ONOOH . The O-O homolytic cleavage of ONOOH to form $\text{HO}^{\cdot}$ and $\cdot\text{NO}_2$ was calculated to be slightly endoergic ($\Delta G_{298\text{K},\text{rxn}} = 2.2$ kcal/mol) (Scheme 2), and while this process is likely to occur in solution, the formation of DMPO-OH , where the $\text{HO}^{\cdot}$ originates from ONOOH , was previously confirmed using ^{17}O -labeled water.⁶⁴ The hfsc's observed for the formation of small amounts of DMPO-ONX ($a_{\text{N}} = 9.4$, $a_{\text{H}} = 19.9$) follow a qualitative trend predicted for the formation of DMPO-OONO (Table 2) in which the a_{H} is more improved (*i.e.*, 14.2 G) in the presence of explicit interaction of water molecules compared to the just considering the bulk dielectric effect of water. However, DMPO-OONO can only be formed *via* inverted spin trapping through ONOO^- addition to $[\text{DMPO}]^{*+}$. The question on how $[\text{DMPO}]^{*+}$ could be formed in solution can be accounted for by the $\cdot\text{NO}_3$ oxidation of DMPO to DMPO^{*+} .

However, one could argue that $\text{HO}^\bullet$ can further react with nitrate impurity (as HNO_3) to yield $^\bullet\text{NO}_3$ ($\Delta G_{298\text{K},\text{rxn}} = -12.9$ kcal/mol), or *via* H-atom abstraction from HOONO and isomerization of the $^\bullet\text{OONO}$ formed to $^\bullet\text{NO}_3$ with a total $\Delta G_{298\text{K},\text{rxn}} = -46.0$ kcal/mol (see Scheme 2). Hence, $^\bullet\text{NO}_3$ can oxidize DMPO to $\text{DMPO}^{\bullet+}$ to form DMPO-OONO *via* inverted spin trapping with ONOO^- . This oxidation process is exoergic with $\Delta G_{298\text{K},\text{rxn}}$ of -12.6 kcal/mol, subsequent addition of ONOO^- to $\text{DMPO}^{\bullet+}$ gave highly favorable $\Delta G_{298\text{K},\text{rxn}}$ of -27.6 kcal/mol (Figure 3) giving a net $\Delta G_{298\text{K},\text{rxn}}$ of -40.2 kcal/mol for the formation of an EPR-detectable DMPO-OONO . The initial endoergic process for the $^\bullet\text{NO}_3$ oxidation of DMPO to $\text{DMPO}^{\bullet+}$ could explain the low yield (4%) of DMPO-OONO formed in solution. Also, the formation of DMPO-N(O)O_2 can be ruled out due to the lack of additional a_N from the EPR spectrum, and because the formation of DMPO-OONO is more exoergic with $\Delta G_{298\text{K},\text{rxn}} = -40.2$ kcal/mol than the formation of DMPO-N(O)O_2 with $\Delta G_{298\text{K},\text{rxn}} = -0.5$ kcal/mol (Scheme 3).

Under inert atmosphere in aqueous solution (Figure 4d), ONOO^- yielded an N-centered radical adduct, DMPO-NX , as the major product (~57%) with $a_N = 14.5$, $a_N^\bullet = 3.55$, $a_H = 0.93$. Formation of a minor product, DMPO-ONO (6%) was also observed with $a_N = 14.4$, $a_H = 11.3$. The formation of the O- and N-centered adducts was further confirmed by GC-MS and showed distinctive peak at ~ 6.2 min (not observed with DMPO or ONOO^- alone) with significantly intense base peaks at 175.05 m/z and 159.02 m/z corresponding to DMPO/ONOO^- and DMPO-ONO adducts (Figure S8 of SI). Time-of-flight mass spectrum also confirmed the formation of a base peak at 175.14 m/z which was not observed with ONOO^- or DMPO alone (Figure S9 of SI). In addition to the formation of O- and N-centered adducts, the same DMPO-R adduct as also observed as in DMSO suggesting that the origin of the DMPO-R is independent of the solvent used. The only C-centered radical source would be from the nitron itself *via* trapping of another nitron. As proposed above, $^\bullet\text{NO}_3$ at high $[\text{ONOO}^-]$ may oxidize DMPO to $\text{DMPO}^{\bullet+}$ leading to the addition of excess DMPO to $\text{DMPO}^{\bullet+}$ to form $[\text{DMPO-DMPO}]^{\bullet+}$ with $\Delta G_{298\text{K},\text{rxn}} = 0.5$ kcal/mol. The predicted hfsc values for the nitronyl-N and peroxyxynitrite-N (*i.e.*, 11.0 and 5.8, respectively) of DMPO-N(O)OO gave reasonable agreement with the experimental, while significant discrepancy was observed between the predicted and experimental hfsc for a_H with 15.7 and 0.9, respectively (Table 2). Optimization of DMPO-N(O)OO with 2 explicit water molecules was problematic but we predict that the hfsc will not change significantly within ~2 G as also observed for the other adducts with explicit water interaction. Therefore, the nature of DMPO-NX warrants further investigation.

Scheme 4 shows the various modes of ONOOH addition to DMPO where two favorable pathways can be seen: 1) the formation of singlet species, nitroso aldehyde and nitrous acid, *via* electrophilic addition of O(3) with highly favorable overall $\Delta G_{298\text{K}}$'s of -45.3 kcal/mol; and 2) the formation of doublet species, nitrogen dioxide and DMPO-OH *via* O-O homolytic cleavage of ONOOH with also highly favorable $\Delta G_{298\text{K},\text{rxn}}$ of -35.9 kcal/mol. Figure 5 shows the various modes of ONOOH addition to DMPO with only the O(3) addition to form DMPO-OH and $^\bullet\text{NO}_2$ being the most favorable pathway.

Using the same spin trapping conditions employed for DMPO (Figure 4b), the reactivity of ONOO^- with EMPO, DEPMPO and AMPO were also explored in an argon-purged 70% DMSO solution. Shown in Figure S1, similar to DMPO, significant amounts of nitron-R were formed from all nitrones, perhaps *via* formation of $^\bullet\text{CH}_3$. While not observed for DEPMPO, nitron-OX adducts were observed for EMPO and AMPO which can be assigned to the formation of DMPO-OONO most likely *via* inverted spin trapping. Figure 3 shows the thermodynamics of nitron-OONO formation from $[\text{nitron}]^{\bullet+}$ and its addition to ONOO^- giving exoergic free energies of ~28 kcal/mol, with the formation of the cis-isomer only slightly more exoergic by < 1 kcal/mol compared to the trans-isomer. The approximated

calculated hfsc's for the ONOO⁻ adducts of DMPO, EMPO, DEPMPO, and AMPO are shown in Table 2 (for DMPO) and S1 (for the rest of the nitrones), and generally shows higher $a_{\beta\text{H}}$ than a_{N} by as much as ~4 G. This qualitative trend in the relative magnitudes of $a_{\beta\text{H}}$ and a_{N} follows the experimentally observed hfsc's with the exception of EMPO where the observed $a_{\beta\text{H}}$ is lower than a_{N} .

Nitrosoperoxycarbonate (ONOOCO₂⁻)

The optimized structure of ONOOCO₂⁻ was consistent with that of previous theoretical studies,⁸³ which show a 1.58 Å bond distance between the carboxylate-C and peroxy-O, and a C-O-O-N dihedral angle of 82° (Figure 6). The $\Delta G_{298\text{K},\text{rxn}}$ of formation of ONOOCO₂⁻ from *trans*-ONOO⁻ is 0.5 kcal mol⁻¹,⁸³ and is more favorable than its formation from *cis*-ONOO⁻ with $\Delta G_{298\text{K},\text{rxn}}$ = 2.8 kcal mol⁻¹. However, the energy required for the conformational change from *cis*-ONOO⁻ to *trans*-ONOO⁻ was lowered upon complexation with CO₂ (the free energies of conformation change for free and CO₂-complexed ONOO⁻ were determined to be 4.2 and 1.9 kcal mol⁻¹, respectively). During its short half-life, the ONOOCO₂⁻ complex can act as a strong oxidant and, as previously discussed, can undergo homolytic cleavage to form CO₃^{•-} and [•]NO₂ species ($\Delta G_{298\text{K},\text{rxn}}$ = -19.0 kcal mol⁻¹), or rearrange to form NO₃⁻ and CO₂ ($\Delta G_{298\text{K},\text{rxn}}$ = -52.3 kcal mol⁻¹) (Scheme 5). These calculations are consistent with previous models in which the yield of radical products was 30%. Thus, in neutral pH to moderately basic solution, ONOO⁻ is energetically able to exist in equilibrium with ONOOCO₂⁻ where it can act as a strong oxidizing agent and spontaneously form radical species. At neutral pH, ONOO⁻ can undergo competing mechanisms for radical formation such as protonation to form ONOOH, or ONOOCO₂⁻ formation. As the pH is lowered, the protonation of ONOO⁻ can dominate, forming ONOOH.

We further investigated the nature of the DMPO-OX EPR signal in Figure 4a. Under normal atmospheric condition, the observed formation of DMPO-OX adduct could originate from DMPOOCO₂⁻. This hypothesis was confirmed by bubbling pure CO₂ through argon-purged DMSO before ONOO⁻ spin trapping (Figure 4c). The spectra obtained from the CO₂ bubbled solution yielded one radical adduct at low ONOO⁻ concentration, which is identified as DMPO-OCO₂⁻ based on our previous reported hfsc's values of a_{N} = 14.32, a_{H} = 10.68, $a_{\beta\text{H}}$ = 1.37 for DMPO-OCO₂⁻.⁸⁴ Figure S2 shows that only when [ONOO⁻] >> [CO₂] the DMPO-R adduct is formed, while at [ONOO⁻] << [CO₂], the DMPO-OCO₂⁻ adduct is more predominant due to the formation of [•]CH₃ from HO[•] via O-O homolytic cleavage of ONOOH as discussed above. Figure 6 shows the optimized geometries of DMPO-ONOOCO₂⁻ arising from the various modes of ONOOCO₂⁻ addition reaction to DMPO. Results suggest that N- and carboxylate-O(4) addition to DMPO gave the most exoergic $\Delta G_{298\text{K},\text{rxn}}$ with -7.5 and -13.4 kcal/mol, respectively. Due to the highly negative charge density of the carboxylate-O(4)'s ONOOCO₂⁻ (-0.74e), its addition to DMPO is therefore nucleophilic in nature and is the most favorable to give DMPO-OCO₂⁻ and [•]NO₂, further confirming the EPR results observed in the presence of CO₂ and ONOO⁻. GC-MS analysis of the sample showed evidence of DMPO-OCO₂ adduct formation where the chromatogram showed a unique peak at RT 3.1 min with a base mass peak 174.91 *m/z* corresponding to [DMPO-OCO₂H + H]⁺ (Figure S10 of SI).

In DMSO, the possibility of forming the decomposition products, CH₃S=O(NO₂) + [•]CH₃ and CH₃S=O(CO₃)⁻ + [•]CH₃, due to oxidation by [•]NO₂ and CO₃^{•-}, respectively, was also considered. However, results show endoergic free energies of 40.5 kcal mol⁻¹ and 22.3 kcal mol⁻¹, for reactions of [•]NO₂ or CO₃^{•-} with DMSO, respectively, compared to the exoergic [•]OH reaction to DMSO to form CH₃S=O(OH) + [•]CH₃ with $\Delta G_{298\text{K},\text{rxn}}$ = -2.3 kcal/mol indicate that [•]CH₃ formation is less likely to occur from [•]NO₂ and CO₃^{•-} oxidation of

DMSO. Furthermore, formation of N_2O_4 from $\cdot\text{NO}_2$ with $\Delta G_{298\text{K},\text{rxn}} = -0.5$ kcal/mol can compete with $\cdot\text{NO}_2$ oxidation of DMSO.

Spin Trapping Studies using S-Nitroso-N-Acetyl-D,L-Penicillamine (SNAP) and 3-Morpholininosydnonimine (SIN-1)

Scheme 6 shows the reaction of SNAP with DMPO. Homolytic cleavage of the S-N bond of SNAP into $\cdot\text{NO}$ and thiol radical ($\text{RS}^\bullet$) was determined to be endoergic, while the ionic cleavage of SNAP into NO^+ and thiol anion (RS^-) was calculated to be highly unfavorable with $\Delta G_{298\text{K},\text{rxn}} = 14.5$ kcal/mol and $\Delta G_{298\text{K},\text{rxn}} = 50.5$ kcal/mol, respectively, suggesting that the only relevant mechanism of SNAP decomposition is the release of $\cdot\text{NO}$ and $\text{RS}^\bullet$. While it is energetically favorable for the free $\text{RS}^\bullet$ to form an adduct with DMPO ($\Delta G_{298\text{K},\text{rxn}} = -1.1$ kcal/mol), it was calculated that the formation of a disulfide from two $\text{RS}^\bullet$ molecules was significantly more favorable with $\Delta G_{298\text{K},\text{rxn}} = -33.6$ kcal/mol.

EPR spin trapping from SNAP decomposition was performed. Solutions of irradiated SNAP in water alone under inert atmosphere and air did not give any quartet EPR signal. The EPR spectra of DMPO in the presence of SNAP under normal atmosphere was determined to be $\text{HO}^\bullet$ adducts (Figure 7a) with slightly different g values (*i.e.*, major species: $a_{\text{N}} = 15.2$, $a_{\text{H}} = 14.8$; and minor; $a_{\text{N}} = 15.8$, $a_{\text{H}} = 15.2$) indicating that the presence O_2 can lead to the formation of DMPO-OH via perhaps the initial reaction of O_2 with the NO to form an oxidizing species. However, under Ar atmosphere, formation of an $\text{RS}^\bullet$ adduct was observed (Figure 7b) with $a_{\text{N}} = 14.6$, $a_{\text{H}} = 16.0$ close to the reported hfsc values for DMPO-SNAP of $a_{\text{N}} = 15.3$, $a_{\text{H}} = 17.7$ and DMPO-SG of $a_{\text{N}} = 15.2$, $a_{\text{H}} = 16.3$.⁸⁵ GC-MS analysis of the irradiated DMPO/SNAP solution gave a significantly intense GC peak at 6.45 min compared to DMPO and SNAP alone with base mass peak of 302.94 m/z (see Figure S11 of SI).

The formation of S-centered radical adducts from SNAP were further confirmed by spin trapping using AMPO, EMPO and DEPMPO, showing hfsc's consistent to that of the formation of S-centered radical (see Figure S3). The signal intensity increased significantly when the experiment was repeated under inert atmosphere (Figure 7b) which suggests $\cdot\text{NO}_2$ formation under atmospheric conditions can annihilate the signal, probably *via* its direct reaction to the DMPO-SR adduct. It has been previously shown that as $\cdot\text{NO}$ concentration increases in the presence of O_2 , the rate of $\cdot\text{NO}_2$ formation increases, such that at 1.0 mM $\cdot\text{NO}$, the half-life of $\cdot\text{NO}$ due to $\cdot\text{NO}_2$ formation is 0.56 s.¹⁹ Calculations suggest that the formation of a DMPO-NO adduct is thermodynamically unfavorable, with $\Delta G_{298\text{K},\text{rxn}} = 14.0$ kcal/mol,⁸⁶ while formation of a DMPO- NO_2 adduct is favorable, with $\Delta G_{298\text{K},\text{rxn}} = -1.4$ – -1.2 kcal/mol. While DMPO may display thermodynamically favorable trapping reactions with $\cdot\text{NO}_2$, the radical characteristic of $\cdot\text{NO}_2$ can be 'hidden' *via* its complexation to form N_2O_4 which can compete with its addition to the nitron, hence, no $\cdot\text{NO}_2$ adduct was observed (see Scheme 6).

As expected, the unfavorable reactivity of $\cdot\text{NO}$ to DMPO only yielded $\text{O}_2^{\bullet-}$ adduct using SIN-1 as shown in Figure S4. While the SIN-1 decomposition mechanism was proposed⁸⁷ to include the formation of a C-centered radical, only DMPO-OOH was observed in this study.

Nitrogen trioxide ($\cdot\text{NO}_3$), Amino ($\cdot\text{NH}_2$), Azidyl ($\cdot\text{N}_3$), and Nitroxyl (HNO)

Figure 8 shows the various radical adducts formed from $\cdot\text{NO}_3$, $\cdot\text{NH}_2$ and $\cdot\text{N}_3$. The formation of O-centered radical adduct, DMPO- ONO_2 , is more favorable than the N-centered radical adduct, DMPO- NO_3 , with $\Delta G_{298\text{K},\text{rxn}}$ of -20.0 kcal/mol and 56.4 kcal/mol, respectively. Due to the higher reduction potential of NO_3 $E^\circ(\cdot\text{NO}_3/\text{NO}_3^-) = 1.9$ – 2.6 V⁸⁸ compared to the oxidation potential of DMPO $E^\circ(\text{DMPO}/\text{DMPO}^+) = 1.6$ V⁸⁹ (consistent with the free energies of reduction shown in Table 1), it is expected that the

mechanism of DMPO-NO₃ formation should proceed *via* inverted-spin trapping where DMPO is oxidized to DMPO^{•+}•NO₃ and subsequent addition of NO₃⁻ to DMPO^{•+}. The high negative charge on the O atoms of NO₃⁻ favored the nucleophilic addition of the radical to the nitron. Spin density distribution on the nitro-N in DMPO-ONO₂ is negligible. EPR spin trapping of •NO₃ only gave spectrum corresponding to DMPOOH. Although the formation of DMPO-ONO₂ was already exoergic, its hydrolysis to DMPO-OH and HNO₃ is also favorable with $\Delta G_{298K,rxn} = -4.1$ kcal/mol (Scheme 7)—significantly more favorable than the highly endoergic formation of DMPO-OH and HOONO with $\Delta G_{298K,rxn} = 28.0$ kcal/mol. Scheme 7 further shows that DMPO-OH and HNO₃ are more likely to be formed *via* O-N homolytic cleavage of DMPO-ONO₂ to form DMPO-O• and •NO₂ with $\Delta G_{298K,rxn} = 19.1$ kcal/mol compared to the O-N heterolytic cleavages resulting in the formation of DMPO-O⁺ and NO₂⁻, or DMPO-O⁻ and NO₂⁺ with $\Delta G_{298K,rxn}$'s of 41.1 and 67.5 kcal/mol, respectively.

Since only DMPO-OH adduct was observed during irradiation of DMPO in the presence of HNO₃ and H₂O₂, the effect of HNO₃ concentration on DMPO-OH formation was further investigated. It is expected that the DMPO-OH concentration will be dependent on HNO₃ concentration at constant H₂O₂ concentration. Results show that lower DMPO-OH is formed at higher HNO₃ concentration which indicates that HNO₃ can be competing with DMPO for reaction with HO• (see Figure S5 of SI). Moreover, DMPO-OH formation was evident even at large excess of HNO₃ over H₂O₂ (*i.e.*, 800 mM HNO₃: 140 mM H₂O₂) which may also indicate •NO₃ addition to DMPO, however, the use of HN¹⁷O₃ would have been more straightforward for the elucidation of the mechanism of DMPO-OH formation. Nevertheless, the calculated thermochemistries and formation of DMPO-OH adduct at large excess of HNO₃ suggest origination of DMPO-OH from DMPO-NO₃.

Formations of DMPO-NH₂ and DMPO-N₃⁻ gave exoergic $\Delta G_{298K,rxn}$'s of -28.3 and -13.4 kcal/mol, respectively, with the most electronegative N-atoms adding to the nitron. Spin densities of the N-atom directly bound to the nitron range from 2%-3%, which is enough to impart additional a_N due to the N-atom as shown in Figure 9. The formation of DMPO-NH₂ was previously reported by Kirino, et al.⁹⁰ ($a_N = 15.9$, $a_H = 19.0$ and $a_{N-amino} = 1.7$) as well as by Chignell, et al.⁹¹ ($a_N = 15.9$, $a_H = 19.3$ and $a_{N-amino} = 1.60$). These values are consistent with the current data of $a_N = 15.8$, $a_H = 19.3$ and $a_{N-amino} = 1.6$. GC-MS analysis showed evidence of DMPO-NH₂ adduct formation where chromatogram showed a unique peak at 3.18 min with a base mass peak at 128.97 m/z (Figure S12 of SI). Generation of DMPO-N₃ and its detection by EPR spin trapping was previously reported⁵¹ giving a quartet of triplets with hfcs values of $a_N = 14.8$, $a_H = 14.2$ and $a_{N-azide} = 3.1$. These values are consistent with the current data of $a_N = 14.8$, $a_H = 14.2$ and $a_{N-amino} = 3.4$. GC-MS analysis showed evidence of DMPO-N₃ adduct formation in which the chromatogram gave a unique peak at 6.21 min with a base mass peak 155.03 m/z . (Figure S13 of SI).

Addition of ³NO⁻ and HNO to DMPO (Figure 10) *via* the N-atom each resulted in ring opening to form the nitroso-oximate and nitroso-oxime, with the latter being exoergic having $\Delta G_{298K,rxn}$ of -2.2 kcal/mol. Previously,⁹² spin trapping studies using DMPO and Angeli's salt only showed formation of DMPO-OH and C-centered adducts as also confirmed in this work (see Figure S6 of SI). GC-MS analysis revealed a unique GC peak at 3.21 min with base mass peak 113.08 m/z (Figure S14 of SI) corresponding to an ene-aldoxime which can be formed from the elimination of the NO from the C-nitroso moiety of the originally formed nitroso-oxime.

Conclusions

EPR and theoretical studies of the reaction between DMPO and $\cdot\text{NO}_2$ suggest formation of an O-centered radical adduct, DMPO-ONO. The formation of an O-centered radical adduct from $\cdot\text{NO}_2$ was also demonstrated using AMPO, EMPO and DEPMPO. The addition reactions of DMPO with ONOO $^-$ /ONOOH primarily result in ring opening to form the nitroso-aldehyde and nitrite/nitrous acid. The formation of DMPO-OH, as detected by EPR, from the O-O homolytic cleavage of ONOOH was also thermodynamically favorable. Decomposition of ONOOCO $_2^-$ in the presence of DMPO yielded CO $_3^{\cdot-}$ adduct at high CO $_2$ concentration. However, alkyl adduct formation in an all PBS system suggests inverted spin trapping of DMPO by DMPO $^{\cdot+}$ which is likely generated from the highly oxidizing $\cdot\text{NO}_3$. The EPR-detectable nitronyl adduct as DMPO-OONO, whose formation can also be rationalized by inverted spin trapping, was shown to be also formed using AMPO and EMPO as well. EPR studies of the NO-donor SNAP in the presence of DMPO show S-centered radical adduct formation under inert atmosphere. Reaction of SIN-1 with DMPO only yielded O $_2^{\cdot-}$, while $\cdot\text{NO}_3$ addition to DMPO initially formed DMPO-ONO $_2$, its subsequent hydrolysis yielded DMPO-OH and HNO $_3$ as the former detected by EPR, consistent with the calculated exoergic free energy. Additions of $\cdot\text{N}_3$ and $\cdot\text{NH}_2$ to DMPO yielded N-centered radicals with significant hyperfine contributions from the N-atoms of the radical moieties. Reaction of HNO to DMPO only yielded a ring opening product to give nitroso-oxime and DMPO-OH. This study shows a myriad of RNS reaction with nitrones, and through the use of spin traps, can provide insights into the mechanisms of RNS-mediated reactions in chemical and biological systems. Moreover, this study can lead to the development of better probe designs for RNS detection, and to the understanding the antioxidative properties of nitrones.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

This publication was made possible by Grant RO1 HL81248 from the NIH National Heart, Lung, and Blood Institute. This work was supported by an allocation of computing time from the Ohio Supercomputer Center.

Abbreviations

EPR	electron paramagnetic resonance
ROS	reactive oxygen species
RNS	reactive nitrogen species
SNAP	<i>S</i> -nitroso- <i>N</i> -acetyl-DL-penicillamine
TEMPONE-H	1-hydroxy-2,2,6,6-tetramethyl-4-oxo-piperidine
HRP	horseradish peroxidase
DMPO	5,5-dimethyl-pyrroline <i>N</i> -oxide
AMPO	5-carbamoyl-5-methyl-pyrroline <i>N</i> -oxide
EMPO	5-ethoxycarbonyl-5-methyl-pyrroline <i>N</i> -oxide
DEPMPO	5-(diethoxyphosphoryl)-5-methyl-1-pyrroline <i>N</i> -oxide

References

1. Bossy-Wetzel E, Schwarzenbacher R, Lipton SA. Molecular pathways to neurodegeneration. *Nat. Med.* 2004; 10(Suppl):S2–9. [PubMed: 15272266]
2. Escobales N, Crespo MJ. Oxidative-nitrosative stress in hypertension. *Curr. Vasc. Pharmacol.* 2005; 3:231–246. [PubMed: 16026320]
3. Kaneki M, Shimizu N, Yamada D, Chang K. Nitrosative stress and pathogenesis of insulin resistance. *Antioxid. Redox Signal.* 2007; 9:319–329. [PubMed: 17184170]
4. Llorens S, Nava E. Cardiovascular diseases and the nitric oxide pathway. *Curr. Vasc. Pharmacol.* 2003; 1:335–346. [PubMed: 15320480]
5. Sugiura H, Ichinose M. Nitrate stress in inflammatory lung diseases. *Nitric Oxide.* 2011; 25:138–144. [PubMed: 21440655]
6. Szabo C. Role of nitrosative stress in the pathogenesis of diabetic vascular dysfunction. *Br. J. Pharmacol.* 2009; 156:713–727. [PubMed: 19210748]
7. Thomas DD, Ridnour LA, Isenberg JS, Flores-Santana W, Switzer CH, Donzelli S, Hussain P, Vecoli C, Paolocci N, Ambs S, Colton CA, Harris CC, Roberts DD, Wink DA. The chemical biology of nitric oxide: implications in cellular signaling. *Free Radical Biol. Med.* 2008; 45:18–31. [PubMed: 18439435]
8. Ferguson LR. Chronic inflammation and mutagenesis. *Mutat. Res.* 2010; 690:3–11. [PubMed: 20223251]
9. Bartesaghi S, Valez V, Trujillo M, Peluffo G, Romero N, Zhang H, Kalyanaraman B, Radi R. Mechanistic studies of peroxynitrite-mediated tyrosine nitration in membranes using the hydrophobic probe *N*-tert-BOC-*L*-tyrosine tert-butyl ester. *Biochemistry.* 2006; 45:6813–6825. [PubMed: 16734418]
10. Kalyanaraman B. Nitrated lipids: a class of cell-signaling molecules. *Proc. Natl. Acad. Sci. U S A.* 2004; 101:11527–11528. [PubMed: 15292510]
11. Lancaster JR Jr. Protein cysteine thiol nitrosation: Maker or marker of reactive nitrogen species-induced non-erythroid cellular signaling? *Nitric Oxide.* 2008; 19:68–72. [PubMed: 18503780]
12. Nuriel T, Hansler A, Gross SS. Protein nitrotryptophan: Formation, significance and identification. *J. Proteomics.* 2011
13. Ridnour LA, Thomas DD, Mancardi D, Espey MG, Miranda KM, Paolocci N, Feelisch M, Fukuto J, Wink DA. The chemistry of nitrosative stress induced by nitric oxide and reactive nitrogen oxide species. Putting perspective on stressful biological situations. *Biol. Chem.* 2004; 385:1–10. [PubMed: 14977040]
14. Shahani N, Sawa A. Nitric oxide signaling and nitrosative stress in neurons: role for S-nitrosylation. *Antioxid. Redox Signal.* 2011; 14:1493–1504. [PubMed: 20812870]
15. Tiago T, Palma PS, Gutierrez-Merino C, Aureliano M. Peroxynitrite-mediated oxidative modifications of myosin and implications on structure and function. *Free Radical Res.* 2010; 44:1317–1327. [PubMed: 20815777]
16. Rosen GM, Tsai P, Pou S. Mechanism of free-radical generation by nitric oxide synthase. *Chem. Rev.* 2002; 102:1191–1200. [PubMed: 11942793]
17. McAninly J, Williams DLH, Askew SC, Butler AR, Russell C. Metal ion catalysis in nitrosothiol (RSNO) decomposition. *Chem. Commun.* 1993; 23:1758–1759.
18. Angelo M, Singel DJ, Stamler JS. An S-nitrosothiol (SNO) synthase function of hemoglobin that utilizes nitrite as a substrate. *Proc. Natl. Acad. Sci. U S A.* 2006; 103:8366–8371. [PubMed: 16717191]
19. Beckman JS, Koppenol WH. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and the ugly. *Am. J. Physiol.* 1996; 271:C1424–C1437. [PubMed: 8944624]
20. Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc. Natl. Acad. Sci. U S A.* 1990; 87:1620–1624. [PubMed: 2154753]
21. Czapski G, Goldstein S. The role of the reactions of NO with superoxide and oxygen in biological systems: a kinetic approach. *Free Radical Biol. Med.* 1995; 19:785–794. [PubMed: 8582651]

22. Alvarez B, Demicheli V, Duran R, Trujillo M, Cervenansky C, Freeman BA, Radi R. Inactivation of human Cu,Zn superoxide dismutase by peroxynitrite and formation of histidinyl radical. *Free Radical Biol. Med.* 2004; 37:813–822. [PubMed: 15304256]
23. Ischiropoulos H, Zhu L, Chen J, Tsai M, Martin JC, Smith CD, Beckman JS. Peroxynitrite-mediated tyrosine nitration catalyzed by superoxide dismutase. *Arch. Biochem. Biophys.* 1992; 298:431–437. [PubMed: 1416974]
24. Pearce LL, Martinez-Bosch S, Manzano EL, Winnica DE, Epperly MW, Peterson J. The resistance of electron-transport chain Fe-S clusters to oxidative damage during the reaction of peroxynitrite with mitochondrial complex II and rat-heart pericardium. *Nitric Oxide.* 2009; 20:135–142. [PubMed: 19118636]
25. Rubbo H, Denicola A, Radi R. Peroxynitrite inactivates thiol-containing enzymes of *Trypanosoma cruzi* energetic metabolism and inhibits cell respiration. *Arch. Biochem. Biophys.* 1994; 308:96–102. [PubMed: 8311481]
26. Yang D, Wang H-L, Sun Z-N, Chung N-W, Shen J-G. A highly selective fluorescent probe for the detection and imaging of peroxynitrite in living cells. *J. Am. Chem. Soc.* 2006; 128:6004–6005. [PubMed: 16669647]
27. Sun Z-N, Wang H-L, Liu F-Q, Chen Y, Tam PKH, Yang D. BODIPY-based fluorescent probe for peroxynitrite detection and imaging in living cells. *Org. Lett.* 2009; 11:1887–1890. [PubMed: 19331349]
28. Lancaster JR Jr. The use of diaminofluorescein for nitric oxide detection: Conceptual and methodological distinction between NO and nitrosation. *Free Radical Biol. Med.* 2010; 49:1145. [PubMed: 20600838]
29. Cortes JS, Granados SG, Ordaz AA, Jimenez JAL, Griveau S, Bedioui F. Electropolymerized manganese tetraaminophthalocyanine thin films onto platinum ultramicroelectrode for the electrochemical detection of peroxynitrite in solution. *Electroanalysis.* 2007; 19:61–64.
30. Lorch S, Lightfoot R, Ohshima H, Virag L, Chen Q, Hertkorn C, Weiss M, Souza J, Ischiropoulos H, Yermilov V, Pignatelli B, Masuda M, Szabo C. Detection of peroxynitrite-induced protein and DNA modifications. *Methods Mol. Biol.* 2002; 196:247–275. [PubMed: 12152205]
31. Ikeda K, Yukihiro Hiraoka B, Iwai H, Matsumoto T, Mineki R, Taka H, Takamori K, Ogawa H, Yamakura F. Detection of 6-nitrotryptophan in proteins by Western blot analysis and its application for peroxynitrite-treated PC12 cells. *Nitric Oxide.* 2006; 16:18–28. [PubMed: 16765071]
32. Kalyanaraman B, Karoui H, Singh RJ, Felix CC. Detection of thiyl radical adducts formed during hydroxyl radical- and peroxynitrite-mediated oxidation of thiols-a high resolution ESR spin-trapping study at Q-band (35 GHz). *Anal. Biochem.* 1996; 241:75–81. [PubMed: 8921168]
33. Goldstein S, Czapski G, Lind J, Merényi G. Tyrosine nitration by simultaneous generation of NO and O₂ under physiological conditions. *J. Biol. Chem.* 2000; 275:3031–3036. [PubMed: 10652282]
34. Tsai P, Porasuphatana S, Pou S, Rosen GM. Investigations into the spin trapping of nitric oxide and superoxide: models to explore free radical generation by nitric oxide synthase. *J. Chem. Soc., Perkin Trans.* 2000; 2:983–988.
35. Sikora A, Zielonka J, Lopez M, Joseph J, Kalyanaraman B. Direct oxidation of boronates by peroxynitrite: Mechanism and implications in fluorescence imaging of peroxynitrite. *Free Radical Biol. Med.* 2009; 47:1401–1407. [PubMed: 19686842]
36. Goldstein S, Lind J, Merenyi G. Chemistry of peroxynitrites as compared to peroxynitrates. *Chem. Rev.* 2005; 105:2457–2470. [PubMed: 15941219]
37. Merényi, G. b.; Lind, J.; Czapski, G.; Goldstein, S. Direct determination of the Gibbs' energy of formation of peroxynitrous acid. *Inorg. Chem.* 2003; 42:3796–3800. [PubMed: 12793816]
38. Goldstein S, Rabani J. Mechanism of nitrite formation by nitrate photolysis in aqueous solutions: the role of peroxynitrite, nitrogen dioxide, and hydroxyl radical. *J. Am. Chem. Soc.* 2007; 129:10597–10601. [PubMed: 17676847]
39. Merenyi G, Lind J, Goldstein S, Czapski G. Mechanism and thermochemistry of peroxynitrite decomposition in water. *J. Phys. Chem.* 1999; 103:5685–5691.

40. Bonini MG, Radi R, Ferrer-Sueta G, Ferreira AMDC, Augusto O. Direct EPR detection of the carbonate radical anion produced from peroxynitrite and carbon dioxide. *J. Biol. Chem.* 1999; 274:10802–10806. [PubMed: 10196155]
41. Meli R, Nauser T, Latal P, Koppenol WH. Reaction of peroxynitrite with carbon dioxide: intermediates and determination of the yield of CO₃ and NO₂. *J. Biol. Inorg. Chem.* 2002; 7:31–36. [PubMed: 11862538]
42. Goldstein S, Czapski G. Viscosity effects on the reaction of peroxynitrite with CO₂: Evidence for radical formation in a solvent cage. *J. Am. Chem. Soc.* 1999; 121:2444–2447.
43. Massari J, Tokikawa R, Zanolli L, Tavares MFM, Assuncao NA, Bechara EJH. Acetyl radical production by the methylglyoxal-peroxynitrite system: A possible route for L-lysine acetylation. *Chem. Res. Toxicol.* 2010; 23:1762–1770. [PubMed: 20923167]
44. Royer LO, Knudsen FS, De Oliveira MA, Tavares MFM, Bechara EJH. Peroxynitrite-initiated oxidation of acetoacetate and 2-methylacetoacetate esters by oxygen: Potential sources of reactive intermediates in keto acidoses. *Chem. Res. Toxicol.* 2004; 17:1725–1732. [PubMed: 15606150]
45. Imaram W, Gersch C, Kim KM, Johnson RJ, Henderson GN, Angerhofer A. Radicals in the reaction between peroxynitrite and uric acid identified by electron spin resonance spectroscopy and liquid chromatography mass spectrometry. *Free Radical Biol. Med.* 2010; 49:275–281. [PubMed: 20406679]
46. Imaram W, Johnson RJ, Angerhofer A. ESR spin trapping of the reaction between urate and peroxynitrite: the hydrogen adduct. *Appl. Magn. Reson.* 2009; 37:463–472.
47. Olson LP, Bartberger MD, Houk KN. Peroxynitrate and peroxynitrite: A complete basis set investigation of similarities and differences between these NO_x species. *J. Am. Chem. Soc.* 2003; 125:3999–4006. [PubMed: 12656637]
48. Shiri M, Zolfigol MA, Kruger HG, Tanbakouchian Z. Advances in the application of N₂O₄/NO₂ in organic reactions. *Tetrahedron.* 2010; 66:9077–9106.
49. Batinic-Haberle I, Spasojevic I, Stevens RD, Hambright P, Neta P, Okado-Matsumoto A, Fridovich I. New class of potent catalysts of O₂.bul.-dismutation. Mn(III) ortho-methoxyethylpyridyl- and di-ortho-methoxyethylimidazolylporphyrins. *Dalton Trans.* 2004:1696–1702. [PubMed: 15252564]
50. Bowler MW, Montgomery MG, Leslie AGW, Walker JE. How azide inhibits ATP hydrolysis by the F-ATPases. *Proc. Natl. Acad. Sci. USA.* 2006; 103:8646–8649.
51. Kalyanaraman B, Janzen EG, Mason RP. Spin trapping of the azidyl radical in azide/catalase/hydrogen peroxide and various azide/peroxidase/hydrogen peroxide peroxidizing systems. *J. Biol. Chem.* 1985; 260:4003–4006. [PubMed: 2984193]
52. Partridge RS, Monroe SM, Parks JK, Johnson K, Parker WD Jr, Eaton GR, Eaton SS. Spin trapping of azidyl and hydroxyl radicals in azide-inhibited rat brain submitochondrial particles. *Arch. Biochem. Biophys.* 1994; 310:210–217.
53. Amadelli R, Maldotti A, Bartocci C, Carassiti V. ESR spin-trapping investigation of azide oxidation on cadmium sulfide and zinc oxide suspension. *J. Phys. Chem.* 1989; 93:6448–6453.
54. Clarke K, Edge R, Johnson V, Land EJ, Navaratnam S, Truscott TG. Direct observation of NH₂ reactions with oxygen, amino acids, and melanins. *J. Phys. Chem. A.* 2008; 112:1234–1237. [PubMed: 18215026]
55. Kemp-Harper BK. Nitroxyl (HNO): a novel redox signaling molecule. *Antioxid. Redox Signal.* 2011; 14:1609–1613. [PubMed: 21299468]
56. Dikalov S, Skatchkov M, Fink B, Bassenger E. Quantification of superoxide radicals and peroxynitrite in vascular cells using oxidation of sterically hindered hydroxylamines and electron spin resonance. *Nitric Oxide.* 1997; 1:423–431. [PubMed: 9441913]
57. Rosen GM, Rauckman I. J. Formation and reduction of a nitroxide radical by liver microsomes. *Biochem. Pharmacol.* 1977; 26:675–678. [PubMed: 856200]
58. Finkelstein E, Rosen GM, Rauckman EJ. Spin trapping. Kinetics of the reaction of superoxide and hydroxyl radicals with nitrones. *J. Am. Chem. Soc.* 1980; 102:4994–4999.
59. Rosen, GM.; Britigan, BE.; Halpern, HJ.; Pou, S. Free radicals: Biology and detection by spin trapping. Oxford University Press; New York, New York: 1999.

60. Rangelova K, Mason RP. The fidelity of spin trapping with DMPO in biological systems. *Magn. Reson. Chem.* 2011; 49:152–158. [PubMed: 21246623]
61. Makino K, Hagiwara T, Hagi A, Nishi M, Murakami A. Cautionary note for DMPO spin trapping in the presence of iron ion. *Biochem. Biophys. Res. Commun.* 1990; 172:1073–1080. [PubMed: 2173913]
62. Ebersson L. Inverted spin trapping. Part III. Further studies on the chemical and photochemical oxidation of spin traps in the presence of nucleophiles. *J. Chem. Soc., Perkin Trans.* 1994; 2:171–176.
63. Astolfi P, Greci L, Panagiotaki M. Spin trapping of nitrogen dioxide and of radicals generated from nitrous acid. *Free Radical Res.* 2005; 39:137–144. [PubMed: 15763961]
64. Gatti RM, Alvarez B, Vasquez-Vivar J, Radi R, Augusto O. Formation of spin trap adducts during the decomposition of peroxynitrite. *Arch. Biochem. Biophys.* 1998; 349:36–46. [PubMed: 9439580]
65. Durand G, Proszak RA, Han Y, Ortial S, Rockenbauer A, Pucci B, Villamena FA. Spin trapping and cytoprotective properties of fluorinated amphiphilic carrier conjugates of cyclic versus linear nitrones. *Chem. Res. Toxicol.* 2009; 22:1570–1581. [PubMed: 19678661]
66. Labanowski, JW.; Andzelm, J. *Density Functional Methods in Chemistry*. Springer; New York, N. Y.: 1991.
67. Firsich, MJ., et al. *Gaussian 03*. Gaussian, Inc.; Pittsburgh, PA: 2004.
68. Scott AP, Radom L. Harmonic vibrational frequencies: An evaluation of Hartree-Fock, Moeller-Plesset, quadratic configuration interaction, density functional theory, and semiempirical scale factors. *J. Phys. Chem.* 1996; 100:16502–16513.
69. Cossi M, Barone V. Analytical second derivatives of the free energy in solution by polarizable continuum models. *J. Chem. Phys.* 1998; 109:6246–6254.
70. Cossi M, Barone V, Cammi R, Tomasi J. Ab initio study of solvated molecules: a new implementation of the polarizable continuum model. *Chem. Phys. Lett.* 1996; 255:327–335.
71. Reed AE, Curtiss LA, Weinhold F. Intermolecular interactions from a natural bond orbital, donor-acceptor viewpoint. *Chem. Rev.* 1988; 88:899–926.
72. Baboul AG, Curtiss LA, Redfern PC. Gaussian-3 theory using density functional geometries and zero-point energies. *J. Chem. Phys.* 1999; 110:7650–7657.
73. Izgorodina EI, Brittain DR, Hodgson JL, Krenske EH, Lin CY, Namazian M, Coote ML. Should contemporary density functional theory methods be used to study the thermodynamics of radical reactions? *J. Phys. Chem. A.* 2007; 111:10754–10768. [PubMed: 17887739]
74. Rockenbauer A, Korecz L. Automatic computer simulations of ESR spectra. *Appl. Magn. Reson.* 1996; 10:29–43.
75. Robinson KM, Beckman JS. Synthesis of peroxynitrite from nitrite and hydrogen peroxide. *Methods in Enzymol.* 2005; 396:207–214. [PubMed: 16291234]
76. Pace MD, Carmichael AJ. Quantitative EPR spin trapping. 1. Nitrogen dioxide radicals and nitrite ions from energetic materials in alkaline aqueous solution. *J. Phys. Chem. A.* 1997; 101:1848–1853.
77. Crow JP, Spruell C, Chen J, Gunn C, Ischiropoulos H, Tsai M, Smith CD, Radi R, Koppenol WH, Beckman JS. On the pH-dependent yield of hydroxyl radical products from peroxynitrite. *Free Radical Biol. Med.* 1994; 16:331–338. [PubMed: 8063196]
78. Villamena FA. Superoxide radical anion adduct of 5,5-dimethyl-1-pyrroline *N*-oxide. 6: Redox properties. *J. Phys. Chem. A.* 2010; 114:1153–1160. [PubMed: 19968309]
79. Henry DJ, Parkinson CJ, Radom L. An assessment of the performance of high-level theoretical procedures in the computation of the heats of formation of small open-shell molecules. *J. Phys. Chem. A.* 2002; 106:7927–7936.
80. Koppenol WH, Liebman JF. The oxidizing nature of the hydroxyl radical. A comparison with the ferryl ion (FeO^{2+}). *J. Phys. Chem. A.* 1984; 88:99–101.
81. Stanbury DM. Reduction potentials involving inorganic free radicals in aqueous solution. *Adv. Inorg. Chem.* 1989; 33:69–138.

82. Rockenbauer A, Györ M, Tüdös F. Spin trapping reactions with nitric oxides IV. Reactions with olefins. *Tetrahedron Lett.* 1986; 27:3425–3428.
83. Houk KN, Condroski KR, Pryor WA. Radical and concerted mechanisms in oxidations of amines, sulfides, and alkenes by peroxynitrite, peroxynitrous acid, and the peroxynitrite-CO₂ adduct: Density functional theory transition structures and energetics. *J. Am. Chem. Soc.* 1996; 118:13002–13006.
84. Villamena FA, Locigno EJ, Rockenbauer A, Hadad CM, Zweier JL. Theoretical and experimental studies of the spin trapping of inorganic radicals by 5,5- dimethyl-1-pyrroline *N*-oxide (DMPO). 2. Carbonate radical anion. *J. Phys. Chem.* 2007; 111:384–391.
85. Singh RJ, Hogg N, Joseph J, Kalyanaraman B. Mechanism of nitric oxide release from S-nitrosothiols. *J. Biol. Chem.* 1996; 271:18596–18603. [PubMed: 8702510]
86. Villamena FA, Hadad CM, Zweier JL. Comparative DFT study of the spin trapping of methyl, mercapto, hydroperoxy, superoxide, and nitric oxide radicals by various substituted cyclic nitrones. *J. Phys. Chem. A.* 2005; 109:1662–1674. [PubMed: 16833491]
87. Rojas Wahl RU. Decomposition mechanism of 3-N-morpholinosydnonimine (SIN-1)-a density functional study on intrinsic structures and reactivities. *J. Mol. Model.* 2004; 10:121–129. [PubMed: 14997366]
88. Dutton AS, Fukuto JM, Houk KN. Theoretical reduction potentials for nitrogen oxides from CBS-QB3 energetics and (C)PCM solvation calculations. *Inorg. Chem.* 2005; 44:4024–4028. [PubMed: 15907131]
89. McIntire GL, Blount HN, Stronks HJ, Shetty RV, Janren EG. Spin trapping in electrochemistry. 2. Aqueous and nonaqueous electrochemical characterizations of spin traps. *J. Phys. Chem. A.* 1980; 84:916–921.
90. Kirino Y, Ohkuma T. a. Kwan T. Spin trapping with 5,5-dimethylpyrroline-N-oxide in aqueous solution. *Chem.Pharm. Bull.* 1981; 29:29–34.
91. Chignell CF, Kalyanaraman B, Sik RH, Mason RP. Spectroscopic studies of cutaneous photosensitizing agents. II. Spin trapping of photolysis products from sulfanilamide and 4-aminobenzoic acid using 5,5-dimethyl-1-pyrroline-1-oxide. *Photochem. Photobiol.* 1981; 34:147–156.
92. Stoyanovsky DA, Clancy R, Cederbaum AI. Decomposition of sodium trioxodinitrate (Angeli's Salt) to hydroxyl radical: An ESR spin-trapping study. *J. Am. Chem. Soc.* 1999; 121:5093–5094.
93. Chignell CF, Kalyanaraman B, Mason RP, Sik RH. Spectroscopic studies of cutaneous photosensitizing agents. I. Spin trapping of photolysis products from sulfanilamide, 4-aminobenzoic acid, and related compounds. *Photochem. Photobiol.* 1980; 32:563–571.

Figure 1. Optimized geometries showing bond lengths (in Å), charge and spin densities (e) (in parentheses), and free energies of $\bullet\text{NO}_2$ addition to DMPO at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory.

Figure 2.

X-Band spectra of adducts generated from $\cdot\text{NO}/\text{O}_2$ DMSO in the presence of: (a) DMPO (30 mM), [DMPO-ONO, 95%, $a_{\text{N}} = 13.8$, $a_{\text{H}} = 11.7$; non-nitroxide, 5%, $a_{\text{H}} = 10.5$, $a_{\text{H}} = 10.5$, $a_{\text{H}} = 3.4$]; (b) AMPO (10 mM), [AMPO-ONO I, 83.4%, $a_{\text{N}} = 13.3$, $a_{\text{H}} = 10.8$, $a_{\text{N}}' = 0.6$; AMPO-ONO II, 16.6%, $a_{\text{N}} = 13.8$, $a_{\text{H}} = 17.7$, $a_{\text{N}}' = 2.4$]; (c) DEPMPO (10 mM), [DEPMPO-ONO I, 94%, $a_{\text{N}} = 13.35$, $a_{\text{P}} = 46.97$, $a_{\text{H}} = 10.93$; DEPMPO-ONO I $a_{\text{N}} = 13.51$, $a_{\text{P}} = 45.53$, $a_{\text{H}} = 12.40$; DEPMPO-R, 6%, $a_{\text{N}} = 13.85$, $a_{\text{P}} = 46.94$, $a_{\text{H}} = 20.05$]; (d) EMPO (5 mM), [EMPO-ONO I, $a_{\text{N}} = 13.22$, $a_{\text{H}} = 11.21$, $a_{\text{H}} = 0.3$; EMPO-ONO II, $a_{\text{N}} = 13.37$, $a_{\text{H}} = 12.24$, $a_{\text{H}} = 1.15$].

Figure 3. Optimized structures of *cis*-ONOO⁻, *trans*-ONOO⁻, DMPO^{•+} and spin adducts formed from DMPO^{•+} and ONOO⁻, showing the charge and spin densities (in parentheses), free energies of reaction, and pertinent bond lengths at the PCM/B3LYP/6-31+G**//B3LYP/6-31* level of theory.

Figure 4.

X-Band spectra of adducts of DMPO generated from (a) ONOO^- in 70% DMSO [$\text{CO}_3^{\bullet-}$ adduct, 80%, $a_{\text{N}} = 14.0$, $a_{\text{H}} = 9.8$, $a_{\text{H}\beta} = 1.8$; alkyl adduct, 20%, $a_{\text{N}} = 15.5$, $a_{\text{H}} = 21.5$], (b) ONOO^- in 70% DMSO under inert atmosphere [alkyl adduct, 80%, $a_{\text{N}} = 15.3$, $a_{\text{H}} = 21.5$; $\bullet\text{OH}$ adduct, 16%, $a_{\text{N}} = 15.4$, $a_{\text{H}} = 15.0$; DMPO-ONX, 4%, $a_{\text{N}} = 9.4$, $a_{\text{H}} = 19.9$], (c) ONOO^- in 70% DMSO under CO_2 atmosphere [$\text{CO}_3^{\bullet-}$ adduct, 93%, $a_{\text{N}} = 14.0$, $a_{\text{H}} = 9.8$, $a_{\text{H}\beta} = 1.8$; alkyl adduct, 7.4%, $a_{\text{N}} = 15.6$, $a_{\text{H}} = 21.2$], (d) ONOO^- in 100% PBS under inert atmosphere [alkyl adduct, 27%, $a_{\text{N}} = 16.0$, $a_{\text{H}} = 22.6$; DMPO-NX, 57%, $a_{\text{N}} = 14.5$, $a_{\text{N}'} = 3.55$, $a_{\text{H}} = 0.93$; DMPO-ONO, 6%, $a_{\text{N}} = 14.4$, $a_{\text{H}} = 11.3$].

Figure 5. Optimized structures of ONOOH and energetics of various modes of ONOOH addition to DMPO, showing the charge densities (in parentheses), and pertinent bond lengths at the PCM/B3LYP/6-31+G**//B3LYP/6-31* level.

Figure 6.

Optimized structures of ONOOCO_2^- and the various spin adducts formed with DMPO showing charge and spin densities (in parentheses), and pertinent bond lengths at the PCM/B3LYP/6-31+G**//B3LYP/6-31* level of theory.

Figure 7.

X-Band spectra generated from visible light irradiation of DMPO (100 mM) and SNAP (65 mM) in aqueous solution: (a) under ambient atmosphere, HO• adducts: (major) $a_N = 15.2$, $a_H = 14.8$; (minor) $a_N = 15.8$, $a_H = 15.2$; (b) under Ar atmosphere, RS• adduct: $a_N = 14.6$, $a_H = 16.0$ (lit. value for DMPO-SNAP: $a_N = 15.3$, $a_H = 17.7$).⁸⁵

Figure 8.

Optimized structures of $\bullet\text{NO}_3$, $\bullet\text{NH}_2$, $\bullet\text{N}_3$ and their respective adducts with DMPO showing the free energies of reaction, charge and spin densities (in parentheses), and pertinent bond lengths at the PCM/B3LYP/6-31+G**//B3LYP/6-31* level of theory.

Figure 9.

X-Band spectra in PBS of DMPO adducts of (a) $\cdot\text{N}_3$: $a_{\text{N}} = 14.75$, $a_{\beta\text{-H}} = 14.18$, $a_{\text{N}'} = 3.4$ (lit. value for DMPO- N_3 : $a_{\text{N}} = 14.8$, $a_{\text{H}} = 14.2$ and $a_{\text{N-azide}} = 3.1$)⁵¹; and (b) $\cdot\text{NH}_2$: $a_{\text{N}} = 15.75$, $a_{\beta\text{-H}} = 19.30$, $a_{\text{N}'} = 1.62$; (lit. value for DMPO- NH_2 : $a_{\text{N}} = 15.9$, $a_{\text{H}} = 19.3$ and $a_{\text{N-amino}} = 1.60$).⁹³

($\Delta G_{\text{rxn}} = 16.2$ kcal/mol)

DMPO- NO^-

($\Delta G_{\text{rxn}} = -2.2$ kcal/mol)

DMPO-NOH

Figure 10.

Optimized structures of ${}^3\text{NO}^-$, HNO and their respective addition products (*i.e.*, nitroso-oximate and nitroso-oxime, respectively) with DMPO showing the free energies of reaction, charge densities (in parentheses), and pertinent bond lengths at the PCM/B3LYP/6-31+G**//B3LYP/6-31* level of theory.

Scheme 1.
Formation of RNS from $\cdot\text{NO}$.

Chem Res Toxicol. Author manuscript; available in PMC 2013 August 20.

Scheme 3.

Proposed reaction mechanisms of ONOO^- addition to DMPO and their respective $\Delta G_{298\text{K}, \text{rxn}}$ (in kcal/mol) calculated at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory.

Scheme 4.

Proposed reaction mechanisms of ONOO^- with DMPO with the respective $\Delta G_{298\text{K},\text{rxn}}$ calculated at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory.

Scheme 5.

Proposed reaction mechanisms for the formation of nitron spin adducts with ONOOCO_2^- , and their respective $\Delta G_{298\text{K},\text{rxn}}$ (in kcal/mol) calculated at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory.

Scheme 6.

Proposed reaction mechanisms for the decomposition of SNAP in the presence of DMPO and their respective $\Delta G_{298K,rxn}$ (in kcal/mol) calculated at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory.

Scheme 7.

Proposed decomposition mechanisms for DMPO-NO₃ and their respective $\Delta G_{298\text{K},\text{rxn}}$ (in kcal/mol) calculated at the PCM/B3LYP/6-31+G**//B3LYP/6-31G* level of theory.

Table 1

Free Energies of Reduction, $\Delta G_{\text{aq},298\text{K}}$ (in kcal/mol), of Selected RNS and ROS at the PCM(water)/B3LYP/6-31+G**//B3LYP/6-31G* and PCM(water)/G3(MP2)//B3LYP/6-31G* (in parentheses) Levels of Theory.

Entry	$\Delta G_{298\text{K},\text{red}}$
$\text{ONOOCO}_2^- + 2\text{e}^- / \text{NO}_2^- + \text{CO}_3^{2-}$	-242.5 (-236.5)
$\text{N}_2\text{O}_4 + 2\text{e}^- / 2\text{NO}_2^-$	-240.3 (-232.1)
$\text{cis-ONOOH} + 2\text{e}^- / \text{HO}^- + \text{NO}_2^-$	-237.1 (230.3)
$\text{trans-ONOOH} + 2\text{e}^- / \text{HO}^- + \text{NO}_2^-$	-235.6 (-231.8)
$\text{trans-ONOO}^- + 2\text{e}^- / \text{O}^{2-} + \text{NO}_2^-$	-170.5 (-158.0)
$\text{cis-ONOO}^- + 2\text{e}^- / \text{O}^{2-} + \text{NO}_2^-$	-166.3 (-153.9)
$\text{NO}_3 + \text{e}^- / \text{NO}_3^-$	-152.3 (-155.0)
$\text{NO}^+ + \text{e}^- / \text{NO}$	-146.6 (-140.7)
$\text{DMPO}^{*+} + \text{e}^- / \text{DMPO}$	-139.7 (-147.8)
$\text{N}_3 + \text{e}^- / \text{N}_3^-$	-124.1 (-121.0)
$\text{NO}_2 + \text{e}^- / \text{NO}_2^-$	-120.4 (-116.8)
$\text{OH} + \text{e}^- / \text{HO}^-$	-118.9 (-120.0)
$\text{CO}_3^{*+} + \text{e}^- / \text{CO}_3^{2-}$	-103.3 (-115.9)
$\text{N}_2\text{O}_4 + \text{e}^- / \text{N}_2\text{O}_4^{*-}$	-102.0 (-115.3) ^a
$\text{HO}_2^* + \text{e}^- / \text{HO}_2^-$	-97.2 (-97.0)
$\text{cis-ONOOH} + \text{e}^- / \text{cis-ONOOH}^{*-}$	-86.5 (-78.6)
$\text{NH}_2 + \text{e}^- / \text{NH}_2^-$	-85.9 (-89.9)
$\text{trans-ONOOH} + \text{e}^- / \text{trans-ONOOH}^{*-}$	-83.3 (-80.0) ^b
$\text{NO} + \text{e}^- / \text{}^3\text{NO}^-$	-82.6 (-69.0)
$\text{ONOOCO}_2^- + \text{e}^- / \text{ONOOCO}_2^{*2-}$	-81.7 (-68.1)
$\text{trans-ONOO}^- + \text{e}^- / \text{trans-ONOO}^{*2-}$	-62.8 (-50.0)
$\text{cis-ONOO}^- + \text{e}^- / \text{cis-ONOO}^{*2-}$	-55.7 (-45.9) ^b
$\text{O}_2^{*+} + \text{e}^- / \text{O}_2^{2-}$	-53.7 (-49.6)

^a Calculation using NO_2^- and $\text{}^*\text{NO}_2$ as products for G3(MP2) only and could be an overestimate.

^b Energies for cis-ONOOH^{*-} and trans-ONOO^{*2-} were used for the G3(MP2) calculation only.

Table 2

Predicted Hyperfine Splitting Constants (G) of RNS Adducts of DMPO and their Respective Free Energies, $\Delta G_{298K,rxn}$ (in kcal/mol), at the PCM(water)/B3LYP/6-31+G**//B3LYP/6-31G* Level of Theory. (Values in Parentheses are in the Presence of Two Explicit Water Molecules Calculated at the Same Level of Theory as Above)^a

Adducts	predicted hyperfine splitting constants (G)						ΔG_{rxn}
	a_N	$a_{\beta H}$	$a_{\gamma H}$	a_N (adduct)	$a_{N''}$ for AMPO	a_P for DEPMPO	
-OONO ^b	11.0 (10.4)	11.8 (14.2)	0.6 (0.7)	<i>c</i>	<i>c</i>		-26.3
-N(O)O ₂ ^b	11.0	15.7	<i>c</i>	5.8			-0.5
-NO	10.0 (10.3)	17.2 (19.7)	0.8 (0.7)	2.6 (7.1)	<i>c</i>		14.1
-ONO	10.7 (10.2)	7.3 (9.3)	1.5 (^c)	<i>c</i>	<i>c</i>		1.2
-NO ₂	9.6 (10.2)	7.8 (7.6)	1.5 (1.9)	8.0 (8.4)	<i>c</i>		-1.4
-NH ₂	11.9 (12.5)	12.1 (11.0)	0.5 (0.9)	1.6 (2.0)			-28.3
-N ₃	11.7 (12.1)	7.4 (8.1)	0.8 (0.7)	3.2 (3.0)	<i>c</i>		-13.4
-SNAP	11.1 (11.8)	9.0 (8.8)	0.9 (1.1)				-1.1

^aFor complete list of hyperfine splitting constants of various adduct of AMPO, EMPO and DEPMPO, see the supplementary information (Table S1).

^bFormed from the inverted spin trapping of ONOO⁻ by [nitronel]^{•+}.

^cNot applicable or not significant.