Published in final edited form as:

Ann N Y Acad Sci. 2011 September ; 1233: 58–63. doi:10.1111/j.1749-6632.2011.06130.x.

The effects of ion channel blockers validate the conductance-based model of saccadic oscillations

Aasef G. Shaikh¹, David S. Zee², Lance M. Optican³, Kenichiro Miura⁴, Stefano Ramat⁵, and R. John Leigh¹

¹Department of Neurology, Case Western Reserve University, Cleveland, Ohio ²Department of Neurology, The Johns Hopkins University, Baltimore, Maryland ³Laboratory of Sensorimotor Research, National Eye Institute, National Institutes of Health, Bethesda, Maryland ⁴Kyoto University, Kyoto, Japan ⁵University of Pavia, Pavia, Italy

Abstract

Conductance-based models of reciprocally inhibiting burst neurons suggest that intrinsic membrane properties and postinhibitory rebound (PIR) determine the amplitude and frequency of saccadic oscillations. Reduction of the low-threshold calcium currents (I_T) in the model decreased the amplitude but increased the frequency of the simulated oscillations. Combined reduction of hyperpolarization-activated cation current (I_h) and I_T in the model abolished the simulated oscillations. We measured the effects of a selective blocker of I_T (ethosuximide) in healthy subjects on the amplitude and frequency of saccadic oscillations evoked by eye closure and of a nonselective blocker of I_h and I_T (propranolol) in a patient with microsaccadic oscillation and limb tremor syndrome (mSOLT). Ethosuximide significantly reduced the amplitude but increased the frequency of the saccadic oscillations during eye closure in healthy subjects. Propranolol abolished saccadic oscillations in the mSOLT patient. These results support the hypothetical role of postinhibitory rebound, I_h , and I_T , in generation of saccadic oscillations and determining their kinematic properties.

Keywords

burst neurons; hyperpolarization-activated cation current; low-threshold calcium current; reciprocal innervations

Introduction

Reciprocal innervation of the agonist-antagonist muscle pair and transient removal of the external inhibition generating high-frequency pulse of the neural discharge (postinhibitory rebound, PIR) are the fundamental drives for high velocities of ballistic movements.¹ For example, during saccades, the omnipause neurons (which inhibit the burst neurons during

© 2011 New York Academy of Sciences.

Address for correspondence: Aasef G. Shaikh, M.D., Ph.D., Department of Neurology, University Hospitals Case Medical Center, 11100 Euclid Avenue, Cleveland, OH 44106-5040. aasefshaikh@gmail.com.

Conflicts of interest

The authors declare no conflicts of interest.

Preferred citation: Shaikh, A.G., D.S. Zee, L.M. Optican, K. Miura, S. Ramat & R.J. Leigh. 2011. The effects of ion channel blockers validate the conductance-based model of saccadic oscillations. In Basic and Clinical Ocular Motor and Vestibular Research. Janet Rucker & David S. Zee, Eds. *Ann. N.Y. Acad. Sci.* **1233**: 58-63.

steady fixation) transiently turn off, allowing the ipsilateral excitatory burst neurons (EBNs) to send an excitatory signal to the motoneurons, which innervates the agonist muscle that rotates the eyes to the ipsilateral side.² Simultaneously, the ipsilateral inhibitory burst neurons (IBNs) inhibit the motoneurons to the antagonist muscle.² It was proposed that PIR generated by the transient inhibition of the OPNs is critical for the bursts of neural discharge in EBNs and IBNs, which provide sufficient force to rapidly turn the globe.³ During steady gaze, OPNs inhibit EBNs and IBNs on both sides and prevent unwanted saccades.

A conductance-based neuromimetic model predicted that increased excitability of the burst neurons could result in saccadic oscillations.³ The excitability could be increased either by the removal of inhibition from the OPNs (during eye closure in healthy subjects or decreased glycinergic inhibition during pathological states) or due to pathological depolarizing shifts of the resting membrane potential of the burst neurons (e.g., intoxication, autoimmune disorders, and degenerative cerebellar disorders).³⁻⁶ Hyperpolarization-activated, inward cation current (I_h) and low-threshold calcium current (I_T) are key determinants of the resting membrane potential and the strength of PIR.⁷⁻¹¹ In the conductance-based model of the burst neurons, the activation kinetics and amplitudes of I_h and I_T determined the amplitudes and the frequencies of saccadic oscillations. An increase in I_T increased the amplitude and decreased the frequency, while increasing I_h increased the frequency of saccadic oscillations (Fig. 1).³

These model predictions motivated two hypotheses: selective blockade of I_T should reduce the amplitude but increase the frequency of the saccadic oscillations, and nonselective blockers of I_h and I_T ^{7,8} should reduce the amplitude and frequency of saccadic oscillations. To test these hypotheses, we measured the effects of ethosuximide, a selective I_T blocker,¹² on the properties of physiological saccadic oscillations during eyelid closure in two healthy subjects and the effects of propranolol, a nonselective blocker of I_h and I_T , in a patient with syndrome of microsaccadic oscillations and limb tremor (mSOLT).³

Methods

The mSOLT patient and healthy subjects were evaluated at the Johns Hopkins vestibular laboratory. All subjects gave informed consent. The eye movements of the healthy subjects were recorded with the magnetic-field system using dual search coils (Skalar Medical, Delft, the Netherlands). A search-coil system and an infrared camera (Ober Consulting, Poland) were used to measure eye movements of the mSOLT patient.

The subjects fixed their gaze upon a target that was projected straight ahead. Eye movements were recorded both when looking at a stationary target and in a completely dark room with no visual target. Eye movements in healthy subjects were then recorded with both eyes closed but while fixating on the imaginary stationary target. Eye position signals were sampled at 1000 Hz resolution. The data were further processed with custom software written in the Matlab® environment.

Data from each axis were processed separately. Saccadic oscillations were best characterized from eye velocity records. Oscillations were separated interactively. Cycle-by-cycle analysis was performed in the following steps. The detrended data were normalized with respect to the mean velocity (i.e., normalized velocity = actual velocity – mean velocity). Normalization allowed the velocity to align on the abscissa with the peaks of the cycles remaining positive and the troughs negative. The points of intersection of the velocity data trace, crossing from a negative value to a positive value with the abscissa were recorded. The x coordinates of these values were called positive zero crossings. The cycle duration was calculated from the time between the positive zero crossings. The inverse of

the cycle duration yields the cycle frequency. The average of the peak and trough values of the cycle was considered as the peak velocity of the given cycle. The mean, median, standard deviation, and 95% confidence intervals of the frequencies and peak velocities of all cycles were computed. Saccades were interactively selected in the eye position trace. Saccadic amplitude, velocity, latency, and gain (actual/desired eye movement) were computed.

Results

Effect of ethosuximide on saccadic oscillations during eyelid closure in healthy subjects

The black traces in Fig. 2 illustrate saccadic oscillations in two subjects with the eyes closed. These oscillations were present around all three axes. Eye movements during attempted fixation on the imaginary target with the eyes closed were recorded in the same subject 45 min after taking 250 mg ethosuximide by mouth (gray traces in Fig. 2). Saccadic oscillations around all three axes were again seen under closed eyelids; however, their amplitude was reduced in horizontal and torsional planes in this example. Frequency was relatively increased.

Figure 3 summarizes the frequency and amplitudes of horizontal, vertical, and torsional saccadic oscillations during eye closure in two healthy subjects. Each gray box-whisker plot represents the effects of ethosuximide on the frequency and amplitude of the saccadic oscillations. The horizontal line in each box and whisker plot represents the median value, while notches represent 95% confidence intervals. The frequency of oscillations increased around all three axes after ethosuximide; however, the difference was not statistically significant (one-way ANOVA, $P > 0.05$). In contrast, there was a statistically significant reduction in the amplitude of horizontal and torsional saccadic oscillations in subject 1 and saccadic oscillations around all three axes in subject 2 (one-way ANOVA, $P < 0.05$).

Effect of propranolol on saccadic oscillations in a patient with mSOLT

A young patient with mSOLT was referred to our clinic for occasional brief episodes of blurred vision and fine tremor of the hand. The symptoms were present since early childhood, and were worsened during periods of stress or anxiety. On bedside examination, intermittent bursts of saccadic oscillations and limb tremor were the only neurological findings. Ophthalmoscopy revealed nearly continuous, small-amplitude, high-frequency oscillations of the optic disc. Subsequently, the patient's mother was evaluated and reported similar episodes of blurred vision. On examination, the mother had the same saccadic oscillations and tremor. During a subsequent visit to see the mother, she reported that her primary care provider prescribed propranolol (60 mg by mouth twice a day) for hypertension. The mother now had relief from her intermittent episodes of blurred vision and hand tremor. At this time, the saccadic oscillations were no longer present as examined during ophthalmoscopy.

The black trace in Fig 4 illustrates saccadic oscillations during attempted steady fixation recorded from the mother prior to beginning propranolol. These oscillations were 23.2 18.3 Hz in frequency and 76.2 ± 29.4 degrees/second $\pm$ in amplitude. The treatment with propranolol abolished these oscillations (gray trace in Fig. 4).

Discussion

We tested the prediction of a neuromimetic model of the saccade generators in which the rapid velocities of saccades are due an abrupt increase in the firing rate, the burst, in reciprocally innervating EBNs and IBNs.^{3,13} The model also predicted that a pause in the external inhibition is critical for the high firing rate during the burst, a mechanism known as

PIR. Simulation of the pathological enhancement of PIR, by increasing I_h and/or I_T , revealed high-frequency back-to-back saccades (i.e., saccadic oscillations). The model also predicted that enhanced I_h would increase the frequency of saccadic oscillations, without affecting the amplitude (Fig. 1). In contrast, enhanced I_T would increase the amplitude of saccadic oscillations but decrease their frequency (Fig. 1).³ Our strategy for testing these model predictions was to determine if a selective blocker of I_T , ethosuximide,¹ reduced the amplitude of physiological saccadic oscillations and increased their frequency. We also determined whether nonselective blockade of I_h and I_T , by propranolol,^{7,8} would reduce the amplitude of oscillations. Our findings were consistent with both model predictions; the ethosuximide reduced the amplitude but increased the frequency of saccadic oscillations in both the normal subjects, and propranolol reduced the amplitude of oscillations in the patient with mSOLT.

The model also predicted that the frequency of oscillations is less sensitive to changes in I_T than is the amplitude. Figure 1 depicts that, in the model, for the given change in I_T , the amplitude of simulated saccadic oscillations is affected more than the frequency. Consistent with this prediction, the amount of change in the amplitude of oscillations was larger and reached statistical significance as compared to the amount of change in the frequency in both subjects.

There was an inter-axis variability in the effects of ethosuximide in both subjects. This variability could be explained, in part, by differences in the expression profile of CaV3 ion-channels (conducting I_T) at anatomically separate brainstem sites of the burst neurons controlling vertical, torsional, and horizontal saccades. The burst neurons for horizontal saccades are found in caudal pons, while those for vertical and torsional movements are located in rostral mesencephalon.¹⁴

The conductance-based model predicted that mSOLT is caused by an inherited abnormality causing increased membrane excitability or reduced external inhibition of the pontine burst neurons.³ Our results provided support for this model, since non-selective blockade of I_h and I_T with propranolol could theoretically reduce the excitability of the burst neurons, and, in turn, account for the reduction of the amplitude of saccadic oscillations in the mSOLT patient.

According to the model predictions, blockade of I_h and I_T should reduce PIR and therefore reduce saccade velocity.^{3,15} However, in both our normal subjects, there was no change in saccade velocity. In our model, the strength of PIR required to simulate oscillations had to be supra-physiological; however, to simulate slow saccades, PIR must be significantly less than physiological. Therefore, it is possible that at the given doses, ethosuximide and propranolol might reduce PIR, but not enough to affect saccade velocity.

The recordings in the healthy subjects were done with scleral search coils, under closed eyelids. Does mechanical friction due to lid closure affect the eye velocity or induce coil slippage? We had addressed this issue in an independent study where we had shown that the acceleration and velocity of the eye movement during vestibulo-ocular reflex evoked by head impulses (the mechanism that is not physiologically affected by eye closure) were the same during open and closed eyelids.¹⁶ In the same study, the peak velocities or accelerations of the eye movements evoked by vestibulo-ocular reflex during head impulses are of comparable speed to those of saccades.¹⁶ Therefore, it is unlikely that eye closure induced mechanical hindrance to the movement of the eye or of the search coil mounted on it.

To summarize, the conductance-based model emphasized that increased excitability or disinhibition of the pontine burst neurons can cause saccadic oscillations. The results of the

current experiments are in accord with the specific model predictions and in particular with the hypothesis that I_h and I_T conductance determines the frequency and amplitude of the saccadic oscillations. There are, however, caveats. The drugs could have effects on other structures related to saccade generation that might influence the frequencies and amplitudes of saccadic oscillations. Electrophysiological experiments investigating the effects of selective ion channel blockers on saccades, injected locally at the location of the burst neurons or omnipause neurons, are needed to further test these ideas.

Acknowledgments

The authors thank Mr. Dale Roberts for assistance. This work was supported by grants from NIH EY01849, EY06717, the Department of Veterans Affairs, and the EvenorArmington Fund.

References

- Sherrington CS. On reciprocal innervation of antagonistic muscles—tenth note. *R. Soc. Proc.* 1907; 52:337–349.
- Leigh, RJ.; Zee, DS. *The Neurology of Eye Movements*. 4th ed. Contemporary neurology series; New York: 2006.
- Shaikh AG, Miura K, Optican LM, et al. A new familial disease of saccadic oscillations and limb tremor provides clues to mechanisms of common tremor disorders. *Brain*. 2007; 130:3020–3031. [PubMed: 17921180]
- Shaikh AG, Ramat S, Optican LM, et al. Saccadic burst cell membrane dysfunction is responsible for saccadic oscillations. *J. Neuroophthalmol.* 2008; 28:329–336. [PubMed: 19145136]
- Shaikh AG, Marti S, Tarnutzer AA, et al. Gaze fixation deficits and their implication in ataxia-telangiectasia. *J. Neurol. Neurosurg. Psych.* 2009; 80:858–864.
- Shaikh AG, Hain TC, Zee DS. Oculomotor disorders in adult-onset Still's disease. *J. Neurol.* 2010; 257:136–138. [PubMed: 19756823]
- McCormick DA, Pape HC. Noradrenergic and serotonergic modulation of a hyperpolarization-activated cation current in thalamic relay neurones. *J. Physiol.* 1990; 431:319–342. [PubMed: 1712844]
- Pape HC, McCormick DA. Noradrenaline and serotonin selectively modulate thalamic burst firing by enhancing a hyperpolarization-activated cation current. *Nature*. 1989; 340:715–718. [PubMed: 2475782]
- Shaikh AG, Finlayson PG. Excitability of auditory brainstem neurons, in vivo, is increased by cyclic-AMP. *Hear Res.* 2005; 201:70–80. [PubMed: 15721562]
- Shaikh AG, Finlayson PG. Hyperpolarization-activated (I_h) conductances affect brainstem auditory neuron excitability. *Hear Res.* 2003; 183:126–136. [PubMed: 13679144]
- Perez-Reyes E. Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol. Rev.* 2003; 83:117–161. [PubMed: 12506128]
- Coulter DA, Huguenard JR, Prince DA. Characterization of ethosuximide reduction of low-threshold calcium current in thalamic neurons. *Ann. Neurol.* 1989; 25:582–593. [PubMed: 2545161]
- Ramat S, Leigh RJ, Zee DS, et al. Ocular oscillations generated by coupling of brainstem excitatory and inhibitory saccadic burst neurons. *Exp. Brain Res.* 2005; 160:89–106. [PubMed: 15289966]
- Buttner-Ennever JA. Mapping the oculomotor system. *Prog. Brain Res.* 2005; 171:3–11. [PubMed: 18718275]
- Miura K, Optican LM. Membrane channel properties of premotor excitatory burst neurons may underlie saccade slowing after lesions of omnipause neurons. *J. Comput. Neurosci.* 2006; 20:25–41. [PubMed: 16511656]
- Shaikh AG, Wong AL, Optican LM, et al. Sustained eye closure slows saccades. *Vis. Res.* 2010; 50:1665–1675. [PubMed: 20573593]

Figure 1.

The effect of I_h and I_T on the frequency and amplitude of the simulated saccadic oscillations. Increasing I_T increases the amplitude but decreases the frequency of simulated saccadic oscillations. For the given change in I_T , the amplitudes of saccadic oscillations are more affected than their frequency. Hence, the amplitude is more sensitive to changes in I_T as compared to I_h . Increasing I_h , however, increases the frequency of saccadic oscillations.

Figure 2.

Horizontal, vertical, and torsional saccadic oscillations evoked by eye closure in two healthy subjects (black traces). The eye velocity is plotted along the y -axis, while the time is plotted on the x -axis. Ethosuximide reduces the amplitude and increases the frequency of these saccadic oscillations in both subjects (gray trace).

Figure 3.

Box-whisker plots summarizing the effects of ethosuximide on cycle-by-cycle frequency and amplitude of the saccadic oscillations. The horizontal line in the center of notch represents the median value, the notch represents the 95% confidence interval, the length of the box depicts the inter-quartile interval, and the whiskers represent the range. Nonoverlapping notches represent statistically significant differences between the medians of the compared populations.

Figure 4.

Horizontal saccadic oscillations as the patient (the mother) with microsaccadic oscillation limb tremor (mSOLT) attempts to fixate on a target (black traces). Eye velocities are plotted on the y -axis, while the time is plotted along the x -axis. Propranolol almost abolished the saccadic oscillations in the patient (the mother) with mSOLT (gray trace).