Published in final edited form as:

J Acad Nutr Diet. 2012 September ; 112(9): 1390–1396. doi:10.1016/j.jand.2012.05.011.

Beverage displacement between elementary and middle school, 2004–2007

Reena Oza-Frank, PhD, RD,

Research Institute at Nationwide Children's Hospital, Department of Pediatrics, The Ohio State University, 700 Children's Drive, W254, Columbus, OH 43205, Phone: (614) 355-3530, Fax: (614) 355-3455

Madeline Zavodny, PhD, and

Department of Economics, Agnes Scott College, 141 E. College Ave. Decatur, GA 30030, Phone: (404) 471-6377, Fax: (404) 471-5478

Solveig A. Cunningham, PhD

Hubert Department of Global Health, Rollins School of Public Health, Emory University, 1518 Clifton Rd, NE, 7045, Atlanta GA 30322, Phone: (404) 727-6486, Fax: (404) 727-4590 (fax)

Reena Oza-Frank: reena.oza-frank@nationwidechildrens.org; Madeline Zavodny: mzavodny@agnesscott.edu; Solveig A. Cunningham: sargese@sph.emory.edu

Abstract

Background—Sweetened beverages of low nutritional quality may be displacing more nutritious beverages, such as 100% fruit juice and milk, from children's diets. However, previous research has not examined changes in beverage intake over time using nationally representative longitudinal data.

Objective—To examine changes in sweetened beverage, milk, and juice consumption between 5th and 8th grade.

Design—Longitudinal analysis of self-reported beverage consumption.

Participants/setting—Data were from the Early Childhood Longitudinal Survey-Kindergarten Cohort 1998–1999 (ECLS-K), a nationally representative study of children followed from kindergarten through 8th grade. This analysis used data from Spring 2004 and 2007, when most children were in 5th and 8th grade, respectively (n=7,445).

Main outcome measures—Changes in consumption of sweetened beverages, milk, and 100% fruit juice last week.

Statistical analyses performed—Survey-adjusted linear regression was used to estimate longitudinal relationships between servings of milk, sweetened beverages, and 100% fruit juice, controlling for child and family characteristics and food consumption.

Results—Children's milk consumption decreased between 5th and 8th grade, and these decreases were larger among children who drank sweetened beverages daily. However, after controlling for demographic characteristics, changes in children's milk consumption were not significantly related to changes in their consumption of sweetened beverages over time ($\beta=0.005$, $p=0.81$),

Correspondence to: Reena Oza-Frank, reena.oza-frank@nationwidechildrens.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

while changes in milk consumption were positively related to changes in juice consumption ($\beta=0.087$, $p<0.01$).

Conclusions—Observed decreases in average milk consumption from 5th to 8th grade were not related to changes in sweetened beverage consumption. They were positively related to changes in fruit juice consumption, so not indicating displacement. Caloric beverages generally tended to increase or decrease in tandem, so focus must be placed on their role in children's entire diet and on balancing them with food and total beverage intake.

Keywords

beverage; displacement; elementary school; middle school; ECLS-K; sweetened beverage; milk; juice; longitudinal

Introduction

Patterns of beverage consumption among children and adolescents in the US suggest that sugar-sweetened beverages of low nutritional quality may be replacing more nutritious beverages, such as 100% fruit juice and milk, in children's diets (1–4). Specifically, national cross-sectional data indicate that milk consumption decreased among children ages 2 to 18 years between 1977 and 2001 while soda consumption more than doubled during that period (5). The simultaneous timing of these changes suggests that sugar-sweetened beverages may have displaced more nutritious drinks in children's diets.

Understanding whether less nutritious beverages are displacing more nutritious beverages is important because such displacement may have negative long-term health consequences. For example, soda consumption has been associated with lower dietary intake of vitamin A and C, calcium, magnesium and riboflavin in children and adolescents (1, 6). Increases in soft drinks and fruit juice in children's diet are linked to increases in their total caloric intake, which can lead to weight gain and increased obesity risks (7–14). Obese children are more likely to become obese adults, (15) increasing their risk of other chronic conditions, including hypertension, type 2 diabetes, and premature cardiovascular disease (15, 16).

Several studies have examined whether sweetened beverages displace milk and 100% fruit juice from children's diets. A study in Louisiana found that 10-year olds who consumed large quantities of sweetened beverages drank less milk than did those who consumed few or no sweetened beverages (17). A longitudinal study that examined annual 3-day food and beverage diaries among children in Iowa from age 1 to 5 years found that milk intake was inversely associated with intake of sweetened beverages at almost every age examined (18). A longitudinal study of 24-hour diet recalls among elementary school children in Nebraska found that, over a 2-year period, sugar-sweetened beverages displaced milk in their diets (8). Another study found that as students in Minnesota moved from elementary to junior high or middle school, their consumption of milk decreased while their consumption of soft drinks increased (19).

These studies are suggestive of beverage displacement in children. However, to date, no study has examined beverage displacement using nationally representative longitudinal data. The objective of this study was to examine longitudinal changes in sweetened beverages, milk, and juice consumption between 5th and 8th grade using a large, nationally representative study, the Early Childhood Longitudinal Study, Kindergarten Class of 1998–99 (ECLS-K). Approximately 90% of children aged 2–19 consume sweetened beverages or fruit juice and at least 70% consume milk, making these beverages the most common caloric beverages consumed by school-aged children (20).

Materials and Methods

Data source

The data for this study are from the Early Childhood Longitudinal Study, Kindergarten Class of 1998–99 (ECLS-K), which followed a nationally representative cohort of almost 20,000 children in the United States from kindergarten to 8th grade. The ECLS-K was developed by the National Center for Education Statistics (NCES) of the US Department of Education (21). Multistage probability sampling was used to select a nationally representative sample of children who were in kindergarten in 1998–99 or in 1st grade in 1999–2000. The primary sampling units were counties and groups of counties, with a 2nd stage selection of schools within the sampled primary sampling units and a 3rd stage unit of students within schools.

The ECLS-K included interviews with parents, teachers, and school administrators and direct anthropometric assessments of the children. In the 5th and 8th grade, children filled out a food consumption questionnaire that had 19 questions about food and beverages developed by the US Department of Agriculture (USDA) based on the Youth Risk Behavior Surveillance System (YRBSS) and California Children's Eating and Exercise Practices Survey (CalCHEEPS). Evidence of validity of these questions has been previously published (22).

This study used data from the 6th and 7th waves of data collection, administered in Spring 2004 and 2007, when most children were in 5th and 8th grade, respectively. Most sample attrition from the cohort resulted from random selection for non-sampling due to survey costs, maintaining the nationally representative nature of the data (23). We used data on all 7,445 children with complete data (no missing responses to beverage consumption questions, age, sex or race/ethnicity) in both the 5th and 8th grade. This study was conducted using publicly available data and was determined to be exempt by Emory University's Institutional Review Board (IRB).

Measures

Children were asked the following questions about their drink consumption over the past 7 days:

- “How many times did you drink Soda pop (EXAMPLES Coke, Pepsi, Mountain Dew), sports drinks (EXAMPLE Gatorade), or fruit drinks that are not 100% fruit juice (EXAMPLES Kool-Aid, Hi-C, Fruitopia, Fruitworks)?”
- “How many glasses of milk did you drink? (Include all types of milk, including cow's milk, soymilk or any other kind of milk; include the milk you drank in a glass or cup, from a carton, or with cereal. Count the half pint of milk served at school as equal to one glass.)”
- “How many times did you drink 100% fruit juices such as orange juice, apple juice, or grape juice? (Do not count punch, Kool-Aid, sports drinks, or other fruit-flavored drinks).”

For the sweetened beverage and juice questions, children chose one of the following: none, 1–3 times during the last week, 4–6 times during the last week, 1 time per day, 2 times per day, 3 times per day, or 4 or more times per day. For the milk question, children chose one of the following: 1–3 glasses during the last week, 4–6 glasses during the last week, 1 glass per day, 2 glasses per day, 3 glasses per day, or 4 or more glasses per day. Based on the responses to the above questions, we created three measures of children's consumption of each type of beverage: whether they reported consuming any, whether they reported

consuming it at least daily, and the number of servings they reported consuming during the last week. We calculated servings last week by using the mean value for categories (e.g., 1–3 times per week was coded as 2 servings per week). For children who reported daily consumption, we multiplied frequencies or servings per day (1 through 4) by 7.

Control variables

The regression models controlled for parent-reported child characteristics that may be related to beverage intake. These characteristics included sex, race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic, Asian or Pacific Islander, American Indian, and multiracial, with non-Hispanic white as the omitted category), and socioeconomic status (SES). SES was based on an ECLS-K-created variable reflecting parental income, education, and occupational prestige, based on parent reports; indicator variables for 4 of 5 quintiles were used. Preliminary analyses indicated associations between sweetened beverage consumption and public versus private school attendance, whether the child eats school lunch regularly, eats school breakfast regularly, or receives free or reduced price lunch at school. These variables therefore were also included. Because food intake is related to beverage intake, the other nutrition indicators collected in the ECLS-K were also included: changes in consumption of salad, potatoes, carrots, other vegetables, fruit, and fast food, as reported by children. Additional models were stratified by sex, race/ethnicity, or poverty status to determine if displacement occurred in specific demographic subgroups. The household poverty variable was created by the NCES by comparing income with Census poverty thresholds, with households whose income fell below the threshold classified as living in poverty. The results reported here were robust to including other potentially relevant characteristics: number of days per week the child is active for at least 20 minutes, the number of hours of television watched per week, body mass index, and family characteristics (whether the mother works full time, whether other children live in the household, the number of days the family eats dinner together per week, and the number of days the family eats breakfast together per week).

Statistical methods

T-tests were used to test differences in demographic characteristics. Ordinary least squares (OLS) regressions with longitudinal data were used to estimate the relationship between changes in children's consumption of milk, sweetened beverages, and 100% fruit juice during the previous week. The results indicate how changes in consumption were related, controlling for other characteristics. P-values < 0.05 are indicated as significant in the text below.

The dummy variable adjustment method was used for the small number of observations with missing data on control variables. This method involves creating for each variable with missing values a dummy variable equal to 1 if an individual had a missing value for that variable and setting the variable itself equal to a constant (24). Observations were weighted using the child-parent weights for the 6th and 7th survey wave panel, which account for attrition and non-response. The empirical methods corrected for the complex sample design of the ECLS-K using the svy command in Stata 11.

Results

Table 1 reports sample means for demographic characteristics and measures of beverage intake by children's frequency of sweetened beverage consumption (none, any, and daily) during the previous week for each grade. Children who drank sweetened beverages at all or daily were more often male and white. Additionally, children who drank sweetened

beverages daily more often attended public school, ate school lunch regularly, ate school breakfast regularly, and received free/reduced price school lunch.

Overall, children's reported caloric drink servings per week fell significantly between 5th and 8th grade from 21.6 to 19.7 servings. Milk consumption fell between 5th and 8th grade (both absolutely and as a share of reported caloric beverage consumption) while consumption of 100% fruit juice increased, regardless of sweetened beverage consumption. Milk consumption fell more, both absolutely and as a proportion of initial servings, among children who drank any sweetened beverages than among children who drank none. The decline in milk consumption was even larger among children who drank sweetened beverages daily: that group drank an average of 2 fewer glasses of milk in 8th grade than in 5th grade.

Table 2 presents regression estimates of how the change in milk consumption is related to the changes in sweetened beverage and juice consumption, controlling for child demographic and nutrition characteristics. The change in children's consumption of milk between 5th and 8th grade was not related to the change in their consumption of sweetened beverages ($\beta=0.005$, $p=0.81$). The change in juice consumption was also not related to the change in children's consumption of sweetened beverages ($\beta=0.039$, $p=0.08$). This pattern indicates that sweetened beverages did not displace other caloric beverages from children's diets. Changes in milk consumption and juice consumption were positively related: children who increased their milk consumption also increased their juice consumption over the three-year period. This pattern indicates that milk and juice are complements, not substitutes, in children's diets.

Table 2 also shows results stratified by population subgroups. We did not find evidence of displacement between caloric beverages among children regardless of their sex, race, or poverty status. Among boys and girls, whites and Hispanics, and regardless of poverty status, the change in children's milk consumption was positively related to the change in their juice consumption but not to the change in their sweetened beverage consumption. Among both boys and children in poverty, changes in juice consumption were positively related to changes in sweetened beverage consumption, indicating that the two were complements. Changes in milk consumption were not significantly related to changes in either juice or sweetened beverage consumption for Blacks or Asians. Thus, analyses of multiple subpopulations of children indicate that milk and juice consumption increased or decreased in tandem for most children. Changes in sweetened beverage consumption were not inversely related to changes in milk or juice consumption among any subgroup.

Discussion

This is the first study to examine displacement of caloric drinks over time in a large, nationally representative study of children. Changes in children's milk and juice consumption over a three-year period were not negatively related to changes in their consumption of sweetened beverages. Thus, the observed decreases in average milk consumption from 5th to 8th grade were not associated with increases in sweetened beverage consumption. Regardless of sweetened beverage consumption, milk and 100% fruit juice were complements in children's diets, with children who increased their consumption of one also increasing their consumption of the other. Among boys and among children living in poverty, juice and sweetened beverage consumption were complements, with children increasing or decreasing their consumption of the two drinks in tandem.

Previous longitudinal studies showed that increases in sweetened beverage consumption accompanied decreases in milk and fruit juice intake from early childhood into adolescence

(18, 25, 26). For example, one study showed that girls who consumed soda at age 5 had higher soda intake, lower milk intake, higher intake of added sugars, and lower nutrient intake from age 5 to 15 years compared with girls who did not consume soda at age 5 (27). Another study using a small, regional sample concluded that consumption of juice and milk declined while consumption of soft drinks increased between 3rd and 8th grade (19). However, these studies lacked generalizability due to small sample sizes, focus on a specific geographic location, or restriction of the sample to specific racial/ethnic groups. It is possible that the difference between those results and the results here are due to cohort differences: this study used more recent data (2004–2007) than previous work, and displacement may have occurred among earlier cohorts of children but not among children born in 1992–1993.

The results indicate that relationships between changes in the consumption of sweetened beverages, milk, and juice between 5th and 8th grade were generally similar across sex, racial, ethnic, and socioeconomic groups. Specifically, milk and juice were complements for all groups, and sweetened beverages were not substitutes for milk or juice in any groups. Juice and sweetened beverages were complements for boys and for children living in poverty. Previous studies documented differences in food, beverage, and activity patterns among children by sex (28) and ethnic groups (11, 20, 27). However, although some of these studies were longitudinal or based on nationally representative data, none of the studies were both.

In addition to finding that changes in milk consumption between 5th and 8th grade were not related to changes in consumption of sweetened beverages, we also found that there was no significant change in patterns of children with any or daily sweetened beverage consumption as opposed to no sweetened beverage consumption. This may be because children's sweetened beverage consumption patterns are already established by 5th grade. Still, it may not indicate that consumption of these beverages did indeed stay constant, as the ECLS-K asked only about the number of times of sweetened beverage were consumed, not about the size of those portions without increases in frequency of consumption.

This study faces several limitations. Like most national surveys, the ECLS-K did not include a measure of water intake (5, 20), so we were not able to determine whether the observed decreases in the total consumption of caloric beverages reflected an increase in water consumption or simply a decrease in beverage intake. However, as sweetened beverages, milk, and juice have been shown to be the dominant beverage consumed by children aged 2–19 (20), our finding that children reduced the frequency of their intake of caloric beverages likely indicates that they reduced their liquid calories. Still, the dietary data were based on a retrospective food questionnaire, not a food diary, so we cannot estimate how children's total caloric intake changed with changes in their beverage consumption. However, as all caloric beverages contribute to overall caloric intake, it is concerning that these main caloric beverages consumed by children -milk, juice, and sweetened beverages - were complementary with each other.

Previous work indicates that the accuracy of children's dietary reporting increase with repeated administration of the same food questionnaire with at least 25 days between questionnaires (30). Therefore, reports in the ECLS-K may have been more accurate at the second administration, in 8th grade, particularly since the children were three years older. However, the ECLS-K asks about frequency of consumption, not serving sizes, which may not provide the same information. While this limitation may introduce noise into our estimates, it does not imply a bias in the results.

Finally, children may have not understood what beverages they were being asked about and may not have known, for example, which juice drinks were 100% juice. Children were not specifically asked to exclude diet drinks from their reports, so it is possible that some children included diet drinks in their reports of sugar-sweetened beverage consumption even though the examples did not include diet drinks. However, it has been shown that less than 8% of children aged 2–19 consume diet beverages (20).

The study offers several strengths. The data are from a large, nationally representative longitudinal dataset. The longitudinal nature of the data allows us to examine changes among the same children over time. This controls for much of the unobserved heterogeneity that can confound cross-sectional analyses. The ECLS-K is the only study of this scale to have collected information on beverages from children of elementary and middle school ages. It also has sufficient sample sizes to examine nationally representative subsets of children by sex, race and ethnicity, and poverty status.

Conclusions

Reducing the intake of caloric beverages is a priority given the high prevalence of overweight and obesity. This focus is especially important since the results presented here indicate that caloric beverages tend to be complementary with each other – as children increase their intake of one caloric beverage they also increase their intake of the others. Caloric beverages must be seen within the context of the entire diet and must be balanced with food intake. It is important for dietetics professionals to help children and their families understand that caloric beverages, even those that are generally healthful, contribute to children's total energy intake and must be moderated as part of a healthy diet.

References

1. Ballew C, Kuester S, Gillespie C. Beverage choices affect adequacy of children's nutrient intakes. *Arch Pediatr Adolesc Med.* 2000; 154(11):1148–52. [PubMed: 11074858]
2. Bowman SA. Beverage choices of young females: changes and impact on nutrient intakes. *J Am Diet Assoc.* 2002; 102(9):1234–9. [PubMed: 12792618]
3. Rampersaud GC, Bailey LB, Kauwell GP. National survey beverage consumption data for children and adolescents indicate the need to encourage a shift toward more nutritive beverages. *J Am Diet Assoc.* 2003; 103(1):97–100. [PubMed: 12525800]
4. Harnack L, Stang J, Story M. Soft drink consumption among US children and adolescents: nutritional consequences. *J Am Diet Assoc.* 1999; 99(4):436–41. [PubMed: 10207395]
5. Nielsen SJ, Popkin BM. Changes in beverage intake between 1977 and 2001. *Am J Prev Med.* 2004; 27(3):205–10. [PubMed: 15450632]
6. Guenther PM. Beverages in the diets of American teenagers. *J Am Diet Assoc.* 1986; 86(4):493–9. [PubMed: 3958400]
7. Bermudez OI, Gao X. Greater consumption of sweetened beverages and added sugars is associated with obesity among US young adults. *Ann Nutr Metab.* 2010; 57(3–4):211–8. [PubMed: 21088388]
8. Blum JW, Jacobsen DJ, Donnelly JE. Beverage consumption patterns in elementary school aged children across a two-year period. *J Am Coll Nutr.* 2005; 24(2):93–8. [PubMed: 15798075]
9. LaRowe TL, Adams AK, Jobe JB, Cronin KA, Vannatter SM, Prince RJ. Dietary intakes and physical activity among preschool-aged children living in rural American Indian communities before a family-based healthy lifestyle intervention. *J Am Diet Assoc.* 2010; 110(7):1049–57. [PubMed: 20630162]
10. Lim S, Zoellner JM, Lee JM, Burt BA, Sandretto AM, Sohn W, et al. Obesity and sugar-sweetened beverages in African-American preschool children: a longitudinal study. *Obesity (Silver Spring).* 2009; 17(6):1262–8. [PubMed: 19197261]

11. Ludwig DS, Peterson KE, Gortmaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet*. 2001; 17;357(9255): 505–8.
12. Papandreou D, Malindretos P, Rousso I. Risk factors for childhood obesity in a Greek paediatric population. *Public Health Nutr*. 2010; 13(10):1535–9. [PubMed: 20074393]
13. Vartanian LR, Schwartz MB, Brownell KD. Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis. *Am J Public Health*. 2007; 97(4):667–75. [PubMed: 17329656]
14. Welsh JA, Cogswell ME, Rogers S, Rockett H, Mei Z, Grummer-Strawn LM. Overweight among low-income preschool children associated with the consumption of sweet drinks: Missouri, 1999–2002. *Pediatrics*. 2005; 115(2):e223–9. [PubMed: 15687430]
15. Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr*. 2007; 150(1):12–7. e2. [PubMed: 17188605]
16. National Institutes of Health. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults--The Evidence Report. *Obes Res*. 1998; 6 (Suppl 2):51S–209S. [PubMed: 9813653]
17. Rajeshwari R, Yang SJ, Nicklas TA, Berenson GS. Secular trends in children's sweetened-beverage consumption (1973 to 1994): the Bogalusa Heart Study. *J Am Diet Assoc*. 2005; 105(2): 208–14. [PubMed: 15668676]
18. Skinner JD, Carruth BR. A longitudinal study of children's juice intake and growth: the juice controversy revisited. *J Am Diet Assoc*. 2001; 101(4):432–7. [PubMed: 11320948]
19. Lytle LA, Seifert S, Greenstein J, McGovern P. How do children's eating patterns and food choices change over time? Results from a cohort study. *Am J Health Promot*. 2000; 14(4):222–8. [PubMed: 10915532]
20. Wang YC, Bleich SN, Gortmaker SL. Increasing caloric contribution from sugar-sweetened beverages and 100% fruit juices among US children and adolescents, 1988–2004. *Pediatrics*. 2008; 121(6):e1604–14. [PubMed: 18519465]
21. U.S. Department of Education - National Center for Education Statistics. ECLS-K Longitudinal Kindergarten-Fifth Grade Public-Use Data File and Electronic Codebook (CD-ROM). Washington, DC: 2006.
22. USDA. Design considerations for developing effective wording and format options for a children's nutrition behavior questionnaire. Final report to Economic Research Service, USDA; Calverton, MD: 2002.
23. Tourangeau, K.; Nord, C.; Le, T.; Pollack, JM.; Atkins-Burnett, S. ECLS-K: Combined user's manual for the ECLS-K fifth-grade data Files and Electronic Codebooks. National Center for Education Statistics; Washington, DC: 2006.
24. Allison, Paul. Missing Data. Vol. 136. Sage Publications, Inc; 2001. Quantitative Applications in the Social Sciences.
25. Fisher JO, Mitchell DC, Smiciklas-Wright H, Mannino ML, Birch LL. Meeting calcium recommendations during middle childhood reflects mother-daughter beverage choices and predicts bone mineral status. *Am J Clin Nutr*. 2004; 79(4):698–706. [PubMed: 15051617]
26. Fiorito LM, Mitchell DC, Smiciklas-Wright H, Birch LL. Dairy and dairy-related nutrient intake during middle childhood. *J Am Diet Assoc*. 2006; 106(4):534–42. [PubMed: 16567149]
27. Fiorito LM, Marini M, Mitchell DC, Smiciklas-Wright H, Birch LL. Girls' early sweetened carbonated beverage intake predicts different patterns of beverage and nutrient intake across childhood and adolescence. *J Am Diet Assoc*. 2010; 110(4):543–50. [PubMed: 20338280]
28. Reedy J, Krebs-Smith SM. Dietary sources of energy, solid fats, and added sugars among children and adolescents in the United States. *J Am Diet Assoc*. 2010; 110(10):1477–84. [PubMed: 20869486]
29. Ogden, CLBKK.; Carroll, MDSP. Consumption of sugar drinks in the United States, 2005–2008. National Center for Health Statistics Data Brief; Hyattsville, MD: 2011.

30. Baxter SD, Smith AF, Nichols MD, Guinn CH, Hardin JW. Children's dietary reporting accuracy over multiple 24-hour recalls varies by body mass index category. *Nutr Res.* 2006; 26(6):241–8. [PubMed: 17541449]

TABLE 1

Demographic characteristics and consumption of milk and 100% fruit juice by frequency of sweetened beverage consumption during the previous week, among U.S. children in 5th and 8th grade

	All		None		Any		Daily	
	5 th	8 th	5 th	8 th	5 th	8 th	5 th	8 th
n	7445	7445	1159	1177	6286	6268	2113	1912
<u>Child characteristics</u>								
Female	49.9%	49.9%	53.5%	56.1%	49.2%	48.7%	46.6%	44.6%
White, non-Hispanic	58.4%	58.4%	50.5%	53.2%	59.9%	59.4%	57.6%	56.7%
Black, non-Hispanic	16.5%	16.5%	21.1%	18.7%	15.6%	16.1%	18.2%	19.6%
Hispanic	18.0%	18.0%	18.0%	17.5%	18.0%	18.0%	18.9%	19.1%
Asian/Pacific Islander, non-Hispanic	3.8%	3.8%	5.4%	6.5%	3.4%	3.2%	2.5%	1.9% *
American Indian, non-Hispanic	1.3%	1.3%	1.7%	1.6%	1.3%	1.3%	1.4%	1.6%
Multiracial	2.0%	2.0%	3.2%	2.4%	1.8%	2.0%	1.5%	1.2%
Attends public school	87.7%	88.7% *	85.7%	87.9%	88.0%	88.8%	90.7%	90.6%
Eats school lunch regularly	78.5%	77.0%	78.7%	71.7% *	78.5%	78.0%	81.6%	80.0%
Eats school breakfast regularly	29.8%	26.6% **	29.0%	27.9%	29.9%	26.4% **	34.4%	31.6%
Living in Poverty	20.4%	19.5%	22.4%	21.7%	20.0%	19.1%	23.5%	21.4%
Receives free/reduced price school lunch	37.9%	33.9% **	42.1%	34.8% *	37.1%	33.8% **	42.8%	38.0% *
<u>Family socioeconomic status</u>								
Bottom quintile	18.9%	19.5%	21.8%	20.6%	18.3%	19.2%	21.3%	22.3%
Second quintile	18.5%	20.6% **	17.0%	17.2%	18.8%	21.2% **	25.2%	24.1%
Third quintile	19.4%	20.1%	20.7%	18.4%	19.1%	20.4%	18.6%	22.5% *
Fourth quintile	20.6%	18.8% *	19.7%	18.8%	20.8%	18.8% *	18.7%	17.7%
Top quintile	22.7%	21.1% **	20.9%	24.9%	23.0%	20.3% **	16.1%	13.5%
<u>Food consumption (frequency per week)</u>								
Green salad	2.2	2.6% **	2.8	2.8	2.1	2.6	2.2	3.0% **
Potatoes (not fries or chips)	2.0	2.1	1.7	2.1	2.1	2.1	2.8	2.7
Carrots	2.8	1.9% **	3.5	2.0	2.7	1.9% **	3.1	2.0% **

	All		None		Any		Daily	
	5 th	8 th	5 th	8 th	5 th	8 th	5 th	8 th
Other vegetables	5.2	4.9 [*]	5.6	5.5	5.1	4.8 [*]	6.3	5.6 [*]
Fruit	7.7	7.2 [*]	8.1	8.0	7.6	7.1 [*]	9.4	8.3 [*]
Fast food	3.2	2.4 ^{**}	2.6	1.5 ^{**}	3.3	2.6 ^{**}	4.9	3.7 ^{**}
Total servings of caloric beverages last week	21.6	19.7 ^{**}	15.1	14.4	22.8	20.6 ^{**}	33.2	29.9 ^{**}
<u>Sweetened beverages</u>								
Drank sweetened beverages at all	83.7%	84.1%	0%	0%	100%	100%	100%	100%
Drank sweetened beverages daily	29.0%	27.0%	0%	0%	34.6%	32.1%	100%	100%
Servings of sweetened beverages last week	6.2	5.5 ^{**}	0	0	7.4	6.5 ^{**}	15.8	14.3 ^{**}
Share of reported caloric beverage consumption	29.1%	28.9%	0%	0%	34.8%	34.4%	51.7%	52.5%
<u>Milk</u>								
Drank milk at all	88.7%	86.5% ^{**}	83.5%	84.0%	89.7%	87.0% ^{**}	87.0%	84.0%
Drank milk daily	53.5%	46.3% ^{**}	51.4%	46.7%	53.9%	46.2% ^{**}	58.9%	49.4% ^{**}
Servings of milk last week	10.3	8.5 ^{**}	9.7	8.5 [*]	10.4	8.5 ^{**}	10.9	8.7 ^{**}
Share of reported caloric beverage consumption	46.3%	42.1% ^{**}	58.6%	56.4%	43.9%	39.4% ^{**}	30.7%	27.3% ^{**}
<u>Juice</u>								
Drank 100% fruit juice at all	75.0%	82.3% ^{**}	72.0%	76.2%	75.6%	83.4% ^{**}	75.1%	81.6% ^{**}
Drank 100% fruit juice daily	24.8%	27.9% ^{**}	28.6%	31.9%	24.0%	27.1% [*]	33.6%	37.2%
Servings of 100% fruit juice last week	5.1	5.6 ^{**}	5.5	5.9	5.1	5.6 [*]	6.5	6.9
Share of reported caloric beverage consumption	23.5%	28.1% ^{**}	34.9%	38.2%	21.3%	26.2% ^{**}	17.6%	20.2% ^{**}

* indicates that 5th and 8th grade means are different at the 0.05 level using t-tests;

** at the 0.01 level.

Notes: Data are from the ECLS-K, with children observed in the 6th and 7th wave of data collection (2003 and 2006).

Survey-adjusted sample means are shown.

“Living in Poverty” indicates that the family is below the federal poverty threshold. Socioeconomic quintile is based on parental income, education, and occupational prestige. Shares of reported caloric beverage consumption are calculated as servings of each beverage type as a percentage of total servings of caloric beverages last week.

TABLE 2

Survey-adjusted linear regression estimates of the association between changes in milk consumption or juice consumption and changes in servings of other caloric beverages between 5th and 8th grade among U.S. children

	All	Girls	Boys	Whites	Blacks	Hispanics	Asians	Living in Poverty	Not Living in Poverty
n	7445	3757	3688	4872	640	1169	456	1112	6333
<u>Outcome: Servings of milk</u>									
Servings of sweetened beverages	0.005 (0.019)	-0.033 (0.031)	0.033 (0.026)	0.014 (0.025)	-0.025 (0.056)	0.004 (0.042)	0.025 (0.119)	0.026 (0.040)	-0.008 (0.023)
Servings of 100% fruit juice	0.087** (0.022)	0.077** (0.029)	0.091** (0.030)	0.081** (0.029)	0.050 (0.054)	0.151** (0.043)	0.097 (0.071)	0.123* (0.042)	0.074** (0.026)
<u>Outcome: Servings of juice</u>									
Servings of sweetened beverages	0.039 (0.022)	0.015 (0.031)	0.060* (0.030)	0.010 (0.026)	0.073 (0.050)	0.051 (0.038)	0.138 (0.075)	0.119* (0.056)	0.008 (0.023)
Servings of milk	0.066** (0.017)	0.061* (0.024)	0.067** (0.021)	0.050** (0.019)	0.056 (0.053)	0.132** (0.038)	0.080 (0.057)	0.120** (0.042)	0.051** (0.019)

* p<0.05;

** p<0.01

Note: Data are from the ECLS-K.

Results shown for all children and stratified by sex, race/ethnicity, and poverty status. Shown are coefficient estimates from linear regression models, with standard errors in parentheses. "Living in Poverty" indicates that the family is below the federal poverty threshold.

Models include sex and race/ethnicity, family socioeconomic status, public/private school attendance, whether the child eats school lunch regularly, eats school breakfast regularly, receives free or reduced price lunch at school, and the change in consumption of salad, potatoes, carrots, other vegetables, fruit, and fast food, as reported by children.