Early Growth Response Protein-1 Promoter-Mediated Synergistic Antitumor Effect of hTERTC27 Gene Therapy and 5-Fluorouracil on Nasopharyngeal Carcinoma

Guimiao Lin,^{1,2} Marie Chia-Mi Lin,³ Suxia Lin,² Hong Yao,³ Shuihong Yu,² Wanxian Yi,² Gaixia Xu,⁴ Samuel Sai-Ming Ng,³ Siping Chen,² Jing Yu,² Xiaomei Wang,^{2*} and Baoxue Yang^{1,5,6*}

Abstract

hTERTC27 is a newly constructed polypeptide that can induce telomere dysfunction. To study the synergistic antitumor effects of the hTERTC27 polypeptide driven by the early growth response protein-1 (Egr-1) promoter and chemotherapeutic 5-fluorouracil (5-FU) on nasopharyngeal carcinoma, a series of *in vitro* and *in vivo* experiments were performed. The results showed that hTERTC27 expression was significantly increased up to 7.21-folds by the 5-FU-activated Egr-1 promoter in C666-1 cells. Overexpressed hTERTC27 made the cells more sensitive to 5-FU, and additionally, inhibited cell proliferation about 20.41%. Combinational therapy of overexpressed hTERTC27 driven by the 5-FU-activated Egr-1 promoter and 5-FU synergistically inhibited cell proliferation and promoted apoptosis of C666-1 cells for about 4.75-fold and 1.76-fold in comparison with a sole therapy of hTERTC27 or 5-FU *in vitro*. *In vivo* experiments showed that overexpressed hTERTC27 driven by 5-FU-activated Egr-1 promoter and 5-FU synergistically reduced tumor volume, tumor weight, and local infiltration, which may be relative to tumor cell apoptosis. These results suggest that combinational therapy of overexpressed hTERTC27, which is driven by the 5-FU-activated Egr-1 promoter, and 5-FU may provide a novel approach to treat nasopharyngeal cancer.

Key words: 5-FU, Egr-1, hTERTC27, nasopharyngeal carcinoma, gene therapy

Introduction

Nasopharyngeal carcinoma (NPC) is one of the most malignant neoplasms in the nasopharynx, especially in Southeastern Asia. 5-fluorouracil (5-FU), the most widely used anticancer agent, has been applied widely to the treatment of NPC.^{1,2} However, 5-FU treatment has only limited benefit to the NPC patients because of severe side effects caused by high dosage and failing preventing recurrence and metastasis due to limited sensitivity.³ Therefore, several strategies of gene therapy have been intensively explored to treat NPC.^{4,5}

Telomerase is an attractive target for cancer gene therapy, because it is highly active in most cancer cells, but it is inactive in most somatic cells.⁶ The primary function of telomerase is to maintain the telomere length and to prolong the life span of cells.⁷ The dysfunction of telomerase will prevent the long-term survival of tumor cells and lead to cell death or senescence.^{8,9} Telomerase reverse transcriptase (TERT), the catalytic protein subunit of telomerase, is critical for the activity of telomerase in tumor cells.¹⁰ Previously, we have reported that a 27-kDa C-terminal polypeptide of hTERT (hTERTC27) is capable of inducing telomere

¹Department of Pathophysiology, Norman Bethune Medical School, Jilin University, Changchun, China.

²The Key Lab of Biomedical Engineering, Department of Physiology, Research Institute of Uropoiesis and Reproduction, School of Medical Sciences, Shenzhen University, Shenzhen, China.

³Division of Neurosurgery, Department of Surgery, Brain Tumor Centre, The Chinese University of Hong Kong, Pokfulam, Hong Kong, China.

⁴Department of Optoelectronic Information Engineering, College of Optoelectronics Engineering, Shenzhen University, Shenzhen, China.

⁵Department of Pharmacology, School of Basic Medical Sciences, Peking University, Beijing, China.

⁶Key Laboratory of Molecular Cardiovascular Sciences, Ministry of Education, Beijing, China.

Address correspondence to: Baoxue Yang; Department of Pharmacology, School of Basic Medical Sciences, Peking University; 38 Xueyuan Road, Haidian District, Beijing, China 100191 or Xiaomei Wang; School of Medical Sciences, Shenzhen University; 3688 Nanshan Road, Nanshan District, Shenzhen, China 518060.

E-mail: baoxue@bjmu.edu.cn or xmwang@szu.edu.cn

dysfunctions, and overexpression of hTERTC27 in human cervical carcinoma HeLa cells slows down the cell proliferation and inhibits the tumorigenicity by significantly increasing anaphase chromosome end-to-end fusion.¹¹ Ectopic overexpression of this polypeptide also makes the HeLa cells more sensitive to H₂O₂-induced oxidative injury.¹² Furthermore, hTERTC27 delivered by recombinant adeno-associated virus or recombinant adeno-associated virus and adenovirus cocktail system suppresses tumor growth *in vivo* in a human U87-MG glioblastoma xenograft model.^{13,14}

The early growth response protein-1 (Egr-1) promoter, which is sensitive to reactive oxygen intermediates,¹⁵ is commonly used as a bridge in gene therapy, because it can be activated by oxygen free radicals generated by radiotherapy or chemotherapy and promotes the high expression of tumor therapeutic genes.^{16,17} Therefore, we hypothesize that the radical agent 5-FU may activate the Egr-1 promoter, which can drive the overexpression of hTERTC27 that makes tumor cells sensitive to the chemotherapy of 5-FU, and have the chemotherapeutic effect on NPC.

In this study, hTERTC27 cDNA was constructed to the downstream of the Egr-1 promoter, which was transfected into the C666-1 cells. The synergistic antitumor effects of the 5-FU and hTERTC27 polypeptide driven by the 5-FU-activated Egr-1 promoter were assayed for the first time. Overexpressed hTERTC27 driven by the Egr-1 promoter made the C666-1 cells more sensitive to 5-FU and synergistically inhibited C666-1 cell proliferation and induced apoptosis with 5-FU treatment. These results suggest that the synergistic effect of 5-FU and Egr-1 promoter-driven hTERTC27 might be a potential clinical strategy for nasopharyngeal cancer therapy.

Materials and Methods

Cell culture

Human NPC C666-1 cells were maintained in our lab and cultured in the Dulbecco's modified Eagle's medium (DMEM; Hyclone), supplemented with 10% fetal bovine serum (FBS; Hyclone), 100 μ g/mL penicillin (Ameresco), and 100 μ g/mL streptomycin (Ameresco). Cells were cultured at 37°C in a humidified atmosphere with 5% CO₂.

Plasmid construction and transfection

The pcDNA3.1-Egr-1 plasmid (pEgr) was a gift from Dr. Jianxiang Liu and Mei Tian. In this plasmid, the cytomegalovirus promoter was substituted with the Egr-1 promoter. cDNAs of a 27-kDa hTERT C-terminal polypeptide (hTERTC27, amino acid residues 882–1132) and enhanced green fluorescent protein (EGFP) were obtained by polymerase chain reaction using the pcDNA3.1-hTERTC27 and pEGFP-N2 (BD Biosciences) as templates, respectively. The hTERTC27 cDNA was subcloned into the pEgr plasmid at *Kpn* I and *Eco*R I sites to generate the pEgr-1-hTERTC27 plasmid (pEgr-C27). EGFP cDNA was then subcloned into the pEgr-C27 plasmid at *Eco*R I and *Xba* I sites to generate a pEgr-C27-GFP plasmid coexpressing hTERTC27 and GFP fusion protein. C666-1 cells were transfected with the pEgr, pEgr-C27, or pEgr-C27-GFP vectors using a Lipofectamine 2000 transfection reagent (Invitrogen), following the manufacturer's instructions, and selected using 600 μ g/mL G418

(Merck) for 2 weeks. The cells stably transfected with the pEgr, pEgr-C27, or pEgr-C27-GFP vectors were called as C666-Con cells, C666-C27 cells, or C666-C27-GFP cells, respectively, and the untransfected C666-1 cells were called as C666-Mock cells.

Detection of EGFP

About 2×10^6 C666-C27-GFP cells were seeded in 10-cm dishes. After 12 hours of incubation, the cells were treated with different concentrations (in μ g/mL) of 5-FU (Sigma) of 0, 3.12, 6.25, 12.5, 25, respectively, for 48 hours. The cells were imaged with a fluorescence microscope (Zeiss). Then, cells were washed with phosphate-buffered saline (PBS) twice, harvested by trypsinization, and fixed in 70% ethanol. GFP expression was further determined by a flow cytometry analysis (Becton Dickinson).

Immunoblotting analysis

Cells were lysed in an NP-40 lysis buffer containing 50 mM Tris (pH 7.4), 150 mM NaCl, 1% NP-40, 20 μ g/mL PMSF, 1 μ g/mL pepstatin A, and 1 μ g/mL leupeptin (pH 7.4). Tumor tissue was homogenized in 250 mM sucrose containing 1 mM ethylenediaminetetraacetic acid, 20 μ g/mL PMSF, 1 μ g/mL pepstatin A, and 1 μ g/mL leupeptin (pH 7.4) with a glass Dounce homogenizer. Total protein was assayed using a DC protein assay kit (BioRad). Proteins (30 μ g) were electrophoresed through 12% (w/v) polyacrylamide and 1% (w/v) sodium dodecyl sulfate gels. Proteins were transferred from the gels by blotting onto a nitrocellulose membrane (Millipore). The membranes were blocked with 0.02 mol/L Tris-HCl (pH 7.0), 0.9% (w/v) NaCl, and 0.1% (v/v) Tween-20 containing 5% (w/v) nonfat milk for 1 hour at room temperature. Antibodies against the hTERT COOH-terminal fragment (1:250; Santa Cruz Biotechnology), cleaved caspase-3 (1:1000; Cell Signaling Technology), cleaved caspase-9 (1:1000; Cell Signaling Technology), cleaved PARP (1:1000; Cell Signaling Technology), and Bcl-2 (1:500; Santa Cruz Biotechnology) were added and incubated overnight at 4°C. Bound antibodies were located by a further incubation with horseradish peroxidase-conjugated goat anti-rabbit IgG (1:5000; Santa Cruz Biotechnology). β -actin (1:500; Santa Cruz Biotechnology) was used as a loading control. Blots were developed with an ECL Plus kit (Pierce) and exposed to Kodak films.

Determination of the 50% inhibitory concentration

About 2×10^3 C666-Mock, C666-Con, or C666-C27 cells were seeded in a 96-well plate. Twelve hours later, the cells were treated with different concentrations (in μ g/mL) of 5-FU of 0, 0.78, 1.56, 3.12, 6.25, 12.5, 25, 50, 100, 200, or 400, respectively. An MTT assay was performed 48 hours later. Briefly, 20 μ L of 5 mg/mL MTT (Sigma) in PBS was added, and the cells were then incubated for 4 hours at 37°C with 5% CO₂. Cells were washed twice with PBS, and the pellet was then solubilized in 150 μ L of 100% dimethyl sulfoxide (Sigma) by shaking for 5 minutes. Absorbance was measured on a microplate reader (BioRad) at a wavelength of 570 nm. The inhibitory rate (IR) and 50% inhibitory concentration (IC₅₀) were calculated using Origin 7.5 software package.

Cell growth and proliferation assay

C666-Con or C666-C27 cells were seeded in six-well plates (2.5×10^4 /well) and a 96-well plate (1.0×10^4 /well). Twelve hours later, the cells were treated with PBS or 5-FU (6.25 or 12.5 $\mu\text{g}/\text{mL}$), respectively. Forty-eight hours later, cells in six-well plate were imaged with a light microscope (Zeiss) and then washed with PBS twice, harvested by trypsinization, and counted using a hemocytometer (Shanghai Jinghong Corporation). Cell proliferation in a 96-well plate was measured by the MTT assay as described above.

Colony formation assay

C666-Con or C666-C27 cells (5×10^2 /well) were seeded in six-well plates. Twelve hours later, cells were exposed to PBS, 6.25, or 12.5 $\mu\text{g}/\text{mL}$ 5-FU for 48 hours and allowed to grow for 14 days, fixed with methanol for 10 minutes, and stained with Giemsa for 15 minutes. The plates were washed and dried, and the colonies were counted. All experiments were done in triplicate.

Flow cytometry analysis of apoptosis

For quantitative analysis of apoptosis, cells were harvested at 48 hour after treating with 5-FU, washed once with ice-cold PBS, incubated with annexin V-fluorescein/PI (Boehringer Mannheim) in a calcium-containing HEPES (Sigma) buffer, and then immediately analyzed with a flow cytometry machine (Becton Dickinson).

TUNEL assay

The presence of apoptotic cells in the C666-1 cells was detected by the terminal deoxynucleotidyl transferase-mediated dUTP nick-end-labeling (TUNEL) assay using the *in situ* cell death detection kit (POD; Roche), according to the procedures recommended by the manufacturer. The apoptotic cells were analyzed under a fluorescence microscope.

Tumor xenograft formation and histological analysis

BALB/C nude mice at 6 weeks of age were purchased from the Medical Laboratory Animal Center of Guangdong

Province, China. Mice were injected with 100 μL (5×10^7 cells/mL in PBS) of C666-Con cell suspension (12 mice) or C666-C27 cell suspension (12 mice) subcutaneously into the mid-back region. One week after injection, the mice inoculated with C666-Con or C666-C27 cells were randomized into two groups, respectively (Con, Con+5-FU and C27, C27+5-FU). Mice were administrated (i.p.) either saline or 5-FU (40 mg/kg daily, three times a day) for 5 days. Tumor xenograft formation was measured on a daily base to make the growth curves. At the end of the treatment, mice were sacrificed. Tumor xenografts were surgically dissected, weighed, measured and then fixed with 4% formaldehyde in PBS, and embedded in paraffin. Sections were stained with hematoxylin and eosin (HE staining). The tumor volume was calculated according to the equation $L \times W^2/2$ with the length (mm) being the longest axis of the tumor. This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the Medical Laboratory Animal Center of Guangdong Province, China (the permit number is SCXK2008-0002).

Statistical analysis

All data were presented as mean $\pm$ SD. The results were compared by an analysis of variance. All statistical calculations were performed with SPSS 11.0 software package. A p -value < 0.05 was regarded as statistically significant difference.

Results

5-FU increased hTERTC27-GFP expression driven by Egr-1 promoter in C666-1 cells

We firstly constructed the pEgr-C27-GFP plasmid, in which a cDNA-encoding hTERTC27-GFP fusion protein was put at downstream of Egr-1 promoter. A fluorescence microscopic assay showed the hTERTC27-GFP fusion protein was expressed in pEgr-C27-GFP stably transfected C666 cells (Fig. 1A). About 3.12 $\mu\text{g}/\text{mL}$ 5-FU obviously increased the expression of hTERTC27-GFP fusion protein assayed by flow cytometry ($p < 0.01$, Fig. 1B). With the increase of 5-FU

FIG. 1. 5-FU activated Egr-1 promoter to increase hTERTC27-GFP expression in C666-1 cells. **(A)** hTERTC27-GFP expression was observed under the fluorescence inverse microscope. Scale bar = 200 μm . **(B)** hTERTC27-GFP expression levels induced by 5-FU determined using a flow cytometry analysis. **(C)** Relative hTERTC27-GFP expression levels in C666-C27-GFP cells with or without 5-FU treatment. Values are means $\pm$ SD, $n = 6$; * $p < 0.01$ versus group without 5-FU treatment. **(D)** Western blot analysis of C666-C27-GFP cells with or without 5-FU treatment using corresponding antibodies against the hTERT COOH-terminal polypeptide. Actin was probed with an anti-actin antibody as a loading control. 5-FU, 5-fluorouracil.

FIG. 2. hTERTC27 increased sensitivity of C666-1 cells to 5-FU. **(A)** Dose-response curves of inhibitory effect of 5-FU on C666-Mock, C666-Con, or C666-C27 cells. Cells were seeded into 96-well plates. Twelve hours later, cells were treated with the indicated concentrations of 5-FU for 48 hours. The inhibitory rate was measured by the MTT assay. **(B)** IC₅₀ values from the MTT assay. Values are means $\pm$ SD, $n=6$; * $p<0.05$ versus group mock; ▲ $p<0.05$ versus group control.

dosage, hTERTC27-GFP expression was increased up to 7.21-fold ($p<0.01$, Fig. 1C), which was confirmed by the Western blot analysis (Fig. 1D). The results indicate that 5-FU can increase hTERTC27 expression by activating the Egr-1 promoter.

hTERTC27 increased sensitivity of C666-1 cells to 5-FU

To examine the effect of overexpressed hTERTC27 on the survival of 5-FU-treated C666-1 cells, growth inhibitions of C666-Mock, C666-Con, or C666-C27 cells were analyzed with the MTT assay after treatment with various concentrations of 5-FU for 48 hours. We found that C666-C27 cells showed a lower proliferation ability than C666-Con cells and C666-Mock cells. The IC₅₀ of 5-FU in C666-C27 cells (15.11 ± 0.92 μg/mL) was significantly lower than those in

C666-Mock cells (22.88 ± 2.45 μg/mL) and C666-Con cells (21.31 ± 1.91 μg/mL) ($p<0.05$). IRs of cells were not significantly different between C666-Mock cells and C666-Con cells (Fig. 2). These results indicate that overexpressed hTERTC27 makes the C666-1 cells more sensitive to 5-FU treatment.

Overexpressed hTERTC27 and 5-FU synergistically inhibited C666-1 cell growth

The synergistic effect of the overexpressed hTERTC27 polypeptide and 5-FU on C666-1 cell proliferation was determined by the MTT assay. As shown in Figure 3A and B, cell proliferation was significantly inhibited in group C27 ($p<0.05$) and groups 5-FU ($p<0.01$) when compared with that in group control. Interestingly, groups C27+5-FU demonstrated lower cell proliferation rates than group C27 ($p<0.01$) and groups 5-FU ($p<0.01$). Consistently, colony formation assay showed that the colony formation rate in group C27 ($71.67 \pm 6.50\%$) was significantly lower than that in group control ($95.00\% \pm 7.23\%$) as showed in Figure 3C and D ($p<0.05$). Moreover, the combined therapeutic group of C27+6.25 μg/mL 5-FU ($0.33\% \pm 0.10\%$) demonstrated a further decrease in the colony formation rate when compared with groups C27 ($71.67\% \pm 6.50\%$, $p<0.01$) or 6.25 μg/mL 5-FU ($8.33\% \pm 2.95\%$, $p<0.01$) alone. There was not any colony found in group 12.5 μg/mL 5-FU only and group 12.5 μg/mL 5-FU+C27 (Fig. 3C, D). These results indicate that overexpressed hTERTC27 and 5-FU synergistically inhibited the proliferation of C666-1 cells.

Overexpressed hTERTC27 and 5-FU synergistically induce apoptosis in C666-1 cells

To determine if overexpressed hTERTC27 and 5-FU cause apoptosis of C666-1 cells, the early apoptotic death (AnnexV + /PI-) was assessed by flow cytometry. As shown in Figure 4A and B, the apoptotic death in group C27

FIG. 3. hTERTC27 and 5-FU synergistically inhibited C666-1 cell growth. **(A)** Representative pictures showing C666-Con or C666-C27 cells treated with 6.25 or 12.5 μg/mL 5-FU for 48 hours. Scale bar = 200 μm. **(B)** Effects of hTERTC27 and 5-FU on C666-1 cell proliferation measured by the MTT assay. **(C)** Representative pictures of colony formation assay. **(D)** Colony formation rates. All values are means $\pm$ SD, $n=6$; * $p<0.01$ versus group control; ▲ $p<0.05$ versus group control; * $p<0.01$ versus group C27; ▲ $p<0.01$ versus group 6.25 μg/mL 5-FU; ◇ $p<0.01$ versus group 12.5 μg/mL 5-FU.

FIG. 4. Synergistic effects of hTERTC27 and 5-FU on apoptosis in C666-1 cells. **(A)** Apoptosis in C666-1 cells transfected with the pEgr vector or pEgr-C27 without or with 6.25 or 12.5 $\mu\text{g}/\text{mL}$ 5-FU for 48 hours. Apoptosis was determined using an Annexin V-PI double-staining flow cytometric technique. **(B)** Percentages of early apoptosis from experiments showed in **(A)**. **(C)** Representative fields of the TUNEL assay; scale bar = 400 μm . **(D)** Percentages of TUNEL-positive cells from experiments showed in **(C)**. All values are means $\pm$ SD, $n = 6$; * $p < 0.01$ versus group control; ** $p < 0.01$ versus group C27; $\Delta p < 0.05$ versus group 6.25 $\mu\text{g}/\text{mL}$ 5-FU; $\diamond p < 0.05$ versus group 12.5 $\mu\text{g}/\text{mL}$ 5-FU.

(7.41% $\pm$ 1.81%) was significantly increased compared to that in group control (2.43% $\pm$ 0.37%, $p < 0.01$). Further, group C27 + 6.25 $\mu\text{g}/\text{mL}$ 5-FU (15.77% $\pm$ 3.98%) demonstrated a significant increase in the apoptotic death compared to group C27 ($p < 0.01$) and group 6.25 $\mu\text{g}/\text{mL}$ 5-FU (10.34% $\pm$ 2.07%, $p < 0.05$), and this trend of difference was also found between group C27 + 12.5 $\mu\text{g}/\text{mL}$ 5-FU (35.15% $\pm$ 5.66%) and group 12.5 $\mu\text{g}/\text{mL}$ 5-FU (26.32% $\pm$ 4.12%). To confirm this result, the TUNEL assay was performed (Fig. 4C, D). It was found that $\sim 9.11\%$ C666-1 cells were TUNEL positive in group C27, which is significantly higher than control (about 3.27%, $p < 0.01$). However, about 36.73% and 61.65% C666-1 cells were TUNEL positive in groups C27 + 6.25 $\mu\text{g}/\text{mL}$ 5-FU and C27 + 12.5 $\mu\text{g}/\text{mL}$ 5-FU, respectively, whereas only about 24.34% and 44.14% C666-1 cells were under apoptosis in groups 6.25 $\mu\text{g}/\text{mL}$ 5-FU and 12.5 $\mu\text{g}/\text{mL}$ 5-FU. The apoptotic rates of C666-1 cells in combined therapeutic groups were remarkably higher than groups C27 ($p < 0.01$) and 5-FU ($p < 0.05$). These data suggest that overexpressed hTERTC27 driven by Egr-1 promoter and 5-FU synergistically promotes the C666-1 cell apoptosis.

5-FU-induced hTERTC27 polypeptide and 5-FU synergistically inhibited the growth of NPC tumor in vivo

BALB/C mice were injected subcutaneously with C666-Con cells or C666-C27 cells, respectively. A week after injection, animals were treated with 40 mg/kg 5-FU daily, and tumor growth was monitored for 5 days. As shown in Figure 5A, tumor growth was significantly inhibited by the hTERTC27 polypeptide and 5-FU. The tumor volume and weight in groups C27 and 5-FU were significantly diminished compared with those in group control (Fig. 5A–C, $p < 0.05$). Furthermore, the volume and weight of tumor in group C27 + 5-FU were significantly lower than in groups C27 and 5-FU (Fig. 5B, C, $p < 0.05$). In addition, local muscle

infiltration was found in all mice of group control, whereas local muscle infiltration was found in only 67.00% of group C27 mice, 50.00% of group 5-FU mice, and 16.70% of group C27 + 5-FU mice, respectively (Fig. 5D).

5-FU-induced hTERTC27 polypeptide and 5-FU promote the cleavage of poly(ADP-ribose) polymerase, the activation of caspases, and down regulation of Bcl-2

It is well known that apoptosis is characterized by the cleavage of poly(ADP-ribose) polymerase (PARP) and activation of caspases. Thus, we examined the effect of hTERTC27 and 5-FU on the cleavage of PARP and the activation of caspases. As shown in Figure 6A and B, the amount of PARP cleavage was markedly enhanced in group C27 + 5-FU when compared to that in group 5-FU. Similar results were also observed in the activation of caspases (Fig. 6A, B). These results consistently showed that the combination of hTERTC27 and 5-FU synergistically induced apoptotic death in NPC tumors. Then, we evaluated changes in the signaling proteins of the mitochondrion-dependent apoptotic pathway. Bcl-2 is a very important protein of the antiapoptotic family. As shown in Figure 6C and D, the Bcl-2 expression in groups C27 and 5-FU remarkably decreased in comparison with that in group control. Further, the Bcl-2 expression in group C27 + 5-FU diminished obviously when compared with that in groups C27 or 5-FU.

Discussion

The motivation of this study was to determine if the Egr-1 promoter can be activated by 5-FU to increase hTERTC27 expression, and if 5-FU can be used as an inducer to increase Egr-promoter-driven hTERTC27 polypeptide overexpression and a chemotherapeutic agent on NPC cancer. Our results showed the Egr-1 promoter, which is sensitive to reactive

FIG. 5. *In vivo* synergistic antitumor effect of hTERTC27 and 5-FU on nasopharyngeal carcinoma tumor. **(A)** Tumor growth curves in C666-1 cell-transplanted mice. Tumors were measured in three dimensions on the indicated days after treatment with 40 mg/kg 5-FU. Values are means $\pm$ SD, $n=6$; * $p < 0.05$. **(B)** Images of tumors. Scale bar = 10 mm. **(C)** Weights of xenograft tumors at day 5 post-treatment. Values are means $\pm$ SD, $n=6$; $\Delta p < 0.05$ versus group control; * $p < 0.01$ versus group control; * $p < 0.05$ versus group C27; $\Delta p < 0.05$ versus group 5-FU. **(D)** Hematoxylin and eosin staining, indicating the local muscle infiltration (arrow) in tumors. Scale bar = 200 μ m.

oxygen intermediates,¹⁸ was activated by 5-FU to drive hTERTC27 overexpression. At appropriate concentrations of 5-FU, we observed the synergistic antitumor effect of the hTERTC27 polypeptide regulated by the Egr-1 promoter and 5-FU on nasopharyngeal carcinoma.

As we know, 5-FU is one of the most widely used drugs in NPC chemotherapy^{1,2} because it can effectively inhibit the activity of thymidylate synthase, block the DNA synthesis,

arrest the cell cycle, and promote apoptosis. However, the severe side effects of 5-FU and its limited sensitivity to cancer cells still bring great obstacles to cancer therapy. Different strategies have been designed to decrease the dosage and increase the antitumor sensitivity of 5-FU. Therefore, it is very important to combine 5-FU chemotherapy and gene therapy to make the tumor cells sensitive to 5-FU treatment and enhance its antitumor effects.^{19–21}

FIG. 6. Immunoblotting analysis of cleaved PARP, caspases, and Bcl-2 protein. **(A)** Immunoblotting analysis for cleaved PARP and caspases in tumor tissues after treatment. Actin was probed as loading controls. **(B)** Quantification of cleaved PARP and caspases in tumor tissues. Values are means $\pm$ SD, $n=3$; * $p < 0.01$ versus group control; $\square p < 0.01$ versus group C27; $\Delta p < 0.05$ versus group 5-FU. **(C)** Representative immunoblotting analysis of Bcl-2 in tumor tissues. **(D)** Quantification of the Bcl-2 expression in tumor tissues. Values are means $\pm$ SD, $n=3$; * $p < 0.01$ versus group control; $\Delta p < 0.05$ versus group control; * $p < 0.01$ versus group C27; $\Delta p < 0.01$ versus group 5-FU. PARP, poly(ADP-ribose) polymerase.

Recently, a variety of therapeutic approaches are being investigated to target telomere or telomerase, since many tumors express high levels of telomerase.^{6,22–24} We have newly constructed the hTERTC27 polypeptide and previously reported that this polypeptide rendered the HeLa cells more sensitive to H₂O₂ treatment. Moreover, our previous study showed that overexpressed hTERTC27 induced telomere dysfunction in HeLa cells and inhibits cervical carcinoma and glioblastoma tumor growth and tumorigenicity.^{16–18} However, hTERTC27 gene therapy alone cannot quickly inhibit the tumor growth because of its limited expression level. Here, our results showed that hTERTC27 remarkably inhibited cell proliferation, promoted apoptosis of C666-1 cells *in vitro*, and obviously slowed down the NPC tumor growth *in vivo* as well. The IC₅₀ value of 5-FU in hTERTC27-overexpressed NPC cells was significantly lower than that in control cells, which indicated that a low concentration of 5-FU treatment in combination with hTERTC27 polypeptide achieves the same antitumor effect as a higher 5-FU concentration. This strategy can decrease the therapeutic dosage and the side effects of 5-FU.

To further optimize the therapeutic strategy of the hTERTC27 gene therapy and 5-FU chemotherapy, we introduced the Egr-1 promoter as a bridge to amplify the antitumor effect of hTERTC27 and 5-FU. The Egr-1 promoter increased hTERTC27 expression in NPC cells due to reactive oxygen intermediates caused by 5-FU, and the high expression of hTERTC27 rendered the cells more sensitive to 5-FU and made the cells generate more reactive oxygen intermediates and further led to a higher hTERTC27 protein level. By recycling, the combined therapy of hTERTC27 driven by Egr-1 promoter and 5-FU exerted a more efficient antitumor effect just as found in our study. Also, the combined treatment of hTERTC27 and 5-FU significantly promoted apoptosis in C666-1 cells. An *in vivo* mouse model also showed that hTERTC27 driven by the Egr-1 promoter and 5-FU synergistically inhibited tumor growth and local infiltration.

The Bcl-2 family proteins have been shown to play important roles in apoptotic cell death.²⁵ Bcl-2 inhibits cell apoptosis. Here, we found that Bcl-2 expression was obviously diminished in tumors of combined treating group. Caspase-3 and caspase-9 are involved in cell apoptosis. A classical substrate of activated caspases is PARP, which is a DNA-repair enzyme needed for normal cells.²⁶ Interestingly, we found that obviously elevated activated caspase-3 and 9 expressions and cleaved PARP with combined therapy of C27 and 5-FU. These results confirmed the synergistic antitumor effects of the hTERTC27 polypeptide driven by Egr-1 promoter and 5-FU on nasopharyngeal carcinoma.

In summary, 5-FU activated the Egr-1 promoter to overexpress hTERTC27 polypeptide, which enhanced the sensitivity of NPC cells to 5-FU treatment. Our study proposes that the combinational therapy of 5-FU-Egr-1-hTERTC27 is adaptable to treating nasopharyngeal cancer.

Acknowledgments

This work was supported by the grants from Shenzhen/Hong Kong Innovation Circle (CXQ2008043), National Natural Science Foundation of China (NSFC) (30870921, 81170632, and 30900335), and Doctoral Training Fund 20100001110047,

International Science & Technology Cooperation Program of China 2012DFA11070.

Disclosure Statement

The authors declare no conflict of interest.

References

1. Zheng J, Wang G, Yang GY, et al. Induction chemotherapy with nedaplatin with 5-FU followed by intensity-modulated radiotherapy concurrent with chemotherapy for locoregionally advanced nasopharyngeal carcinoma. *Jpn J Clin Oncol* 2010;40:425.
2. Lee N, Harris J, Garden AS, et al. Intensity-modulated radiation therapy with or without chemotherapy for nasopharyngeal carcinoma: Radiation therapy oncology group phase II trial 0225. *J Clin Oncol* 2009;27:3684.
3. Tham IW, Lu JJ. Controversies and challenges in the current management of nasopharyngeal cancer. *Expert Rev Anticancer Ther* 2010;10:1439.
4. Pan JJ, Zhang SW, Chen CB, et al. Effect of recombinant adenovirus-p53 combined with radiotherapy on long-term prognosis of advanced nasopharyngeal carcinoma. *J Clin Oncol* 2009;27:799.
5. Song Y, Dong MM, Yang HF. Effects of RNA interference targeting four different genes on the growth and proliferation of nasopharyngeal carcinoma CNE-2Z cells. *Cancer Gene Ther* 2011;18:297.
6. Tian X, Chen B, Liu X. Telomere and telomerase as targets for cancer therapy. *Appl Biochem Biotechnol* 2010;160:1460.
7. Blackburn EH. Structure and function of telomeres. *Nature* 1991;350:569.
8. Aubert G, Lansdorp PM. Telomeres and aging. *Physiol Rev* 2008;88:557.
9. Shamas MA. Telomeres, lifestyle, cancer, and aging. *Curr Opin Clin Nutr Metab Care* 2011;14:28.
10. Osterhage JL, Friedman KL. Chromosome end maintenance by telomerase. *J Biol Chem* 2009;284:16061.
11. Huang JJ, Lin MC, Bai YX, et al. Ectopic expression of a COOH-terminal fragment of the human telomerase reverse transcriptase leads to telomere dysfunction and reduction of growth and tumorigenicity in HeLa cells. *Cancer Res* 2002;62:3226.
12. Huang J, Bai YX, Han SW, et al. A human TERT C-terminal polypeptide sensitizes HeLa cells to H₂O₂-induced senescence without affecting telomerase enzymatic activity. *Biochem Biophys Res Commun* 2003;301:627.
13. Ng SS, Gao Y, Chau DH, et al. A novel glioblastoma cancer gene therapy using aav-mediated long-term expression of human TERT C-terminal polypeptide. *Cancer Gene Ther* 2007;14:561.
14. Gao Y, Ng SS, Chau DH, et al. Development of recombinant adeno-associated virus and adenovirus cocktail system for efficient hTERT27 polypeptide-mediated cancer gene therapy. *Cancer Gene Ther* 2008;15:723.
15. Datta R, Rubin E, Sukhatme V, et al. Ionizing radiation activates transcription of the EGR1 gene via CarG elements. *Proc Natl Acad Sci U S A* 1992;89:10149.
16. Du N, Pei XT, Xiao WH, et al. Effect of fluorouracil-inducible GM-CSF gene therapy regulated by EGR-1 promoter on chemotherapeutic hematopoietic damage of tumor-bearing mice. *Zhonghua Yi Xue Za Zhi* 2009;89:1281.
17. Mezhr JJ, Smith KD, Posner MC, et al. Ionizing radiation: A genetic switch for cancer therapy. *Cancer Gene Ther* 2006; 13:1.

18. Datta R, Taneja N, Sukhatme VP, et al. Reactive oxygen intermediates target CC(A/T)6GG sequences to mediate activation of the early growth response 1 transcription factor gene by ionizing radiation. *Proc Natl Acad Sci U S A* 1993;90:2419.
19. Li XP, Li CY, Li X, et al. Inhibition of human nasopharyngeal carcinoma growth and metastasis in mice by adenovirus-associated virus-mediated expression of human endostatin. *Mol Cancer Ther* 2006;5:1290.
20. Qin L, Zhang X, Zhang L, et al. Downregulation of BMI-1 enhances 5-fluorouracil-induced apoptosis in nasopharyngeal carcinoma cells. *Biochem Biophys Res Commun* 2008;371:531.
21. Wu J, Xiao X, Zhao P, et al. Minicircle-IFN γ induces antiproliferative and antitumoral effects in human nasopharyngeal carcinoma. *Clin Cancer Res* 2006;12:4702.
22. Hsu YH, Lin JJ. Telomere and telomerase as targets for anti-cancer and regeneration therapies. *Acta Pharmacol Sin* 2005;26:513.
23. Zhang H, Yang MH, Zhao JJ, et al. Inhibition of tankyrase 1 in human gastric cancer cells enhances telomere shortening by telomerase inhibitors. *Oncol Rep* 2010;24:1059.
24. Beisner J, Dong M, Taetz S, et al. Nanoparticle mediated delivery of 2'-O-methyl-RNA leads to efficient telomerase inhibition and telomere shortening in human lung cancer cells. *Lung Cancer-J Iaslc* 2010;68:346.
25. Elmore S. Apoptosis: A review of programmed cell death. *Toxicol Pathol* 2007;35:495.
26. Rosen A, Casciola-Rosen L. Macromolecular substrates for the ICE-like proteases during apoptosis. *J Cell Biochem* 1997;64:50.