Fractionation and characterization of satellite DNAs of the kangaroo rat (*Dipodomys ordii*)

F.T.Hatch, J.A.Mazrimas

Biomedical Division, Lawrence Livermore Laboratory, University of California, Livermore, California 94550, USA

Received 6 February 1974

ABSTRACT

Nuclear DNA from liver cells of the kangaroo rat species *Dipodomys ordii* was fractionated and characterized with the aid of buoyant density gradients in neutral and alkaline CsCl and in Ag^+ - Cs_2SO_4 . More than one-half of the DNA was present in three density satellites, a greater proportion than in any other species yet reported; the purified satellite DNAs were denser than principal DNA. All satellite fractions revealed sharp isopycnic bands and narrow denaturation profiles. Two had identical buoyant densities but differed substantially in T_m , base composition, and reassociation kinetics. In alkaline CsCl all three satellites, as well as a shoulder of intermediate repetitive DNA on the heavy side of the principal band, revealed unique strand densities. The most highly repetitive satellite was unusually rich in (G + C) and contained 6.7% of 5-methylcytosine. A survey of internal organs and spermatozoa of an adult male revealed no significant differences in distribution of the satellites among tissues.

INTRODUCTION

The nuclei of eucaryotic cells contain much more DNA than is accounted for by current knowledge of the requirements for coding and metabolic regulation. A significant fraction of the excess DNA consists of reiterated series of similar copies of base sequences¹. The most highly repetitive sequences, designated satellite DNAs, differ from the remainder of nuclear DNA in buoyant density in salt concentration gradients. Satellite DNA has become the focus of intensive study because of its relative ease of isolation and because few clues have been found to explain its function in the cell.

In studies of animals living in a tritiated desert ecosystem² we discovered that the most prevalent small mammal *Dipodomys ordii* exhibited more satellite DNA than any species reported up to that time. This paper describes the fractionation and physiochemical characterization of three satellites from this species, which together comprise more than one-half of

the total cell DNA. Previous communications described the density gradient pattern and an unusual thermal ultraviolet transition curve³, the enrichment of the satellites in the nucleolar and heavy chromatin fractions derived from the cell nuclei⁴, and a remarkable variability in satellite distributions among other species of the genus⁵. Hybridization of complementary RNA to DNA in situ revealed that one satellite was localized to the centromeres of most chromosomes. The other two satellites were located throughout the short arms of the same chromosomes and in the centromeres only of about three pairs of chromosomes that lacked the first satellite⁶. These locations corresponded heterochromatin as revealed by C-banding. Replication of the satellites in cultured cells occurred asynchronously in the second half of S period, when the replication of principal DNA was essentially completed⁷. Finally, a basic sequence of ten nucleotide pairs, with minor variations, has been determined for the centromeric satellite⁸.

MATERIALS AND METHODS

(a) Preparation of DNA solutions

Dipodomys ordii (presumed subspecies monoensis) were trapped alive in various locations at the Nevada Test Site of the U.S. Atomic Energy Commission. Livers were excised after cervical dislocation, and the DNA was isolated according to Kay, Simmons & Dounce⁹ with the modifications described previously³. The DNA samples were dissolved in 0.01 M EDTA, pH 8.0 by shaking overnight in a mechanical shaker. After filtration through glass-fiber prefilters (Millipore Filter Corp., Bedford, Mass.) to remove dust and any undissolved DNA, the solutions were dialyzed twice against 0.01 M KCl-0.0005 M EDTA, pH 8.0.

The samples to be subjected to thermal denaturation and measurement of reassociation kinetics were dialyzed against 0.06 M KCl-0.0005 M EDTA and sheared by repeated passage through a 27 gauge hypodermic needle or by two passages through a French Press at 20,000 psi.

(b) Survey of tissues for satellite DNAs

To find out whether the distribution of satellite DNAs varied among

different cell types, the following tissues were assayed: heart, kidney, skeletal muscle, brain, spleen, liver, lung, thymus gland, and a pool of mesenteric and inguinal lymph nodes. About 100 mg of frozen tissue was thawed and homogenized in 2 ml of 0.15 M NaCl-0.015 M Sodium citrate (SSC) and centrifuged to eliminate cell debris, blood components, and RNA; the pelleted chromatin material was again homogenized and centrifuged. The resulting pellet was homogenized in 1.6 ml of 0.15 M NaCl, and 0.16 ml of 5% sodium dodecyl sulfate and 50 μ l of DNAase-free Pronase solution (2 mg/ml) were added. The mixture was shaken for 3 hr at room temperature. A sample of this solution was then submitted to analytical CsCl centrifugation as described in the next section.

Spermatozoa were collected from the epididymis and vas deferens of male animals. After the spermatozoa concentration was determined by counting in a hemacytometer, aliquots of sperm were pelleted by centrifugation and resuspended in 0.5 ml 0.01 M KCl-0.0005 M EDTA; sodium N-lauroyl sarcosinate (5% solution) was added to a final concentration of 0.5%, together with DNAase-free Pronase, 40 μ g. This mixture, at room temperature, was stirred vigorously at intervals for one hour. The resulting solution was submitted to analytical CsCl centrifugation as described below. Contamination of the sperm preparations by diploid cells was estimated from the hemacytometer counts to be less than 4%.

(c) Fractionation by preparative density-gradient centrifugation (Fig. 1)

The DNA samples were prepared for Cs_2SO_4 - Ag^+ gradients by dialysis against a solution containing 0.01 M $\text{Na}_2\text{B}_4\text{O}_7$ and 0.01 M Na_2SO_4 ; precautions were taken to exclude traces of Cl^- . To 4.15 ml of 0.01 M borate buffer at pH 8.0 were added 4.67 g Cs_2SO_4 (Harshaw Chemical Co. optical grade), 1.0 ml of the dialyzed DNA solution containing 100 to 250 μ g DNA, and 1.0 ml of freshly prepared solution containing 0.175 μ g AgNO_3 per μ g of DNA. The order of reagent addition was critical. The final conditions were: volume 7.0 ml, density 1.54 g/ml, pH 9.0-9.3, and molar ratio of Ag^+ to DNA-phosphorus 0.33. Centrifugation was in a 50 Ti rotor in a Spinco Model L2 ultracentrifuge for 48 hr at 38,000 rpm. Fractions of

0.2 ml were collected after puncture of the tube bottoms and absorbance was measured at 260 nm. Peaks A, B and C in Fig. 1 are lettered in order of removal from the tubes. Removal of Ag^+ was necessary prior to further use of solvents containing Cl^- ; therefore, the pooled fractions were dialyzed successively against large volumes of 0.01 M NaCN and 0.01 M KCl-0.0005 M EDTA.

Approximately 50- μg samples of DNA from each of the pooled fractions then were centrifuged in 7 ml of neutral CsCl solution (Harshaw radiotracer grade) of initial density 1.71 g/ml. Centrifugation was performed as above for 60 hr at 35,000 rpm. Fractions of 0.1 ml were collected and measured as above. For optimal purification of the DNA fractions, preparative neutral CsCl gradient centrifugation was repeated once or twice. In each case appropriate fractions were pooled, dialyzed, and monitored by neutral analytical CsCl gradients (Fig. 1) as described below. The final products were also examined in alkaline CsCl gradients.

(d) Analytical density gradient ultracentrifugation

Approximately 2 μg of the whole DNA or its fractions were submitted to neutral isopycnic CsCl (Harshaw optical grade) centrifugation in an AnG rotor. The initial density ρ_0 of the solution was 1.710 g/ml. Poly(dAT-dAT) ($\rho = 1.679$ g/ml) and in some cases DNA from Micrococcus lysodeikticus ($\rho = 1.731$ g/ml) were added singly or together as density markers for calculating relative densities as described by Ifft, Voet & Vinograd¹⁰. All densities values are expressed on a scale in which E. coli DNA is assigned a density of 1.710 g/ml. After 24 hr at 40,000 rpm in a Spinco Model E Ultracentrifuge, the cells were scanned at 265 nm with an ultraviolet multiplex scanner.

Alkaline CsCl centrifugations were performed with the AnF rotor at 52,000 rpm as described above, except that ρ_0 of the solutions was 1.76 to 1.78 g/ml. The pH of the solutions was adjusted to 12.5 to 12.8 with 1 M NaOH, and EDTA was added to a final concentration of 0.001 M. Poly(dAT-dAT) was used as a marker ($\rho = 1.722$ g/ml in alkali) for the calculations of relative density.

Fig. 1. Flow chart showing the analytical buoyant density gradient patterns developed during the fractionation and characterization of DNA from livers of *D. ordii*. (a) Initial pattern of whole DNA in neutral CsCl. (b) Ag⁺-CsSO₄ gradient separation. (c) Neutral CsCl patterns for each peak from the Ag⁺-Cs₂SO₄ gradient, before further purification. (d) Neutral CsCl gradients for the final products. (e) Alkaline CsCl gradients (pH 12.5) for each fraction.

Integrations of the areas and percentages of each peak were performed on the Model 310 Curve Resolver (E. I. duPont de Nemours Co., Wilmington, Delaware) as previously described, except that measurements were made directly on the scanner tracings³. A minimum number of gaussian peaks was employed to outline the total area of density distribution of DNA. For unfractionated DNA samples four components were required at positions corresponding to densities of 1.698, 1.702, 1.707, and 1.713 g/ml in neutral CsCl.

(e) Determination of thermal denaturation profiles

Thermal denaturation analysis was done in a Gilford Model 2400 recording spectrophotometer (Gilford Instrument Co., Oberlin, Ohio) modified for digital recording of both temperature and absorbance readings. Melting profiles were monitored at 260 nm, and all data were corrected for thermal expansion of the solvent. The calculations and plots were made with the aid of a computer. Normalized first derivatives of the thermal denaturation curves were obtained by sequentially fitting a parabola to each successive series of seven data points, with determination of the midpoint tangents by a computer subroutine.

(f) Reassociation kinetics

The rate of reassociation of the separated strands of sheared heat-denatured DNA fractions was measured by following the return of hypochromicity at 260 nm with the recording spectrophotometer. The data were analyzed with a computer program based on the methods of Wetmur and Davidson¹¹ and Thrower and Peacocke¹².

(g) Base composition

About 50 µg of DNA in 0.3 ml of 0.05 M Tris-HCl buffer (pH 7.1) was incubated at room temperature with 10 µl of DNAase I solution (5 mg/ml, Worthington Biochemical Corp. code DPFF) plus 10 µl of 0.25 M MgCl₂ and 10 µl of 0.625 M CaCl₂ for 2 hr. The pH was raised to 9.5 by the addition of 1 N NaOH; 30 µl of venom phosphodiesterase I (8 mg/ml, Worthington code VPH in solution containing 10 mg/ml of bovine albumin) was added and mixed,

and the mixture was incubated for three hr at 37°C. Then an additional aliquot of enzyme was added and digestion was allowed to continue for 1 hr. The pH of the digest was adjusted to about 4.0 with 1 N HCl and 1 ml of 0.001 M formic acid was added with about 20 mg of partially deactivated charcoal^{13,26}. After thorough mixing and standing for 10 min, the digest was centrifuged at 1200 xg in a clinical centrifuge for 2 min; the supernatant was removed and one ml of fresh formic acid solution was added to the charcoal residue. After the charcoal was washed and centrifuged twice more, 1 ml of ethanol-ammonia solution (3 ml conc. ammonia in 100 ml 50% ethanol) was added. The solution was mixed and allowed to stand 15 min. After 5 min of centrifugation at high speed, the almost clear supernatant was carefully removed and evaporated under nitrogen at 50°C to about 0.25 ml. This concentrated nucleotide solution was centrifuged again at high speed to remove the remaining charcoal fines and evaporated to dryness. Then 50 µl of phosphate buffer (0.002 M KH_2PO_4 , pH 4.15) was added in preparation for column chromatography. The recovery of nucleotides was about 70%. Significant losses are customarily encountered in this procedure; however, recovery of standard mixtures indicated that no bias was introduced into the base composition.

For separation of the 5'-mononucleotides 5 µl of the concentrated nucleotide preparation was injected into a Model 830 Dupont high-pressure liquid chromatograph fitted with a one-meter anion exchange column (Dupont AAX); elution was with 0.002 M KH_2PO_4 at pH 4.15. Separation was effected in about 25 minutes at 1200 psi. Resolution of the 5'-deoxymononucleotides was complete except for partial overlapping of 5-methyldeoxycytidylate and deoxycytidylate. Integration of peaks on the chart was by means of an attached electronic chart integrator. The response factor of each nucleotide was determined with pure standards purchased from P-L Biochemicals, Inc. The published base compositions of mouse satellite and

guinea-pig satellite DNAs¹ were verified using this system.

(h) Nomenclature of the satellite DNAs (based on *D. ordii*

In a previous report³, we showed that in neutral CsCl gradients, *D. ordii* DNA is resolved into two satellite peaks, designated MS (1.707 g/ml) and HS (1.713 g/ml). As shown below, we have further fractionated the HS peak on Ag^+ - Cs_2SO_4 and alkaline CsCl gradients and have shown it to consist of two components. To be consistent with the nomenclature of Flamm, Walker & McCallum¹⁴ for guinea pig satellites, the component with the greater compositional bias between strands (in alkaline CsCl) was assigned the symbol α ; the component with the lesser bias was assigned the symbol β .

RESULTS

(a) Isopycnic density gradients

(i) Neutral CsCl

Four subspecies of *D. ordii* from different geographic locations, (Nev., Ariz., Tex., Colo.) gave very similar density gradient patterns. The differences did not exceed those observed among different preparations of the Nevada subspecies, and depended somewhat on the duration of frozen storage of liver before DNA isolation. Likewise, no significant differences were noted in the patterns of DNA from liver, lung, thymus, spleen, lymph nodes, kidney, heart, skeletal muscle, brain, and spermatozoa of a single animal.

(ii) Ag^+ - Cs_2SO_4 Gradients

Liver DNA in this special preparative gradient was fairly completely resolved into three peaks (Fig. 1b) when the ratio of Ag^+ to DNA was optimal. When, after dialysis, the three components were recentrifuged in neutral CsCl (Fig. 1c), the heavier peak (A) was found to consist of approximately equal quantities of satellite DNA banding at 1.707 g/ml and 1.713 g/ml and to be contaminated with some lighter DNA; the large middle peak (B) consisted of principal DNA banding at 1.698 g/ml, another major component at 1.702 g/ml which we have called "intermediate" DNA, together with a

small amount of satellite DNA at 1.707 g/ml; the small light peak (C) consisted of satellite banding at 1.713 g/ml contaminated with a small amount of satellite at 1.707 g/ml.

(b) Fractionation of satellite, intermediate and principal DNA

The general scheme for fractionation of D. ordii DNA is given in the flow chart of Fig. 1. After the initial Ag^+ - Cs_2SO_4 gradient centrifugation described above, the three separated peaks (Fig. 1b) were run individually in analytical (Fig. 1c) and in preparative gradients of neutral CsCl. The preparative gradient derived from the heavy peak (A) was separated into two pools essentially in the middle; that derived from the light peak (C) was simply improved by discarding small amounts from both sides of the peak. The fractions thus selected were subjected to one or two additional preparative gradients in neutral CsCl, monitored by analytical gradients. The final products (Fig. 1d), one satellite (MS) of density 1.707 g/ml and two satellites (HS- α and HS- β) of density 1.713 g/ml, were estimated to be 90-95 percent pure.

The preparative CsCl gradient derived from the middle peak (B) was broad and devoid of fine structure. Further fractionation was much less satisfactory than in the case of the satellites. In order to achieve the best possible separation of the two major components at densities 1.698 and 1.702 g/ml that appeared in the analytical monitoring gradient, single fractions estimated to be closest to those densities in the preparative gradient (shaded bands in Fig. 1c) were subjected to further preparative gradients as above, with the result shown in Fig. 1d. The component banding chiefly at 1.698 g/ml, which we have called principal DNA by analogy with other mammalian DNAs and because it is constantly present in all Dipodomys species we have examined, is significantly contaminated with the 1.702 g/ml component⁵. The component banding chiefly at 1.702 g/ml, designated intermediate DNA which is variable among Dipodomys species, contains DNA at both 1.698 and 1.707 g/ml that was not eliminated by repeated CsCl gradients.

On the basis of the relative areas of ultraviolet absorbance at 265 nm

in the analytical gradient patterns the proportions of total DNA contributed by each component were estimated to be: HS- α , 19%; HS- β , 11%; MS, 22%; intermediate DNA, 15%; principal DNA, 33%. Quantitative recovery of fractions throughout the series of preparative gradients was impossible because of intentional sacrifice of yield in favor of purity.

(c) Properties of DNA fractions

(i) Buoyant density

In neutral CsCl gradients the buoyant densities of each DNA fraction obtained were as shown in the flow chart (Fig. 1). In Ag^+ - Cs_2SO_4 gradients, buoyant densities are critically dependent on the molar ratio of Ag^+ to DNA phosphorus. Our data are qualitative in this respect, indicating that HS- α and MS bound more Ag^+ than principal and intermediate DNA, and that HS- β bound less Ag^+ or perhaps none (Fig. 1b). The buoyant densities in alkaline CsCl for the single strands of each fraction are shown in Fig. 1e. The satellites show distinct resolution into unlike strands. This is possibly true of intermediate DNA; however, the principal DNA, in common with that of other mammals, shows no dissimilarity of single strands.

(ii) Thermal denaturation profiles

Table I gives the melting data for the purified DNA fractions that had been sheared in preparation for reassociation experiments, and for two comparison DNAs. The midpoints of the thermal transitions T_m increased progressively with increased buoyant density, except that the T_m for HS- β was about 6°C higher than that for HS- α of the same density. Indications of the sharpness of the transitions are given by the maxima in the derivative curves, expressed as change in molar extinction at 260 nm per °C, and by the width in °C of the transitions between 0.25 and 0.75 of completion. As expected, the satellite fractions melted more sharply than the more heterogeneous principal and intermediate DNAs. The thermal transition curve for HS- α revealed a small second transition at about 86°C, which was indicative of minor contamination with HS- β . It is possible that some fragments of DNA isolated in density gradients contain segments of both satellite sequences.

Table 1

Thermal ultraviolet transition data on sheared principal, intermediate and satellite DNA fractions from *Dipodomys ordii* and comparison species^a

Fraction	T _m	Max d ε/dT ^b	σ .25 - .75 ^c
<u><i>Dipodomys ordii</i> fractions</u>			
Principal (1.698 g/ml)	78.4	327	4.6
Intermediate (1.702 g/ml)	78.9	309	4.5
MS (1.707 g/ml)	79.5	794	1.4
HS-α (1.713 g/ml)	80.4	728	1.3
HS-β (1.713 g/ml)	86.6	452	2.9
<u>Comparison DNAs</u>			
Mouse satellite	76.3	717	1.6
Guinea pig β + γ satellites ^d	79.6	552	2.2

^aSolvent: 0.06 M KCl-0.0005 M EDTA.

^bMaximum rate of melting, expressed as change of molar extinction at 260 nm per °C.

^cBreadth in °C between 0.25 and 0.75 of completion of transition.

^dUnresolved mixture of the two satellites, which have similar properties.

(iii) Preliminary information on reassociation kinetics

Our studies of the reassociation of the DNA fractions with reiterated sequences from the kangaroo rat indicate that at least two parameters are important in characterizing the process. Both the rate and the final extent of reassociation differed among the fractions. In addition we found that the fractions are affected differently by the stringency of reaction conditions i. e., temperature and ionic strength, and that the correction factors for ionic strength given by Britten & Smith¹⁵ do not apply to our material. Therefore, we have elected to report the observed second-order rate constants and the fractional completion of the spectrophotometric reaction (Table 2). Data are given also for comparison DNAs measured under the

same conditions. The conditions selected, namely, 0.06 M ionic strength and 62°C, combine practical utility and rather high stringency. The temperature of reassociation was approximately 17°C below the T_m of all fractions except HS- β (25°C); however, for HS- β there was little difference in rate and extent of reassociation over the range of 15-25°C below T_m . The satellite HS- β reassociated nearly three times faster than the mouse and guinea pig satellites, but to a slightly lesser final extent. The HS- α and MS satellites of kangaroo rat reassociated about one order of magnitude more slowly and to a substantially lesser final extent than HS- β or the comparison samples. Further study of the effects of varying DNA molecular weight, ionic strength and temperature on reassociation rates is in progress.

Table 2

Reassociation kinetic data from sheared heat-denatured DNA fractions^a

Fraction	k_2^b	Extent of Reaction ^c
<u>Dipodomys ordii</u> fractions	$M^{-1} sec^{-1}$	%
Principal (1.698 g/ml)	30	14
Intermediate (1.702 g/ml)	20	17
MS (1.707 g/ml)	37	39
HS- α (1.713 g/ml)	36	30
HS- β (1.713 g/ml)	490	59
Comparison DNAs		
Mouse satellite	170	70
Guinea pig $\beta + \gamma^d$	182	70
T2 Bacteriophage	0.18	103

^aSolvent: 0.06 M KCl-0.0005 EDTA. Temperature 62°C.

^bSecond-order rate constant.

^cPercent of restoration of original hypochromicity (at 260 nm) of the DNA at the extrapolated termination of reassociation.

^dUnresolved mixture of the two satellites, which have similar properties.

(iv) Base composition

The base composition data on whole DNA and its fraction from *D. ordii* are presented in Table 3. Two findings are noteworthy: first, the G + C + 5 mC content of the HS- β satellite (66.1%) was very much higher than that of HS- α (43.4%) despite their identical buoyant densities, and second, the 5-methylcytosine content of HS- β was 6.7%, which is probably the highest value yet observed in an animal DNA^{16,17}.

Table 3

Base composition of DNA fractions from *D. ordii*

	5dMeCMP	dCMP	dTMP	dAMP	dGMP	G + C + 5mC ^b
Whole DNA (100) ^a	1.5	19.7	29.7	27.3	21.6	42.8
1.698 g/ml (33)	0.9	18.9	31.9	31.9	16.2	36.0
1.702 g/ml (15)	0.5	20.1	30.2	28.5	20.6	41.2
HS- α (19)						
H-strand	1.0	3.0	34.8	21.8	39.2	
L-strand	0.9	37.7	23.1	33.3	5.0	
duplex	1.0	20.3	28.9	27.5	22.1	43.4
HS- β (11)						
duplex	6.7	25.8	16.5	17.4	33.6	66.1
MS (22)						
H-strand	0.3	26.4	51.2	8.5	13.8	
L-strand	0.3	12.6	11.7	49.6	25.7	
duplex	0.3	19.5	31.4	29.0	19.7	39.5

^aFigures in parentheses are the percent of whole DNA represented by each fraction, estimated by curve-resolver analysis of tracings from analytical buoyant-density gradients.

^bG = dGMP, C = dCMP, and 5 mC = 5dMeCMP.

DISCUSSION

More than one-half of the nuclear DNA of the kangaroo rat *Dipodomys ordii* is distributed among two heavy satellites of identical buoyant density (HS- α and HS- β , 1.713 g/ml) and one satellite of intermediate density (MS, 1.707 g/ml). The remaining DNA can be divided into partially separable components, an "intermediate" DNA at 1.702 g/ml and principal DNA at 1.698 g/ml. This distribution applied to four subspecies of *D. ordii* obtained

in different locations, and to a variety of internal organs, including sperm. However, an anomalous and possibly questionable subspecies D. ordii compactus showed a different pattern (Mazrimas, Hatch & Best, unpublished data), and other species of the genus showed a remarkable variability that was correlated with phylogenetic and karyotypic characteristics⁵.

Fractionation of the three satellites to rather high purity and partial separation of the intermediate and principal DNAs could be accomplished only by successive preparative centrifugations in isopycnic gradients containing Ag^+ - Cs_2SO_4 and CsCl . Separation was aided particularly by considerable binding of Ag^+ to HS- α and MS, an intermediate degree of binding to principal and intermediate DNAs, and minimal or absent binding to HS- β . The variations in binding of Ag^+ are not related in an obvious way to the overall base compositions of the fractions.

The physicochemical properties of the satellites were typical of those observed previously in other animals, for example the mouse and guinea pig used for comparison in this study¹. Banding at equilibrium in neutral buoyant density gradients was sharp; two bands of different densities appeared in alkaline CsCl gradients; thermal transitions were sharp; reassociation of complementary strands from the denatured state under appropriate conditions was rapid. The foregoing properties are consistent with a high degree of homogeneity of fragments of the satellites arising from numerous iterations of a short basic sequence. Indeed, a basic sequence of only ten nucleotide pairs, with minor variations has been determined for HS- β by Fry *et al.*⁸. The basic sequence is as follows:

5' - - - A C A C A G C G G G - - - 3' Light strand

3' - - - T G T G T C G C C C - - - 5' Heavy strand

Unlike the sequence proposed by Southern¹⁸ for the guinea pig α satellite, this sequence contains no nonsense codons. It is not known whether the HS- β satellite is transcribed *in vivo*, but if there is a random interspersion of the observed 10, 11, and 12 nucleotide variants of the basic sequence it is difficult to see how it could code for a useful protein. Furthermore, one strand of the HS- β satellite lacks thymine and the other strand lacks adenine.

This drastically limits the possible amino acids that either strand could code for if it were transcribed. The bulk of available evidence on other satellites does not indicate transcription in vivo of the very highly repetitive sequences¹.

Previously some satellite DNAs have been shown to deviate from empirically observed relationships between T_m and base composition¹⁹ and between buoyant density and base composition²⁰ as reviewed by Flamm²¹ (1972). Among the satellites of D. ordii there are obvious deviations in the two satellites HS- α and HS-B, which have the same density but very different T_m and base composition (see ref. 22 for a similar example in Crustacea). For all three of the kangaroo rat satellites, the prediction of base composition (G + C) from buoyant density was grossly in error. The predictions from T_m at three different ionic strengths were in error for HS- β , but in reasonable agreement for HA- α and MS. The physicochemical basis for these discrepancies is unknown. Although 5-methylcytosine alters both the T_m and the density of synthetic polynucleotides, the proportion of the base in these satellites is insufficient to account for the observed deviations (J. Gill, personal communication). A number of departures from normal DNA structure as revealed by X-ray diffraction have been noted in the regularly repeating sequences of synthetic polynucleotides by Langridge²³. Alterations in helical pitch, base-pair alignment, and nearest-neighbor frequencies can reasonably affect the hydrodynamic properties and the strength of strand interactions; however, the molecular mechanics of the postulated anomalies and any consequent implications for the role of repetitive sequences in chromatin are as yet unknown.

The properties of intermediate DNA are under investigation. Purification is difficult and incomplete at this time. Enriched material appears to separate into two strands in alkaline CsCl and to contain some proportion of repetitive sequences. One hypothesis that we are testing by further experiments is that intermediate DNA consists of tandem combinations of covalently linked principal and satellite DNAs. Progressive scission of these molecules in the CsCl gradients may account for the continuing heterogeneity of observed densities.

Neither the evolutionary origin nor the present-day function of satellite and other highly repetitive DNA sequences are satisfactorily understood. There is increasing evidence of their strong association with nuclear heterochromatin²⁴, but this information does little to elucidate their roles. The availability of a mammal containing major quantities of three different satellites and the demonstration that these occupy large blocks in the genome is expected to promote progress in the field. Determination of analogies that may exist between the extraordinary situation in D. ordii and the less spectacular anomalies of DNA in most higher organisms, including man²⁵, seems to be a necessary and promising extension of this research.

Acknowledgments

This work was performed under the auspices of the U.S. Atomic Energy Commission. Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Atomic Energy Commission to the exclusion of others that may be suitable.

We acknowledge gratefully the assistance of Mr. M. L. Stuart in trapping some of the kangaroo rats, and the advice of Prof. V. M. Schumaker on density gradients and that of Mr. S. P. Krotz on ultracentrifugation.

REFERENCES

- 1 Walker, P. M. B. (1971) Prog. Biophys. Molec. Biol. 23, 145-190.
- 2 Hatch, F. T., Mazrimas, J. A. Koranda, J. J. and Martin, J. R. (1970) Radiation Res. 44, 97-107.
- 3 Hatch, F. T. and Mazrimas, J. A. (1970) Biochim. Biophys. Acta 224, 291-294.
- 4 Mazrimas, J. A. and Hatch, F. T. (1970) Exptl. Cell Res. 63, 462-466.
- 5 Mazrimas, J. A. and Hatch, F. T. (1972) Nature New Biol. 240, 102-105.
- 6 Prescott, D. M., Bostock, C. J., Hatch, F. T. and Mazrimas, J. A. (1973) Chromosoma 42, 205-213.
- 7 Bostock, C. J., Prescott, D. M. and Hatch, F. T. (1972) Exptl. Cell Res. 74, 487-495.
- 8 Fry, K., Poon, R., Whitcome, P., Idriss, J., Salser, W., Mazrimas, J. and Hatch, F. (1973) Proc. Nat. Acad. Sci. U.S. 70, 2642-2646.
- 9 Kay, E. R. M., Simmons, N. S. and Dounce, A. L. (1952) J. Am. Chem. Soc. 74, 1724-1726.
- 10 Ifft, J. B., Voet, D. H. and Vinograd, J. (1961) J. Phys. Chem. 65, 1138-1145.
- 11 Wetmur, J. G. and Davidson, N. (1968) J. Mol. Biol. 31, 349-370.
- 12 Thrower, K. J. and Peacocke, A. R. (1968) Biochem. J. 109, 543-557.
- 13 Symons, R. H. (1970) Biochim. Biophys. Acta 209, 296-305.
- 14 Flamm, W. G., Walker, P. M. B. and McCallum, M. (1969) J. Mol. Biol. 42, 441-455.
- 15 Britten, R. J. and Smith, J. (1970) Carnegie Inst. Washington Yearbook

68, 378-386.

16 Salomon, R., Kaye, A.M. and Herzberg, M. (1969) J. Mol. Biol. 43, 581-592.

17 Vanyushin, B.F., Tkacheva, S.G. and Belozersky, A.N. (1970) Nature 225, 948-949.

18 Southern, E.M. (1970) Nature 227, 794-798.

19 Marmur, J. and Doty, P. (1962) J. Mol. Biol. 5, 109-118.

20 Schildkraut, C.L., Marmur, J. and Doty, P. (1962) J. Mol. Biol. 4, 430-443.

21 Flamm, W.G. (1972) Int. Rev. Cytol. 32, 1-51.

22 Skinner, D.M., Beattie, W.G., Kerr, M.S. and Graham, D.E. (1970) Nature 227, 837-839.

23 Langridge, R. (1969) J. Cell Physiol. 74, Suppl. 1, 1-20.

24 Yunis, J.J. and Yasmineh, W.G. (1972) Adv. Cell Molec. Biol. 2, 1-46.

25 Hsu, T. C. Arrighi, F.E. and Saunders, G.F. (1972) Proc. Nat. Acad. Sci., (U.S.) 69, 1464-1466.

26 We are indebted to Mr. Kirk Fry and Dr. Winston Salser, Department of Zoology, University of California, Los Angeles, for gifts of partially deactivated charcoal and advice on the desalting of nucleotides.