Primary structure of tRNA^{Lys} of *E. coli* BKalpana Chakraborty, Assaf Steinschneider*, Richard V. Case[†], Alan H. Mehler*

Department of Biochemistry, Medical College of Wisconsin, Milwaukee, WI 53233, USA

Received 27 August 1975

ABSTRACT.

The primary structure of tRNA^{Lys} of *E. coli* was determined by use of [³²P]-tRNA. The sequence is pGGGUCGUUAGCUCAGDDGGDAGAGCAGUUGACUmam⁵-s²-UUU-t⁶AAΨCAAUUGm⁷GXCAGAGGTΨCGAAUCCUGCAGACCCACCA. No s⁴-U was detected in position 8. No other lysine tRNA was detected but the existence of another species has not been ruled out.

INTRODUCTION.

During the past 10 years the primary structures of many species of tRNA have been determined¹. Although it has not yet been possible to define completely recognition sites, binding sites or other features that affect enzyme activity, many interesting approaches to analyzing the interaction between aminoacyl-tRNA synthetases and tRNA have been made possible by knowledge of the tRNA structure². Since each enzyme of this series is obviously different from the others in many respects³, it is necessary to determine the structure of the cognate tRNA to try to understand the properties of the aminoacyl-tRNA synthetases. We have previously purified and studied some properties of the lysyl-tRNA synthetase of *E. coli*⁴ and now we describe the structure of the corresponding tRNA.

MATERIALS AND METHODS

tRNA^{Lys} was prepared by modifications of the method of Gillam and Tener⁵ from unlabeled *E. coli* B in early studies⁶. For subsequent analyses [³²P]-tRNA was prepared by methods modified from those described previously.⁷ The bacteria were harvested in late log phase after cooling the flasks in ice water. After the phenol treatments, the RNA was extracted with 1 M NaCl in the cold. Stripping of amino acids was carried out in 0.1 M Tris, pH 8.0, 0.5 M NaCl, 10⁻²M MgCl₂ and 10⁻²M mercaptoethanol for 60 min at room temperature. "Sham" phenoxyacetylation was

carried out on the stripped RNA before passage through a BD cellulose column. The lysine tRNA was charged with a purified synthetase and the preparation was treated with 10^{-2} M periodate for 20 min at 0° in the dark. Excess periodate was destroyed with 4 μ l of ethyleneglycol and the RNA was precipitated with ethanol. The precipitate was suspended in 4 ml of 50% (V/V) Sepharose 4B modified with hydrazine⁸ and shaken 60-90 min at room temperature. The gel was transferred to a Pasteur pipette, drained and washed with 5 ml of 1 M NaCl. The combined effluents contained 10-14% of the applied radioactivity and 80-90% of tRNA^{Lys}. The NaCl was diluted to 0.3 M and the solution was applied to a second BD cellulose column, which was eluted with 1 M NaCl, then with a gradient of 1 M NaCl to 2.5 M NaCl containing 25% ethanol. tRNA^{Lys} precipitated from the major peak with 67% ethanol was at least 95% pure as judged from ribonuclease T1 fingerprints, acrylamide gel electrophoresis and lysine acceptance.

The approach of Holley *et al.*⁹ was used initially with fragments of tRNA separated according to Tomlinson and Tener¹⁰. The methods of Sanger as developed by Barrell¹¹ and Brownlee¹² were used for analysis of [³²P] tRNA. Modified nucleotides were identified by thin-layer chromatography¹³.

RESULTS.

The purity of the tRNA prepared from unlabeled cells was evaluated by the content of esterified lysine, which ranged from 1300 to 1650 pmoles per OD₂₆₀ unit. Rechromatography of the stripped nucleic acid on BD cellulose showed 95% of the absorbancy to elute in a single symmetrical peak. [³²P] tRNA was of similar purity; lysine acceptance was approximately 1 mole per mole of tRNA and the fingerprints of digests with ribonuclease A and ribonuclease T1 (Fig. 1) are characteristic of a single species without significant contamination.

The identity of each spot on the fingerprints is reported in Table 1. Where oligonucleotides are not separated, as in the case of spot 5 of the ribonuclease A digest, subsequent analysis of the components following additional hydrolysis revealed the composition of the mixture. For example, ribonuclease T1 digestion of spot 5 produced two equivalents each of Gp and ApCp for each equivalent of ApGp and Cp. Similar analyses resolved the mixture in spot 9. Some faint spots, such as spot 10, appear to represent analogues of major fragments with incomplete modification of minor bases. The sum of spot 10 and spot 14 equals one equivalent of trinucleotide. The minor spot 8 is related to spot 11 according to TLC chromatography of total hydrolyzates.

Fig. 1 Fingerprints of Ribonuclease A and Ribonuclease T1 Digests of tRNA^{Phe} A, Radioautogram of ribonuclease A digest; B, radiogram of ribonuclease T1 digest.

Spot 16 of the ribonuclease T1 digest usually appears as a double spot. We have not been able to characterize the two portions but it seems likely that the same incomplete modification of a base (probably the t⁶A) responsible for the appearance of spot 10 in the ribonuclease A digest is responsible for the second component of spot 16.

The small size of most fragments and the large number of free mononucleotides liberated limit the amount of ordering that can be done on the basis of overlapping sequences in the two digests. Therefore, it was necessary to use partial digestion of the tRNA to prepare larger fragments that contain longer sequences. Fig. 2 shows the unique linear sequence deduced from analyses of the overlapping regions. The conventional clover-leaf folding of tRNA molecule is shown in Fig. 3.

TABLE I

Fragments Released from tRNA^{Lys} by Ribonucleases

Spot No.	Ribonuclease A	Ribonuclease T1
1	U (8)	G (4)
1a	ψ	
1b	U!	
1c	mam ⁵ -s ² -U	
1d	D	
2	C (9)	AG
3	AC (2)	CAG (2)
4	GC (2)	CACG
5	AGC (1), GAC (2)	ACCCACCA
6	AAU	pG
7	GU	DAG
8		UCG
9	AGD, AGU	m ⁷ GXCG
10	-	CUCAG
11	Gm ⁷ GXC	DDG
12	AGAGC	UUG
13	GAAU	UUAG
14	t ⁶ AAψ	TψCG
15	AGGT	AAUCCUG
16	GGD	AC (Umam ⁵ -s ² -UUU) t ⁶ AAψCAAUUG
17	pGGGU	

Numbers in parentheses indicate relative quantities released. X appears to be 3-(3-amino-3-carboxypropyl)uridine according to thin layer chromatography.

Fig. 2 The arrows (↑) indicate the sites of ribonuclease T1 cleavage. Only some of the ribonuclease A breaks are indicated (↑) in order to show the fragments that were important in obtaining an unambiguous sequence. The lines above the sequence shows the useful fragments obtained after partial hydrolysis by both ribonuclease A and T1.

Fig. 3 Cloverleaf structure of tRNA^{Lys}.

DISCUSSION.

The sequence deduced above is a unique structure compatible with the structures of the fragments isolated although the order of the series of uracil residues in the anticodon remains to be confirmed. It is very probable, however, that the modification of U to mam⁵-s²-U occurs at the first position of the anticodon as shown by Ohashi *et al.*¹⁴ for this base in glutamic acid tRNA and generalized by Nishimura¹³ for thiobases in the anticodon. Ishikura *et al.*¹⁵ have shown that tRNA^{Lys} of *E. coli* contains t⁶A, which we have shown to be adjacent to the anticodon, as predicted.

The methods of the Sanger group for determining the sequences of most of the spots were straightforward but the pentadecanucleotide containing the anticodon region (spot 16 in the ribonuclease T1 digest) presented special problems. Most of the sequence of this oligonucleotide could be determined from the fragments produced by ribonuclease U2. Some uncertainty remains about the position of the modified U found with three unmodified U's in the anticodon region. Attempts to obtain interpretable fragments of the anticodon region following digestion of intact tRNA^{Lys} by ribonuclease S1 or treatment of an oligonucleotide with silkworm nuclease have not been successful.

4-Thiouridine is found widely among tRNA species of *E. coli* and has been reported to exist in tRNA^{Lys}. Our early analyses on unlabeled tRNA were compatible with s⁴-U in position 8: some ultraviolet absorption was seen at 335 nm and the yield of UUAG (the fragment including U8) from the ribonuclease T1 digest was low⁶. Using the [³²P]tRNA our evidence is against the existence of s⁴-U: we have not detected s⁴-U among the total digestion products, no new spot was seen on fingerprints after irradiation of the intact tRNA with ultraviolet light under conditions that formed cross-linked material in tRNA^{Arg}, which contains s⁴-U in position 8¹⁶, and the yield of UUAG was not diminished.

Previous studies have indicated that at least two separable forms of tRNA^{Lys} exist in *E. coli*¹⁷ but the coding and chemical properties of the material in the separate peaks were identical. In the present work, in which precautions were taken to minimize alteration of the nucleic acid, no evidence was found for a second structure and, as suggested earlier⁷, it is possible that only one species of tRNA^{Lys} is transcribed in this bacterium. The existence of two codons for lysine, on the other hand, evokes caution in concluding that a single tRNA for lysine exists. Separate species that recognize AAA and AAG codons, respectively, have been obtained from rat liver (U.L. RajBhandary, personal communication). Preliminary binding studies using synthetic triplets (Miles Laboratory, kindly given to us by Dr. Barbara Vold) showed about one quarter as much tRNA^{Lys} bound to ribosomes with AAG as with AAA. These results do not allow a definite conclusion to be drawn about the ability of one tRNA to satisfy the needs of *E. coli* to translate both lysine codons. It is possible that the structure determined here is capable of binding to both AAA and AAG but it is also possible that part of the preparation has no sulfur in the first position of the anticodon and that only the sulfur-deficient species interacts with AAG.

Two lysine tRNA structures from yeast have been described^{18,19}. The species obtained from a haploid strain shows very few similarities to the structure presented above. The other species, however, shows many features in common, including the entire dihydrouridine and anticodon loops. The dihydrouridine stem differs only in the methylation of G10. The extra loop contains m⁷G followed by modified pyrimidines. Many differences exist, however, in the other stems, including the 5'-terminus. The similarities and differences may be useful in determining the nature of interaction between nucleic acids and their cognate synthetases.

ACKNOWLEDGEMENTS.

We are grateful for helpful technical advice from Dr. U. L. RajBhandary and for suggestions on purification of the tRNA from Dr. I. Sela. This work was supported by grants GM-13037 from the National Institutes of Health and GM-27440 from the National Science Foundation.

* Present address: Department of Biochemistry, University of Illinois Medical School, Chicago, Illinois

+ Present address: P-L Biochemicals, Inc., Milwaukee, Wis.

REFERENCES.

- 1 Barrell, B.G. and Clark, B.F.C. (1974) Handbook of Nucleic Acid Sequences, Joynson-Bruvvers, England.
- 2 Söll, D. and Schimmel, P.R. (1974) in The Enzymes, Vol. X, pp. 489-538, Academic Press, New York.
- 3 Mehler, A.H. and Chakraburttty, K. (1971) in Advances in Enzymology, Vol. 35, pp. 443-501, edited by A. Meister, John Wiley & Sons, Inc.
- 4 Stern, R. and Mehler, A.H. (1965) Biochem. Z. **342**, 400-409.
- 5 Gillam, I., Blew, D., Warrington, R.C., von Tigerstrom, M. and Tener, G.M. (1968), Biochemistry, **7**, 3459-3468.
- 6 Case, R.V., (1971) Dissertation, Marquette University.
- 7 Steinschneider, A., (1975), J. Chromatography, **103**, 355-363.
- 8 Remy, P., Birmele, C. and Ebel, J.P., (1972) FEBS Letter, **27**, 134-138.
- 9 Holley, R.W., Apgar, J., Everett, G.A., Madison, J.T., Marquise, M., Merrill, S.H., Penswick, J.R. and Zamir, Z., (1965) Science, **147**, 1462-1465.
- 10 Tomlinson, R.V. and Tener, G.M. (1963), Biochemistry **2**, 697-702.
- 11 Barrell, B.G. (1971) in Procedures in Nucleic Acid Research Vol. II pp. 751-811, Edited by G.L. Contoni and D.R. Davies, Harper & Row, New York.
- 12 Brownlee, G.G. (1972) in Determination of Sequences in RNA, American Elsevier, Publishing Co., New York.
- 13 Nishimura, S. (1972) in Progress in Nucleic Acid Research and Molecular Biology Vol. 12, pp. 49-85 Edited by J. N. Davidson and Waldo E. Cohn, Academic Press, New York.
- 14 Ohashi, Z., Saneyoshi, M., Harada, F., Hara, H. and Nishimura, S. (1970), Biochem. Biophys. Res. Comm. **40**, 866-872.
- 15 Ishikura, H., Yamada, Y., Murao, K., Saneyoshi, M. and Nishimura, S. (1969) Biochem. Biophys. Res. Comm. **37**, 990-995.
- 16 Chakraburttty, K. Nucleic Acid Research, in press.
- 17 Söll, D. and RajBhandary, U.L. (1967) J. Mol. Biol. **39**, 113-124.
- 18 Smith, C.J., Ley, A.N., Dobrenan, P. and Mitra, S.K. (1971) J. Biol. Chem. **246**, 7817-7819.
- 19 Madison, J.T., Boguslawski, S.J. and Teetor, G.H., (1972), Science, **176**, 687-689.