Published in final edited form as:

Cell Mol Neurobiol. 1993 June ; 13(3): 247–261.

Chronic Caffeine Alters the Density of Adenosine, Adrenergic, Cholinergic, GABA, and Serotonin Receptors and Calcium Channels in Mouse Brain

Dan Shi¹, Olga Nikodijević¹, Kenneth A. Jacobson¹, and John W. Daly¹

¹ Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892.

SUMMARY

1. Chronic ingestion of caffeine by male NIH strain mice alters the density of a variety of central receptors.
2. The density of cortical A₁ adenosine receptors is increased by 20%, while the density of striatal A_{2A} adenosine receptors is unaltered.
3. The densities of cortical β_1 and cerebellar β_2 adrenergic receptors are reduced by *ca.* 25%, while the densities of cortical α_1 and α_2 adrenergic receptors are not significantly altered. Densities of striatal D₁ and D₂ dopaminergic receptors are unaltered. The densities of cortical 5 HT₁ and 5 HT₂ serotonergic receptors are increased by 26–30%. Densities of cortical muscarinic and nicotinic receptors are increased by 40–50%. The density of cortical benzodiazepine-binding sites associated with GABA_A receptors is increased by 65%, and the affinity appears slightly decreased. The density of cortical MK-801 sites associated with NMDA-glutaminergic receptors appear unaltered.
4. The density of cortical nitrendipine-binding sites associated with calcium channels is increased by 18%.
5. The results indicate that chronic ingestion of caffeine equivalent to about 100 mg/kg/day in mice causes a wide range of biochemical alterations in the central nervous system.

Keywords

caffeine; adenosine receptors; adrenergic receptors; cholinergic receptors; serotonin receptors; GABA receptors; calcium channels; dopamine receptors; NMDA receptors

INTRODUCTION

Chronic caffeine can cause tolerance in animals and humans, and subsequent abstinence from caffeine can then lead to withdrawal syndromes. The underlying mechanisms are poorly understood (Nehlig *et al.*, 1992; Daly, 1993). Blockade of central adenosine receptors resulting in an up-regulation of adenosine receptors has received attention as one explanation of the chronic effects of caffeine. Chronic treatment with caffeine does result in an apparent up-regulation of A₁ adenosine receptors in rodents (Fredholm, 1982; Boulenger *et al.*, 1983; Wu and Coffin, 1984; Green and Stiles, 1986; Zielke and Zielke, 1987; Hawkins *et al.*, 1988; Ramkumar *et al.*, 1988; Daval *et al.*, 1989; Rudolphi *et al.*, 1989; see also references in Nehlig *et al.*, 1992; Daly, 1993), and an enhanced sensitivity to behavioral effects of adenosine analogs (Ahlijanian *et al.*, 1986; Nikodijević *et al.*, 1993). The effects of chronic caffeine on other central receptors has received less attention. β -Adrenergic

receptors appear down-regulated, while α -adrenergic receptors appear unchanged (Goldberg *et al.*, 1982; Green and Stiles, 1986). Caffeine does alter turnover of not only norepinephrine, but also of dopamine and serotonin (Berkowitz *et al.*, 1970; Berkowitz and Spector, 1971; Corrodi *et al.*, 1972; Hadfield and Milio, 1989). The effects of chronic caffeine on levels of dopamine and serotonin receptors do not appear to have been assessed. Similarly, while behavioral effects of cholinergic agonists and antagonists are altered after chronic caffeine (Nikodijevi *et al.*, 1992), changes in muscarinic and nicotinic receptors have not been assessed. The density of benzodiazepine sites associated with GABA_A-receptors appear increased after chronic caffeine (Wu and Coffin, 1984; Wu and Phillis, 1986).

The effect of chronic ingestion of caffeine on a variety of central receptors and on calcium channels has now been probed using radio ligand binding assays with male NIH-Swiss strain mice. The results provide evidence for significant alterations in adenosine, adrenergic, serotonergic, cholinergic and GABAergic systems. Such changes may underlie the changes in behavioral responses of such mice seen after chronic caffeine (Nikodijevi *et al.*, 1993).

METHODS

Materials

[³H] CGS21680 (specific activity, 47.2 Ci/mmol), [³H]N⁶ cyclohexyladenosine (30.2 Ci/mmol), [³H] dihydroalprenolol (57.5 Ci/mmol), [³H] SCH 23390 (71.3 Ci/mmol), [³H]nicotine (75 Ci/mmol), [³H]quinuclidinylbenzilate (45.5 Ci/mmol), [³H]nitrendipine (71.2 Ci/mmol), [³H]diazepam (84.5 Ci/mmol), [³H]5-hydroxytryptamine (24.4 Ci/mmol), [³H]ketanserin (64.1 Ci/mmol), [³H]clonidine (60.0 Ci/mmol), [³H]prazosin (18.1 Ci/mmol), [³H]MK-801 (22.5 Ci/mmol) and [³H]spiperone (23.1 Ci/mmol) were from New England Nuclear (Boston, MA). Caffeine (free base) was from Matheson, Coleman and Bell (Cincinnati, OH). R-N⁶-Phenylisopropyladenosine, 2-chloroadenosine, clonidine, phentolamine, MK-801, nicotine, pargyline, and methysergide were from Biochemicals, Inc. (Natick, MA). Propranolol, diisopropylfluoro-phosphate, polyethylenimine, and butaclamol were from Sigma (St. Louis, MO). Atropine was from Merck & Co., Inc. (Rahway, NJ). Nifedipine was from Delbay Pharmaceuticals Inc. (Bloomfield, NJ). Serotonin was from Regis Chemical Co. (Chicago, IL). R015-1788 was from Hoffmann La Roche, Inc. (NJ).

Chronic Caffeine Administration

White male mice of the NIH Swiss strain, weight 25–30 g, were divided into two groups. The control group was given free access to water and food, while the chronic caffeine group was given access to water containing 0.1 % caffeine for 4 days and to food. Caffeine ingestion was continued up to two to four hours before sacrifice. Based on water consumption, mice ingested ~100 mg/kg of caffeine per day. For further details and behavioral studies with such mice see Nikodijevi *et al.* (1992).

Membrane Preparation

The mice were killed by cervical fracture and the brains rapidly removed into ice-cold 50 mM Tris-HCl buffer (pH 7.4). Cerebral cortex (5 mice), cerebellum (4 mice) or striatum (10 mice) were dissected from mice of either control or caffeine groups. Tissue was homogenized in 3 ml of ice-cold 50 mM Tris-HCl buffer (pH 7.4), using a polytron at setting number 6 for 10 seconds. The homogenate was adjusted in volume to 7 ml using ice-cold 50 mM Tris-HCl (pH 7.4) and centrifuged at 35000 × g for 15 min at 4°C. The pellet was suspended with polytron and recentrifuged in the same volume of buffer. The final pellet was resuspended in the appropriate incubation buffer at protein concentration of 2.3 mg/ml. Aliquots were used immediately for receptor-binding assays. Protein concentrations

were determined by the BCA protein assay reagents (Pierce Chemical Co., Rockford, IL) using bovine albumin as standard.

[³H]N⁶-Cyclohexyladenosine (CHA) Binding

[³H]CHA binding to cortical membranes was measured by a modification of the method of Bruns *et al.* (1980). The final pellet had been resuspended in 50 mM Tris-HCl buffer (pH 7.4), containing 8 μg/ml adenosine deaminase. Incubations were in the same buffer containing 100 μl membrane suspension and [³H]CHA from 0.5–16 nM. The final volume was 1 ml. Each assay (90 min, 25°C) was performed in triplicate. Nonspecific binding was defined in the presence of 10 μM 2-chloroadenosine and was *ca.* 5% of the total binding. Binding reactions were terminated by filtration through Whatman GF/B filters using a Brandel M24R Cell Harvester (Brandel, Gaithersburg, MD). Filters were washed two times with 5 ml ice-cold buffer and placed in scintillation vials with 5 ml of Hydrofluor scintillation fluid, followed by counting for tritium.

[³H]CGS 21680 Binding

[³H]CGS 21680 binding to striatal membranes was measured by a modification of the method of Jarvis *et al.* (1989). The final pellet had been resuspended in 50 mM Tris-HCl (pH 7.4) buffer containing 8 μg/ml adenosine deaminase and 10 mM MgCl₂. Incubations were in the same buffer containing 100 μl membrane suspension and [³H]CGS 21680 from 0.5–16 nM. The final volume was 1 ml. Each assay (90 min, 25°C) was performed in triplicate. Nonspecific binding was defined in the presence of 30 μM N⁶-R-phenylisopropyladenosine and was *ca.* 20% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]Prazosin Binding

[³H]Prazosin binding to cortical membranes was measured by a modification of the method of Glossman and Hornung (1980). The final pellet had been resuspended in 50 mM Tris HCl buffer (pH 7.4). Incubations were in the same buffer, containing 100 μl membrane suspension and [³H]prazosin from 0.05–0.8 nM. The final volume was 1 ml. Each assay (30 min, 25°C) was performed in triplicate. Nonspecific binding was defined in the presence of 1 μM phentolamine and was *ca.* 10% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]Clonidine Binding

[³H]Clonidine binding to cortical membranes was measured by a modification of the method described by U'Prichard *et al.* (1977). The final pellet had been resuspended in 50 mM Tris HCl buffer (pH 7.4). Incubations were in the same buffer containing 100 μl membrane suspension and [³H]clonidine from 0.25–16 nM. The final volume was 1 ml. Each assay (30 min, 25°C) was performed in triplicate. Nonspecific binding was defined in the presence of 10 μM clonidine and was *ca.* 10% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]Dihydroalprenolol (DHA) Binding

[³H]DHA binding to cortical membranes for β₁ adrenergic receptors and to cerebellum membranes for β₂ adrenergic receptors was measured by a modification of the method of Bylund and Snyder (1976). The final pellet had been resuspended in 50 mM Tris HCl buffer (pH 7.4), containing 10 mM MgCl₂ and 1 mM EDTA. Incubations were in the same buffer containing 100 μl membrane suspension and [³H]DHA from 0.125–8 nM. The final volume was 1 ml. Each assay (30 min, 30°C) was performed in triplicate. Nonspecific binding was

defined in the presence of 10 μ M propranolol and was *ca.* 30% of total binding. Filtration, washing and scintillation spectroscopy were as described for [3 H]CHA.

[3 H]SCH 23390 Binding

[3 H]SCH 23390 binding to the D₁ dopaminergic receptor in striatal membranes was measured essentially as described (Ferretti, *et al.*, 1992). The final pellet had been resuspended in 50 mM Tris HCl buffer (pH 7.4), containing 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, and 1 mM MgCl₂. Incubations were in the same buffer containing 100 μ l membrane suspension and [3 H]SCH 23390 from 0.05–3.2 nM. The final volume was 1 ml. Each assay (30 min, 37°C) was performed in triplicate. Nonspecific binding was defined in the presence of 10 μ M butaclamol and was *ca.* 5% of total binding. Filtration, washing and scintillation spectroscopy were as described for [3 H]CHA.

[3 H]Spiperone Binding

[3 H]Spiperone binding to D₂-dopaminergic receptors in striatal membranes was measured essentially as described (Michaluk *et al.*, 1982). The final pellet had been resuspended in 50 mM Tris HCl buffer (pH 7.4), containing 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, 10 μ M pargyline, and 0.1% ascorbic acid. Incubations were in the same buffer containing 100 μ l membrane suspension and [3 H]spiperone from 0.05–3.2 nM. Each assay (25 min, 37°C) was performed in triplicate. Nonspecific binding was defined in the presence of 1 μ M butaclamol and was *ca.* 10% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [3 H]CHA.

[3 H]Nicotine Binding

[3 H]Nicotine binding to cortical membranes was measured by a modification of described procedures (Martino Garrows and Kellar, 1986; Marks *et al.*, 1986). The final pellet had been resuspended in 20 mM Hepes buffer (pH 7.4), containing 1 mM MgCl₂, 120 nM NaCl, 5 mM KCl, and 2 mM CaCl₂. Incubations were in the same buffer containing 100 μ l membrane suspension, 200 μ M diisopropyl-fluorophosphate and [3 H]nicotine from 0.125–8.0 nM. The final volume was 1 ml. Each assay (120 min, 0–4°C) was performed in triplicate. Nonspecific binding was defined in the presence of 1 μ M nicotine and was *ca.* 20% of total binding. Filtration, washing and scintillation spectroscopy were as described for [3 H]CHA. The filters were presoaked in cold 0.3% polyethylenimine to reduce nonspecific binding.

[3 H]Quinuclidinylbeuzilate (QNB)

[3 H]QNB binding to muscarinic receptors in cortical membranes was measured as follows: (see Yamamura and Snyder, 1979; Hammer *et al.*, 1980). The final pellet had been resuspended in 20 mM Hepes buffer (pH 7.4), containing 100 mM NaCl and 10 mM MgCl₂. Incubations were in the same buffer containing 100 μ l membrane suspension and [3 H]QNB from 0.05–0.8 nM. The final volume was 1 ml. Each assay (30 min, 37°C) was performed in triplicate. Nonspecific binding was defined in the presence of 1 μ M atropine and was *ca.* 25% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [3 H]CHA. The filters were presoaked in 0.3% polyethylenimine to reduce nonspecific binding to filter.

[3 H]Diazepam Binding

[3 H]Diazepam binding to benzodiazepine sites on GABA_A receptors in cortical membranes was measured essentially as described (Wu and Coffin, 1984). The final pellet had been resuspended in 50 mM Tris-HCl buffer (pH 7.4). Incubations were in the same buffer containing 100 μ l membrane suspension and [3 H]diazepam from 0.5–16 nM. The final

volume was 1 ml. Each assay (60 min, 0–4°C) was performed in triplicate. Nonspecific binding was defined in the presence of 20 μ M Ro15-1788 and was *ca.* 10% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]5-Hydroxytryptamine Binding

[³H]5-Hydroxytryptamine binding to 5-HT₁ receptors in cortical membranes was measured by a modification of the method described by Pedigo *et al.* (1981). The final pellet had been resuspended in 50 mM Tris-HCl buffer (pH 7.4), containing 4 mM CaCl₂. Incubations were in the same buffer containing 10 μ M pargyline, 100 μ l membrane suspension and [³H]5 hydroxytryptamine from 0.5–16 nM. The final volume was 1 ml. Each assay (10 min, 37°C) was performed in triplicate. Nonspecific binding was defined in the presence of 10 μ M 5 hydroxytryptamine and was *ca.* 30% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]Ketanserin Binding

[³H]Ketanserin binding to 5-HT₂ receptors in cortical membranes was measured by a modification of the method described by Leysen *et al.* (1986). The final pellet had been resuspended in 50 mM Tris-HCl buffer (pH 7.4). Incubations were in the same buffer, containing 100 μ l membrane suspension and [³H]ketanserin from 0.25–8 nM. The final volume was 1 ml. Each assay (15 min, 37°C) was performed in triplicate. Nonspecific binding was defined in the presence of 1 μ M methysergide and was *ca.* 20% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]MK-801 Binding

[³H]MK-801 binding to sites on NMDA receptors in cortical membranes was measured by a modification of the method described by Wong *et al.* (1986). The final pellet had been resuspended in 5 mM Tris-HCl buffer (pH 7.4). Incubations were in the same buffer, containing 100 μ l membrane suspension and [³H]MK 801 from 0.5–16 nM. The final volume was 1 ml. Each assay (45 min, 23°C) was performed in triplicate. Nonspecific binding was defined in the presence of 100 μ M MK-801 and was *ca.* 5% of the total binding. Filtration, washing and scintillation spectroscopy were as described for [³H]CHA.

[³H]Nitrendipine Binding

[³H]Nitrendipine binding to cortical membranes was measured by a modification of the method of Ehlert *et al.* (1982). The final pellet had been resuspended in 50 mM Tris-HCl buffer (pH 7.4). Incubations were in the same buffer, containing 100 μ l membrane suspension and [³H]nitrendipine from 0.05–1 nM. The final volume was 1 ml. Each assay (90 min, 25°C) was performed in triplicate. Nonspecific binding was defined in the presence of 1 μ M and was *ca.* 10% of total binding. Filtration, washing and scintillation were as described for [³H]CHA.

Data Analysis

The dissociation constant (K_D) and the maximal number of binding sites (B_{max}) were analyzed with the computer program LIGAND (Munson and Rodbard, 1980) for each experiment and for pooled data from at least three experiments and are reported as means $\pm$ SEM. The statistical evaluations were with the Student's t-test.

RESULTS

Chronic ingestion of caffeine for a period of 4 days by male NIH Swiss strain mice results in significant changes in densities of a variety of cortical or cerebellar receptors and in density

of cortical nitrendipine binding sites, presumed to be associated with L-type voltage sensitive calcium channels (Table I). Significant increases in $B_{\max}$ values in cortical membranes occur for the following: A_1 adenosine receptors (20%), $5HT_1$ and $5HT_2$ serotonergic receptors (26–30%), nicotinic and muscarinic receptors (40–50%), benzodiazepine binding sites associated with $GABA_A$ receptors (65%), and nitrendipine binding sites (18%) (Fig. 1). Only in the case of [3H]diazepam binding to benzodiazepine modulatory sites was a significant change in K_D observed, namely a slight decrease in affinity (Table 1, $P < 0.05$). Significant decreases in $B_{\max}$ values occur for cortical β_1 -adrenergic receptors (25%) and for cerebellar β_2 adrenergic receptors (27%). The $B_{\max}$ values for striatal A_{2A} adenosine receptors and D_1 and D_2 dopaminergic receptors and for cortical α_1 and α_2 adrenergic receptors and for [3H]MK-801-channel sites associated with NMDA receptors were unchanged after chronic caffeine ingestion in male NIH Swiss strain mice.

DISCUSSION

The increase in density of cortical A_1 adenosine receptors after chronic blockade of such receptors by caffeine (Table 1) is not unexpected. Indeed, such have been previously reported by many laboratories primarily for rats (see INTRODUCTION). One report of no change in A_1 -adenosine receptors in cerebral cortex after chronic caffeine has appeared (Holtzman *et al.*, 1991). In contrast to the up-regulation of A_1 -adenosine receptors, there was no increase density of striatal A_{2A} -adenosine receptors after chronic blockade of such receptors by caffeine (Table 1). One study had reported no increase in striatal A_{2A} -adenosine receptors as measured by autoradiography (Lupica *et al.*, 1991), while another study reported a slight increase as measured by membrane binding assays (Hawkins *et al.*, 1988). In vitro in striatal slices A_1 , but not A_{2A} -adenosine receptors undergo desensitization (Abbraccio *et al.*, 1992). In contrast, in vivo in rats an adenosine analog, NECA, down-regulated striatal A_{2A} -adenosine receptors, while having no effect on A_1 adenosine receptors (Portal *et al.*, 1998).

The increase in A_1 adenosine receptors in male NIH Swiss strain mice (Table 1) is accompanied by an increase in sensitivity to the depressant effects of adenosine analogs on open-field locomotor activity (Nikodijevi *et al.*, 1992). Remarkably, this applies to both A_1 and A_2 selective analogs. There are synergistic effects on locomotor activity of A_1 and A_2 selective agonists (Nikodijevi *et al.*, 1991) and thus endogenous adenosine through the A_1 , receptors could enhance behavioral effects of A_2 selective adenosine analogs in control and chronic caffeine treated mice. It should be noted that male NIH Swiss strain mice do not become tolerant to the stimulatory effects of caffeine after chronic caffeine-ingestion (Nikodijevi *et al.*, 1993).

Caffeine is not known to block extracellular receptors other than adenosine receptors at the dosage level (~100 mg/kg/day) employed in the present study. Thus, effects of chronic caffeine on levels of other central receptors are likely indirect. Certainly, blockade by chronic caffeine of the tonic input of endogenous adenosine to presynaptic receptors that are inhibitory to neurotransmitter release (Williams, 1987) would be expected to alter turnover and hence postsynaptic activation of receptors for a variety of central neurotransmitters, including norepinephrine, dopamine, serotonin, acetylcholine, γ -aminobutyric acid and glutamate. Homeostatic regulation in such receptors or the second messenger systems subserved by such receptors might then be expected. The present results indicate that densities of some, but not all of such possible neurotransmitter receptors do significantly change as a result of 4 days of chronic caffeine ingestion (~100mg/kg/day) in male NIH Swiss strain mice.

The reduction on density of both β_1 and β_2 adrenergic receptors in cerebral cortex and cerebellum, respectively (Table 1), suggests a caffeine induced increase in turnover and action of norepinephrine at such receptors. Caffeine has been reported to increase norepinephrine turnover (Berkowitz *et al.*, 1970; Corrodi *et al.*, 1972; Hadfield and Milio, 1989). A down-regulation of cortical β_1 adrenergic receptors has been previously reported for rats after chronic caffeine (Goldberg *et al.*, 1982; Green and Stiles, 1986) or theophylline (Fredholm *et al.* 1984). The lack of change in α_1 and α_2 adrenergic receptors after chronic caffeine ingestion (Table 1) suggests either that these receptors exist mainly at sites or pathways whose noradrenergic input is unaffected by caffeine or that such receptors are not under homeostatic control in this strain of mice. In rat brain the densities of α_1 , and α_2 adrenergic receptors were unaltered after chronic caffeine (Goldberg *et al.*, 1982). A single injection of pentoxifylline (3, 7-dimethyl-1-(5-oxohexyl)xanthine) caused a decrease in densities of both α_2 and β_1 adrenergic receptors in cortex of rats (Lowenstein *et al.*, 1982).

The lack of effect of chronic caffeine ingestion on dopamine receptors (Table 1) is remarkable in view of many reports, indicating a close interrelationship of adenosine and dopamine systems in striatum (see review by Ferré *et al.*, 1992). Thus, blockade of striatal adenosine receptors by caffeine enhances dopaminergic function (*cf.* Josselyn and Beninger, 1991 and references therein). Conversely, chronic central infusion of adenosine analogs in rats causes an attenuation in both A_{2A} adenosine and D₁ dopamine receptor elicited stimulation of striatal adenylate cyclase (Porter *et al.*, 1988). Other treatments, such as dopamine receptor blockade or denervation, do result in alterations in sensitivity to dopaminergic agonists (Yarbrough 1975; Siggins *et al.*, 1984; Chipkin *et al.*, 1984). Behaviorally, chronic ingestion of caffeine in the male NIH Swiss strain mice has no significant effect on stimulation of locomotor activity elicited by amphetamine, but does increase the maximal percent stimulation elicited by cocaine (Nikodijevi *et al.*, 1993), agents thought to act by increasing dopaminergic function.

The densities of cortical 5HT₁ and 5HT₂-serotonin receptors are increased after chronic caffeine ingestion of male NIH Swiss strain mice (Table 1), suggesting a reduction in serotonergic input and a resultant up regulation of receptors. However, caffeine has been reported in some studies to *increase* rather than decrease serotonin turnover in rats (Berkowitz and Spector, 1971; Corrodi *et al.*, 1972) and mice (Valzelli *et al.*, 1973). Thus, a down regulation of serotonin receptors might have been expected. Behaviorally, effects of serotonergic agents do not appear to have been investigated after chronic caffeine.

The densities of cholinergic receptors, both nicotinic and muscarinic, appear increased after chronic caffeine ingestion, suggesting a decrease in acetylcholine function and a resultant up regulation of receptors. However, chronic treatment of rats with nicotine increases the density of nicotinic receptors (Ksir *et al.*, 1985; Hillard and Pound, 1993). Thus, caffeine might be increasing nicotinic receptors by enhancing acetylcholine release, rather than by decreasing release. Acute caffeine does increase acetylcholine release from rat cerebral cortex (Phillis *et al.*, 1980). Behaviorally, it appears that muscarinic function is enhanced in NIH Swiss strain mice after chronic caffeine ingestion, since higher doses of a muscarinic antagonist, scopolamine, are required to stimulate openfield locomotor activity compared to control mice (Nikodijevi *et al.*, 1993). Both nicotinic (nicotine) and muscarinic (oxotremorine) agonists reduce openfield locomotor activity in control NIH Swiss strain mice, while being ineffective (nicotine) or less effective (oxotremorine) after chronic caffeine (Nikodijevi *et al.*, 1993). This too could be interpreted as due to increase in cholinergic function, so that any further depressant effects of nicotine or oxotremorine are blunted. The mice do exhibit a marked long term reduction in openfield locomotor activity after chronic caffeine (Nikodijevi *et al.*, 1993), which might reflect enhanced function of the up-regulated cholinergic receptors. Further studies directed towards such hypotheses are

required. Chronic caffeine has been reported to reduce the excitatory effects of acetylcholine on rat cerebral cortical neurons (Lin and Phillis, 1990). Chronic nicotine results in tolerance to depressant effects of nicotine in spite of regulation of nicotinic receptors (Ksir *et al.*, 1985).

The density of GABA_A receptors, as assessed by binding of the GABA_A modulator diazepam, is increased markedly in NIH strain mice (Table 1). There also appears to be a small but significant decrease in the affinity of diazepam; the nature of this decrease requires further investigation. Such an increase suggests an up-regulation of GABA_A channels as a result of a decrease in GABAergic function during chronic caffeine treatment. An up regulation of benzodiazepine sites after chronic caffeine has been previously reported for rats (Wu and Coffin, (1984; Wu and Phillis, 1986). Chronic caffeine at a low dose of 10 mg/kg/day has little effect on diazepam-elicited changes in glucose utilization in brains of rat (Nehlig *et al.*, 1987). Chronic theophylline does decrease the convulsant activity of various agents in rats (Szot *et al.*, 1987), but whether this is due to up-regulation of adenosine receptors or an up regulation of GABA_A receptors or is unknown.

The density of the NMDA class of glutamate receptors, as assessed by binding of the NMDA-channel blocker MK-1801, appears unchanged after chronic caffeine (Table 1). This is perhaps surprising in view of reports of functional : interrelationships of NMDA and adenosine receptors (Bartrup and Stone, 1990; Nagaoka *et al.*, 1993).

The density of L-type calcium channels, as assessed by binding of the L-type calcium channel antagonist nitrendipine is significantly increased in cortical membranes in male NIH Swiss strain mice after chronic caffeine ingestion (Table 1) The mechanism is unknown. Nitrendipine binding is also reported to be increased after chronic ethanol (Dolin *et al.*, 1987). Behavioral effects of calcium channel agents after chronic caffeine do not appear to have been assessed.

In summary, chronic ingestion of caffeine in male NIH Swiss strain mice results in increases in densities of A₁, but not A_{2A} adenosine receptors commensurate at least for the A₁ receptors with direct chronic blockade of such receptors by caffeine. Other G protein linked receptors also show changes in densities probably due to indirect effects of the blockade of adenosine receptors by caffeine on the tonic input of neurotransmitters to such G protein linked receptors. The β_1 and β_2 adrenergic receptors are down-regulated, while 5HT₁ and 5HT₂ serotonin receptors and muscarinic receptors are up-regulated. α_1 and α_2 Adrenergic receptors and D₁ and D₂ dopaminergic receptors are unaffected. Receptors that are part of a receptor-ion channel complex are also affected by chronic caffeine ingestion. The densities of the nicotinic receptor-channel and the GABA_A receptor channel are increased, while the NMDA receptor channel appears unchanged. The density of L-type calcium channels, as assessed by nitrendipine binding is increased by chronic caffeine ingestion. In rats, the density of forskolin-binding sites, presumably associated with adenylate cyclase, is increased after chronic caffeine (Daval *et al.*, 1989).

It appears likely that most alterations in receptors, ion channels and effector systems after chronic caffeine will prove related to blockade of tonic adenosine input by caffeine, rather than to blockade of phosphodiesterases or release of calcium from intracellular sites, since the latter two effects become significant only at much higher concentrations than the expected 10 μ M brain concentrations of caffeine attained during chronic ingestion of caffeine (100 mg/kg/day) (see Ahljianian and Takemori, 1986). It should be noted that phosphodiesterase inhibitors, including xanthines much more potent than caffeine do cause down-regulation of β -adrenergic receptors (Lowenstein *et al.*, 1982; Schultz and Schmidt, 1986). Further studies on behavioral responses of caffeine treated NIH Swiss strain mice (*cf.*

Nikodijevi *et al.*, 1993) will be required to correlate changes in receptors and ion channels with caffeine induced tolerance, alteration in drug responsiveness, and withdrawal

Acknowledgments

The research on caffeine was supported by a grant to D.S. from the International Life Science Institute.

REFERENCES

- Abbracchio MP, Fogliatto G, Paoletti AM, Rovati GE, Cattabeni F. Prolonged in vitro exposure of rat brain slices to adenosine analogues: selective desensitization of adenosine A₁ but not A₂ receptors. *Eur. J. Pharmacol.* 1992; 227:317–324. [PubMed: 1473554]
- Ahlijanian MK, Takemori AE. Cross-tolerance studies between caffeine and (–)-N⁶-(phenylisopropyl)-adenosine (PIA) in mice. *Life Sci.* 1986; 88:577–588. [PubMed: 3003486]
- Bartrup JT, Stone TW. Activation of NMDA receptor coupled channels suppresses the inhibitory action of adenosine on hippocampal slices. *Brain Res.* 1990; 530:330–334. [PubMed: 1702344]
- Berkowitz BA, Spector S. The effect of caffeine and theophylline on the disposition of brain serotonin in the rat. *Eur. J. Pharmacol.* 1971; 16:322–325.
- Berkowitz BA, Tarver JH, Spector S. Release of norepinephrine in the central nervous system by theophylline and caffeine. *Eur. J. Pharmacol.* 1970; 10:64–71.
- Boulenger J-P, Patel J, Post RM, Parma AM, Marangos PJ. Chronic caffeine consumption increases the number of brain adenosine receptors. *Life Sci.* 1983; 32:1135–1142. [PubMed: 6298543]
- Bruns RF, Daly JW, Snyder SH. Adenosine receptor in brain membranes: binding of N⁶-cyclohexyl-[³H]adenosine and 1,3-diethyl-8-[³H]phenylxanthine. *Proc. Natl. Acad. Sci. USA.* 1980; 77:5547–5551. [PubMed: 6254090]
- Bylund DB, Snyder SH. Beta adrenergic receptor binding in membrane preparations from mammalian brain. *Mol. Pharmacol.* 1976; 12:568–580. [PubMed: 8699]
- Chipkin RE, McQuade RD, Iorio LC. D₁ and D₂ dopamine binding site up-regulation and apomorphine-induced stereotypy. *Pharmacol. Biochem. Behav.* 1987; 28:477–482. [PubMed: 2893388]
- Corrodi H, Fuxe K, Jonsson G. Effects of caffeine on central monoamine neurons. *J. Pharm. Pharmacol.* 1972; 24:155–158.
- Daly, JW. Mechanism of action of caffeine.. In: Garattini, S., editor. *Caffeine, Coffee and Health*. Raven Press Ltd.; New York: 1993. p. 97-149.
- Daval JL, Deckert J, Weiss SRB, Post RM, Marangos PJ. Upregulation of adenosine A₁ receptors and forskolin binding sites following chronic treatment with caffeine or carbamazepine: a quantitative autoradiographic study. *Epilepsia.* 1989; 30:26–33. [PubMed: 2912715]
- Dolin S, Little H, Hudspeth M, Pagonis C, Littleton J. Increased dihydropyridine-sensitive calcium channels in rat brain may underlie ethanol physical dependence. *Neuropharmacology.* 1987; 26:275–279. [PubMed: 2438583]
- Ehlert FJ, Roeske WR, Itoga E, Yamamura HI. The binding of [³H]nitrendipine to receptors for calcium channel antagonists in the heart, cerebral cortex, and ileum of rats. *Life Sci.* 1982; 30:2192–2202.
- Ferré S, Fuxe K, Von Euler G, Johansson B, Fredholm BB. Adenosine-dopamine interactions in the brain. *Neuroscience.* 1992; 51:501–512. [PubMed: 1488111]
- Ferretti C, Blengio M, Vigna I, Ghi P, Genazzini E. Effects of estradiol on the ontogenesis of striatal D₁ and D₂ receptor sites in male and female rats. *Brain Res.* 1992; 571:212–217. [PubMed: 1535267]
- Fredholm BB. Adenosine actions and adenosine receptors after 1 week treatment with caffeine. *Acta Physiol. Scand.* 1982; 115:283–286. [PubMed: 6291335]
- Fredholm BB, Jonzon B, Lindgren E. Changes in noradrenaline release and in beta receptor number in rat hippocampus following long-term treatment with theophylline or L-phenylisopropyladenosine. *Acta Physiol. Scand.* 1984; 122:55–59. [PubMed: 6095597]

- Glossmann H, Hornung R. α -Adrenoceptors in rat brain: sodium changes the affinity of agonists for prazosin sites. *Eur. J. Pharmacol.* 1980; 61:407–408. [PubMed: 6245895]
- Goldberg MR, Curatolo PW, Tung C-S, Robertson D. Caffeine down-regulates β adrenoreceptors in rat forebrain. *Neuroscience Lett.* 1982; 31:47–52.
- Green RM, Stiles GL. Chronic caffeine ingestion sensitized the A₁ adenosine receptor-adenylate cyclase system in rat cerebral cortex. *J. Clin. Invest.* 1986; 77:222–227. [PubMed: 3003150]
- Hadfield MG, Milio C. Caffeine and regional brain monoamine utilization in mice. *Life Sci.* 1989; 45:2637–2644. [PubMed: 2575694]
- Hammer R, Berrie CP, Birdsall NJM, Burgeri ASV, Hulme. Pirenzepine distinguishes between different subclasses of muscarinic receptors. *Nature.* 1980; 283:90–93. [PubMed: 7350532]
- Hawkins M, Dugich MM, Porter NM, Urbancic M, Radulovacki M. Effects of chronic administration of caffeine on adenosine A₁ and A₂ receptors in rat brain. *Brain Res. Bull.* 1988; 21:479–482. [PubMed: 3214753]
- Hillard CJ, Pounds JJ. Effects of chronic nicotine treatment on the accumulation of [³H]tetraphenylphosphonium by cerebral cortical synaptosomes. *J. Neurochem.* 1993; 60:687–695. [PubMed: 8419544]
- Holtzman SG, Mante S, Minneman KP. Role of adenosine receptors in caffeine tolerance. *J. Pharmacol. Exp. Ther.* 1991; 256:62–68. [PubMed: 1846425]
- Jarvis MF, Schulz K, Hutchison AJ, Do UH, Sills MA, Williams M. [³H]CGS 21680: A selective A₂ adenosine receptor agonist directly labels A₂ receptors in rat brain. *J. Pharmacol. Exp. Ther.* 1989; 251:888–893. [PubMed: 2600819]
- Josselyn SA, Beninger RJ. Behavioral effects of intrastriatal caffeine mediated by adenosinergic modulation of dopamine. *Pharmacol. Biochem. Behav.* 1991; 39:97–103. [PubMed: 1924519]
- Ksir C, Hakan R, Hall DP, Kellar KJ. Exposure to nicotine enhances the behavioral stimulant effect of nicotine and increases the binding of [³H]acetylcholine to nicotine receptors. *Neuropharmacology.* 1985; 24:527–531. [PubMed: 4022267]
- Leysen JE, Gompel PV, Gommeren W, Woestenborghs R, Janssen PAJ. Down regulation of serotonin-S₂ receptor sites in rat brain by chronic treatment with the serotonin-S₂ antagonists: ritanserin and setoperone. *Psychopharmacology.* 1986; 88:434–444. [PubMed: 3010361]
- Lin Y, Phillis JW. Chronic caffeine exposure reduces the excitant action of acetylcholine on cerebral cortical neurons. *Brain Res.* 1990; 524:316–318. [PubMed: 2292013]
- Lowenstein PR, Vacas MI, Cardinali DP. Effect of pentoxifyline on α - and β -adrenoceptor sites in cerebral cortex, medial basal hypothalamus and pineal gland of the rat. *Neuropharmacology.* 1982; 21:243–248. [PubMed: 6280095]
- Lupica CR, Berman FR, Jarvis MF. Chronic theophylline treatment increases adenosine A₁, but not A₂, receptor binding in the rat brain; an autoradiographic study. *Synapse.* 1991; 9:95–102. [PubMed: 1821488]
- Marks MJ, Stitzel JA, Romm E, Wehner JM, Collins AC. Nicotine binding sites in rat and mouse brain: comparison of acetylcholine, nicotine, and α -bungarotoxin. *Mol. Pharmacol.* 1986; 30:427–436. [PubMed: 3534542]
- Martino-Barows AM, Kellar KJ. [³H]Acetylcholine and [³H](–)nicotine label the same recognition site in rat brain. *Mol. Pharmacol.* 1986; 31:169–174.
- Michaluk J, Antkiewicz-Michaluk L, Roskosz-Pelc A, Sansone M, Oliver A, Vetulani J. Dopamine receptor in the striatum and limbic system of various strains of mice: Relation to difference in responses to apomorphine. *Pharmacol. Biochem. Behav.* 1982; 17:1115–1118.
- Munson PJ, Rodbard D. LIGAND: A versatile computerized approach for characterization of ligand-binding systems. *Anal. Biochem.* 1980; 107:220–239. [PubMed: 6254391]
- Nagaoka H, Sakurada S, Sakurada T, Takeda S, Nakagawa Y, Kisara K, Arai Y. Theophylline-induced nociceptive behavioral response in mice: possible indirect interaction with spinal N-methyl-D-aspartate receptors. *Neurochem. Int.* 1993; 22:69–74. [PubMed: 7680261]
- Nehlig A, Daval J-L, Debry G. Caffeine and the central nervous system: Mechanisms of action, biochemical, metabolic, and psychostimulant effects. *Brain Res. Rev.* 1992; 17:139–170. [PubMed: 1356551]

- Nehlig A, Daval J-L, Pereira de Vasconcelos A, Boyet. Caffeine-diazepam interaction and local cerebral glucose utilization in the conscious rat. *Brain Res. Rev.* 1987; 419:272–278.
- Nikodijevi O, Jacobson KA, Daly JW. Locomotor activity in mice during chronic treatment with caffeine and withdrawal. *Pharmacol. Biochem. Behav.* 1993; 44:199–2160. [PubMed: 7679219]
- Nikodijevi O, Sarges R, Daly JW, Jacobson KA. Behavioral effects of A₁- and A₂-selective adenosine agonists and antagonists: Evidence for synergism and antagonism. *J. Pharmacol. Exp. Therap.* 1991; 259:286–294. [PubMed: 1920121]
- Pedigo NW, Yamamura HI, Nelson DL. Discrimination of multiple [³H]5-hydroxytryptamine binding sites by the neuroleptic spiperone in rat brain. *J. Neurochem.* 1981; 36:220–226. [PubMed: 7463047]
- Phillis JW, Jiang ZG, Chelack BJ, Wu PR. The effect of morphine on purine and acetylcholine release from rat cerebral cortex: Evidence for a purinergic component in morphine's action. *Pharmacol. Biochem. Behav.* 1980; 13:421–4270.
- Porter NM, Radulovacki M, Green RD. Desensitization of adenosine and dopamine receptors in rat brain after treatment with adenosine analogs. *J. Pharmacol. Exp. Therap.* 1988; 244:218–225.
- Ramkumar V, Bumgarner JR, Jacobson KA, Stiles GL. Multiple components of the A₁ adenosine-adenylate cyclase system are regulated in rat cerebral cortex by chronic caffeine ingestion. *J. Clin. Invest.* 1988; 82:242–247.
- Rudolphi KK, Keli M, Fastbom J, Fredholm BB. Ischaemic damage in gerbil hippocampus is reduced following up regulation of adenosine (A₁) receptors by caffeine treatment. *Neurosci. Lett.* 1989; 103:275–280. [PubMed: 2812514]
- Schultz JE, Schmidt BH. Rolipram, a stereospecific inhibitor of calmodulin independent phosphodiesterase, causes β -adrenoceptor subsensitivity in rat cerebral cortex. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 1986; 333:23–30. [PubMed: 3016560]
- Siggins GR, Hoffer BJ, Ungerstedt U. Electrophysiological evidence for the involvement of cyclic adenosine monophosphate in dopamine responses of caudate neurons. *Life Sci.* 1974; 16:779–792. [PubMed: 4378085]
- Szot P, Sanders RC, Murray TF. Theophylline-induced upregulation of A₁-adenosine receptors associated with reduced sensitivity to convulsants. *Neuropharmacology.* 1987; 26:1173–1180. [PubMed: 3658121]
- U'Prichard DC, Greenberg DA, Snyder SH. Binding characteristics of a radiolabeled agonist and antagonist at central nervous system alpha noradrenergic receptors. *Mol. Pharmacol.* 1977; 13:454–4730. [PubMed: 17827]
- Valzelli L, Bernasconi S. Behavioral and neurochemical effects of caffeine in normal and aggressive mice. *Pharmacol. Biochem. Behav.* 1973; 1:251–254. [PubMed: 4798095]
- Williams M. Purine receptors in mammalian tissues: pharmacology and functional significance. *Ann. Rev. Pharmacol. Toxicol.* 1987; 27:315–345.
- Wong EHF, Knight AR, Woodruff GN. [³H]MK-801 labels a site on the N-methyl-D-aspartate receptor channel complex in rat brain membranes. *J. Neurochem.* 1988; 50:274–281. [PubMed: 2826686]
- Wu PH, Coffin VL. Up-regulation of brain [³H]diazepam binding sites in chronic caffeine-treated rats. *Brain Res.* 1984; 294:186–1890. [PubMed: 6697237]
- Wu PH, Phillis JW. Up-regulation of brain [³H]diazepam binding sites in chronic caffeine-treated rats. *Gen. Pharmacol.* 1986; 17:501–503. [PubMed: 3019827]
- Yamamura HI, Snyder SH. Muscarinic cholinergic binding in rat brain. *Proc. Nat. Acad. Sci. USA.* 1974; 71:1725–1729. [PubMed: 4151898]
- Yarborough GG. Supersensitivity of caudate neurons after repeated administration of haloperidol. *Eur. J. Pharmacol.* 1975; 31:367–369. [PubMed: 1149799]
- Zielke CL, Zielke HR. Chronic exposure to subcutaneously implanted methylxanthines Differential elevation of A₁-adenosine receptors in mouse cerebellar and cerebral cortical membranes. *Biochem. Pharmacol.* 1987; 36:2533–2538. [PubMed: 3606653]

Fig. 1.

Scatchard Analyses for Alterations in Binding of Radioligands after Chronic Caffeine in NIH Swiss Strain Mice. Control, open circles; chronic caffeine, closed circles. A. Cortical A_1 -adenosine receptor with [3H]CHA. B. Cortical $B_{1,}$ -adrenergic receptor with [3H]DHA. C. Cerebellar β_2 -adrenergic receptor with [3H]DHA. D. Cortical 5-HT $_1$ -serotonin receptor with [3H]5-hydroxytryptamine. E. Cortical 5-HT $_2$ -serotonin receptor with [3H]5-ketanserin. F. Cortical nicotinic receptor with [3H]nicotine. G. Cortical muscarinic receptor with [3H]QNB. H. GABA $_A$ receptor with [3H]diazepam. I. Calcium channel with [3H]nitrendipine. Scatchard Analyses for the following were unchanged and are not shown: Striatal A_{2A} -adenosine receptor with [3H]CGS 21680; Cortical α_1 -adrenergic receptor with [3H]prazosin; Cortical α_2 -adrenergic receptor with [3H]clonidine; Striatal D $_1$ -dopaminergic receptor with [3H]SCH 23390; Striatal D $_2$ -dopaminergic receptor with [3H]spiperone; Cortical NMDA receptor with [3H]MK-801.

Table I

Effect of Chronic Caffeine Ingestion on Receptors and Ion Channels in Brain Membranes from Male NIH Swiss Strain Mice.

	K_d (nM)		B_{max}(fmol/mg protein)	
	Control	Chronic Caffeine	Control	Chronic Caffeine
A ₁ -Adenosine	1.95 ± 0.33	2.00 ± 0.45	911 ± 23	1089 ± 39 ^{**}
A _{2A} -Adenosine	10.26 ± 0.66	9.98 ± 0.19	872 ± 57	884 ± 44
α ₁ -Adrenergic	0.049 ± 0.003	0.053 ± 0.012	175 ± 7	189 ± 12
α ₂ -Adrenergic	1.59 ± 0.20	1.37 ± 0.20	200 ± 3	193 ± 2
β ₁ -Adrenergic	1.15 ± 0.08	1.07 ± 0.14	224 ± 9	167 ± 5 [*]
β ₂ - Adrenergic	1.57 ± 0.08	1.56 ± 0.10	158 ± 12	115 ± 11 [*]
D ₁ -Dopamine	0.53 ± 0.11	0.48 ± 0.06	3097 ± 81	3165 ± 66
D ₂ -Dopamine	0.108 ± 0.021	0.084 ± 0.018	729 ± 21	725 ± 55
5-HT ₁	2.93 ± 0.37	3.35 ± 0.58	361 ± 14	474 ± 48 ^{**}
5-HT ₂	0.33 ± 0.09	0.35 ± 0.06	275 ± 11	347 ± 11 [*]
Nicotinic	0.77 ± 0.05	0.95 ± 0.05	34 ± 2	50 ± 3 [*]
Muscarinic	0.17 ± 0.02	0.17 ± 0.03	1153 ± 56	1509 ± 47 ^{**}
NMDA (MK-801)	2.76 ± 0.29	2.66 ± 0.17	2653 ± 97	2588 ± 46
GABA _A (diazepam)	2.82 ± 0.26	4.69 ± 0.50 ^{**}	1061 ± 69	1741 ± 100 [*]
Ca ²⁺ Channel (nitrendipine)	0.16 ± 0.03	0.17 ± 0.02	314 ± 6	369 ± 12 ^{**}

Kinetic analysis of binding of radioligands to cortical, cerebellar or striatal membranes from control mice and chronic caffeine-ingesting mice was carried out as described in METHODS. Values for K_D and B_{max} are means ± S.E.M. (n = 3 or greater).

* P < 0.01

** P < 0.05.