Arterioportal Fistulas in Liver Transplant Recipients

Wael E. A. Saad, M.D., F.S.I.R.¹

¹ Division of Vascular Interventional Radiology, Department of Radiology and Medical Imaging, University of Virginia Health System, Charlottesville, Virginia

Semin Intervent Radiol 2012;29:105–110

Address for correspondence and reprint requests Wael E. A. Saad, M.D., F.S.I.R., Division of Vascular Interventional Radiology, Department of Radiology and Medical Imaging, University of Virginia Health System, Box 800170, 1215 Lee Street, Charlottesville, VA 22908 (e-mail: wspikes@yahoo.com).

Abstract

Keywords

- transplant
- liver
- vascular complications
- endovascular
- portal
- arterial
- fistula
- embolization

Arterioportal fistulas (APFs) are classified into intrahepatic (>75% of all reported) and extrahepatic (<25% of all reported). Anecdotally, investigators are more likely to report more sensational cases (typically extrahepatic APFs), so the actual prevalence of intrahepatic APFs is probably much higher (likely >90% of APFs). All reported APFs in liver transplant recipients have been intrahepatic. Hemodynamically significant APFs in liver transplant recipients are rare, occurring in 0.2%; however, APFs (hemodynamically significant or not) are not uncommonly seen in hepatic angiograms of liver transplant recipients (up to 5.4% of hepatic arteriograms in transplants). Interestingly, hemodynamically significant APFs warranting endovascular treatment are reported more commonly in the literature in native compared with transplanted livers ($n > 280$ –300 versus $n = 13$, respectively). This article discusses APFs that are specific to liver transplant recipients; their incidence, etiology, pathogenesis, natural history, clinical presentation, and endovascular management are discussed in detail.

Objectives: On completion of this article, the reader will be able to identify the pathophysiology, clinical presentation, treatment options, and outcomes of liver transplant recipients with arterioportal fistulas.

Accreditation: Tufts University School of Medicine is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit: Tufts University School of Medicine designates this journal-based CME activity for a maximum of **1 AMA PRA Category 1 Credit™**. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Arterioportal fistulae (APFs) are classified into intrahepatic (>75% of all reported APFs) and extrahepatic (<25% of all reported APFs).¹ Anecdotally, physicians are more likely to report more sensational cases (extrahepatic APFs), so the actual prevalence of intrahepatic APFs is probably much higher (>90%). Reportedly, all APFs in liver transplant recipients have been intrahepatic APFs.^{1–3} APFs are further classi-

fied into hemodynamically significant and hemodynamically insignificant. The determinant of hemodynamic significance is Doppler ultrasound findings suggestive of involvement of the main, right, or left portal vein.^{1–4}

This article discusses APFs that are specific to liver transplant recipients. Their incidence, etiology, pathogenesis, natural history, clinical presentation, imaging evaluation, and endovascular management are discussed in detail.

Incidence and Etiology

Incidence

Hemodynamically significant APFs in liver transplant recipients are rare, occurring in 0.2% of cases.² There appears to be no difference in occurrence between living related (split livers) and deceased donor (split or whole grafts).² However, APFs (hemodynamically significant or not) are not uncommonly seen in hepatic angiograms of liver transplant recipients, reported in up to 5.4% of hepatic arteriograms in this patient population.^{1,5–7} Interestingly, hemodynamically significant

APFs warranting endovascular treatment have been reported more frequently in the literature in native (nontransplanted) compared with transplanted livers ($n > 280$ –300 versus $n = 13$, respectively).^{1,8,9} This is despite the fact that liver transplant recipients undergo numerous percutaneous procedures (liver biopsies and percutaneous transhepatic cholangiograms) and are under constant imaging surveillance.^{1–3} This may shed light as to the pathogenesis of hemodynamically significant APFs.^{1–3,10}

Etiology and Pathogenesis

APFs are considered arterial injuries leading to a connection between the high-pressure hepatic artery and an adjacent lower-pressure portal vein branch.^{1–3,8,11–15} APFs are believed to be a product of percutaneous transhepatic procedures such as percutaneous liver biopsies and percutaneous transhepatic cholangiograms (PTCs) with or without percutaneous biliary drain placement, hence the presumed etiology of iatrogenic vascular injury.^{1–3,13–15} Of the 13 APFs in liver transplants evaluated by Saad and coworkers, 11 (85%) had clearly described prior biopsies (with or without PTCs), averaging 103 days (range: 1–450 days) prior to the diagnosis of biopsy-associated APFs.^{2,8,10,16–21} Eight of the 11 cases of APFs (73%) had biopsies within 90 days from the APF diagnosis.^{2,8,10,16–21} Arterial injuries caused by percutaneous transhepatic procedures have been classified into four types: parenchymal contusions, active arterial bleeding, intrahepatic pseudoaneurysms, and arteriovenous fistulas.^{3,9,12–15} Over time, non-APF “acute” injuries (pseudoaneurysms, contusions, active bleeding) either heal or lead to death. Thus because of its enduring and insidious nature, the most prevalent arterial injury is the APF. As a result, APFs represent 37%, 86%, and 100% of all arterial injuries immediately after, 1 week after, and 4 weeks after percutaneous biopsies, respectively.^{12–14}

Certain APFs close spontaneously, whereas others may progress to large hemodynamically significant APFs.^{2,5–7,16,17} The pathogenesis of this is poorly understood and highly speculative. However, the discrepancy between the incidence of APFs in native versus transplanted livers has been a point of discussion and speculation regarding the pathogenesis of large hemodynamically significant APFs.^{1–3,10} It has been surmised that the relatively compromised arterial flow and the relative poor compliance of transplanted livers (stiff graft) are a hindrance to the progression of APFs from small hemodynamically insignificant APFs to those that are hemodynamically significant.^{1–3}

Clinical Presentation

Most APFs, hemodynamically significant or not, are asymptomatic.^{1–3,8,12,13} In those that are symptomatic, APFs have been the cause of portal hypertension (especially extrahepatic and chronic central intrahepatic APFs), pain, sepsis, hemobilia, biliary obstruction, and, rarely, pulmonary hypertension.^{1–3,8,10,16–23} Unlike large arteriovenous fistulas in the systemic circulation, hemodynamically significant APFs have not been described as causing congestive heart failure.^{2,8}

This is probably because the outflow in APFs is not systemic and the “short-circuit” between the systemic arterial to the systemic venous components is not present.²

In an analysis by Saad and coworkers,² of 11 APFs clearly described in detail in the literature, the most common presentation for APFs (73% of cases) was a dysfunctional liver transplant that had abnormal liver function tests (LFTs) only. Another common presentation was gastrointestinal bleeding (45%), with hemobilia as the most common type (27%). Pain was also a common complaint that occurred in 36% of liver transplant recipients with APFs.² Hepatic graft dysfunction and pain are a product of hemodynamically significant APFs; however, bleeding can be caused by both large and small APFs.²

Imaging Diagnosis

The gold standard imaging modality is to diagnose APF with catheter-directed digital subtraction angiography (DSA).^{1–3} However, noninvasive imaging such as contrast-enhanced computed tomography (CT) or contrast-enhanced magnetic resonance imaging (MRI) and Doppler ultrasound are important preprocedural diagnostic modalities and posttreatment follow-up modalities. Doppler ultrasound is particularly useful because it is specific to hemodynamically significant APFs.^{1–3}

Doppler Ultrasound

Doppler ultrasound may be the most important noninvasive imaging modality for diagnosing and following APFs in liver transplant recipients. In a study by Saad and coworkers, Doppler was able to detect only 50% of APFs; however, it detected all hemodynamically significant APFs.¹ The advantages of Doppler ultrasound are that it is noninvasive, the main screening tool for all vascular complications in liver transplants, and most importantly is the only noninvasive imaging modality that can hemodynamically assess the hepatic artery, portal vein, and APF.^{1–4} In fact, the definition of hemodynamic significance is based on either Doppler ultrasound criteria or conventional angiography.^{1–3}

Doppler ultrasound findings of APFs include visualizing a focus of turbidity and aliasing at the site of the APFs (►Fig. 1A), reduced arterial resistive indices (RIs) of the feeding hepatic artery, and reversal of flow with or without arterialization of venous flow in the receiving portal venous branch (►Fig. 1B).^{1,3} A reduced arterial resistive index is defined as either nominally low (<0.50) or an arterial RI that is decreased from a prior Doppler baseline examination.^{1,3} Hemodynamic significance is defined by identifying the portal venous findings in the main portal vein of the hepatic graft or its first-order branch.^{1–3} If the graft is a whole hepatic graft, the main portal vein is the main portal vein proper and its first-order branch is the right or left portal vein. If the graft is a split graft, the main portal vein of the graft is the right portal vein with its first-order branches the right posterior or anterior branch. Similarly, in a left lobe split graft, the main portal branch is the left portal vein with its first-order branches the segmental branches to Couinaud segments I, II, or IV (if IV exists, and it usually does).¹

Figure 1 (A) Doppler ultrasound of the left hepatic lobe in a liver transplant recipient who is status post liver biopsy months earlier. There is a focus of turbidity with aliasing (open arrow) with reversal of flow in the involved left portal vein branch (white arrow). (B) Doppler ultrasound with spectral waveform analysis of the involved left portal vein branch, demonstrating reversal and arterialization of flow in the left portal vein (open arrow). (C, D) Contrast-enhanced T1-weighted axial magnetic resonance (MR) images of the same patient as (B) and (C) in the arterial phase of the enhancement. The images demonstrate the involved left hepatic lobe in a portal venous phase parenchymal of enhancement, which precedes the remainder of the liver that is normal in its enhancement. The dashed arrows show the demarcation between the abnormally enhancing left lobe and the normally enhancing right hepatic lobe. Also noted is a dilated left hepatic artery (solid white arrow) when compared with the right hepatic artery (open white arrow) (D). (E, F) Two images in sequence of the initial diagnostic digitally subtracted angiogram demonstrating a portal vein branch (solid arrow) coursing adjacent to the feeding artery (open arrow) seen in the early arterial phase of the hepatic angiogram. By definition, this is an arterioportal fistula (APF). The asterisk identifies the vicinity of the APF. (G) Two contrast-enhanced T1-weighted axial MR images in the arterial phase of the enhancement before (G-i) and after (G-ii) APF embolization. The images demonstrate that the abnormal left lobe enhancement is out of phase (G-i) with the rest of the liver but becomes in phase with the rest of the liver after embolization (G-ii). The demarcation seen in (G-i) (white arrows) is no longer visualized on the postembolization image. (H) Digitally subtracted angiogram through a microcatheter (open black arrow) that has been placed into the feeding artery. Again seen is the early portal vein drainage (open white arrow). The asterisk signifies the vicinity of the APF. (I) Digitally subtracted angiogram through a microcatheter after coil embolization of the main artery (dashed arrows) supplying the APF (open arrow). Embolizing the main arterial supply and negating its sump effect has unmasked a smaller second arterial branch (solid black arrows) that has fistulized with the same portal vein branch (asterisk). (J, K) Two digitally subtracted angiogram images in two separate oblique projections better visualizing the smaller artery (solid black arrows) that fistulizes (asterisk) with the portal vein branch (solid white arrows). Coils are seen in the embolized main arterial supply (open arrow). The operator at this point used polyvinyl alcohol to embolize this small branch. (L) Two digitally subtracted angiograms of the same frame rate, same time after injection and same injection rate before (L-i) and after embolization (open arrow, L-ii). The APF (asterisk, L-i) has been completely occluded (L-ii); the early portal vein branch (white arrows) is no longer visualized. Notice that there is better filling of the hepatic artery due to the loss of the sump effect (steal effect) of the APF.

Contrast-Enhanced Computed Tomography or Magnetic Resonance Imaging

Contrast-enhanced CT and/or contrast-enhanced MRI, with or without angiography, probably are the most sensitive noninvasive modalities in identifying all sizes of APFs. CT and MRI, however, are usually not specific to borderline hemodynamically significant APFs. Typically, on the arterial

phase of a CT or MR, the involved hepatic segment(s) (the segment supplied by the “fistulized” portal vein branch) demonstrates portovenous parenchymal enhancement that is “ahead” and mistimed with the remainder of the liver enhancement, which is still in the arterial phase of the contrast enhancement (►Fig. 1C, 1D).¹ A large abnormally enhancing hepatic vascular territory usually indicates central

Figure 1 Continued.

APFs but not necessarily large hemodynamically significant APFs; having said that, central APFs have a higher likelihood of developing into a hemodynamically significant APF. Occasionally, a dilated feeding hepatic artery and/or a dilated receiving/draining portal vein radical is visualized, signifying a hemodynamically significant APF (→Fig. 1D).

Digital Subtraction Angiography

DSA is the gold standard examination for diagnosing APFs of all sizes, as well as identifying hemodynamically significant APFs.¹⁻³ It is a very sensitive and specific modality for both. Even the smallest APFs are defined as visualizing a portal vein branch adjacent to a hepatic arterial branch in the early arterial phase of the DSA.¹⁻³ The fistulous tract does not have to be seen, and in fact is rarely seen, to make the diagnosis.¹⁻³ Hemodynamically significant APFs are defined as visualization/contrast filling of the main portal vein of the graft or its first-order-branch.¹⁻³

Postembolization Imaging Follow-up

Post-successful embolization imaging follow-up is performed to evaluate for the reversal of signs of hemodynamic significance. It should be primarily performed by Doppler ultrasound, which is the best noninvasive imaging modality to assess and evaluate for the hemodynamic significance of APFs.¹⁻³ On follow-up imaging, one should notice reversal of the portal venous Doppler ultrasound findings that signify hemodynamic significance.¹⁻³ By contrast-enhanced CT or MR, the out-of-phase enhancement pattern of the involved

hepatic segment, described earlier, should return to a normal enhancement pattern (→Fig. 1G).¹

Management

Traditionally, only symptomatic APFs (e.g., causing ascites, graft failure, bleeding) warrant treatment.¹⁻³ However, with the advent of minimally invasive endovascular therapy, with its relative safety and effectiveness, asymptomatic but hemodynamically significant APFs with hepatic graft dysfunction (abnormal LFTs) have been treated. In addition, APFs that show progressive enlargement have also been treated.¹⁻³ Anecdotally, it is unfortunately not always easy to determine definitively whether an APF is the cause of abnormal LFTs. Abnormal LFTs/dysfunctional grafts are nonspecific, and embolizing APFs may be a “diagnostic-therapeutic” approach to a problem (i.e., “Fix it and see if that alleviates the dysfunction”). This approach, however, is still controversial, and APF cases in transplants should be managed on a case-by-case basis.

Surgical Management

The management of APFs historically is surgical, by ligation of the involved hepatic artery segment, hepatic graft segmentectomy, lobectomy, or even retransplantation.^{1,3,13,16,24,25} Being surgically minimalistic by performing intrahepatic surgical ligation is actually not the simplest of surgeries; it requires tedious methods of tissue dissection and vascular reconstruction, and it may involve massive blood loss.²³

Endovascular Management

There has been a paradigm shift in the last decade toward the endovascular management of APFs in liver transplant recipients.^{8,10,16–22} This was attributed by Saad and coworkers to the inadequate and invasive surgical techniques due to the (usually) deeply located APFs within the hepatic parenchyma that cause major blood loss or result in a significant size segmentectomy.² However, endovascular management (almost always with embolization) is also difficult and not without risk.²

Endovascular management of APFs in transplant recipients is riskier than endovascular management of APFs in native (nontransplanted) livers. This is because the hepatic graft is a far less forgiving organ than a native liver in terms of hepatic arterial supply, and it is particularly susceptible to ischemia.^{1–3,26} Any hepatic artery thrombosis as a complication of embolization may have devastating consequences.^{1–3} Furthermore, the fistulous tract is very short and often not precisely visualized, and thus the embolization is actually of the more proximal feeding artery; this results in a much larger potential zone of ischemia. The larger the APF and the larger and more central the feeding artery, the higher the risk that a successful embolization of the feeding artery will also devascularize a large territory of hepatic graft parenchyma from the arterial flow.¹ This is the reason why some authors advocate treating rapidly growing asymptomatic APFs before they grow to an extent where larger arterial branches are required to be embolized.¹ It is not uncommon (15% of APF cases) to find progression of the hemodynamic significance of APFs over 2 to 9 months.^{2,20} Regarding growth and centrality of APFs, endovascular embolization procedures have been classified as either low ischemic risk procedures (small peripheral feeding vessels that can be sacrificed with minimal collateral devascularization) and high ischemic risk procedures (larger more central feeding vessels that when sacrificed will have significant collateral devascularization due to a large vascular territory that has been embolized).²

Embolization of APFs are now typically performed with microcatheters and microcoil techniques in an attempt to be as superselective as possible. As mentioned earlier, the fistulous tract is usually not definitively visualized, and the aim is usually to embolize the feeding artery as close to the APF as possible (►Fig. 1H, 1I). The angiographic end point is neither pristine angiographic images nor complete obliteration of the APF. The successful end point is a reduction of the APF from hemodynamically significant to hemodynamically insignificant.^{1–3} As a result, not uncommonly (20 to 25% of APFs) may require more than one session for adequate embolization/obliteration of the APF.² In fact, some authors are content with partial embolization as long as the portal venous findings of hemodynamic significance are alleviated by follow-up Doppler ultrasound.¹

Particulate embolization with polyvinyl alcohol and even Gelfoam (Upjohn, Inc., Kalamazoo, MI) may be helpful in reaching the desired end point (►Fig. 1I–K). An interesting angiographic finding is the phenomenon of having multiple feeding arteries, seen in up to 20 to 25% of APF cases in liver transplants, which has led some authors to dub these cases

“complex APFs.”^{1,3} Unfortunately, not all APF fistulas present themselves on the initial DSA. Similar to embolizing an arteriovenous malformation, embolizing the main feeding artery may unmask and expose the secondary feeding vessels (►Fig. 1I–K). For this reason, it is important that operators perform a less selective hepatic angiogram after initial embolization to identify additional secondary feeders/APFs (►Fig. 1L).

Endovascular management of APF overall is safe and effective.² Approximately 20 to 25% of cases require more than one session to reach the desired effect/hemodynamic end point.² The overall technical success rate is 63 to 77%, with an anticipated complication rate of 12 to 15%. However, the intent-to-treat graft survival at 6 months has been reported to be up to 36%.²

Conclusion

APFs in liver transplants are not uncommon; however, most are incidentally noted and of questionable significance; hemodynamically significant APFs are rare, occurring in ~0.2% of transplants. Doppler ultrasound and digital subtraction angiography are the two modalities that can define hemodynamic significance. It remains controversial whether to treat asymptomatic APFs, although treatment is performed for patients with abnormal LFTs in the presence of a hemodynamically significant APF. Endovascular therapy is safe and effective, and now considered the treatment of choice.

References

- 1 Saad WE, Davies MG, Rubens DJ, et al. Endoluminal management of arterioportal fistulae in liver transplant recipients: a single-center experience. *Vasc Endovascular Surg* 2006;40(6):451–459
- 2 Saad WE, Lippert AJ, Davies MG, et al. Prevalence, presentation, and endovascular management of hemodynamically or clinically significant arterio-portal fistulae in living- and cadaveric-donor liver transplant recipients. *Clin Transpl* In press
- 3 Saad WE. Management of nonocclusive hepatic artery complications after liver transplantation. *Tech Vasc Interv Radiol* 2007;10(3):221–232
- 4 Saad WE, Lin E, Ormanoski M, Darcy MD, Rubens DJ. Noninvasive imaging of liver transplant complications. *Tech Vasc Interv Radiol* 2007;10(3):191–206
- 5 Wozney P, Zajko AB, Bron KM, Point S, Starzl TE. Vascular complications after liver transplantation: a 5-year experience. *AJR Am J Roentgenol* 1986;147(4):657–663
- 6 Zajko AB, Bron KM, Starzl TE, et al. Angiography of liver transplantation patients. *Radiology* 1985;157(2):305–311
- 7 Karatzas T, Lykaki-Karatzas E, Webb M, et al. Vascular complications, treatment, and outcome following orthotopic liver transplantation. *Transplant Proc* 1997;29(7):2853–2855
- 8 Jabbour N, Reyes J, Zajko A, et al. Arterioportal fistula following liver biopsy. Three cases occurring in liver transplant recipients. *Dig Dis Sci* 1995;40(5):1041–1044
- 9 Portal A. Mémoires sur quelques maladies du foie, qu'on attribue à d'autres organes et sur des maladies dont on fixe ordinairement le siège dans le foie quoi qu'il n'y soit pas. *Histoire de l'Académie Royale Sciences*, année 1777, 1780:601
- 10 Figueras J, Soubrane O, Pariente D, Devictor D, Houssin D. Fatal hemobilia after liver graft biopsy in a transplanted child. *Gastroenterol Clin Biol* 1994;18(8–9):786–788

- 11 Saad WE. Management of nonocclusive hepatic artery complications after liver transplantation. *Tech Vasc Interv Radiol* 2007; 10(3):221–232
- 12 Saad WE, Davies MG, Ryan CK, et al. Incidence of arterial injuries detected by arteriography following percutaneous right-lobe ultrasound-guided core liver biopsies in human subjects. *Am J Gastroenterol* 2006;101(11):2641–2645
- 13 Hellekant C. Vascular complications following needle puncture of the liver. *Clinical angiography. Acta Radiol Diagn (Stockh)* 1976;17(2):209–222
- 14 Okuda K, Musha H, Nakajima Y, et al. Frequency of intrahepatic arteriovenous fistula as a sequela to percutaneous needle puncture of the liver. *Gastroenterology* 1978;74(6):1204–1207
- 15 Hellekant C, Olin T. Vascular complications following needle puncture of the liver. An angiographic investigation in the rabbit. *Acta Radiol Diagn (Stockh)* 1973;14(5):577–582
- 16 Chavan A, Harms J, Pichlmayr R, Galanski M. Transcatheter coil occlusion of an intrahepatic arterioportal fistula in a transplanted liver. *Bildgebung* 1993;60(4):215–218
- 17 Vivas S, Palacio MA, Lomo J, et al. Arterioportal fistula and hemobilia in a patient with hepatic transplant [in Spanish]. *Gastroenterol Hepatol* 1998;21(2):88–89
- 18 Charco R, Margarit C, López-Talavera JC, et al. Outcome and hepatic hemodynamics in liver transplant patients with portal vein arterIALIZATION. *Am J Transplant* 2001;1(2):146–151
- 19 Botelberge T, Van Vlierberghe H, Voet D, Defreyne L. Detachable balloon embolization of an arterioportal fistula following liver biopsy in a liver transplant recipient: a case report and review of literature. *Cardiovasc Intervent Radiol* 2005;28(6): 832–835
- 20 Detry O, De Roover A, Delwaide J, Dondelinger RF, Meurisse M, Honoré P. Selective coil occlusion of a large arterioportal fistula in a liver graft. *Liver Transpl* 2006;12(5):888–889
- 21 Otobe Y, Hashimoto T, Shimizu Y, et al. Formation of a fatal arterioportal fistula following needle liver biopsy in a child with a living-related liver transplant: report of a case. *Surg Today* 1995;25(10):916–919
- 22 Chua DY, Pavillion G, Tay KH, Tan JL, Chung AY. Pulmonary hypertension: an unusual presentation of an iatrogenic hepatic arterioportal fistula and its successful resolution post-embolotherapy. *Am Surg* 2009;75(6):511–514
- 23 Strodel WE, Eckhauser FE, Lemmer JH, Whitehouse WM Jr, Williams DM. Presentation and perioperative management of arterioportal fistulas. *Arch Surg* 1987;122(5):563–571
- 24 Agha FP, Raji MR. Successful transcatheter embolic control of significant arterioportal fistula: a serious complication of liver biopsy. *Br J Radiol* 1983;56(664):277–280
- 25 Saad WE. Management of hepatic artery steno-occlusive complications after liver transplantation. *Tech Vasc Interv Radiol* 2007; 10(3):207–220
- 26 Saad WE, Davies MG, Sahler LG, et al. Hepatic artery stenosis in liver transplant recipients: primary treatment with percutaneous transluminal angioplasty. *J Vasc Interv Radiol* 2005;16(6): 795–805