GIANT SUPRAGLOTTIC SCHWANNOMA

Rajani Kanth. T.V.,* R. Nanayya,** Y. Satyaprabhakara,* K. Blandeena,*** P.S.N. Murthy****

Abstract: Schwannomas arising from the supraglottis are uncommon, if not rare. They are known to arise from the head and neck region but oropharyngeal presentation is not often seen. Few cases of pedunculated schwannoma arising from the aryepiglottic fold presenting as giant tumours in the oropharynx have been reported so far. The tumour in this case has been excised via intra-oral approach, with nil residual morbidity. A follow up period of one year was uneventful without any recurrence.

Keywords: Schwannoma, Supraglottis.

INTRODUCTION

Schwannomas are benign, encapsulated, peripheral nerve tumours that arise from the Schwann cell. Approximately 45% of them occur in the head and neck region¹. The lateral cervical region and the mouth are the most common sites. Schwannomas tend to affect women more often than men and are most frequently seen in the third to sixth decades of life². Neurogenic tumours are uncommon in the larynx comprising less than 0.1% of benign laryngeal neoplasms. In the larynx, these tumours tend to arise either from the aryepiglottic fold or the true vocal cord and are submucosal. Only a few cases of pedunculated schwannomas arising from the aryepiglottic fold presenting as oropharyngeal masses have been reported so far and we are reporting one such case.

CASE REPORT

A thirty year old male presented with the history of a feeling of lump in the throat, difficulty in swallowing and change in voice for the last one year. The symptoms were insidious in onset and progressively worsened with time. For the last couple of months, the patient experienced difficulty in breathing, especially when he was lying down supine. He was a nonsmoker and was not an alcoholic.

On examination, he was found to be poorly nourished and had a mild degree of stridor. His speech was plummy with an element

of hyponasality. Clinical examination of the throat revealed a smooth, globular, pedunculated, pinkish mass occupying almost the entire oropharynx. It seemed to arise from below, the superior aspect of the mass being free. On palpation, it was firm, nontender and the site of the attachment could not be determined. Indirect laryngoscopy was not possible because of the oropharyngeal mass. No significant cervical lymphadenopathy was noted. An X-ray film of lateral view of the neck showed a well rounded soft tissue shadow filling up the oropharynx and impinging on the laryngeal inlet. CT scan of the neck showed the same with the mass occupying more than 90% of the air column in the maximum diameter. There was partial enhancement with contrast.

Tracheostomy was done to relieve him of his respiratory difficulty. Punch biopsy of the lesion revealed only chronic inflammatory tissue on histopathological examination. We further decided to do an excisional biopsy of the tumour. After routine investigations and obtaining fitness for surgery, the patient was taken up for operative procedure. Under general anesthesia, a Boyle-Davis mouth gag was introduced and the oral cavity accessed. The pedicle of the tumour was identified at the right aryepiglottic fold and was ligated. Hemostasis was achieved with the help of cautery. The specimen was sent for histopathological examination. The patient was started on oral feeds on the first postoperative day and decannulation of tracheostomy was done on the third postoperative day. He was subsequently discharged on the fifth postoperative day. There has been no evidence of recurrence or any other complaint for the last one year.

Pathological examination showed a cellular schwannoma composed of compact (Antoni A) areas and loose meshed (Antoni B) areas. The tumour was made of spindle cells with wavy nuclei showing nuclear pleomorphism with thick

Fig 1: Oropharyngeal part of the tumour seen on oral examination

Fig 2: X-ray film of lateral view of the neck showing a well-defined soft tissue swelling

*Assistant Professor, ** Professor, ***Associate Professor, NRIAMS, Vijayawada, **** Professor, PSIMS, Vijayawada

Fig III: CT Scan film of the tumour filling up the Oropharynx.

Fig IV: The excised Tumour specimen

Fig V: Histopathology showing Antoni A, Antoni B Areas.

walled blood vessels. Few lymphocytes were scattered in between the tumour cells.

DISCUSSION

Schwannomas are a type of benign peripheral nerve sheath tumors (BPNSTs), also known as neurilemmomas. They were first described by Verocay³. Schwannomas usually arise from the cranial (10%) and spinal (25%) nerve roots. They can arise at any site of the peripheral nervous system, with the exception of the olfactory and optic nerves.

Among the neurogenic tumours in the head and neck region, parapharyngeal space is the most commonly involved. Isolated schwannoma of the larynx is rare. New and Erich⁴ found only one case of schwannoma of 722 benign laryngeal tumours. Holinger and Johnston⁵ found only one case of Schwannoma among 1197 benign lesions of the larynx. Dysphagia and shortness of breath are the usual presenting symptoms of laryngeal schwannomas mentioned in the literature⁶. Laryngeal schwannomas are nearly always solitary, submucosal, well circumscribed and occasionally painful. Schwannoma of the larynx usually involves the aryepiglottic folds⁷ but the involvement of the ventricular folds⁸ arytenoids⁹ and vocal cords⁴ have also been described. Surgery is the treatment of choice. An endoscopic approach can be attempted for smaller tumours but for large tumours, external approaches like medial thyrotomy, lateral pharyngotomy and lateral thyrotomy are used. In this case, where the tumour was exceptionally large reaching the oropharynx, the pedicle was narrow and we were able to ligate it and remove the tumour by transoral approach.

Histopathological examination reveals alternating cellular and edematous regions referred to as Antoni A and Antoni B areas. In cellular schwannomas, increased mitotic activity is seen. Immunohistochemistry of Schwannoma shows reactivity for S-100 protein. Since laryngeal schwannomas are supraglottic, the internal branch of superior laryngeal nerve is supposed to be the nerve of origin¹⁰.

CONCLUSION

Schwannomas arising from the supraglottic region usually present

early because of symptoms such as dysphagia and change in voice. Rarely they may present at a later stage when they occupy the oropharynx. When they are small, they can be removed transorally with the aid of endoscopy. Large tumours usually need an external approach. This case, we feel, is an exception than the rule.

REFERENCES

1. Hawkins DB, Luxford WM. Schwannoma of the Head and Neck in children. *Laryngoscope* 1980; 12; 1921-1926.
2. Thurner D., Quint C., Parmer J., et al - *Archives of Otolaryngology and Head and Neck Surgery*, 128 (7) : 850-852, 2002.
3. Decker P. J., Haidar A. Neurilemmoma of the larynx. *British Journal of Clinical Practice*. 1994; 48 (3) ; 159-160
4. New GB, Erich JB. Benign tumours of the larynx ; a study of 722 cases . *Arch Otolaryngol* 1983 ; 28 ; 841-910.
5. Holinger PH, Johnston KC. Benign tumours of the larynx. *Ann Otol Rhinol Laryngol* 1951 ; 60; 496-509.
6. Mehta S. Neurilemmoma of the larynx. *Ear Nose Throat Journal*. 1991 ; 70(2); 72-73.
7. Ogura JR, Biller HF. Cysts and tumors of the larynx . In; Paperella MM, Schumrick DA , eds *Otolaryngology*. Vol.3. Philadelphia; WB Saunders, 1973, 658-665.
8. Palva T, Jokenon K, Karja J. Neurilemmoma of the larynx. *Journal of Laryngology and Otolaryngology* 1975; 89 (2); 203-207.
9. Hippel K, Chmielewski G. Schwannom des Larynx bei Morbus Recklinghausen. *Differentialdiagnostische Erwagungen und immunohistochemischer Nachweis*. *Laryngorhinootologie* 1989; 68 ; 611-613.
10. Nanson EM. Neurilemmoma of the larynx ; a case study. *Head Neck Surg* 1978 ; 1; 69-74.

Address for correspondence

Dr. Rajani Kanth T.V.,
60-27-3/5, Behind Pinnamaneni Polyclinic,
Sidharthanagar, Vijayawada 520 010.
Phone- 9398324153