Pathological Fracture of the Mandible, Secondary to Eosinophilic Granuloma in a Medically Compromised Patient

M. J. Baliga, Prof. and Head, **E. Darvish**, Resident, **L. Mallya**, Asst. Prof.,
G. K. Singh, Post Graduate Orthopaedics,

Dept. of Oral and Maxillofacial Surgery, College of Dental Surgery, K.M.C. , Chandra Building
Balmatta, Mangalore -575 001, India

A case of eosinophilic granuloma of the mandible with submandibular lymph node involvement in a medically compromised patient is presented. The lesion caused the pathologic fracture of the mandible. It was treated radically with the resection of the mandible between the angle posteriorly and the midline anteriorly and immediate reconstruction with a 3DBDB AO reconstruction plate. This was followed by telecobalt therapy of 2000 cGys in 10 fractions over 20 days.

The combined surgical and radiation therapy should be reserved for patients having : (1) unremitting pain following curettage (2) Limitation of activity (3) a medically compromised patient or (3) in whom radiographic findings suggest an aggressive lesion involving adjacent tissues and/or causing pathologic fractures.

Key Words

Eosinophilic Granuloma, Histiocytosis X, Pathological Fracture

INTRODUCTION

Eosinophilic granuloma account for more than 50% of all cases pertaining to Histiocytosis X^{1, 2, 3} : It occurs most frequently in the first two decades of life, with males being affected twice as often as females⁴. This reticuloendothelial disorder shows great variations with respect to its presentation. The presenting symptoms vary from none to pain, swelling and tenderness, accompanied by general malaise and fever.

This case report describes a case of eosinophilic

granuloma of the mandible with submandibular lymph node involvement in a medically compromised patient. The purpose of this presentation is to report a case where in eosinophilic granuloma led to a pathologic fracture of the involving bone ; and to formulate criterion for selection of treatment modality to be instituted in patients suffering from this disease.

CASE REPORT

A 45 year old female patient complaining of pain in the left posterior region of the mandible, was treated for apical periodontitis of the molar, followed by extraction of the offending teeth, by her family dentist. The pain resolved and reappeared after 4 weeks. No Radiographic examination was done at this stage but the patient was advised to undergo medical treatment. Three months later the patient reported to a medical centre where she was investigated and treated. On the first of March 1993, the patient reported to the maxillofacial clinic at Mangalore, with complaints of pain and tenderness at the left lower border of the mandible.

The patient's medical history was significant as she was found to be a diabetic. Extraoral examination showed no facial asymmetry, with palpable left submandibular lymph node, which was nontender and firm in consistency. Intra-orally the left lower alveolus was found to be obliterated with no obvious pathology. A panoramic radiograph revealed a radiolucent lesion with ill defined borders, extending to the inferior border of the mandible and antero posteriorly from the angle (left) to the midline

anteriorly. A pathologic fracture of the body of the mandible was noticed in the premolar region (Fig.1).

Contact with the previous medical centre revealed that the patient had undergone curettage in the same region, which was diagnosed as eosinophilic granuloma. To exclude the polycystic form of the disease, a bone marrow biopsy and ultra sound of the abdomen were done along with

patient was put on plain insulin prior to surgical intervention.

Under naso-endotracheal anaesthesia the mandible was resected from the left angle to the midline with the clearance of the soft tissues in the submandibular triangle. Immediate reconstruction of the surgical defect was undertaken with a 3DBDB AO reconstruction plate (Fig.2). The patient had an uneventful

Fig.1

Panoramic radiograph showing a radiolucent lesion of the left mandibular body. Note the pathological fracture of the inferior cortical plate

other conventional osseous radiographs. Other investigations done included a total hemoglobin count, urine analysis and E.C.G. which were found to be within normal limits. A fine needle aspiration biopsy of the left submandibular lymph node revealed an infiltrate of histiocytes and eosinophils.

Following admission, consultation with the endocrinology and oncology units, were obtained. Since the patient's blood glucose was found to be very high (fasting 210 mg/dl), the

recovery. This was followed by telecobalt therapy of 2000 cGys in 10 fractions over 20 days. The patient has been followed up, showing good relief and no recurrence to date.

Microscopically, the lesion was composed primarily of histiocytes, some of which were found to contain lipid material and variable infiltrate of inflammatory cells including eosinophils and multinucleated giant cells (Fig.3). This confirmed the diagnosis of eosinophilic granuloma.

DISCUSSION

Eosinophilic granuloma primarily affects children and young adults, although it may occur at any

The treatment of this disease varies from surgical curettage, local irradiation, chemotherapy to local steroid injections. The treatment modality

Fig.2

Post surgical panoramic radiograph showing reconstruction of surgical defect using – 3 DBDB AO reconstruction plate.

age. In a review done by Slater and Swarm¹, 70% of the affected patients were younger than 20 years, with males being affected more often than females. It is characterised by solitary or multiple lesions of bone, involving the calvarium, femur, mandible or the ribs, with or without pulmonary involvement and usually associated with a good prognosis. Monostatic form of the disease occurs more frequently than the eopolystatic form in most series¹⁻³.

The etiology still remains uncertain. The various possible causes mentioned in the literature include – (1) a genetic predisposition⁵, (2) infections⁶⁻⁸, (3) immunodeficiency or (4) an abnormal response of the immune system to certain antigens⁹.

Fig.3

Photomicrograph of the excised lesion showing abundance of histiocytes and a variable infiltrate of inflammatory cells H&E stain 400 X.

depends upon the age and medical status of the patient, extent of the disease, and the sites of involvement.

The use of intralesionary injections of steroids was tried at various times, based upon the hypothesis that the drug destroys the granulomatous tissue and induces osteogenesis. Curettage of the lesion has been suggested but complete removal of the lesion is not necessary in most cases. Response to biopsy alone exists in literature¹⁰. Some authors suggest that curettage followed by filling the defect with cancellous chips prevents further complications like pathological fractures.

Radiation therapy should be reserved for inaccessible lesions and should not be contemplated in children, because of the

development of secondary malignancies. The combined surgical and radiation therapy should be reserved for patients having unremitting pain following curettage limitations of activity, a medically compromised patient or in whom radiographic findings suggest an aggressive lesion involving adjacent tissues and/or causing pathologic fractures. Reports of pathologic fracture of the mandible as a complication of eosinophilic granuloma are rare², but since this case is associated with majority of the symptoms, mentioned above, indicating a more radical treatment plan, a combined surgical and radiation therapy was undertaken. It is also essential to undertake a full body CT-scan or osseous radiographs to rule out the disseminated form of the disease prior to any form of surgical intervention.

References

1. Slater JM, (1980) : Swarm OJ. Eosinophilic Granuloma of Bone. *Med. Pediatr. oncol* 8 : 151.
2. Hartman KS, (1980) : Histiocytosis X : A review of 114 cases with oral involvement *Oral Surg.* 49 : 38.
3. Wester SM, Krishnan VK, Beabou JW, et al (1982) : Langerhans cell Granulomatosis (Histiocytosis X) of bone in adults. (1982) : 413.
4. Oberman HA, (1961) : Idiopathic Histiocytosis : a clinicopathologic study of 40 cases and review of the literature on Eosinophilic granuloma of Bone, Hand – Schuller-Christian Disease, And Letterer - Siwe Disease. (1961) : *Pediatrics* 28 : 307.
5. Zinkham WH, (1976) : Multifocal Eosinophilic Granuloma. *Am J MED* 60 : 457.
6. Rapis AD, (1978) : Histiocytosis X : Analysis of 50 cases. *Inj J Oral Surg* 7 : 76
7. Deman JCH, (1968) : Rod-like Tubular Structures in the cytoplasm of Histiocytes in "Histiocytosis x" *J. Pathol Bacteriol* 95 : 123.
8. Krutchkoff DJ, Jones CR. (1984) : Multifocal Eosinophilic Granuloma A Clinical Pathologic Conference. *J. Oral Pathol.* 13(5) : 472-88.
9. Granda FM, Mc Daniel RK, (1982) : Multiple Progressive Eosinophilic granuloma of the jaws. *J Oral Maxillofac Surg.* 40(3) : 174-8.
10. McCullough CJ, (1980) : Eosinophilic granuloma of Bone. *Arch Orthop Scand* 51 : 389.

Congratulation !

Dr. S. K. Dey for publishing 6th edition of Fundamentals of Ear, Nose and Throat and Head-Neck Diseases.