Lead Article

Abstract

A study was undertaken to evaluate the role of unilateral/bilateral submucous resection of the inferior turbinates in fifty cases of chronic hypertrophic rhinitis. Patients associated with deflected nasal septum or sinus infections were excluded from the study. Decongestion of turbinates was done to exclude the cases with predominantly mucosal hypertrophy. Gertner (1984) plate method was used to assess the nasal patency pre-operatively and then post-operatively at 1, 3 and 6 months follow up. The analysis of observations made revealed SMR of inferior turbinate to be a very effective modality for relieving nasal obstruction due to bony turbinal hypertrophy. The procedure has least interference with nasal mucosal integrity and functions and complications associated with the procedure have been found to be minimal. Histopathological examination of mucosa and of ostentid tissue revealed infiltration by chronic inflammatory cells suggesting chronic nonspecific inflammation either due to non-specific infections and / or allergy.

S M R of Inferior Turbinate in Chronic Hypertrophic Rhinitis

Karan Sharma, Senior Lecturer, K. K. Duggal, Professor Incharge,
Jagdeep S. Hundal, Junior Resident, Department of E.N.T.,
Govt. Medical College, Amritsar, India
Pin-143001

Turbinate hypertrophy is one of the major causes of nasal obstruction. Usually it is the inferior turbinate which often gets enlarged as a result of deflected nasal septum, Allergic or Non allergic rhinitis, Rhinitis medicamentosa, oral medications like antihypertensive and oral contraceptives etc. While the symptoms of rhinorrhoea and sneezing often respond to medical treatment, nasal obstruction secondary to chronic hypertrophy doesn't. Many local procedure have been mentioned in the literature for the relief of perennial nasal obstruction due to hypertrophied inferior turbinate. These include intratubinal steroid injection, electric cauterization cryosurgery, sub-mucosal diathermy, turbinate out-fracture, partial or total inferior turbinectomy and submucosal resection of conchal bone. Most of these procedures give temporary relief and if carried out too extensively or repeated too frequently result in fibrosis and altered physiology. Partial or total turbinectomy may offer complete relief from nasal obstruction but do carry a definite risk of profound physiological changes often resulting in mucosal atrophy and ozaena. Thus with the aim of permanent relief without any altered nasal physiology, we have carried out unilateral or bilateral submucosal turbinectomy to evaluate its results.

Material and Methods

Fifty patients with unilateral or bilateral nasal obstruction as a result of chronic hypertrophic rhinitis were taken up for surgery in this study, conducted in the E.N.T. Department of Government Medical College, Amritsar. Patients having symptomatic Deflected Nasal Septum and sinus infections refractory to conventional surgical/medical treatment were outrightly excluded from this study. The shrinkage of turbinates were performed in all cases to differentiate mucosal hypertrophy from bony turbinal enlargement and patients with thick turbinal bone were only included in this study. Nasal smear examination was done in all the cases for the presence of eosinophils or mast cells. Nasal patency was measured by the plate method of Gertner (1984) based on the hygrometric principle i.e. by measuring the total area of plate undergoing vapour condensation.

Surgical Technique

Premedication with intramuscular 0.6 mg Atropine and 10 mg Diazepam was given half an hour prior to surgery. Nasal cavities were packed with xylocaine 4% first and then after 10-15 minutes, the inferior turbinate was infiltrated with xylocaine 2% with adrenaline. An elliptical incision was made on the anterior end of inferior turbinate to circumscribe a crescent of hyperplastic mucosa. The mucoperiosteum was elevated from all the surfaces of roughened turbinal bone

by sharp dissection and then the bone was out as laterally as possible. Little of mucosa and a piece of excised turbinal bone was subjected to histopathological examination. The incision given was approximated properly and the nasal cavity/cavities were gently packed above and below the turbinate with an antibiotic soaked gauze pack. The packs removed after 48 hours and antibiotics along with antihistaminics decongestants and analgesics were given orally for 5-7 days. Patients were followed at 1, 3 and 6 months interval. Our technique differed from House (1951) in a way that almost complete turbinal bone was removed, of course, just medial to its lateral attachment so as to preserve future support for left periosteomucous memberane and to prevent its collapse. Also our incision was large and was left unstitched, though apposed properly, avoiding any haematoma formation post operatively.

Measurement of Nasal Patency

A chromium coated plate of 10 cm x 12 cm size, marked with arches 1 cm apart was held by the patient under the nostrils in a horizontal position in such a way that the vertical axis of the plate is 90° towards the upper lip and straight under the collumella (Fig. 1). The patient was asked to breathe out through nose slowly without force and with mouth closed. The area undergoing fogging on each side was calculated by the formula :

$S = \Pi ab$ where

a = half vertical diameter of fogging area

b = half transverse diameter of fogging area.

Observations

Sixty percent of the patients were in third decade of life. The youngest patient was fifteen years old whereas the eldest one was 37 years old. 76% of patients in this study were males. All the patients had chief complaint of almost continuous nasal obstruction which was unilateral in 28% and bilateral in rest 72%. Intermittent or continuous mucoid to muco-purulent anterior nasal discharge was complained off by all the patients. 80% also complained off posterior nasal discharge. In addition, 36% patients had the chief complaint of troublesome sneezing. Hyposmia was observed in one fifth of the cases. Nose examination revealed bilateral inferior turbinate hypertrophy in 72% whereas in rest

Fig. 1. The patient, 10 years old boy with swelling of left side of midface.

28%, it was unilateral. Signs and symptoms of otitis media were seen in 32% of patients out of which half were having otitis media with effusion whereas the rest were having frank tube-tympanic type of suppurative otitis media. Recurrent attacks of sore throat was complained by 18% and history of headache was seen in 8% of the patients in this study.

Pre-operatively, nasal patency test revealed total area of vapour condensation on the plate to be 13.54 sq.cms (Range 9.9-21.44) in bilateral cases whereas it was found to be 6.06 sq.cms. (Range 4.71-7.54) on the side of obstruction in unilateral enlargement cases. Post-operatively in bilateral SMR of turbinate cases, total area of vapour condensation was found to be 15.13 sq. cms. (Range 11. 56-23.50), 19.49 sq. cms. (Range 15.77-26.15) and 20.81 sq. cms. (Range 12.44-28.34) at 1 month, 3 months and 6 months follow-ups respectively. Similarly in cases of unilateral SMR of inferior turbinate the mean area of vapour condensation on the operated side was 6.88 sq.cms. (Range 5.59-8.02), 7.65 sq.cms. (Range 7.46-10.70) and 9.73 sq.cms. (Range 7.91-11.78) at three consecutive follow ups of 1, 3 and 6 months respectively. (Fig. 2).

Percentage improvement was calculated by dividing the area of vapour condensation post operatively by the area of vapour condensation pre-operatively and then multiplying it by 100. This percentage improvement was calculated separately at each follow up and it was found to be maximum at 3rd follow up. It was in the range of 60-80% in 60% cases and more than 80% in 8% of the cases. The range of improvement

Figure II
Graph showing mean area of vapour condensation (in sq.cm.) Preoperatively and postoperatively at follow up of one month, three months and six months in Bilateral and unilateral sub-mucous turbinectomy groups.

was between 40-60% in 24% of the cases and was less than 40% in another 8% of the cases in this study.

Nasal Smear examination revealed 4-5 eosinophils per high power field in 28% of the patients and mast cells in none. Histopathological examination of the mucosa and turbinate bone was consistent with chronic non-specific inflammatory changes. Soft tissue section showed pseudostratified columnar epithelium with prominent basement membrane and numerous seromucinous glands with intervening stromal tissue infiltrated by lymphocytes, plasma cells and macrophages. Section of turbinate bone also showed infiltration of osteoid tissue with chronic inflammatory cells. None of the cases showed any signs of chronic specific rhinitis.

Excessive intraoperative bleeding was seen in 4% cases which of course was managed by tight anterior basal packing, removed after 72 hours. None of our cases had reactionary or secondary haemorrhage. Synechiae formation was also observed in 4% cases at first follow up and was treated. None of our cases had any tendency towards crust formation.

Discussion

Many modalities in the treatment of hypertrophied turbinates have been tried but none so far has been established as the method of choice. Total inferior turbinectomy was first reported in 1895 by Jones. Since then many new methods have been advocated and sub-mucous resection of inferior turbinate is one of them. It was House in 1951 who recommended this procedure for uncontrollable perennial inferior turbinal enlargement and advocated removal of only anterior one-third of inferior turbinate bone sub-mucosally without fracturing of its remaining two-third. In the pathology of chronic rhinitis, although the mucosa throughout the nasal cavity and paranasal sinuses is involved yet the areas severely affected are the ones heavily endowed with sinusoidal erectile tissue and mucous secreting glandular elements (Principato 1979). These are primarily seen over the entire surface of the inferior turbinate lower one-third to one-half of the middle turbinate and an area of septal mucosa adjacent to anterior one-third of inferior turbinate. Moreover the inferior turbinate is the one which bears the main brunt during the physiology of nasal respiration. Thus sub-

mucosal resection of the inferior turbinate is thought to be an ideal procedure for reducing the size of the inferior turbinate with a minimal interference to its physiological functions.

For confirming and evaluating the subjective complaint of nasal obstruction Zwardemaker (1889) first described a simple hygrometric method of assessing the nasal patency by using a cold mirror. Using the same principle of hygrometry, Gertner et al (1984) evolved the plate method for assessing the nasal patency. During this study, we have found metal plate method to be a very simple and easy to perform method of rhinometry. It is comfortable for the patient and is clinically applicable as an office procedure. This method is not dependent upon the temperature and humidity of the surrounding air and above all, it does not interfere with the anatomy of the nose at all.

Following surgery all of our patients in this study showed an increase in nasal patency, though the percentage improvement varied from 32.83 to 87.41 (Mean 53.69) in bilateral turbinectomies and from 48.54 to 82.16 (Mean 60.0) in cases with unilateral turbinectomies. On an analysis of questionnaire, put to all patients, 76% were found to be fully satisfied with the relief from nasal obstruction whereas in rest 24% although there was an objective improvement in their nasal patency yet they were not satisfied subjectively. Results of other subjective parameters i.e. nasal discharge, sneezing etc for assessing the efficacy of this procedure were less encouraging. Measurement of total area of vapour condensation at each follow up has showed that with passage of time it gradually increased with the gradual subsidence of post-operative oedema and scarification of cavernous spaces. It is consistent with the findings of Shahinian (1953) who stated that it is best to wait for several months after doing submucous turbinectomy as this procedure concomitantly reduced vasomotor intermescence in some cases by scarification of cavernous space. Sherman (1977) too in a study of nasal airway function post-operatively reported increased improvement with passage of time and found it to be maximum (80.26%) at two to 12 months follow ups. House (1951) reported subjective improvement in nasal airway patency in 78% and the extent of relief was from 50% to complete relief. Martinez (1983), in his study, had reported relief of nasal obstruction in 96%

of cases following turbinectomy whereas Richard (1984) reported relief of symptoms in 94% cases. Odentoyinbo (1987) reported permanent relief of nasal obstruction in 90% with relief of rhinorrhoea and anosmia to be 36% and 61% in his study of inferior turbinectomies in 39 patients. Ophir et al (1992) reported relief from nasal obstruction in 82% cases of inferior turbinectomy with rhinorrhoea persisting in 34% cases.

The study of complications associated with this procedure revealed primary haemorrhage in 4% of cases and was probably due to subclinical infection. None of the cases had any post-operative reactionary or secondary haemorrhage. Post-operatively 4% cases develop synechiae which was excised without any further problem. None of our patients at any stage complained of nasal crusting. House (1951), Shahinian (1953), Principato (1979) and others who did submucous turbinectomy had reported similar complications of this procedure. Richard (1984) and Davies (1987) have reported the incidence of primary haemorrhage in their cases of turbinectomy to be 8.6% and 8.8% respectively. Woodhead et al (1989) and William et al (1991) have reported the complication of a vascular necrosis of the turbinate bone in cases who underwent submucous diathermy.

Conclusion

Inferior turbinate hypertrophy whether unilateral or bilateral is one of main causes of nasal obstruction. The turbinate hypertrophy may be of mucosa or bone and it is very important to differentiate it by simply decongesting the turbinate. It is the bony hypertrophy cases who should be subjected to submucous resection of the turbinate whereas mucosal hypertrophy cases should be treated with electric or cryocautery. SMR of the turbinate has been found to be a very affective procedure for relieving the nasal obstruction as a result of irreversible, perennial bony inferior turbinal enlargement. Complications associated with this procedure are minimal when compared with other surgical procedures available for reducing the size of the turbinates. Moreover the procedure has least interference with the physiology of the turbinal tissue. Also during this study, Gertner's plate method has been found to be a very simple and easy method of measuring the nasal patency.

References

1. Davies, PJD (1987) : Complications of inferior turbinectomy. *Journal of Laryngology & Otology*, 101 : 1136-1139
2. Gertner, R. ; Pondoshin, L and Fradus, M (1984) : A simple method of measuring the nasal airway in clinical work. *Journal of Laryngology & Otology*, 98 : 351-355.
3. House, H P. (1951) : Submucous resection of inferior turbinal bone. *Laryngoscope*, 61 : 637-646.
4. Jones, M (1895) : Turbinal hypertrophy. *Lancet*, 2 : 879.
5. Martinez, S. A; Nissen, AJ; Stock, CR et al (1983) : Nasal turbinate resection for relief of nasal obstruction. *Laryngoscope*, 93 : 871-875.
6. Odentoyinbo, O (1987) : Complications following total inferior turbinectomy : Facts or myths. *Clin. Otolaryngol*, 1987 : Oct. 12 (5) : 361-363.
7. Ophir, D; Schindel, D; Halpering, D and Marshak, G (1992) : Long term follow up of the effectiveness and safety of inferior turbinectomy. *Plastic Reconstructive Surgery*, 90 (6) : 980-984.
8. Principato, JJ (1979) : Chronic vasomotor rhinitis, cryogenic and other surgical modes of treatment. *Laryngoscope*, 89 : 619-639
9. Richard, AP and Rodney JR (1984) : Inferior turbinate surgery. *Plastic and Reconstructive Surgery. Jour.*, 74 (2) : 227-233.
10. Shahinian, L (1953) . Chronic vasomotor rhinitis, treated by submucous diathermic coagulation. *Arch. Otolaryngol* , 57 . 475-489.
11. Sherman AH (1977) : A study of nasal airway function in the post-operative period of nasal surgery. *Laryngoscope* March 1977 (3) : 229-304.
12. William, HO; Fisher, EW, Goldingwood, DG (1991) : Two stage turbinectomy . sequestration of the inferior turbinate following submucosal diathermy. *Journal of Laryngology & Otology*, 105 (1) : 14-16.
13. Woodhead, CH; Wickham, MH; Smett, JC and Macdonald, AW (1983) : Some observations on submucosal diathermy. *Journal or Laryngology & Otology*, 103 : 1047-1049.
14. Zwaared emaker, M (1889) : Cited by M. Uddstromer in *Acta Otolaryngologica Supplement*, 42, p. 69, 1940.