USE OF SPECIFIC IgE TO FRACTIONS OF GYNANDROPSIS GYNANDRA POLLEN IN *IN VITRO* DIAGNOSTIC TEST

G. Suman Latha, V. Vijaya Lakshmi, H. Surekha Rani and K.J.R. Murthy

Bhagwan Mahavir Hospital and Research Centre, Hyderabad--500 004 (A.P), India.

ABSTRACT

The study was aimed at presence of specific IgE antibody levels *invitro* to the identified antigen. Based on positive skin test with Gynandropsis gynandra and elevated levels of total IgE (>325 IU/ml) 104 patients were selected. Healthy, asymptomatic individuals (25) with low total IgE (<325 IU/ml) were included as controls. The mean OD values by ELISA for specific IgE were 0.67 ± 0.21 , 0.57 ± 0.18 and 0.56 ± 0.18 with whole pollen antigen, 46-37 kD fraction and 36-32 kD fraction, respectively. The specificity and sensitivity between skin test positivity with whole pollen antigen verses fraction with mol.wt 46-37 kD was 90% and 90% and for fraction with mol.wt 36-32 kD was found to be 81.1% and 89.4%. The clusters with molecular weights 46-37 kD and 36-32 kD may be useful in *in vitro* diagnostic test. Fractions within these clusters need to be identified for a higher specificity.

KEY WORDS

ELISA, *Gynandropsis gynandra*, IgE, Skin test, Specificity, Sensitivity.

INTRODUCTION

Asthma is a chronic airway disease affecting 100 million people world wide with increasing prevalence each year. In Indian population the prevalence of asthma is 10%-15%, where as in Hyderabad 30%-50% are suffering with respiratory allergy symptoms in Urban areas. In semi-urban areas, 29.7% are suffering with chronic bronchitis and 18.8% with asthma. (1)

Currently the reagents which are used for diagnosis are crude extracts of the allergen. These reagents are non-specific in nature, lack sensitivity, specificity and cross reactivity. Specific and appropriate major allergen(s) may be useful as monospecific reagent(s) for diagnosis. Fractions with molecular weights 46-37 kD and 36-32 kD of *Gynandropsis gynandra* pollen were demonstrated to be major allergens in a study conducted at this center (2). The present study was aimed at estimation of specific IgE antibody levels *in vitro* to the identified antigens.

Author for Correspondence :

Dr. K.J.R. Murthy, MD, FCCP
Director PPM DOTS,
Research Co-ordinator,
10/1/1, Bhagwan Mahavir Hospital and
Research Center, A.C. Guards,
Hyderabad-500 004 (A.P.), India
Fax : 0091-40-7560638
E-mail: kollurijrm@hotmail.com

MATERIAL AND METHODS

Subjects

Criteria

Of the patients attending the Allergy Clinic at Mahavir Hospital and Research Center, 104 of them who had a confirmed diagnosis of bronchial asthma were studied. The criteria used to confirm atopic bronchial asthma in these patients were clinical symptoms combined with elevated total serum IgE (sIgE) levels and positive skin test responses. Patients below 12 years, above 60 years and pregnant women were excluded from the study. The consent of every patient was taken prior to skin testing and serum collection (3).

Skin test

Pollen antigens selected based on the local aerobiological calendar were used for skin testing by intradermal injections (Cure well India Ltd). Intradermal injection of buffer saline acted as negative control and histamine phosphatase as positive control. Immediate and late phase cutaneous responses were recorded at 20 min and 6-8 hours after allergen challenge, respectively (4).

Clinical studies

Based on intradermal skin test positivity of *G. gynandra* pollen, 104 patients were selected for the present study. For 25 asymptomatic healthy individuals skin test was not performed. Serum samples were collected from the patients and healthy individuals and were

stored at -20°C until use.

Pollen Extraction

The proteins were extracted from the pollens of *G. gynandra* as described by Sridhara *et al.* (5) Pollens collected from flowers of *G. gynandra* were defatted with diethyl ether. The extract was prepared in 0.05 M ammonium bicarbonate buffer, pH 8.0 by continuous stirring for 18 hours on a magnetic stirrer at 4°C. The extract was centrifuged (10,000 rpm, 4°C) and the supernatant brought to 90% saturation with ammonium sulphate. The precipitate was dissolved in distilled water and dialysed using "Spectropor" membrane. (Molecular weight cut off 2,000). The dialysed extract was passed through millipore filter membrane (0.45 µm), lyophilized in aliquots and stored at -20°C.

Total IgE

The Total IgE levels were estimated by ELISA using a commercial kit (General Biologicals Corp, Taiwan). Anti-IgE coated wells were used to detect the levels of total IgE. Patient's sera 20 µl and controls were added into appropriate wells. One hundred microlitre of zero buffer was dispensed in each well and mixed thoroughly for 10 seconds and incubated at room temperature for 30 minutes. Washing was carried out 5 times with distilled water and 150 µl of anti-IgE enzyme conjugate was added and gently mixed and incubated for 30 minutes at room temperature, washed the wells 5 times and added 100 µl of tetramethylbenzidine (TMB) solution and incubated for 20 minutes. Reaction was stopped by adding 50 µl of stop solution in each well and read the optical density at 490 nm with a micro titer plate reader within 15 minutes.

Specific IgE

ELISA was performed to measure the *G. gynandra*-specific IgE levels in individual patients sera with whole pollen antigen and also with its fractions by the method of Voller *et al.* (6). Antigenic extract were diluted in carbonate bicarbonate coating buffer, pH 9.6 and added, 100 µl per well (0.1 µg/100 µl). This was followed by overnight incubation at 4°C. After washing thrice with phosphate buffer saline (PBS) tween 20, the plates were incubated with 1% BSA for 2 hrs at 37°C. After washing thrice with PBS tween 20, the plates were incubated with 100 µl (1:100 diluted in PBS pH 7.4) individual patient's sera for two hrs at 37°C. As a control, normal healthy human serum was also tested. After washing thrice with PBS tween 20. The Anti-IgE conjugated to horse radish peroxidase-HRP (Sigma Co) was added to each well (1:2000 diluted in PBS pH 7.4) incubated for two hours at 37°C followed by washing thrice with PBS tween20, 100 µl of enzyme substrate O-phenylenediamine dihydrochloride was

added in 0.05M phosphate citrate buffer containing 30% H₂O₂ was added to each well. The plates were incubated for 30 minutes at room temperature. The colour development was stopped by adding 50 µl of 5 M H₂SO₄ to each well of microtiter plate. Absorbance was measured at 490 n.m. Sera showing ≥ 3 times the O.D. compared to negative control was considered as ELISA positive.

Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE)

SDS-PAGE was performed with the extracted proteins, using a vertical slab gel apparatus (Bio-rad, laboratories) as described by Laemmli (7). Electrophoresis was carried out in a discontinuous buffer (0.025M Tris, 0.2 M glycine and 0.1% SDS) system. Antigen samples were diluted in sample buffer and boiled at 100°C for 3 min. Extract was loaded in concentration of 5 µg/well on 6%, 15% stacking gel and resolving gel. Samples were allowed to run for 5-6 hrs at 100 volts and 25 m Amps. Molecular weights were estimated with the use of standard molecular weight protein markers (Genei).

Electro elution

Portions of the gel containing the separated fractions were cut and processed for electro elution by employing the method described by Andersen (8). Electro elution was done using a 2mm phosphate buffer, pH 6.8. This buffer was chosen because it provided effective elution associated with a minimal generation of heat and found non-toxic for pollen antigen. Gels were equilibrated for 40 min in equilibrium buffer after the termination of the electrophoretic run during which removal of toxic products SDS, Tris, Glycine and also swelling of the gel occurred. The electro elution of the gel was performed for 20 min, with a 40 V current. After the termination of the electro elution, the separated components were harvested. Fractions were kept frozen at -20°C until the use.

Statistical Analysis

The mean, standard deviation, specificity and sensitivity (9) were calculated for the above parameters.

Results

The specific serum IgE levels were performed by ELISA method. The mean O.D values for specific serum IgE levels in 104 patients with whole pollen antigen of *Gynandropsis gynandra* and its fractions with molecular weights 46 to 37 kD and 36 to 32 kD was observed to be 0.67±0.21, 0.57±0.18, 0.56±0.18 and in normals it was found to be 0.13±0.03 respectively (Table 1).

Table 1. The mean specific IgE levels with whole pollen antigen and fractions in patients allergic to *Gynandropsis gynandra* pollen antigen by ELISA

IgE	Mean $\pm$ S.D
Specific IgE (Whole pollen antigen) N=104	0.670 $\pm$ 0.21
46-37kD (Fraction) N=104	0.57 $\pm$ 0.18
36-32kD (Fraction) N=104	0.56 $\pm$ 0.18
Specific IgE (Asymptomatics) N=25	0.13 $\pm$ 0.03

The specificity and sensitivity between the skin test positivity with whole pollen antigen verses fractions were performed and it was observed that for fraction with mol.wt 46-37 kD, was 90% and 90% and for fraction with mol.wt 36-32 kD was found to be 81.1% and 89.4% respectively.

These above results are supported by western blot analysis, which showed a positive band at 43 kD in the allergic patients. (10)

DISCUSSION

IgE triggers the release of chemicals that are responsible for inducing the symptoms of asthma. A genetically predisposed individual can develop an IgE immune response only after environmental exposure to the offending antigen.

Currently diagnosis of allergic diseases is based on the typical history of allergic symptoms, clinical and serodiagnostic tests. It is the major isotype of anaphylactic antibodies, and although theoretically IgG₄ can also be a reagin, its clinical importance is not significant. Thus *in vivo* and *in vitro* tests used in the diagnosis of allergic diseases are directed towards the detection of free or cell-bound IgE.

Immediate hypersensitivity skin test used to identify specific IgE sensitization is influenced by various factors such as systemic and topical antihistamines, which may suppress the weal and flare reaction; other medications like corticosteroids can also interfere with the test. The test must be performed on a regular basis to maintain reliability and interpretation of test results is subjective moreover is not standardized. There may be false-positive and false-negative reactions,

intermittent difficulty in performing the test and anaphylaxis is a potential risk. The most important factors to be concentrated are sensitivity and reproducibility of the test (11,12).

In vitro tests are better and invaluable in many allergy studies. Though total IgE is the original screening test for allergy, it is not 100% specific for allergic disease and has been superseded by newer specific tests. Whereas specific IgE is not influenced by any concurrent drug treatment, it can be measured *in vitro*, the specificity of the test being as high as 90% for inhaled allergens. Therefore, by screening the allergy patients for specific allergen IgE sensitivity may help for better diagnosis.

In the present study, patients had high IgE antibody titers towards the fractions 46-37 kD and 36-32 kD of *G. gynandra*. Similar observations were reported in other pollens like *Prosopis juliflora* and *Amaranthus spinosus*, fraction with mol.wt. 20kD of *Prosopis juliflora* pollen is allergenic among six fractions identified (13). While in *Amaranthus spinosus* pollen, seven antigenic protein fractions were identified by Singh *et al.* (14).

In the present study, the specificity and sensitivity of skin test positivity with whole pollen antigen verses fraction with mol.wt. 46-37 kD were observed to be 90.4% and 90% and for fraction with mol.wt. (36-32 kD) was found to be 81.1% and 89.4% respectively. Quiralte *et al.* has studied the Olive allergen specific IgE responses in patients with different allergenic fractions of Olive and found that ole e 1 and ole e2 as the major allergens, which can be used for both *invitro* and *in vivo* as diagnostic purposes (15).

On this pollen *Gynandropsis gynandra*, there are no reports available.

In vitro testing for the diagnosis of allergy was acceptable as frontline diagnostic tools (16). Hence the results of this preliminary study on pollen fractions with molecular weights 46-37 kD and 36-32 kD distinctly demonstrates that these fractions may be used as *invitro* diagnostic reagent preparations. However, there is a need to confirm this observation by testing patients positive to pollens other than *Gynandropsis gynandra* and also to identify the specific fraction(s) within these clusters.

ACKNOWLEDGEMENTS

We are grateful to Lady Tata Memorial Trust & Council of Scientific and Industrial Research (CSIR) for the financial support during the study. We thank Dr.Akbar Yazdani, Dr. Nafees Sultana & Ms. R. Eswari for the clinical and technical help rendered during the study.

REFERENCES

1. Satish Kumar, K., Prasad, C.E. and Uma Maheswara Reddy, P. (1999) Prevalence of asthma in a sample population of Hyderabad. Indian Biologist Vol.XXXI, No. 1, 62-66.
2. Suman Latha, G., Vijaya Lakshmi, V., Surekha Rani, H. and Murthy K.J.R. (2004) Use of pollen fractions to assess the effect of hyposensitization with Gynandropsis Pollen. The Indian Journal of Allergy and Applied Immunology, Vol.18-Issue 1.
3. Madhuri, Jaju, Vijayalakshmi, V., Reddy, P.S., Jain, S.N. and Murthy, K.J.R. (1992) Intradermal skin testing, total IgE levels and atopic status in asthmatics. Lung India X, No.2, 61-64.
4. Shivpuri, D.N., Kartar Singh (1971) Studies in yet unknown pollens of Delhi state metropolitan (Clinical aspects). Ind. J. Med. Res., 59, 1411-1419.
5. Sridhara, S., Singh, B.P., Arora, Naveen, Verma, Jyotsna and Gangal, S.V. (1992) A study of antigenic and allergenic changes during storage in three different biological extracts. Asian Pacific J. Allergy Immunol. 10, 33-38.
6. Voller, A., Bidwell, D., Barlett, A. (1980) Enzyme linked immunosorbent assay; in Rose NR, Friedman H, (eds): Manual of Clinical Immunology. Washington, American Society for Microbiology.
7. Laemmli, U.K. (1970) Cleavage of structural proteins during assembly of the head of the bacteriophage T₄. Nature 227, 680-685.
8. Andersen, P. and Heron. I. (1993) Simultaneous electroelution of whole SDS-polyacrylamide gels for the direct cellular analysis of complex protein mixtures. J. Immunol. 161, 29-39.
9. Wayne, W. Daniel (2000) Biostatistics: A foundation for analysis in the health sciences. Seventh Edition, Pg 71-75.
10. Suman Latha, G., Lakshmi Kiran, A., Vijayalakshmi, V., Surekha Rani, H. and Murthy, K.J.R. (2001) Immunoblot Responses in Patients Allergic to Gynandropsis gynandra pollen. The Indian Journal of Allergy and Applied Immunology 14 (2), 53-59.
11. Anon, J.B. (1993) Introduction to *in vivo* allergy testing. Otolaryngol Head Neck Surg. 109 (3 Pt 2), 593-600.
12. Sousa, C.A. and Norton, A.L. (1990) Advances in methodology for diagnosis of allergic skin disease. Vet. Clin. North Am. Small Anim. Pract. 20 (6), 1419-1427.
13. Thakur, I.S. (1989) Fractionation and analysis of allergenicity of allergens from Prosopis juliflora pollen. Int. Arch. Allergy Appl. Immunol. 90 (2), 124-129.
14. Singh, A.B. and Dahiya, P. (2002) Antigenic and allergenic properties of Amaranthus Spinosa pollen—a commonly growing weed in India. Ann. Agric. Environ. Med. 9 (2), 147-151.
15. Quiralte, J., Florido, F., Arias de Saavedra, J.M., Gomez, A., Saenz de San Pedro, B., Gonzalez, E. and Rodriguez, R. (2002) Olive allergen-specific IgE responses in patients with Olea europaea pollinosis. Allergy 57 Suppl 71, 47-52.
16. Poon, A.W., Goodman, C.S. and Rubin, R.J. (1998) *In vitro* and skin testing for allergy: comparable clinical utility and costs. Am. J. Manag. Care. 4 (7), 969-985.